

Appendix H: Tanner crab molt increment data

William T. Stockhausen

05 September, 2018

Contents

Tanner crab growth data	2
Mean growth	3
Comparison with the 2016 assessment model	4

List of Tables

1	Estimated growth parameters for the EBS molt increment data with post-molt size as a power law of pre-molt size.	3
2	2016 assessment model mean growth parameters.	3
3	Growth parameters based on Kodiak data, used as prior means for parameters in the assessment model.	3

List of Figures

1	Tanner crab molt increment data, by region and sex.	2
2	Tanner crab growth data, by region and sex. Colored lines indicate mean growth, by sex, as determined by the assessment model.	4




Figure 1: Tanner crab molt increment data, by region and sex.

Tanner crab growth data

Figure 1 shows molt increment data collected from crab near Kodiak Island in the Gulf of Alaska and in the eastern Bering Sea (EBS). The Kodiak data was collected over a 20+ year period during opportunistic surveys and caged grow-out experiments. The EBS data was collected in 2014, 2015, and 2016 through cooperative research conducted by the AFSC/NMFS and the Bering Sea Research Foundation (BSFRF).

Mean growth

Sex-specific parameters for post-molt size as a power function of pre-molt size ($z_{post} = e^a \cdot z_{pre}^b$) were estimated in R using the `glm` function from the EBS data on the log-scale using the regression formula $\ln[z_{post}] = a + b \cdot \ln[z_{pre}]$. The resulting estimates

Table 1: Estimated growth parameters for the EBS molt increment data with post-molt size as a power lae of pre-molt size..

parameter	males	females
a	0.2708370	0.6106653
b	0.9922623	0.8975509

Sex-specific parameters from the 2016 assessment model reflecting estimated mean growth are listed in Table 2, where $z_{post} = e^a \cdot z_{pre}^b$.

Table 2: 2016 assessment model mean growth parameters.

parameter	males	females
a	0.4220295	0.6999999
b	0.9721004	0.8850577

Growth parameters estimated from the Kodiak data, used as prior mean values for parameters in the assessment model are listed in Table 3.

Table 3: Growth parameters based on Kodiak data, used as prior means for parameters in the assessment model.

parameter	males	females
a	0.437941	0.5656024
b	0.948700	0.9132661




Figure 2: Tanner crab growth data, by region and sex. Colored lines indicate mean growth, by sex, as determined by the assessment model.

Comparison with the 2016 assessment model

The 2016 assessment model estimated mean growth parameters from size composition data. Priors were placed on the growth parameters based on a previous analysis by Rugolo and Turnock of molt increment data from Kodiak Island in the Gulf of Alaska. The estimated mean growth curves from the assessment over-predict post-molt size at larger pre-molt sizes for both males and females. The molt increment data from the EBS does not appear to be radically different from that collected at Kodiak. In the current assessment, only the EBS data will be included to fit.