

8. Aleutian Islands Golden King Crab Stock Assessment

May 2020 Crab SAFE Draft Report

M.S.M. Siddeek¹, J. Zheng¹, C. Siddon¹, B. Daly², M.J. Westphal³, and L. Hulbert¹

¹Alaska Department of Fish and Game, Division of Commercial Fisheries, P.O. Box 115526,
Juneau, Alaska 99811

²Alaska Department of Fish and Game, Division of Commercial Fisheries, 351 Research Ct.,
Kodiak, Alaska 99615

³Alaska Department of Fish and Game, Division of Commercial Fisheries, PO Box 920587,
Dutch Harbor, Alaska 99692.

Executive Summary

1. Stock

Golden king crab, *Lithodes aequispinus*, Aleutian Islands, east of 174° W longitude (**EAG**) and west of 174° W longitude (**WAG**).

2. Catches

The Aleutian Islands golden king crab (AIGKC) commercial fishery has been prosecuted every year since 1981/82. Retained catch peaked in 1986/87 at 2,686 t (5,922,425 lb) and 3,999 t (8,816,319 lb), respectively, for **EAG** and **WAG**, but the retained catch dropped sharply from 1989/90 to 1990/91. The fishery has been managed separately east (**EAG**) and west (**WAG**) of 174° W longitude since 1996/97, and Guideline Harvest Levels (GHLs) of 1,452 t (3,200,000 lb) for **EAG** and 1,225 t (2,700,000 lb) for **WAG** were introduced into management for the first time in 1996/97. The GHL was subsequently reduced to 1,361 t (3,000,000 lb) beginning in 1998/99 for **EAG**. The reduced GHLs remained at 1,361 t (3,000,000 lb) for **EAG** and 1,225t (2,700,000 lb) for **WAG** through 2007/08, but were increased to 1,429 t (3,150,000 lb) for **EAG** and 1,294 t (2,835,000 lb) for **WAG** beginning with the 2008/09 fishing season following an Alaska Board of Fisheries (BOF) decision. The management specification changed from GHL to TAC (Total Allowable Catch) with adoption of the Crab Rationalization Program in 2005/06 (NPFMC 2007b). The TACs were increased by another BOF decision to 1,501 t (3,310,000 lb) for **EAG** and 1,352 t (2,980,000 lb) for **WAG** beginning with the 2012/13 fishing season. The below par fishery performance in **WAG** in recent years lead to reduction in TAC to 1,014 t (2,235,000 lb), which reflected a 25% reduction in the TAC for **WAG**, while the TAC for **EAG** was kept at the same level 1,501 t (3,310,000 lb) for the 2015/16 through 2017/18 fishing seasons. With the improved fishery performance and stock status in 2017/18, the TACs were further increased to 1,134 t (2,500,000 lb) for **WAG** and 1,749 t (3,856,000lb) for **EAG** beginning with the 2018/19 fishing season. With the implementation of a revised state harvest strategy in 2019, the TACs were further increased to 1,302 t (2,870,000 lb) for **WAG** and 1,955 t (4,310,000 lb) for **EAG**. The **EAG** fishery achieved 100% of TAC while the **WAG** fishery is ongoing with 96% of TAC harvested for the 2019/20 fishing season at the time of this assessment.

Catches have been steady under the GHL/TAC and the fishery has harvested close to allowable levels since 1996/97. These TAC levels were set below the ABCs determined under Tier 5 criteria

(considering 1991–1995 mean catch for the whole Aleutian Islands region, 3,145 t (6,933,822 lb), as the limit catch) under the most recent crab management plan. A new harvest strategy based on model estimated mature male abundance was accepted by the BOF in March 2019, specifying a 15% maximum harvest rate for **EAG** and 20% maximum harvest rate for **WAG**, and implemented during the 2019/20 fishery. In addition to the retained catch allotted as TAC, there was retained catch in a cost-recovery fishery towards a \$300,000 goal in 2013/14 and 2014/15 to fund an onboard observer program, and towards a \$500,000 goal in 2015/16 to 2019/20 in order to fund an onboard observer program and stock survey.

Total mortality of Aleutian Islands golden king crab includes retained catch in the directed fishery, mortality of discarded catch, and bycatch in fixed-gear and trawl groundfish fisheries, though bycatch in other fisheries is low compared to mortality in the directed fishery. Total retained catch in the post-rationalized fishery (2005/06–2019/20) has ranged from 2,498 t (5,508,100 lb) to 3,274 t (7,218,545 lb). Total mortality ranged from 2,506 t (5,525,000 lb) to 3,693 t (8,141,000 lb) for the same period. Total retained catch in 2019/20 was 3,274 t (7,218,545 lb): 2,031 t (4,476,775 lb) from the **EAG** fishery (which included cost-recovery catch), and 1,244 t (2,741,770 lb) from the **WAG** fishery. Discarded (non-retained) catch occurs mainly during the directed fishery. Although low levels of discarded catch can occur during other crab fisheries, there have been no such fisheries prosecuted since 2004/05, except as surveys for red king crab conducted under an Alaska Department of Fish and Game (ADF&G) Commissioner’s Permit (and no golden king crab were caught during the cooperative red king crab survey performed by industry and ADF&G in the Adak area in September 2015; Hilsinger et al. 2016). Estimates of the bycatch mortality during crab fisheries decreased during 1995/96–2005/06, both in absolute value and relative to the retained catch weight and stabilized during 2005/06–2014/15. Total estimated bycatch mortality during crab fisheries in 2019/20 was 275 t (607,000 lb) for **EAG** and 116 t (256,000 lb) for **WAG**. Discarded catch also occurs during fixed-gear and trawl groundfish fisheries but is small relative to the directed fishery. Groundfish fisheries are a minor contributor to total fishery discard mortality, 23 t (52,000 lb) for **EAG** and 3 t (8,000 lb) for **WAG** in 2019/20.

Catch per unit effort (CPUE, i.e., catch per pot lift) of retained legal males decreased from the 1980s into the mid-1990s, but increased after 1994/95, particularly with the initiation of the Crab Rationalization Program in 2005/06. Although CPUE for the two areas showed similar trends through 2010/11, during 2011/12–2014/15 CPUE trends have diverged (increasing for **EAG** and decreasing for **WAG**).

A cooperative golden king crab survey was performed by the Aleutian Islands King Crab Foundation (an industry group) and ADF&G in the **EAG** and **WAG** (for the first time in August 2018) fisheries, by vessels that were quota fishing (i.e., each vessel fishing an allotted share of total allowable catch). For the purpose of catch accounting for 2019/20, it was assumed that bycatch mortality that occurred during the survey was accounted for by reported discards for the 2019/20 fishery.

3. Stock biomass

Estimated mature male biomass (MMB) for **EAG** under all scenarios decreased from the 1980s to the 1990s, then increased during the 2000s and sharply increased since 2014. Estimated MMB for **WAG** decreased during the late 1980s and 1990s, increased during the 2000s, decreased for several years since 2009 and has increased since 2014. The low levels of MMB for **EAG** were observed

in 1995–1997 and in 1990s for **WAG**. Stock trends reflected the fishery standardized CPUE trends in both regions.

4. Recruitment

The numbers of recruits to the model size groups under all scenarios have fluctuated in both **EAG** and **WAG**. For **EAG**, model recruitment was high in 2016, highest in 2017; and lowest in 1986. The model recruitment for **WAG** was high during 1984 to 1986, highest in 1985, and lowest in 2011. A slightly increasing trend in recruitment was observed since 2011 in **WAG**.

5. Management performance

The size-based assessment model was accepted at the September 2016 CPT and October 2016 SSC meetings for OFL determination for the 2017/18 fishery cycle. In addition, the CPT in January 2017 and SSC in February 2017 recommended using the Tier 3 method to compute OFL and ABC. The assessment model was first used for setting OFL and ABC for the 2017/18 fishing season. This was followed since. The CPT in May 2017 and SSC in June 2017 accepted the authors' recommendation of using scenario 9 (i.e., model using the knife-edge maturity to determine MMB) for OFL and ABC calculation. During the May 2017 meeting, the CPT noted that a single OFL and ABC are defined for Aleutian Islands golden king crab (AIGKC). However, separate models are available by area. Hence, following previous assessments, OFLs and ABCs by area were summed to calculate OFL and ABC for the entire stock.

All models for **EAG** and **WAG** considered the previous season's fishery information (i.e., 2019/20 fishery, concluded in **EAG** and almost 96% of TAC achieved in **WAG**). We recommend two models from the common four models for **EAG** and **WAG**: model 20_1b Ver 2 (re-evaluation of observer CPUE indices after reducing the number of gear codes; selection of a fixed period, 1987–2012, for mean number of recruits calculation for reference points estimation; and standardization of fishery CPUE by the negative binomial generalized linear model); and model 20_2 (consideration of year and area interaction factor for observer CPUE standardization).

Model 20_1 is the base model (accepted model 19_1 in 2019) with the knife-edge male maturity at 111 mm CL, an M of 0.21yr^{-1} , and the addition of 2019/20 data. Models 20_1b, 20_1b Ver 2, 20_1c, 20_1d, 20_2, and 20_2b are modifications from the base model.

The total catch, 3.693 t, did not exceed OFL, 5.249 t, in 2019/20; therefore, overfishing did not occur.

The mature male biomass, 16.323 t, is above MSST, 5.909 t, in 2019/20; hence, the stock was not overfished.

Status and catch specifications (1000 t) of Aleutian Islands golden king crab

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch ^a	OFL	ABC ^b
2016/17	N/A	N/A	2.515	2.593	2.947	5.69	4.26
2017/18	6.044	14.205	2.515	2.585	2.942	6.048	4.536
2018/19	5.880	17.848	2.883	2.965	3.355	5.514	4.136
2019/20	5.909 ^c	16.323 ^c	3.257	3.274 ^d	3.693 ^d	5.249	3.937
2020/21 ^e		14.760				4.793	3.595
2020/21 ^f		15.106				4.993	3.745
2020/21 ^g		14.774				4.798	3.599

Status and catch specifications (million lb) of Aleutian Islands golden king crab

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch ^a	OFL	ABC ^b
2016/17	N/A	N/A	5.545	5.716	6.497	12.53	9.40
2017/18	13.325	31.315	5.545	5.699	6.487	13.333	10.000
2018/19	12.964	39.348	6.356	6.536	7.396	12.157	9.118
2019/20	13.027 ^c	35.985 ^c	7.180	7.219 ^d	8.141 ^d	11.572	8.679
2020/21 ^e		32.540				10.566	7.925
2020/21 ^f		33.303				11.008	8.256
2020/21 ^g		32.571				10.579	7.934

- Total retained catch plus estimated bycatch mortality of discarded bycatch during crab fisheries and groundfish fisheries.
- 25% buffer was applied to total catch OFL to determine ABC.
- MSST and MMB determined by Model 20_1b Ver 2
- 100% TAC was achieved in **EAG**, but over 96% TAC was achieved in **WAG** at the time of this assessment. The **WAG** fishery is ongoing.
- Model 20_1b, up to 2019/20 data, mean number of recruit calculation time period for **EAG**: 1986–2017 and for **WAG**: 1987–2018.
- Model 20_2, up to 2019/20 data.
- Model 20_1b Ver 2, up to 2019/20 data, mean number of recruit calculation time period for **EAG** and **WAG**: 1987–2012.

6. Basis for the OFL

The length-based model developed for the Tier 3 analysis estimated mature male biomass (MMB) on February 15 each year for the period 1986 through 2020. The terminal year mature male biomass was projected by an additional year to determine OFL and ABC for the 2020/21 season. The Tier 3 approach uses a constant annual natural mortality (M), knife-edge maturity size, and the mean number of recruits for different time periods for OFL and ABC calculation. Previously derived M of 0.21 yr⁻¹ from the combined data and a knife-edge maturity size of 111 mm carapace length (CL) from the **EAG** and **WAG** data were used (Siddeek *et al.* 2018).

We provide the OFL and ABC estimates for **EAG** and **WAG** separately and combined (i.e., for the entire Aleutian Islands; **AI**) from seven models, 20_1, 20_1b, 20_1b Ver 2, 20_1c, 20_1d, 20_2, and 20_2b, for **EAG**; and from four models, 20_1, 20_1b, 20_1b Ver 2, and 20_2, for **WAG** and for **AI** in the following six tables. We treat model 20_1 as the base model.

EAG (Tier 3):

Biomass, total OFL, and ABC for the next fishing season in millions of pounds. Current MMB = MMB on 15 Feb. 2021.

Model	Tier	<i>MMB</i> _{35%}	Current	MMB/	<i>F</i> _{OFL}	Recruitment	<i>F</i> _{35%}	OFL	ABC	ABC
			MMB	<i>MMB</i> _{35%}		Years to define			(P*=0.49)	(0.75*OFL)
EAG20_1	3a	14.553	18.809	1.29	0.61	1987–2012	0.61	6.648	6.609	4.986
EAG20_1b	3a	14.935	18.674	1.25	0.61	1986–2017	0.61	6.583	6.544	4.937
EAG20_1bVer2	3a	14.547	18.694	1.29	0.61	1986–2012	0.61	6.592	6.553	4.944
EAG20_1c	3a	14.481	15.293	1.06	0.61	1986–2017	0.61	4.977	4.939	3.733
EAG20_1d	3a	14.724	17.173	1.17	0.61	1986–2017	0.61	5.850	5.826	4.387
EAG20_2	3a	14.979	19.104	1.28	0.61	1986–2017	0.61	6.908	6.869	5.181
EAG20_2b	3a	14.579	16.177	1.11	0.61	1986–2017	0.61	5.478	5.438	4.109

Biomass in 1000 t; total OFL and ABC for the next fishing season in t.

Model	Tier	<i>MMB</i> _{35%}	Current	MMB/	<i>F</i> _{OFL}	Recruitment	<i>F</i> _{35%}	OFL	ABC	ABC
			MMB	<i>MMB</i> _{35%}		Years to Define			(P*=0.49)	(0.75*OFL)
EAG20_1	3a	6.601	8.532	1.29	0.61	1987–2012	0.61	3,015.592	2,997.858	2,261.694
EAG20_1b	3a	6.774	8.470	1.25	0.61	1986–2017	0.61	2,985.928	2,968.143	2,239.446
EAG20_1bVer2	3a	6.599	8.480	1.29	0.61	1987–2012	0.61	2,990.063	2,972.283	2,242.547
EAG20_1c	3a	6.568	6.937	1.06	0.61	1986–2017	0.61	2,260.998	2,504.178	1,695.748
EAG20_1d	3a	6.679	7.790	1.17	0.61	1986–2017	0.61	2,653.436	2,642.813	1,990.077
EAG20_2	3a	6.794	8.665	1.28	0.61	1986–2017	0.61	3,133.485	3,115.767	2,350.114
EAG20_2b	3a	6.613	7.338	1.11	0.61	1986–2017	0.61	2,484.903	2,466.646	1,863.677

7.

WAG (Tier 3):

Biomass, total OFL, and ABC for the next fishing season in millions of pounds. Current MMB = MMB on 15 Feb. 2021.

Model	Tier	<i>MMB</i> _{35%}	Current MMB	MMB/ <i>MMB</i> _{35%}	<i>F</i> _{OFL}	Recruitment		OFL	ABC (P*=0.49)	ABC (0.75*OFL)
						Years to Define <i>MMB</i> _{35%}	<i>F</i> _{35%}			
WAG20_1	3a	11.473	13.844	1.21	0.56	1987–2012	0.56	3.974	3.958	2.981
WAG20_1b	3a	11.725	13.867	1.18	0.56	1987–2018	0.56	3.983	3.968	2.988
WAG20_1bVer2	3a	11.507	13.877	1.21	0.56	1987–2012	0.56	3.987	3.971	2.990
WAG20_2	3a	11.778	14.199	1.21	0.56	1987–2018	0.56	4.100	4.084	3.075

8.
9.

Biomass in 1000 t; total OFL and ABC for the next fishing season in t.

Model	Tier	<i>MMB</i> _{35%}	Current MMB	MMB / <i>MMB</i> _{35%}	<i>F</i> _{OFL}	Recruitment Years		OFL	ABC (P*=0.49)	ABC (0.75*OFL)
						to Define <i>MMB</i> _{35%}	<i>F</i> _{35%}			
WAG20_1	3a	5.204	6.279	1.21	0.56	1987–2012	0.56	1,802.747	1,795.486	1,352.060
WAG20_1b	3a	5.319	6.290	1.18	0.56	1987–2018	0.56	1,806.903	1,799.775	1,355.177
WAG20_1bVer2	3a	5.220	6.295	1.21	0.56	1987–2012	0.56	1,808.318	1,801.190	1,356.239
WAG20_2	3a	5.343	6.441	1.21	0.56	1987–2018	0.56	1,859.828	1,852.480	1,394.871

Aleutian Islands (AI)

Total OFL and ABC for the next fishing season in millions of pounds.

Model	OFL	ABC (P*=0.49)	ABC (0.75*OFL)
20_1	10.622	10.567	7.967
20_1b	10.566	10.512	7.925
20_1bVer2	10.579	10.524	7.934
20_2	11.008	10.953	8.256

Aleutian Islands (AI)

Total OFL and ABC for the next fishing season in t.

Model	OFL	ABC (P*=0.49)	ABC (0.75*OFL)
20_1	4,818.34	4,793.34	3,613.75
20_1b	4,792.83	4,767.92	3,594.62
20_1bVer2	4,798.38	4,773.47	3,598.79
20_2	4,993.31	4,968.25	3,744.99

7. Probability density functions of the OFL

Assuming a lognormal distribution of total OFL, we determined the cumulative distributions of OFL and selected the median as the OFL.

8. Basis for the ABC recommendation

An x proportion buffer on the OFL; i.e., $ABC = (1.0 - x) * OFL$.

The CPT recommended $x = 0.25$.

See also the section G on ABC.

9. A summary of the results of any rebuilding analysis:

Not applicable.

A. Summary of Major Changes

1. Changes (if any) to management of the fishery

In 2019, a new state harvest strategy was implemented.

2. Changes to input data

Commercial fisheries data were updated with values from the most recent observer and fish ticket data for 2019/20: retained catch for the directed fishery and discarded catch estimates for the directed fishery, non-directed crab fisheries, and groundfish fisheries. Thus, the time series of data used in the model are retained catch (1981/82–2019/20), total catch (1990/91–2019/20), and groundfish bycatch (1989/90–2019/20) biomass and size compositions.

Fish ticket retained CPUE were standardized by the generalized linear model (GLM) with the lognormal and negative binomial link functions for the 1985/86–1998/98 period.

Observer pot sample legal size crab CPUE data were standardized by the GLM with the negative binomial link function with variable selection by CAIC (modified AIC) followed by R square criterion, separately for 1995/96–2004/05 and 2005/06–2019/20 periods. A Year and Area interaction factor was considered in one model to estimate a set of CPUE indices. The habitat areas were determined from observer historical pot locations as fishing footprints (see Appendix B).

3. Changes to assessment methodology

None

4. Changes to assessment results

As expected, the addition of the 2019/20 data changed the OFL and ABC estimates, but changes in parameter or abundance estimates were not dramatic.

B. Response to SSC and CPT comments

January 2020 CPT Comments

Comment# 1: The CPT reiterates the SSC request for a brief description of the cooperative survey in the assessment document, including the area sampled, size composition and a summary of results.

Response:

This is an evolving project to collect AIGKC data by active fishing vessels, following a designed two-stage sampling. The data collection covers species, sex, count of crab by size, by pot, by string, and by vessel. Additional data such as depth of fishing, soak time, bait type, mesh size, and pot size are also collected. We use the number of legal-size male crabs at the vessel/string/pot level to estimate the CPUE by a hierarchical random effects model. A brief explanation of the method is provided in Appendix C.

*We have completed the cooperative surveys for five fishing seasons (2015/16, 2016/17, 2017/18, 2018/19, and 2019/20) in the **EAG** region. We also extended the survey for the first time in the*

WAG region in 2018/19. The data series is too short to obtain meaningful results. However, we used the *EAG* CPUE indices in some model scenarios in this analysis to get some feedback.

Comment# 2: Revised approach to select mean recruitment: The proposed approach sets mean recruitment to the average over the years for which the standard deviations of the recruitment estimates is 70% of the assumed standard deviation of inter-annual variability in recruitment. The choice of 70% is the lowest percentage at which a contiguous set of years would be selected. The CPT agrees with the general approach, and requests that the authors include the basis for the 70% in the next report.

Response:

The 70% value is an arbitrary choice satisfying the need to remove a few years from the tail end of the recruitment time series. Instead of using 70% of the fixed R_{σ} , we used the 90th percentile cutoff level based on 1986 to 2020 recruit standard errors estimated by the base model 20_1 to exclude years with high recruit standard deviations. The 90th percentile choice is also an arbitrary level but uses the actual recruitment standard errors to obtain the cutoff level instead of R_{σ}

Comment# 3: Revised approach for standardizing the fishery catch-rate data for 1995/96 – 2019/20. The CPT notes that basis for the specific blocks chosen for Year and Area interaction needs to be more clearly documented. The weight assigned to each block needs to be the total number of $1^0 \times 1^0$ cells ever fished. One potential problem with this approach is that there are blocks x years with no (or very few) data. The CPT made two suggestions:

a. Fit a model of the form $B_{i,j} = A_i + C_j$ where $B_{i,j}$ is the index of biomass for year i and block j , A_i is a year factor, and C_j is a block factor, and use this model to infer the biomass index for blocks x years with no (or very limited) data.

b. The variance of the total biomass index should be computed as:

$$\text{Var}(B_i) = \sum_j N_{\text{ever},j}^2 \text{var}(CPUE_{i,j})$$

where $N_{\text{ever},j}$ is the total number of $1^0 \times 1^0$ cells ever fished in block j , and $CPUE_{i,j}$ is the expected CPUE index for year i and block j .

Response:

We followed both suggestions. We used a GLM procedure to fit the year and area factors to available $B_{i,j}$ indices and used the fitted model to fill the gap for missing year by block values. We also estimated the variance of the biomass index using the suggested formula (Appendix B).

Comment# 4: Analysis of the cooperative survey data. The use of a mixed-effects model is appropriate. However, the choice of covariates needs additional justification. For example, it was not clear that vessel * pot number should be treated as a fixed effect rather than pot number random within vessel. Similarly, a hierarchical structure for strings * block should be considered, such as string random within block, which is itself random. In general, the model for the analysis of the survey data should be more closely aligned with the design of the survey. One possible model would be:

Sumcatch ~ Year + (1|vessel/pot number) + ns(soakdays,ns=9) + ns(Depth,df=6) + (1|block/string).

Response:

We followed the hierarchical random effects model structure suggested by the CPT to analyze the cooperative survey data (Appendix C).

Comment# 5: The CPT recommended the following models for exploration for the May 2020 CPT meeting:

- **Model 19.1b.** As for model 19.1 but with revised periods of years for defining mean recruitment (EAG: 1985-2016; WAG: 1987-2016) and the fish ticket CPUE data standardized assuming a negative binomial distribution.
- **Model 19.1c.** As for model 19.1b except that the EAG 2015-2019 cooperative survey CPUE index is included in the assessment.
- **Model 19.2.** As for model 19.1b, except that the 1995/96 – 2018/19 CPUE data are standardized using year*area interactions.
- **Model. 19.2b.** As for model 19.2, except that the EAG 2015-2019 cooperative survey CPUE index is included in the assessment.

Response:

We considered all suggested models in this report (see Table T1).

January 2020 SSC comments:

Comment# 1: The SSC reiterates for a description of the cooperative survey in the assessment document, including the area sampled and size compositions.

Response:

Please refer to our response to CPT comment#1.

Comment# 2: SSC supports exploration of treating pot as a random effect nested within vessel, or possibly string, and encourages alternative random effects model structures that align with assumptions of the cooperative survey design.

Response:

We followed the random effects approach to analyze the cooperative survey data because of the two-stage sampling design. As per CPT suggestion#3, we used the pot within vessel and string within block structures in the random effects model analysis for this report. The exploration is continuing.

Comment# 3:

The SSC also reiterates the CPT request on the rationale for the 0.7 Sigma_R criterion for recruitments included in the estimation of reference points as this does not seem justified at this point.

Response:

The R_{σ} value is user enforced, came from an arbitrary weight specified to the recruit likelihood. We made it non-subjective by setting the cutoff recruit deviation value at 90th percentile of the

model-estimated recruitment standard deviations for the whole time series. Recruitments with standard deviations less than the cutoff value are included for reference point estimation.

Comment# 4:

The SSC supports the CPT recommendation to explore the given set of models (CPT comments#5) for the May CPT meeting that explore new recruitment time series, different formulations of CPUE standardization, and the inclusion of cooperative survey CPUE.

Response:

We did in this report.

C. Introduction

1. Scientific name:

Golden king crab, *Lithodes aequispinus* J.E. Benedict, 1895.

2. Distribution:

General distribution of golden king crab is summarized by NMFS (2004). Golden king crab, also called brown king crab, occur from the Japan Sea to the northern Bering Sea (ca. 61° N latitude), around the Aleutian Islands, generally in high-relief habitat such as inter-island passes, on various sea mounts, and as far south as northern British Columbia (Alice Arm) (Jewett *et al.* 1985). They are typically found on the continental slope at depths of 300–1,000 m on extremely rough bottom. They are frequently found on coral bottom.

The Aleutian Islands king crab stock boundary is defined by the boundaries of the Aleutian Islands king crab Registration Area O (Figure 1). In this chapter, “Aleutian Islands Area” means the area described by the current definition of Aleutian Islands king crab Registration Area O. Leon *et al.* (2017) define the boundaries of Aleutian Islands king crab Registration Area O:

The Aleutian Islands king crab management area’s eastern boundary is the longitude of Scotch Cap Light (164°44.72'W long), the northern boundary is a line from Cape Sarichef (54°36'N lat) to 171°W long, north to 55°30'N lat, and the western boundary the Maritime Boundary Agreement Line as described in the Maritime Boundary Agreement between the United States and the Union of Soviet Socialist Republics signed in Washington, June 1, 1990 (Figure 1-1 in Leon et al. 2017). Area O encompasses territorial waters of the state of Alaska (0–3 nautical miles) and waters of the Exclusive Economic Zone (3–200 nautical miles).

During 1984/85–1995/96, the Aleutian Islands king crab populations had been managed using the Adak and Dutch Harbor Registration Areas, which were divided at 171° W longitude (Figure 2), but from the 1996/97 season to present the fishery has been managed using a division at 174° W longitude (Figure 2). In March 1996 the Alaska Board of Fisheries (BOF) replaced the Adak and Dutch Harbor areas with the newly created Aleutian Islands Registration Area O and directed ADF&G to manage the golden king crab fishery in the areas east and west of 174°W longitude as two distinct stocks. That re-designation of management areas was intended to more accurately reflect golden king crab stock distribution, coherent with the longitudinal pattern in fishery production prior to 1996/97 (Figure 3). The longitudinal pattern in fishery production relative to 174° W longitude since 1996/97 is similar to that observed prior to the change in management area

definition, although there have been some changes in the longitudinal pattern in fishery production within the areas east and west of 174° W longitude (Figure 4).

Commercial fishing for golden king crab in the Aleutian Islands Area typically occurs at depths of 100–275 fathoms (183–503 m). Pots sampled by at-sea fishery observers in 2013/14 were fished at an average depth of 176 fathoms (322 m; N=499) in the area east of 174° W longitude and 158 fathoms (289 m; N=1,223) for the area west of 174° W longitude (Gaeuman 2014).

3. Evidence of stock structure:

Given the expansiveness of the Aleutian Islands Area and the existence of deep (>1,000 m) canyons between some islands, at least some weak structuring of the stock within the area would be expected. Data for making inferences on stock structure of golden king crab within the Aleutian Islands are largely limited to the geographic distribution of commercial fishery catch and effort. Catch data by statistical area from fish tickets and catch data by location from pots sampled by observers suggest that habitat for legal-sized males may be continuous throughout the waters adjacent to the islands in the Aleutian chain. However, regions of low fishery catch suggest that availability of suitable habitat, in which golden king crab are present at only low densities, may vary longitudinally. Catch has been low in the fishery in the area between 174° W longitude and 176° W longitude (the Adak Island area, Figures 3 and 4) in comparison to adjacent areas, a pattern that is consistent with low CPUE for golden king crab between 174° W longitude and 176° W longitude (Figure 5) during the 2002, 2004, 2006, 2010, and 2012 NMFS Aleutian Islands bottom trawl surveys (von Szalay et al. 2011). In addition to longitudinal variation in density, there is also a gap in fishery catch and effort between the Petrel Bank-Petrel Spur area and the Bowers Bank area; both of those areas, which are separated by Bowers Canyon, have reported effort and catch. Recoveries during commercial fisheries of golden king crab tagged during ADF&G surveys (Blau and Pengilly 1994; Blau *et al.* 1998; Watson and Gish 2002; Watson 2004, 2007) provided no evidence of substantial movements by crab in the size classes that were tagged (males and females ≥ 90 -mm carapace length [CL]). Maximum straight-line distance between release and recovery location of 90 golden king crab released prior to the 1991/92 fishery and recovered through the 1992/93 fishery was 61.2 km (Blau and Pengilly 1994). Of the 4,567 recoveries reported through 12 April 2016 for the male and female golden king crab tagged and released between 170.5° W longitude and 171.5° W longitude during the 1991, 1997, 2000, 2003, and 2006 ADF&G Aleutian Island golden king pot surveys, none of the 3,807 with recovery locations specified by latitude and longitude were recovered west of 173° W longitude and only fifteen were recovered west of 172° W longitude (V. Vanek, ADF&G, Kodiak, pers. comm.). Similarly, of 139 recoveries in which only the statistical area of recovery was reported, none were recovered in statistical areas west of 173° W longitude and only one was in a statistical area west of 172° W longitude.

4. Life history characteristics relevant to management:

There is a paucity of information on golden king crab life history characteristics due in part to the deep depth distribution (~200–1000 m) and the asynchronous nature of life history events (Otto and Cummiskey 1985; Somerton and Otto 1986). The reproductive cycle is thought to last approximately 24 months and at any time of year, ovigerous females can be found carrying egg clutches in highly disparate developmental states (Otto and Cummiskey 1985). Females carry large, yolk-rich, eggs, which hatch into lecithotrophic (i.e., the larvae can develop successfully to juvenile crab without eating; Shirley and Zhou 1997) larvae that are negatively phototactic (Adams

and Paul 1999). Molting and mating are also asynchronous and protracted (Otto and Cummiskey 1985; Shirley and Zhou 1997) with some indications of seasonality (Hiramoto 1985). Molt increment for large males (adults) in Southeast Alaska is 16.3 mm CL per molt (Koeneman and Buchanan 1985) and was estimated at 14.4 mm CL for legal males in the **EAG** (Watson *et al.* 2002). Annual molting probability of males decreases with increasing size, which results in a protracted inter-molt period and creates difficulty in determining annual molt probability (Watson *et al.* 2002). Male size-at-maturity varies among stocks (Webb 2014) and declines with increasing latitude from about 130 mm CL in the Aleutian Islands to 90 mm CL in Saint Matthew Island section (Somerton and Otto 1986). Along with a lack of annual survey data, limited stock-specific life history stock information prevents development of the standard length-based assessment model.

5. Brief summary of management history:

A complete summary of the management history through 2015/16 is provided in Leon *et al.* (2017). The first commercial landing of golden king crab in the Aleutian Islands was in 1975/76 but directed fishing did not occur until 1981/82.

The Aleutian Islands golden king crab fishery was restructured beginning in 1996/97 to replace the Adak and Dutch Harbor areas with the newly created Aleutian Islands Registration Area O and golden king crab in the areas east and west of 174° W longitude were managed separately as two stocks (ADF&G 2002). Hereafter, the east of 174° W longitude stock segment is referred to as **EAG** and the west of 174° W longitude stock segment is referred to as **WAG**. Table 1 provides the historical summary of number of vessels, GHL/TAC, harvest, effort, CPUE and average weight in the Aleutian Islands golden king crab fishery.

The fisheries in 1996/97–1997/98 were managed for GHLs of 1,452 t (3,200,000 lb) in **EAG** and 1,225 t (2,700,000 lb) in **WAG** (Table 1). During 1998/99–2004/05 the fisheries were managed with GHLs of 1,361 t (3,000,000 lb) for **EAG** and 1,225 t (2,700,000 lb) for **WAG**. During 2005/06–2007/08 the fisheries were managed with a total allowable catch (TAC) of 1,361 t (3,000,000 lb) for **EAG** and a TAC of 1,225 t (2,700,000 lb) for **WAG**. By state regulation (5 AAC 34.612), TAC for the Aleutian Islands golden king crab fishery during 2008/09–2011/12 was 1,429 t (3,150,000 lb) for **EAG** and 1,286 t (2,835,000 lb) for **WAG**. In March 2012 the BOF changed 5 AAC 34.612 so that the TAC beginning in 2012/13 would be 1,501 t (3,310,000 lb) for the **EAG** and 1,352 t (2,980,000 lb) for **WAG**. Additionally, the BOF added a provision to 5 AAC 34.612 that allows ADF&G to lower the TAC below the specified level if conservation concerns arise. The TAC for 2016/17 (and 2017/18) was reduced by 25% for **WAG** to 1,014 t (2,235,000 lb) while keeping the TAC for **EAG** at the same level as the previous season.

During 1996/97–2019/20 the annual retained catch during commercial fishing (including cost-recovery fishing that occurred during 2013/14–2019/20) has averaged 2% below the annual GHL/TACs. During 1996/97–2019/20, the retained catch has been as much as 13% below (1998/99) and as much as 6% above (2000/01) the GHL/TAC.

A summary of other relevant State of Alaska fishery regulations and management actions pertaining to the Aleutian Islands golden king crab fishery is provided below:

Beginning in 2005/06 the Aleutian Islands golden king crab fishery has been prosecuted under the Crab Rationalization Program. Accompanying the adoption of crab rationalization program was implementation of a community development quota (CDQ) fishery for golden king crab in the

eastern Aleutians (i.e., **EAG**) and the Adak Community Allocation (ACA) fishery for golden king crab in the western Aleutians (i.e., **WAG**; Hartill 2012). The CDQ fishery in the eastern Aleutians is allocated 10% of the golden king crab TAC for the area east of 174° W longitude and the ACA fishery in the western Aleutians is allocated 10% of the golden king crab TAC for the area west of 174° W longitude. The CDQ fishery and the ACA fishery are managed by ADF&G and prosecuted concurrently with the individual fisheries quota fishery.

Golden king crab may be commercially fished only with king crab pots (defined in state regulation *5 AAC 34.050*). Pots used to fish for golden king crab in the Aleutian Islands Area must be operated from a shellfish longline and, since 1996, each pot must have at least four escape rings of five and one-half inches minimum inside diameter installed on the vertical plane or at least one-third of one vertical surface of the pot composed of not less than nine-inch stretched mesh webbing to permit escapement of undersized golden king crab (*5 AAC 34.625 (b)*). Prior to the regulation requiring an escape mechanism on pots, some participants in the Aleutian Islands golden king crab fishery voluntarily sewed escape rings (typically 139 mm [5.5 inches]) into their gear or, more rarely, included panels with escape mesh (Beers 1992). Regarding the gear used since the establishment of *5 AAC 34.625 (b)* in 1996, Linda Kozak, a representative of the industry, reported in a 19 September 2008 email to the Crab Plan Team, "... the golden king crab fleet has modified their gear to allow for small crab sorting," and provided a written statement from Lance Nylander, of Dungeness Gear Works in Seattle, who "believes he makes all the gear for the golden king crab harvesting fleet," saying that, "Since 1999, DGW has installed 9[-inch] escape web on the door of over 95% of Golden Crab pot orders we manufactured." A study to estimate the contact-selection curve for male golden king crab was conducted aboard one vessel commercial fishing for golden king crab during the 2012/13 season and found gear and fishing practices used by that vessel were highly effective in reducing bycatch of sublegal-sized males and females (Vanek *et al.* 2013). In March 2011 (effective for 2011/12), the BOF amended *5 AAC 34.625 (b)* to relax the "biotwine" specification for pots used in the Aleutian Islands golden king crab fishery relative to the requirement in *5 AAC 39.145* that "(1) a sidewall ...of all shellfish and bottomfish pots must contain an opening equal to or exceeding 18 inches in length... The opening must be laced, sewn, or secured together by a single length of untreated, 100 percent cotton twine, no larger than 30 thread." Regulation *5 AAC 34.625 (b)(1)* allows the opening described in *5 AAC 39.145 (1)* to be "laced, sewn, or secured together by a single length of untreated, 100 percent cotton twine, no larger than 60 [rather than 30] thread."

Regulation (*5 AAC 34.610 (b)*) sets the commercial fishing season for golden king crab in the Aleutian Islands Area as 1 August through 30 April. That regulatory fishing season became effective in 2015/16 (the commercial fishing season was set in regulation as 15 August through 15 May during 2005/06–2014/15).

Current regulations (*5 AAC 39.645 (d)(4)(A)*) stipulate that onboard observers are required on catcher vessels during the time that at least 50% of the retained catch is captured in each of the three trimesters of the 9-month fishing season. Onboard observers are always required on catcher-processor vessels during the fishing season.

In addition, the commercial golden king crab fishery in the Aleutian Islands Area may only retain at least 6.0-inches (152.4 mm) carapace width (CW), including spines (*5 AAC 34.620 (b)*), which is at least one annual molt increment larger than the 50% maturity length of 120.8 mm CL for males estimated by Otto and Cummiskey (1985). A carapace length (CL) ≥ 136 mm is used to identify legal-size males when CW measurements are not available (Table 3-5 in NPFMC 2007b).

Note the size limit for golden king crab has been 6-inches (152.4 mm) CW for the entire Aleutian Islands Area since the 1985/86 season. Prior to the 1985/86 season, the legal-size limit was 6.5-inches (165.1 mm) CW for at least one of the now-defunct Adak or Dutch Harbor Registration Areas.

We re-evaluated the male maturity size using 1991 pot survey measurements of carapace length and chela height in **EAG** and 1984 NMFS measurements in **WAG** (Siddeek *et al.* 2018). Bootstrap analysis of chela height and carapace length data provided the median 50% male maturity length estimates of 107.02 mm CL in **EAG** and 107.85 mm CL in **WAG**. We used a knife-edge 50% maturity length of 111.0 mm CL, which is the lower limit of the next upper size bin, for mature male biomass (MMB) estimation.

Daily catch and catch-per-unit effort (CPUE) are determined in-season to monitor fishery performance and progress towards the respective TACs. Figures 6 to 8 provide the 1985/86–2018/19 time series of catches, CPUE, and the geographic distribution of catch during the 2018/19 fishing season. Increases in CPUE were observed during the late 1990s through the early 2000s, and with the implementation of crab rationalization in 2005. This is likely due to changes in gear configurations in the late 1990s (crab harvesters, personal communication, 1 July 2008) and, after rationalization, to increased soak time (Siddeek *et al.* 2015), and decreased competition owing to the reduced number of vessels fishing. Decreased competition could allow crab vessels to target only the most productive fishing areas. Trends in fishery CPUE within the areas **EAG** and **WAG** generally paralleled each other during 1985/86–2010/11 but diverged during 2011/12–2019/20 (an increasing trend in **EAG** and a decreasing followed by increasing trends in **WAG**). Sharp increases in CPUE were observed since 2016/17 in **WAG** and 2017/18 in **EAG**, with moderate declines in 2019/20.

6. Brief description of the annual ADF&G harvest strategy:

In March 2019, the BOF adopted a revised harvest strategy (Daly *et al.* 2019). The annual TAC is set by state regulation, 5 AAC 34.612 (Harvest Levels for Golden King Crab in Registration Area O), per:

- (a) In that portion of the Registration Area O east of 174° W. long., the total allowable catch level shall be established as follows:
 - (1) if MMA_E is less than 25 percent of $MMA_{E,(1985-2017)}$, the fishery will not open;
 - (2) if MMA_E is at least 25 percent but not greater than 100 percent of $MMA_{E,(1985-2017)}$, the number of legal male golden king crab available for harvest will be computed as $(0.15) \times (MMA_E / MMA_{E,(1985-2017)}) \times (MMA_E)$ or 25 percent of LMA_E , whichever is less; and
 - (3) if MMA_E is greater than 100 percent of $MMA_{E,(1985-2017)}$, the number of legal male golden king crab available for harvest will be computed as $(0.15) \times (MMA_E)$ or 25 percent of LMA_E , whichever is less.
- (b) In that portion of the Registration Area O west of 174° W. long., the total allowable catch level shall be established as follows:
 - (1) if MMA_W is less than 25 percent of $MMA_{W,(1985-2017)}$, the fishery will not open
 - (2) if MMA_W is at least 25 percent but not greater than 100 percent of $MMA_{W,(1985-2017)}$, the number of legal male golden king crab available for harvest will be

- computed as $(0.20) \times (\text{MMA}_W / \text{MMA}_{W,(1985-2017)}) \times \text{MMA}_W$ or 25 percent of LMA_W , whichever is less; and
- (3) if MMA_W is greater than 100 percent of $\text{MMA}_{W,(1985-2017)}$, the number of legal male golden king crab available for harvest will be computed as $(0.20) \times (\text{MMA}_W)$ or 25 percent of LMA_W , whichever is less.
- (c) In implementing this harvest strategy, the department shall consider the reliability of estimates of golden king crab, the manageability of the fishery, and other factors the department determines necessary to be consistent with sustained yield principles and to use the best scientific information available and consider all sources of uncertainty as necessary to avoid overfishing.
- (d) In this section,
- (1) MMA_E means the abundance of male golden king crab in the portion of the Aleutian Islands Management Area O east of 174° W. long that are greater than or equal to 111 millimeters in carapace length estimated by the stock assessment model for the time prior to the start of the fishery;
 - (2) $\text{MMA}_{E,(1985-2017)}$ means the mean value of the abundance of male golden king crab in the portion of the Aleutian Islands Management Area O east of 174° W. long that are greater than or equal to 111 millimeters in carapace length estimated by the stock assessment model for the time prior to the start of the fishery for the period 1985 – 2017;
 - (3) LMA_E means the abundance of male golden king crab in the portion of the Aleutian Islands Management Area O east of 174° W. long that are greater than or equal to 136 millimeters in carapace length estimated by the stock assessment model for the time prior to the start of the fishery;
 - (4) MMA_W means the abundance of male golden king crab in the portion of the Aleutian Islands Management Area O west of 174° W. long that are greater than or equal to 111 millimeters in carapace length estimated by the stock assessment model for the time prior to the start of the fishery;
 - (5) $\text{MMA}_{W,(1985-2017)}$ means the mean value of the abundance of male golden king crab in the portion of the Aleutian Islands Management Area O west of 174° W. long that are greater than or equal to 111 millimeters in carapace length estimated by the stock assessment model for the time prior to the start of the fishery for the period 1985 – 2017;
 - (6) LMA_W means the abundance of male golden king crab in the portion of the Aleutian Islands Management Area O west of 174° W. long that are greater than or equal to 136 millimeters in carapace length estimated by the stock assessment model for the time prior to the start of the fishery.

In addition to the retained catch that is limited by the TAC established by ADF&G under 5 AAC 34.612, ADF&G has authority to annually receive receipts up to \$500,000 through cost-recovery fishing on Aleutian Islands golden king crab. The retained catch from that cost-recovery fishing is not counted against attainment of the annually established TAC.

7. Summary of the history of the basis and estimates of MMBMSY or proxy MMBMSY:

We estimated the proxy MMB_{MSY} as $\text{MMB}_{35\%}$ using the Tier 3 estimation procedure, which is explained in a subsequent section.

D. Data

1. Summary of new information:

(a) Commercial fishery retained catch by size, estimated total catch by size, groundfish male discard catch by size, observer CPUE index, and commercial fishery CPUE index were updated to include 2019/20 information. Available data by year are shown below

Year	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	.	.	17	18	19	
Ret.C. & Size Comp.	[Blue arrow spanning from 81 to 19]																																					
Total C. & Size Comp.	[Blue arrow spanning from 90 to 19]																																					
Ground fish ByC. & Size Comp.	[Blue arrow spanning from 90 to 19]																																					
Observ. CPUE	[Blue arrow spanning from 96 to 19]																																					
Fishery CPUE	[Blue arrow spanning from 85 to 19]																																					
Tag release										◆			◆			◆			◆			◆																
Tag Recovery	[Blue arrow spanning from 90 to 19]																																					

2. Data presented as time series:

a. Total Catch:

Fish ticket data on retained catch weight, catch numbers, effort (pot lifts), CPUE, and average weight of retained catch for 1981/82–2019/20 (Table 1). Estimated total catch weight for 1990/91–2019/20 (Table 2a).

b. Bycatch and discards:

Retained catch, bycatch mortality (male and female of all sizes) separated by the crab fishery and groundfish fishery, and total fishery mortality for 1981/82–2019/20 (Table 2). Crab fishery discards are available after observer sampling was established in 1988/89. Observer data for the 1988/89–1989/90 seasons are not considered reliable. Table 2 provides crab fishery discards and groundfish fishery bycatch for 1991/92–2019/20 seasons.

c. Catch-per-unit-effort:

Pot fishery and observer nominal retained and total CPUE, pot fishery effort, observer sample size, and estimated observer CPUE index delineated by **EAG** and **WAG** for 1985/86–2019/20 (Table 3).

Estimated commercial fishery CPUE index with coefficient of variation (Table 4 for **EAG** and Table 13 for **WAG**). The estimation methods, and CPUE fits are described in Appendix B.

d. Catch-at-length:

Information on length compositions are provided (Figures 9 to 11 for **EAG**; and 27 to 29 for **WAG**).

e. Survey biomass estimates:

Estimates are not available for the area because no systematic surveys, covering the entire fishing area, have occurred.

f. Survey catch–at–length:

Not available.

g. Other time series data: None.

3. Data which may be aggregated over time:

Molt and size transition matrix: Tag release – recapture –time at liberty records from 1991, 1997, 2000, 2003, and 2006 male tag crab releases were aggregated by year at liberty to determine the molt increment and size transition matrix by the integrated model.

Weight-at-length: Male length-weight relationship: $W = aL^b$ where $a = 3.7255 \cdot 10^{-4}$, $b = 3.0896$ (updated estimates).

Natural mortality: A previous model estimated fixed natural mortality value of 0.21 yr⁻¹, was used in the assessment.

4. Information on any data sources that were available, but were excluded from the assessment:

Data from triennial ADF&G pot surveys for Aleutian Islands golden king crab in a limited area in **EAG** (between 170° 21' and 171° 33' W longitude) that were performed during 1997 (Blau *et al.* 1998), 2000 (Watson and Gish 2002), 2003 (Watson 2004), and 2006 (Watson 2007) are available, but were not used in this assessment. However, the tag release and recapture data from these surveys were used.

Data from the cooperative pot surveys conducted during 2015 to 2019 are available but is limited in time span for full usage. The **EAG** survey covers the full time series but **WAG** survey started only in 2018. We incorporate the **EAG** data in a model scenario as a test run in this assessment.

E. Analytic Approach

1. History of modeling approaches for this stock:

A size structured assessment model based on only fisheries data was under development for several years for the **EAG** and **WAG** golden king crab stocks and accepted in 2016 for OFL and ABC setting for the 2017/18 season. The CPT in January 2017 and SSC in February 2017 recommended using the Tier 3 procedure to set the OFL and ABC. They also suggested using the maturity data to estimate the male mature biomass (MMB). We followed these suggestions in this report to estimate the model based OFL and ABC.

2. Model Description:

a. Description of overall modeling approach:

The underlying population dynamics model is male-only and length-based (Appendix A). This model combines commercial retained catch, total catch, groundfish fishery discarded catch, standardized observer legal size catch-per-unit-effort (CPUE) indices, fishery retained catch size composition, total catch size composition, and tag recaptures by release-recapture length to estimate stock assessment parameters. The tagging data were used to calculate the size transition matrix. To estimate the MMB, we used the knife-edge 50% maturity based on the chela height and carapace length data analysis. To include a long time series of CPUE indices for stock abundance contrast, we also considered the 1985/86–1998/99 legal size standardized CPUE indices as a separate likelihood component in all scenarios (see Table T1).

There were significant changes in fishing practice associated with changes in management regulations (e.g., constant TAC since 1996/97 and crab rationalization since 2005/06), pot configuration (escape web on the pot door increased to 9-inch since 1999), and improved observer recording in Aleutian Islands golden king crab fisheries since 1998. These changes prompted us to consider two sets of catchability and total selectivity parameters with only one set of retention parameters for the periods 1985/86–2004/05 and 2005/06–2019/20.

We fitted the observer and commercial fishery CPUE indices with estimated (by GLM) standard errors and an additional model estimated constant variance. The assessment model predicted total and retained CPUEs. However, we compared only the predicted retained CPUE with the observer legal size crab CPUE indices in the likelihood function because observer recordings of legal-size crabs are reliable.

The data series ranges used for the **WAG** are the same as those for **EAG**.

b. Software:

AD Model Builder (Fournier *et al.* 2012).

c.–f. Details are given in Appendix A.

g. Critical assumptions and consequences of assumption failures:

Because of the lack of an annual stock survey, we relied heavily on standardized CPUE indices (Appendix B) and catch and size composition information to determine the stock abundance trends in both regions. We assumed that the observer and fish ticket CPUE indices are linearly related to exploitable abundance. We kept M constant at 0.21

yr⁻¹ and knife-edge maturity size at 111 mm CL (Siddeek *et al.* 2018). We assumed directed pot fishery discard mortality at 0.20 yr⁻¹, overall groundfish fishery mortality at 0.65 yr⁻¹ (mean of groundfish pot fishery mortality [0.5 yr⁻¹] and groundfish trawl fishery mortality [0.8 yr⁻¹]), groundfish fishery selectivity at full selection for all length classes (selectivity = 1.0). Any discard of legal-size males in the directed pot fishery was not considered in this analysis. These fixed values invariably reduced the number of model parameters to be estimated and helped in convergence. We assumed different q 's (scaling parameter for standardized CPUE in the model, Equation A.13) and logistic selectivity patterns (Equation A.9) for different periods for the pot fishery.

h. Changes to any of the above since the previous assessment:

None.

i. Model code has been checked and validated.

The codes have been checked at various times by independent reviewers and the current codes are available from the first author.

3. Model Selection and Evaluation

a. Description of alternative model configurations:

We considered seven models for **EAG** and four for **WAG** (Table T1). We presented OFL and ABC results for all models separately for **EAG**, **WAG**, and the entire **AI** in the executive summary tables. We considered model 20_1 as the base model. It considers:

- i) Initial abundance by the equilibrium condition considering the mean number of recruits for 1987–2012: The equilibrium abundance was determined for 1960, projected forward with only M and annual recruits until 1980, then retained catches removed during 1981–1984 and projected to obtain the initial abundance in 1985 (see Equations A.4 and A.5).
- ii) Observer CPUE indices for 1995/96–2019/20.
- iii) Fishery CPUE indices for 1985/86–1998/99.
- iv) Initial (Stage-1) weighting of effective sample sizes: number of vessel-days for retained and total catch size compositions, and number of fishing trips for groundfish discard size composition (the groundfish size composition was not used in the model fitting); and (Stage-2) iterative re-weighting of effective sample sizes by the Francis method.
- v) Two catchabilities and two sets of logistic total selectivities for the periods 1985/86–2004/05 and 2005/06–2019/20, and a single set of logistic retention curve parameters.
- vi) Full selectivity (selectivity = 1.0) for groundfish fishery bycatch.
- vii) Knife-edge 50% maturity size of 111 mm CL.
- viii) Stock dynamics $M = 0.21$ yr⁻¹, pot fishery handling mortality = 0.2 yr⁻¹, and mean groundfish bycatch handling mortality = 0.65 yr⁻¹.

- ix) Size transition matrix using tagging data estimated by the normal probability function with the logistic molt probability sub-model. The tag-recaptures were treated as Bernoulli trials (i.e., Stage-1 weighting).
- x) The time period, 1987–2012, was used to determine the mean number of recruits for $MMB_{35\%}$ (a proxy for MMB_{MSY}) estimation under Tier 3.

The salient features and variations from the base scenario of all other scenarios are listed in Table T1. The list of fixed and estimable parameters is provided in Table A1 and detail weights with coefficient of variations (CVs) assigned to each type of data are listed in Table A2.

Best estimates of parameter values for models 20_1b and 20_2 were jittered to confirm model global convergence. The results indicated that global convergence was achieved for most runs (Appendix D).

Table T1. Features of all model scenarios: Initial condition was estimated in year 1960 by the equilibrium condition; two catchability and two sets of logistic total selectivity curves were used for the pre- and post-rationalization periods; a single retention curve was used for the whole period; a knife-edge minimum maturity size of 111 mm CL was used for MMB calculation; and a common M of 0.21 yr⁻¹ was used. The effective sample sizes for size compositions were estimated in two stages: Stage-1: as the number of vessel days/trips and Stage-2: as the Francis re-iteration method. Changes in model specifications are highlighted by the shaded text.

Model	CPUE Data Type	Time Period for Mean Number of Recruit Calculation for (a) Initial Equilibrium Abundance Composition and (b) Reference Points Estimations
20_1 (accepted model in May 2019, implemented with up to 2019/20 data)	Observer data from 1995/96–2019/20 Fish ticket data from 1985/86–1998/99. Observer CPUE standardization by negative binomial and Fish ticket CPUE standardization by lognormal models	1987–2012
20_1b	20_1+ Fish ticket CPUE standardization by negative binomial	EAG:1986–2017; WAG:1987–2018
20_1b Ver2	20_1b+	EAG & WAG:1987–2012
20_1c	20_1b+ cooperative survey CPUE indices for 2015–2019.	EAG:1986–2017
20_1d	20_1c+ restrict cooperative survey CPUE indices to 2015–2018	EAG:1986–2017
20_2	20_1b+ Year:Area interaction for observer CPUE standardization.	EAG:1986–2017; WAG:1987–2018
20_2b	20_2+ cooperative survey CPUE indices for 2015–2019	EAG:1986–2017

b. Progression of results:

The OFL and ABC estimates are similar to estimates by the 2019 model.

c. Label the approved model from the previous year as model:

We used the notation 20_1 for the base model which came from the last year accepted assessment model, 19_1.

d. Evidence of search for balance between realistic and simpler models:

Unlike annually surveyed stocks, Aleutian Islands golden king crab stock biomass is difficult to track, and several biological parameters are assumed based on knowledge from red king crab (e.g., handling mortality rate of 0.2 yr^{-1}) due to a lack of species/stock specific information. We fixed several model parameters after initially running the model with free parameters to reduce the number of parameters to be estimated (e.g., groundfish bycatch selectivity parameters were fixed). In CPUE standardization, instead of using the traditional AIC we used the Consistent Akaike Information Criteria (Bozdogan 1987) that considers number of parameters and data points used for fitting when selecting the final model. The models also considered different configuration of parameters to select parsimonious models. The detailed results of all models are provided in tables and figures.

e. Convergence status and criteria:

ADMB default convergence criteria were used.

f. Table of the sample sizes assumed for the size compositional data:

We estimated the initial input effective sample sizes (i.e., Stage-1) either as number of vessel-days for retained and total catch compositions or number of fishing trips for groundfish size composition (note: we did not use the groundfish size composition in the model fit) for all model scenarios. Then we estimated the Stage-2 effective sample sizes iteratively from Stage-1 input effective sample sizes using the Francis' (2011, 2017) mean length-based method.

We provide the initial input sample sizes (Stage-1) and Stage-2 effective sample sizes for models 20_1, 20_1b, and 20_2 in Tables 5 to 7 for **EAG** and Tables 14 to 16 for **WAG**.

g. Provide the basis for data weighting, including whether the input effective sample sizes are tuned, and the survey CV adjusted:

Described previously (f).

h. Do parameter estimates make sense and are they credible?

The estimated parameter values are within the bounds and various plots suggest that the parameter values are reasonable for a fixed M value for the golden king crab stocks.

i. Model selection criteria:

We used several diagnostic criteria to select the appropriate models for our recommendation: CPUE fits, observed vs. predicted tag recapture numbers by time at large and release size, retained and total catch, and groundfish bycatch fits. Figures are provided for all model scenarios in the Results section.

j. **Residual analysis:**

We illustrated residual fits by bubble plots for retained and total catch size composition predictions in various figures in the Results section.

k. **Model evaluation:**

Only one model with several model scenarios is presented and the evaluations are presented in the Results section below.

4. Results

1. List of effective sample sizes and weighting factors:

The Stage-1 and Stage-2 effective sample sizes are listed for various models in Tables 5 to 7 for **EAG** and Tables 14 to 16 for **WAG**. The weights, with the corresponding coefficient of variations specifications, for different data sets are provided in Table A2 for various models for both **EAG** and **WAG**. These weights (with the corresponding coefficient of variations) adequately fitted the length compositions and no further changes were examined.

We used weighting factors for catch biomass, recruitment deviation, pot fishery F, and groundfish fishery F. We set the retained catch biomass weight to an arbitrarily large value (500.0) because retained catches are more reliable than any other data sets. We scaled the total catch biomass weight in accordance with the observer annual sample sizes (number of pots) with a maximum of 250.0. The total catches were derived from observer nominal total CPUE and effort. In some years, observer sample sizes were low (Tables 3). We chose a small groundfish bycatch weight (0.2) based on the September 2015 CPT suggestion for a lower its weight. We used the best fit criteria to choose the lower weight for the groundfish bycatch. Groundfish bycatch of Aleutian Islands golden king crab is very low (Table 2). We set the CPUE weights to 1.0 for all models. We included a constant (model estimated) variance in addition to input CPUE variance for the CPUE fit. We used the Burnham *et al.* (1987) suggested formula for $\ln(\text{CPUE})$ (and $\ln(\text{MMB})$) variance estimation (Equation A.14). However, the estimated additional variance values were small for both observer and fish ticket CPUE indices for the two regions. Nevertheless, the CPUE index variances estimated from the negative binomial and lognormal GLMs were adequate to fit the model, as confirmed by the fit diagnostics (Fox and Weisberg 2011). Parameter estimates are provided in Tables 8 for **EAG** and 17 for **WAG** for all models. The numbers of estimable parameters are listed in Table A1.

2. Include tables showing differences in likelihood:

Tables 12 and 21 list the total and component negative log likelihood values for **EAG** and **WAG**, respectively.

3. Tables of estimates:

- a. The parameter estimates with coefficient of variation for models 20_1, 20_1b, 20_1b Ver 2, and 20_2 are summarized in Tables 8 and 17 for **EAG** and **WAG**, respectively. We have also provided the boundaries for parameter searches in those tables. All parameter estimates were within the bounds.

- b. All models considered molt probability parameters in addition to the linear growth increment and normally distributed growth variability parameters to determine the size transition matrix.
- c. The mature male and legal male abundance time series for selected models (20_1, 20_1b, and 20_2) are summarized in Tables 9 to 11 for **EAG** and Tables 18 to 20 for **WAG**.
- d. The recruitment estimates for those models are summarized in Tables 9 to 11 for **EAG** and Tables 18 to 20 for **WAG**.
- e. The negative log-likelihood component values and total negative log-likelihood values for models 20_1, 20_1b, 20_1b Ver 2, and 20_2 are summarized in Table 12 for **EAG** and Table 21 for **WAG**. Model 20_2 has the minimum total negative log likelihood for **EAG** whereas model 20_1 has the minimum for **WAG**. However, the total negative log likelihood values for the four models for **WAG** were not very different. We may conclude that the input observer CPUE indices with Year and Area interaction appears to have positively influenced the overall fit.

4. Graphs of estimates:

a. Selectivity:

Total selectivity and retention curves of the pre- and post-rationalization periods for selected models are illustrated in Figure 12 for **EAG** and Figure 30 for **WAG**. Total selectivity for the pre-rationalization period was used in the tagging model. The groundfish bycatch selectivity appeared flat in the preliminary analysis, indicating that all size groups were vulnerable to the gear. This is also shown in the size compositions of groundfish bycatch (Figures 11 and 29 for **EAG** and **WAG**, respectively). Thus, we set the groundfish bycatch selectivity to 1.0 for all length-classes in the subsequent analysis.

b. Mature male biomass:

The mature male biomass time series for selected models are depicted in Figures 26 for **EAG** (for seven models) and **WAG** (for four models). Mature male biomass tracked the CPUE trends well for selected models for **EAG** and **WAG**. The biomass variance was estimated using the Burnham *et al.* (1987) suggested formula (Equation A.14). We determined the mature male biomass values on 15 February each year and considered varying time series of recruits (see Table T1) for estimating mean number of recruits for the $MMB_{35\%}$ calculation under a Tier 3 approach.

c. Fishing mortality:

The full selection pot fishery F over time for selected models is shown in Figure 25 for **EAG** (for seven models) and **WAG** (for four models). The F peaked in late 1980s and early to mid-1990s and systematically declined in the **EAG**. Slight increases in F were observed from 2014 to 2016, followed by a decline in the **EAG**. On the other hand, the F in the **WAG** peaked in late 1980s, 1990s and early 2000s, declined in late 2000s, and slightly increased in 2013–2014 before declining.

- d. **F vs. MMB:**

We provide these plots for models 20_1b Ver2 and 20_2 for **EAG** and **WAG** in Figure 43. The 2019 F was below the overfishing levels in both regions.
 - e. **Stock-Recruitment relationship:** None.
 - f. **Recruitment:**

The temporal changes in total number of recruits to the modeled population are illustrated in Figure 14 for **EAG** (for six models) and in Figure 32 for **WAG** (for four models). The recruitment distribution to the model size group (101–185 mm CL) is shown in Figures 15 and 33 for **EAG** and **WAG**, respectively for the respective number of models.
- 5. Evaluation of the fit to the data:**
- g. **Fits to catches:**

The fishery retained and total catch, and groundfish bycatch (observed vs. estimated) plots are illustrated in Figure 17 for **EAG** (for six models) and in Figure 35 for **WAG** (for four models). The 1981/82–1984//85 retained catch plots for respective number of models are depicted in Figures 18 and 36 for **EAG** and **WAG**, respectively. All predicted fits were very close to observed values, especially for retained catch and groundfish bycatch mortality. However, pre-1995 total catch data did not fit well.
 - h. **Survey data plot:**

We provide some cooperative pot survey data plots in Appendix C.
 - i. **CPUE index data:**

The model predicted CPUE vs. input CPUE indices for six models are shown in Figure 24 for **EAG** and for four models in Figure 42 for **WAG**. The CPUE variance was estimated using the Burnham *et al.* (1987) suggested formula (Equation A.14). These figures compare the effects of different CPUE indices input to models.
 - j. **Tagging data:**

The predicted vs. observed tag recaptures by length-class for years 1 to 6 post tagging are depicted in Figure 13 for **EAG** and Figure 31 for **WAG**. The predictions appear reasonable. Note that we used the **EAG** tagging information for size transition matrix estimation for both stocks (**EAG** and **WAG**). The size transition matrices estimated using **EAG** tagging data in the **EAG** and **WAG** models were similar.
 - k. **Molt probability:**

The predicted molt probabilities vs. CL are depicted for six models in Figures 16 for **EAG** and for four models in Figure 34 for **WAG**. The fitted curves appear to be satisfactory.

l. Fit to catch size compositions:

Retained, total, and groundfish discard length compositions are shown in Figures 9 to 11 for **EAG** and 27 to 29 for **WAG**. The retained and total catch size composition fits appear satisfactory. But, the fits to groundfish bycatch size compositions are bad. Note that we did not use the groundfish size composition in any of the model scenario fits.

We illustrate the standardized residual plots as bubble plots of size composition over time for retained catch (Figures 19 and 21 for **EAG**, and 37 and 39 for **WAG**) and for total catch (Figures 20 and 22 for **EAG**, and 38 and 40 for **WAG**) for two models (20_1b and 20_2). The retained catch bubble plots do not appear to exhibit major pronounced patterns among residuals for the selected models.

m. Marginal distributions for the fits to the composition data:

We did not provide this plot in this report.

n. Plots of implied versus input effective sample sizes and time series of implied effective sample sizes:

We did not provide the plots or table values of implied vs. input effective sample sizes in this report. However, we provide the Stage-1 and the re-weighted Stage-2 effective sample sizes in Tables 5 to 7 for **EAG** and in Tables 14 to 16 for **WAG**, respectively for models 20_1, 20_1b, and 20_2.

o. Tables of RMSEs for the indices:

We did not provide this table in this report.

p. Quantile-quantile (Q-Q) plots:

We did not provide these plots for model fits in this report. However, we provide a Q-Q plot for cooperative survey CPUE fit in Appendix C.

6. Retrospective and historical analysis:

The retrospective fits for scenarios 20_1, 20_1b, 20_1b Ver 2, and 20_2 are shown in Figure 23 for **EAG** and in Figure 41 for **WAG**. The retrospective fits, prepared for the whole time series 1961 to 2019, did not show severe departure when five terminal years' data were sequentially removed, especially for **WAG**, and hence the current formulation of the model appears stable. The modified Mohn rho (1999) values are also given in the figures.

Mohn rho (ρ) formula, modified by Deroba (2014), is:

$$Mohn \rho = \frac{\sum_{n=1}^x \frac{[\widehat{MMB}_{y=T-n,T-n} - \widehat{MMB}_{y=T-n,T}]}{\widehat{MMB}_{y=T-n,T}}}{x}$$

where, $\widehat{MMB}_{y=T-n,T-n}$ is the MMB estimated for year T-n (left subscript) using data up to T-n years (right subscript), T is the terminal year of the entire data, x is the total number of peels, most recent year's data is "peeled off" recursively n times, where n =1, 2, 3. ...x. We used five peels (x=5) and our T =2019.

The low values ($\ll 1.0$) of Mohn rho indicate no severe model misspecification, especially for **WAG**. A severe drop in modeled biomass from the initial MMB occurred when the fishery time series started in 1981.

7. Uncertainty and sensitivity analysis:

The main task was to determine a plausible size transition matrix to project the population over time. In a previous study, we investigated the sensitivity of the model to determining the size transition matrix by using or not using a molt probability function (Siddeek *et al.* 2016a). The model fit improved when molt probability model is included. Therefore, we included a molt probability sub-model for the size transition matrix calculation in all models.

8. Conduct ‘jitter analysis’:

We conducted jitter analysis on models 20_1b and 20_2 (Appendix D). The results indicated that global convergence was achieved for most runs.

F. Calculation of the OFL

1. Specification of the Tier level:

In the following section, we provide the Tier 3 method to determine OFL and ABC.

2. List of parameter and stock size estimates (or best available proxies thereof) required by limit and target control rules specified in the fishery management plan:

The critical assumptions for MMB_{MSY} reference point estimation of Aleutian Islands golden king crab are:

- a. Natural mortality is constant.
- b. A fixed growth transition matrix is adequately estimated from tagging data and a molt probability sub-model.
- c. Total fishery selectivity and retention curves are length-dependent and the 2005/06–2019/20 period selectivity estimates are applicable.
- d. Groundfish bycatch fishery selectivity is kept constant at 1.0 for all length groups.
- e. Model estimated recruits (in millions of crab) are valid for different time periods considered on chosen given model.
- f. Model estimated groundfish bycatch mortality values are appropriately averaged for the period 2010/11–2019/20 (10 years).
- g. A knife-edge 50% maturity size of 111 mm CL, as used for MMB estimation, is correct.

Method:

We simulated the population abundance starting from the model estimated terminal year stock size by length, model estimated parameter values, a fishing mortality value (F), and a constant number of annual recruits. Once stock dynamics stabilized (we used the 99th year estimates) for an F, we calculated the MMB/R for that F.

We computed the relative MMB/R in percentage, $\left(\frac{MMB}{R}\right)_{x\%}$ (where $x\% = \frac{MMB_F}{\frac{R}{MMB_0}} \times 100$ and MMB_0/R is the virgin MMB/R for different F values.

$F_{35\%}$ is the F value producing an MMB/R value equal to 35% of MMB_0/R .

$MMB_{35\%}$ is estimated using the following formula:

$MMB_{35\%} = \left(\frac{MMB}{R}\right)_{35} \times \bar{R}$, where \bar{R} is the mean number of model estimated recruits for a selected period.

3. Specification of the OFL:

a. Provide the equations (from Amendment 24) on which the OFL is to be based:

F_{OFL} uses Equation A.28. The OFL is estimated by an iterative procedure accounting for intervening total removals (see Appendix A).

b. Basis for projecting MMB to the time of mating:

We followed the NPFMC 2007a guideline.

c. Specification of F_{OFL} , OFL, and other applicable measures (if any) relevant to determining whether the stock is overfished or if overfishing is occurring:

The 2019/20 fishery data indicated that overfishing did not occur (Total Catch < OFL) and the stock did not reach the overfished status ($MMB > MSST$). See Management Performance table below. The OFL and ABC values for 2020/21 in the table below are the recommended values. The TACs for 2015/16–2016/17 in the table below do not include landings towards a cost-recovery fishery goal, but the catches towards cost-recovery fishing are included in the retained and total catches.

Status and catch specifications (1000 t) of Aleutian Islands golden king crab

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch ^a	OFL	ABC ^b
2016/17	N/A	N/A	2.515	2.593	2.947	5.69	4.26
2017/18	6.044	14.205	2.515	2.585	2.942	6.048	4.536
2018/19	5.880	17.848	2.883	2.965	3.355	5.514	4.136
2019/20	5.909 ^c	16.323 ^c	3.257	3.274 ^d	3.693 ^d	5.249	3.937
2020/21 ^e		14.760				4.793	3.595
2020/21 ^f		15.106				4.993	3.745
2020/21 ^g		14.774				4.798	3.599

Status and catch specifications (million lb) of Aleutian Islands golden king crab

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch ^a	OFL	ABC ^b
2016/17	N/A	N/A	5.545	5.716	6.497	12.53	9.40
2017/18	13.325	31.315	5.545	5.699	6.487	13.333	10.000
2018/19	12.964	39.348	6.356	6.536	7.396	12.157	9.118
2019/20	13.027 ^c	35.985 ^c	7.180	7.219 ^d	8.141 ^d	11.572	8.679
2020/21 ^e		32.540				10.566	7.925
2020/21 ^f		33.303				11.008	8.256
2020/21 ^g		32.571				10.579	7.934

- Total retained catch plus estimated bycatch mortality of discarded bycatch during crab fisheries and groundfish fisheries.
- 25% buffer was applied to total catch OFL to determine ABC.
- MSST and MMB determined by Model 20_1b Ver 2
- 100% TAC was achieved in **EAG**, but over 96% TAC was achieved in **WAG** at the time of this assessment. The **WAG** fishery is ongoing.
- Model 20_1b, up to 2019/20 data, mean number of recruit calculation time period for **EAG**: 1986–2017 and for **WAG**: 1987–2018.
- Model 20_2, up to 2019/20 data.
- Model 20_1b Ver 2, up to 2019/20 data, mean number of recruit calculation time period for **EAG** and **WAG**: 1987–2012.

4. Specification of the retained portion of the total catch OFL:

The retained catch portion of the total-catch OFL for **EAG**, **WAG**, and the entire Aleutian Islands (**AI = EAG + WAG**) stock were calculated for the three models (20_1, 20_1b, and 20_2):

Model 20_1:

EAG: 2,899 t (6.391 million lb)

WAG: 1,693 t (3.732 million lb)

AI: 4,592 t (10.123 million lb).

Model 20_1b:

EAG: 2,870 t (6.327 million lb)

WAG: 1,697 t (3.741 million lb)

AI: 4,567 t (10.068 million lb).

Model 20_2:

EAG: 3,011 t (6.638 million lb)

WAG: 1,748 t (3.853 million lb)

AI: 4,759 t (10.491 million lb).

G. Calculation of ABC

We estimated the cumulative probability distribution of OFL assuming a log normal distribution of OFL. We calculated the OFL at the 0.5 probability and the maximum ABC at the 0.49 probability and considered an additional buffer by setting $ABC = 0.75 * OFL$

We provide the ABC estimates with the 25% buffer for **EAG**, **WAG**, and **AI** considering models 20_1, 20_1b, and 20_2:

Model 20_1:

EAG: ABC = 2,262 t (4.986 million lb)

WAG: ABC = 1,352 t (2.981 million lb)

AI: ABC = 3,614 t (7.967 million lb).

Model 20_1b:

EAG: ABC = 2,239 t (4.937 million lb)

WAG: ABC = 1,355 t (2.988 million lb)

AI: ABC = 3,594 t (7.925 million lb).

Model 20_2:

EAG: ABC = 2,350 t (5.181 million lb)

WAG: ABC = 1,395 t (3.075 million lb)

AI: ABC = 3.745 t (8.256 million lb).

1. List of variables related to scientific uncertainty:

- Models rely largely on fisheries data.
- Observer and fisheries CPUE indices played a major role in the assessment model.
- Natural mortality, 0.21 yr^{-1} , was estimated in the previous model and not independently estimated here.
- The time period to compute the average number of recruits relative to the assumption that this represents “a time period determined to be representative of the production potential of the stock.”
- Fixed bycatch mortality rates were used in each fishery (crab fishery and the groundfish fishery) that discarded golden king crab.
- Discarded catch and bycatch mortality for each fishery that bycatch occurred during 1981/82–1989/90 were not available.

2. List of additional uncertainties for alternative sigma-b.

We recommend a buffer of 25% to account for additional uncertainties.

3. Author recommended ABC:

Authors recommend two ABC options based on 25% buffer on the OFL under scenarios 20_1bVer2 and 20_2.

H. Data Gaps and Research Priorities

1. Recruit abundances were tied to commercial catch sampling data. The implicit assumption in the analysis was that the estimated recruits come solely from the same exploited stock through growth and mortality. The current analysis did not consider that additional recruitment may occur through immigration from neighboring areas and possibly separate sub-stocks. The analysis also did not consider emigration from the study area, which would result in an assumption of increased M or a reduced estimate of recruits. Extensive tagging experiments or resource surveys are needed to investigate stock distributions.
2. We estimated M in the model. However, an independent estimate of M is needed for comparison, which could be achieved with tagging experiments.
3. An extensive tagging study may provide independent estimates of molting probability and growth. We used historical tagging data to determine the size transition matrix.
4. An arbitrary 20% handling mortality rate on discarded males was used, which was obtained from the red king crab literature (Kruse *et al.* 2000; Siddeek 2002). An experimental-based independent estimate of handling mortality is needed for Aleutian Islands golden king crab.
5. The Aleutian King Crab Research Foundation recently initiated crab survey programs in the Aleutian Islands. This program needs to be strengthened and continued for golden king crab research to address some of the data gaps and establish a fishery independent data source.
6. We have been using a length-weight relationship established based on late 1990s data for golden king crab. It is unclear how the recent changes in environmental conditions in the Bering Sea will affect golden king crab growth and survival. Length-weight data from the cooperative 2018 survey were used in the current assessment; however, more measurements are needed to increase the sample size to refine the length-weight model.
7. We have recently added male maturity data in the model to determine a maturity curve for MMB estimation. These maturity data were collected in 1984 and 1991 and need to be updated. More data and more recent data are needed. The ADF&G observer sampling, dock side sampling, and independent survey programs collected male maturity data during the 2018/19 fishery. Preliminary analysis on these data was presented at the January 2020 CPT meeting. The CPT recommended to collect additional data on small size crab (sublegal) to evaluate the maturity fit. ADF&G and cooperative survey are continuing to collect additional data.
8. Morphometric measurements provide size at maturity. Ideally, an experimental study under natural environment condition is needed to collect male size at functional maturity data to determine functional maturity size.

I. Acknowledgments

We thank ADF&G personnel in Kodiak, Dutch Harbor, and Juneau for preparing/providing various fisheries and biological data and plots for this assessment; We appreciate technical and editorial input from CPT and SSC members, and industry personnel.

J. Literature Cited

- Adams, C.F., and A.J. Paul. 1999. Phototaxis and geotaxis of light-adapted zoeae of the golden king crab *Lithodes aequispinus* (Anomura: Lithodidae) in the laboratory. *Journal of Crustacean Biology*. 19(1): 106-110.
- ADF&G (Alaska Department of Fish and Game). 2002. Annual management report for the shellfish fisheries of the Westward Region, 2001. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K02-54, Kodiak, Alaska.
- Barnard, D.R., and R. Burt. 2004. Summary of the 2002 mandatory shellfish observer program database for the general and CDQ fisheries. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K04-27, Kodiak, Alaska.
- Barnard, D.R., R. Burt, and H. Moore. 2001. Summary of the 2000 mandatory shellfish observer program database for the open access fisheries. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K01-39, Kodiak, Alaska.
- Beers, D.E. 1992. Annual biological summary of the Westward Region shellfish observer database, 1991. Alaska Department of Fish and game, Division of Commercial Fisheries, Regional Information Report 4K92-33, Kodiak.
- Blau, S.F., and D. Pengilly. 1994. Findings from the 1991 Aleutian Islands golden king crab survey in the Dutch Harbor and Adak management areas including analysis of recovered tagged crabs. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 4K94-35, Kodiak.
- Blau, S.F., L.J. Watson, and I. Vining. 1998. The 1997 Aleutian Islands golden king crab survey. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 4K98-30, Kodiak.
- Bowers, F.R., M. Schwenzfeier, S. Coleman, B.J. Failor-Rounds, K. Milani, K. Herring, M. Salmon, and M. Albert. 2008. Annual management report for the commercial and subsistence shellfish fisheries of the Aleutian Islands, Bering Sea, and the Westward Region's shellfish observer program, 2006/07. Alaska Department of Fish and Game, Divisions of Sport Fish and Commercial Fisheries, Fishery Management Report No. 08-02, Anchorage, Alaska.
- Bowers, F.R., M. Schwenzfeier, K. Herring, M. Salmon, J. Shaishnikoff, H. Fitch, J. Alas, and B. Baechler. 2011. Annual management report for the commercial and subsistence shellfish fisheries of the Aleutian Islands, Bering Sea, and the Westward Region's shellfish observer program, 2009/10. Alaska Department of Fish and Game, Divisions of Sport Fish and Commercial Fisheries, Fishery Management Report No. 11-05, Anchorage, Alaska.
- Bozdogan, H. 1987. Model selection and Akaike's Information Criterion (AIC): The general theory and its analytical extensions. *Psychometrika*, 52, 345-370.

- Burnham, K.P., D.R. Anderson, G.C. White, C. Brownie, and K.H. Pollock. 1987. Design and analysis methods for fish survival experiments based on release-recapture. American Fisheries Society, Monograph 5, 437p.
- Burnham, K.P. and D.R. Anderson. 2002. Model Selection and Multimodal Inference, A practical Information- Theoretic Approach. 2nd edition. Springer-Verlag, NY, 488p.
- Daly, B., M.S.M. Siddeek, M. Stichert, S. Martell, and J. Zheng. 2019. Recommended harvest strategy for Aleutian Islands golden king crab. Alaska Department of Fish and Game, Fishery Manuscript Series No. 19-03, Anchorage.
- Deroba, J.J. 2014. Evaluating the consequences of adjusting fish stock assessment estimates of biomass for retrospective patterns using Mohn's rho. *North American Journal of Fisheries Management* 34:380-390.
- Feenstra, J., A. Linnane, M. Haddon, and A. Punt. (unpublished, 2019). Impacts on CPUE from vessel fleet composition changes in an Australian lobster (*Jasus edwardsii*) fishery. New Zealand Journal of Marine and Freshwater Research.
- Fournier, D.A., H.J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J. Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim. Methods Softw.* 27:233-249.
- Fox, J., and S. Weisberg. 2011. An R Companion to Applied Regression. Second edition. Sage Publications, Inc. 449 p.
- Francis, R.I.C.C. 2011. Data weighting in statistical fisheries stock assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* 68: 1124–1138.
- Francis, R.I.C.C. 2017. Revisiting data weighting in fisheries stock assessment models. *Fisheries Research* 192: 5-15.
- Gaeuman, W.B. 2011. Summary of the 2009/10 mandatory crab observer program database for the BSAI commercial crab fisheries. Fishery Data Series No. 11-04. Alaska Department of Fish and Game, Kodiak.
- Gaeuman, W.B. 2014. Summary of the 2013/2014 Mandatory Crab Observer Program Database for the Bering Sea/Aleutian Islands commercial crab fisheries. Alaska Department of Fish and Game, Fishery Data Series No. 14-49, Anchorage.
- Hartill, T. 2012. Annual management report for the community development quota and Adak Community Allocation crab fisheries in the Bering Sea and Aleutian Islands, 2010/11. Pages 177–194 in Fitch, H., M. Schwenzfeier, B. Baechler, T. Hartill, M. Salmon, M. Deiman, E. Evans, E. Henry, L. Wald, J. Shaishnikoff, and K. Herring. Annual management report for the commercial and subsistence shellfish fisheries of the Aleutian Islands, Bering Sea and the Westward Region's Shellfish Observer Program, 2010/11. Alaska Department of Fish and Game, Fishery Management Report No. 12-22, Anchorage.
- Hilsinger, J., C. Siddon, and L. Hulbert. 2016. Cooperative red king crab survey in the Adak area, 2015. Anchorage., Alaska Department of Fish and Game, Fishery Data Series No. 16-18.
- Hiramoto, K. 1985. Overview of the golden king crab, *Lithodes aequispina*, fishery and its fishery biology in the Pacific waters of Central Japan. Pages 297-315, In: Proceedings of the

- International King Crab Symposium. Alaska Sea Grant College Program, AK-SG-85-12, Fairbanks, Alaska.
- Jewett, S.C., N.A. Sloan, and D.A. Somerton. 1985. Size at sexual maturity and fecundity of the fjord-dwelling golden king crab *Lithodes aequispina* Benedict from northern British Columbia. *Journal of Crustacean Biology* 5: 377–385.
- Koeneman, T.M., and D.V. Buchanan. 1985. Growth of the golden king crab, *Lithodes aequispina*, in Southeast Alaskan waters. Pages 281-297, In: Proceedings of the International King Crab Symposium. Alaska Sea Grant College Program, AK-SG-85-12, Fairbanks, Alaska.
- Kruse, G.H., L.C. Byrne, F.C. Funk, S.C. Matulich, and J. Zheng. 2000. Analysis of minimum size limit for the red king crab fishery in Bristol Bay, Alaska. *N. Am. J. Fish. Manage.* 20:307-319.
- Leon, J. M., J. Shaishnikoff, E. Nichols, and M. Westphal. 2017. Annual management report for shellfish fisheries of the Bering Sea–Aleutian Islands management area, 2015/16. Alaska Department of Fish and Game, Fishery Management Report No. 17-10, Anchorage.
- Maunder, M.N., and A.E. Punt. 2004. Standardizing catch and effort data: a review of recent approaches. *Fisheries Research* 70: 141-159.
- Mohn, R. 1999. The retrospective problem in sequential population analysis: An investigation using cod fishery and simulated data. *ICES Journal of Marine Science* 56:473-488.
- Moore, H., L.C. Byrne, and M.C. Schwenzfeier. 2000. Summary of the 1999 mandatory shellfish observer program database for the open access fisheries. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K00–50, Kodiak, Alaska.
- Morrison, R., R.K. Gish, and M. Ruccio. 1998. Annual management report for the shellfish fisheries of the Aleutian Islands. Pages 82–139 in ADF&G. Annual management report for the shellfish fisheries of the Westward Region. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K98-39, Kodiak.
- National Marine Fisheries Service (NMFS). 2004. Bering Sea Aleutian Islands Crab Fisheries Final Environmental Impact Statement. National Marine Fisheries Service, Alaska Region, Juneau, August 2004.
- North Pacific Fishery Management Council (NPFMC). 2007a. Initial Review Draft: Environmental Assessment for proposed Amendment 24 to the Fishery Management Plan for Bering Sea and Aleutian Islands King and Tanner Crabs to Revise Overfishing Definitions. 17 January 2007. North Pacific Fishery Management Council, Anchorage.
- North Pacific Fishery Management Council (NPFMC). 2007b. Public Review Draft: Environmental Assessment for proposed Amendment 24 to the Fishery Management Plan for Bering Sea and Aleutian Islands King and Tanner Crabs to Revise Overfishing Definitions. 14 November 2007. North Pacific Fishery Management Council, Anchorage.
- Otto, R.S., and P.A. Cummiskey. 1985. Observations on the reproductive biology of golden king crab (*Lithodes aequispina*) in the Bering Sea and Aleutian Islands. Pages 123-135 In: Proceedings of the International King Crab Symposium. Alaska Sea Grant College Program, AK-SG-85-12, Fairbanks, Alaska.
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

- Shirley, T.C., and S. Zhou. 1997. Lecithotrophic development of the golden king crab *Lithodes aequispinus* (Anomura: Lithodidae). *J. Crust. Biol.* 17(2):207-216.
- Siddeek, M.S.M. 2002. Review of biological reference points used in Bering Sea and Aleutian Islands (king and Tanner) crab management. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 5J02-06, Juneau, Alaska.
- Siddeek, M.S.M., D.R. Barnard, L.J. Watson, and R.K. Gish. 2005. A modified catch-length analysis model for golden king crab (*Lithodes aequispinus*) stock assessment in the eastern Aleutian Islands. Pages 783-805 in Fisheries assessment and management in data limited situations, Alaska Sea Grant College Program, AK-SG-05-02, Fairbanks, Alaska.
- Siddeek, M.S.M., J. Zheng, and D. Pengilly. 2015. Aleutian Islands Golden King Crab (*Lithodes aequispinus*) Model-Based Stock Assessment in Fall 2015. Draft report submitted for the September 2015 Crab Plan Team Meeting. North Pacific Fishery Management Council, Anchorage, Alaska.
- Siddeek, M.S.M., J. Zheng, A.E. Punt, and V. Vanek. 2016a. Estimation of size-transition matrices with and without moult probability for Alaska golden king crab using tag-recapture data. *Fisheries Research* 180:161-168.
- Siddeek, M.S.M., J. Zheng, and D. Pengilly 2016b. Standardizing CPUE from the Aleutian Islands golden king crab observer data. In: T.J. Quinn II, J.L. Armstrong, M.R. Baker, J. Heifetz, and D. Witherell (eds.), *Assessing and Managing Data-Limited Fish Stocks*. Alaska Sea Grant, University of Alaska Fairbanks, Alaska, USA, pp. 97-116.
- Siddeek, M.S.M., J. Zheng, C. Siddon, B. Daly, J. Runnebaum, and M.J. Westphal. 2018. Aleutian Islands Golden king crab model-based stock assessment in Spring 2018. CRAB2018SAFE chapter. North Pacific Fishery Management Council, Anchorage, Alaska.
- Somerton, D.A., and R.S. Otto. 1986. Distribution and reproductive biology of the golden king crab, *Lithodes aequispina*, in the Eastern Bering Sea. *Fishery Bulletin* 81(3): 571-584.
- Starr, P.J. 2012. Standardized CPUE analysis exploration: using the rock lobster voluntary logbook and observer catch sampling programmes. New Zealand Fisheries Assessment Report 2012/34, 75 p.
- Vanek, V., D. Pengilly, and M.S.M. Siddeek. 2013. A study of commercial fishing gear selectivity during the 2012/13 Aleutian Islands golden king crab fishery east of 174° W longitude. Alaska Department of Fish and Game, Division of Sport Fish and Commercial Fisheries, Fishery Data Series No. 13-41.
- Von Szalay, P.G., C.N. Roper, N.W. Raring, and M.H. Martin. 2011. Data report: 2010 Aleutian Islands bottom trawl survey. U.S. Dep. Commerce., NOAA Technical Memorandum NMFS-AFSC-215.
- Watson, L.J. 2004. The 2003 triennial Aleutian Islands golden king crab survey and comparisons to the 1997 and 2000 surveys (revised October 17, 2005). Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K04-42, Kodiak. [Revised 10/17/2005].
- Watson, L.J. 2007. The 2006 triennial Aleutian Islands golden king crab survey. Alaska Department of Fish and Game, Fishery Management Report No. 07-07, Anchorage.

- Watson, L.J., and R.K. Gish. 2002. The 2000 Aleutian Islands golden king crab survey and recoveries of tagged crabs in the 1997–1999 and 2000–2002 fishing seasons. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 4K02-6, Kodiak.
- Watson, L.J., D. Pengilly, and S.F. Blau. 2002. Growth and molting of golden king crabs (*Lithodes aequispinus*) in the eastern Aleutian Islands, Alaska. Pages 169-187 in Crabs in cold water regions: biology, management, and economics, Alaska Sea Grant College Program, AK-SG-02-01, Fairbanks, Alaska.
- Webb, J. 2014. Reproductive ecology of commercially important Lithodid crabs. Pages 285-314 in B.G. Stevens (ed.). King Crabs of the World: Biology and Fisheries Management. CRC Press, Taylor & Francis Group, New York.

Tables

Table 1. Commercial fishery history for the Aleutian Islands golden king crab fishery 1981/82–2019/20: number of vessels, guideline harvest level (GHL; established in lb, converted to t) for 1996/97 – 2004/05, total allowable catch (TAC; established in lb, converted to t) for 2005/06–2019/20, weight of retained catch (harvest; t), number of retained crab, pot lifts, fishery catch-per-unit- effort (CPUE; retained crab per pot lift), and average weight (kg) of landed crab. The values are separated by **EAG** and **WAG** beginning in 1996/97.

Crab Fishing Season	Vessels	GHL/TAC	Harvest ^a	Crab ^b	Pot Lifts	CPUE ^b	Average Weight ^c
1981/82	14–20	–	599	240,458	27,533	9	2.5 ^d
1982/83	99–148	–	4,169	1,737,109	179,472	10	2.4 ^d
1983/84	157–204	–	4,508	1,773,262	256,393	7	2.5 ^d
1984/85	38–51	–	2,132	971,274	88,821	11	2.2 ^e
1985/86	53	–	5,776	2,816,313	236,601	12	2.1 ^f
1986/87	64	–	6,685	3,345,680	433,870	8	2.0 ^f
1987/88	66	–	4,199	2,177,229	307,130	7	1.9 ^f
1988/89	76	–	4,820	2,488,433	321,927	8	1.9 ^f
1989/90	68	–	5,453	2,902,913	357,803	8	1.9 ^f
1990/91	24	–	3,153	1,707,618	215,840	8	1.9 ^f
1991/92	20	–	3,494	1,847,398	234,857	8	1.9 ^f
1992/93	22	–	2,854	1,528,328	203,221	8	1.9 ^f
1993/94	21	–	2,518	1,397,530	234,654	6	1.8 ^f
1994/95	35	–	3,687	1,924,271	386,593	5	1.9 ^f
1995/96	28	–	3,157	1,582,333	293,021	5	2.0 ^f

Information for subsequent seasons is presented separately for EAG, WAG in the rows below

Table 1. (continued)

Crab Fishing Season	Vessels		GHL/TAC		Harvest ^a		Crab ^b		Pot Lifts		CPUE ^b		Average Weight ^c	
	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG
1996/97	14	13	1,452	1,225	1,493	1,145	731,909	602,968	113,460	99,267	7	6	2.04 ^f	1.91 ^f
1997/98	13	9	1,452	1,225	1,588	1,109	780,610	569,550	106,403	86,811	7	7	2.04 ^f	1.95 ^f
1998/99	14	3	1,361	1,225	1,473	768	740,011	410,018	83,378	35,975	9	11	2.00 ^f	1.86 ^f
1999/00	15	15	1,361	1,225	1,392	1,256	709,332	676,558	79,129	107,040	9	6	1.95 ^f	1.86 ^f
2000/01	15	12	1,361	1,225	1,422	1,308	704,702	705,613	71,551	101,239	10	7	2.00 ^f	1.86 ^f
2001/02	19	9	1,361	1,225	1,442	1,243	730,030	686,738	62,639	105,512	12	7	2.00 ^f	1.81 ^f
2002/03	19	6	1,361	1,225	1,280	1,198	643,886	664,823	52,042	78,979	12	8	2.00 ^f	1.81 ^f
2003/04	18	6	1,361	1,225	1,350	1,220	643,074	676,633	58,883	66,236	11	10	2.09 ^f	1.81 ^f
2004/05	19	6	1,361	1,225	1,309	1,219	637,536	685,465	34,848	56,846	18	12	2.04 ^f	1.77 ^f
2005/06	7	3	1,361	1,225	1,300	1,204	623,971	639,368	24,569	30,116	25	21	2.09 ^f	1.91 ^f
2006/07	6	4	1,361	1,225	1,357	1,030	650,587	527,734	26,195	26,870	25	20	2.09 ^f	1.95 ^f
2007/08	4	3	1,361	1,225	1,356	1,142	633,253	600,595	22,653	29,950	28	20	2.13 ^f	1.91 ^f
2008/09	3	3	1,361	1,286	1,426	1,150	666,946	587,661	24,466	26,200	27	22	2.13 ^f	1.95 ^f
2009/10	3	3	1,429	1,286	1,429	1,253	679,886	628,332	29,298	26,489	26	24	2.09 ^f	2.00 ^f
2010/11	3	3	1,429	1,286	1,428	1,279	670,983	626,246	25,851	29,994	26	21	2.13 ^f	2.04 ^f
2011/12	3	3	1,429	1,286	1,429	1,276	668,828	616,118	17,915	26,326	37	23	2.13 ^f	2.09 ^f
2012/13	3	3	1,501	1,352	1,504	1,339	687,666	672,916	20,827	32,716	33	21	2.18 ^f	2.00 ^f
2013/14	3	3	1,501	1,352	1,546	1,347	720,220	686,883	21,388	41,835	34	16	2.13 ^f	1.95 ^f
2014/15	3	2	1,501	1,352	1,554	1,217	719,064	635,312	17,002	41,548	42	15	2.18 ^f	1.91 ^f
2015/16	3	2	1,501	1,352	1,590	1,139	763,604	615,355	19,376	41,108	39	15	2.09 ^f	1.85 ^f

Crab Fishing Season	Vessels		GHL/TAC		Harvest ^a		Crab ^b		Pot Lifts		CPUE ^b		Average Weight ^c	
	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG
2016/17	3	3	1,501	1,014	1,578	1,015	793,983	543,796	24,470	38,118	32	14	1.99 ^f	1.87 ^f
2017/18	3	3	1,501	1,014	1,571	1,014	802,610	519,051	25,516	30,885	31	17	1.96 ^f	1.95 ^f
2018/19	3	3	1,749	1,134	1,830	1,135	940,336	578,221	25,553	29,156	37	20	1.95 ^f	1.96 ^f
2019/20	3	3	1,955	1,302	2,031	1,244	1,057,464	626,735	30,998	38,733	34	16	1.92 ^f	1.98 ^f

Note:

- a. Includes deadloss.
- b. Number of crab per pot lift.
- c. Average weight of landed crab, including dead loss.
- d. Managed with 6.5" carapace width (CW) minimum size limit.
- e. Managed with 6.5" CW minimum size limit west of 171° W longitude and 6.0" minimum size limit east of 171° W longitude.
- f. Managed with 6.0" minimum size limit.

Catch and effort data include cost recovery fishery.

Table 2. Annual weight of total fishery mortality to Aleutian Islands golden king crab, 1981/82 – 2019/20, partitioned by source of mortality: retained catch, bycatch mortality during crab fisheries, and bycatch mortality during groundfish fisheries. For bycatch in the federal groundfish fisheries, historical data (1991–2008) are not available for areas east and west of 174W, and are listed for federal groundfish reporting areas 541, 542, and 543 combined. The 2009– present data are available by separate **EAG** and **WAG** fisheries and are listed as such. A mortality rate of 20% was applied for crab fisheries bycatch, and a mortality rate of 50% for groundfish pot fisheries and 80% for the trawl fisheries were applied.

Season	Bycatch Mortality by Fishery								
	Retained Catch		Bycatch Mortality by Fishery Type (t)				Total Fishery Mortality		Entire AI
	(t)		Crab	Groundfish		(t)			
EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG		
1981/82	490	95							585
1982/83	1,260	2,655							3,914
1983/84	1,554	2,991							4,545
1984/85	1,839	424							2,263
1985/86	2,677	1,996							4,673
1986/87	2,798	4,200							6,998
1987/88	1,882	2,496							4,379
1988/89	2,382	2,441							4,823
1989/90	2,738	3,028							5,766
1990/91	1,623	1,621							3,244
1991/92	2,035	1,397	515	344		0			4,291
1992/93	2,112	1,025	1,206	373		0			4,716
1993/94	1,439	686	383	258		4			2,770
1994/95	2,044	1,540	687	823		1			5,095
1995/96	2,259	1,203	725	530		2			4,719
1996/97	1,738	1,259	485	439		5			3,926
1997/98	1,588	1,083	441	343		1			3,455
1998/99	1,473	955	434	285		1			3,149
1999/00	1,392	1,222	313	385		3			3,316
2000/01	1,422	1,342	82	437		2			3,285
2001/02	1,442	1,243	74	387		0			3,146
2002/03	1,280	1,198	52	303		18			2,850
2003/04	1,350	1,220	53	148		20			2,792
2004/05	1,309	1,219	41	143		1			2,715
2005/06	1,300	1,204	22	73		2			2,601
2006/07	1,357	1,022	28	81		18			2,506
2007/08	1,356	1,142	24	114		59			2,695
2008/09	1,426	1,150	61	102		33			2,772
2009/10	1,429	1,253	111	108	18	5	1,558	1,366	2,923
2010/11	1,428	1,279	123	124	49	3	1,600	1,407	3,006
2011/12	1,429	1,276	106	117	25	4	1,560	1,398	2,957
2012/13	1,504	1,339	118	145	9	6	1,631	1,491	3,122

2013/14	1,546	1,347	113	174	5	7	1,665	1,528	3,192
2014/15	1,554	1,217	127	175	9	5	1,691	1,397	3,088
2015/16	1,590	1,139	165	157	23	2	1,778	1,298	3,076
2016/17	1,578	1,015	203	145	3	3	1,785	1,163	2,947
2017/18	1,571	1,014	219	126	10	2	1,801	1,142	2,942
2018/19	1,830	1,135	240	140	8	2	2,078	1,277	3,355
2019/20	2,031	1,244	275	116	23	3	2,239	1,363	3,693

Table 2a. Time series of estimated total male catch (weight of crabs on the deck without applying any handling mortality) for the **EAG** and **WAG** golden king crab stocks (1990/91–2019/20). The crab weights are for the size range ≥ 101 mm CL and a length-weight formula was used to predict weight at the mid-point of each size bin. NA: no observer sampling to compute catch.

Year	Total Catch Biomass (t)	
	EAG	WAG
1990/91	1,391	3,626
1991/92	5,813	2,537
1992/93	5,484	1,496
1993/94	NA	2,783
1994/95	1,950	4,872
1995/96	3,681	2,099
1996/97	2,037	1,740
1997/98	2,521	1,777
1998/99	2,762	1,070
1999/00	2,260	2,063
2000/01	2,537	2,197
2001/02	2,086	2,107
2002/03	1,796	1,865
2003/04	1,815	1,845
2004/05	1,621	1,859
2005/06	1,731	1,783
2006/07	1,631	1,546
2007/08	1,814	1,602
2008/09	1,811	1,726
2009/10	1,766	1,681
2010/11	1,750	1,592
2011/12	1,765	1,519
2012/13	1,943	1,825
2013/14	1,834	1,910
2014/15	1,962	1,586
2015/16	2,120	1,551
2016/17	2,224	1,544
2017/18	2,031	1,155
2018/19	2,639	1,507
2019/20	2,985	1,714

Table 3. Time series of nominal annual pot fishery retained, observer retained, and observer total catch-per-unit-effort (CPUE, number of crabs per pot lift), total pot fishing effort (number of pot lifts), observer sample size (number of sampled pots), and GLM estimated observer CPUE Index (for Model 20_1) for the **EAG** and **WAG** golden king crab stocks, 1985/86–2019/20. Observer retained CPUE includes retained and non-retained legal-size crabs.

Year	Pot Fishery Nominal Retained CPUE		Obs. Nominal Retained CPUE		Obs. Nominal Total CPUE		Pot Fishery Effort (no.pot lifts)		Obs. Sample Size (no.pot lifts)		Obs. CPUE Index	
	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG	EAG	WAG
	1985/86	11.90	11.90					117,718	118,563			
1986/87	8.42	7.32					155,240	277,780				
1987/88	7.03	7.15					146,501	160,229				
1988/89	7.52	7.93					155,518	166,409				
1989/90	8.49	7.83					155,262	202,541				
1990/91	8.90	7.00	6.84	8.34	13.00	26.67	106,281	108,533	138	340		
1991/92	8.20	7.40	9.84	6.14	36.91	19.17	133,428	101,429	377	857		
1992/93	8.40	5.90	10.44	4.26	38.52	16.83	133,778	69,443	199	690		
1993/94	7.80	4.40	5.91	12.75	20.81	17.23	106,890	127,764	31	174		
1994/95	5.90	4.10	4.66	6.62	12.91	19.23	191,455	195,138	127	1,270		
1995/96	5.90	4.70	6.03	6.03	16.98	14.28	177,773	115,248	6,388	5,598	1.00	1.17
1996/97	6.50	6.10	6.02	5.90	13.81	13.54	113,460	99,267	8,360	7,194	0.94	0.98
1997/98	7.30	6.60	7.99	6.72	18.25	15.03	106,403	86,811	4,670	3,985	0.87	0.98
1998/99	8.90	11.40	9.82	9.43	25.77	23.09	83,378	35,975	3,616	1,876	1.00	1.09
1999/00	9.00	6.30	10.28	6.09	20.77	14.49	79,129	107,040	3,851	4,523	0.92	0.91
2000/01	9.90	7.00	10.40	6.46	25.39	16.64	71,551	101,239	5,043	4,740	0.82	0.84
2001/02	11.70	6.50	11.73	6.04	22.48	14.66	62,639	105,512	4,626	4,454	1.04	0.82
2002/03	12.40	8.40	12.70	7.47	22.59	17.37	52,042	78,979	3,980	2,509	1.10	0.91
2003/04	10.90	10.20	11.34	9.33	19.43	18.17	58,883	66,236	3,960	3,334	0.97	1.16
2004/05	18.30	12.10	18.34	11.14	28.48	22.45	34,848	56,846	2,206	2,619	1.44	1.24
2005/06	25.40	21.20	29.52	23.89	38.55	36.23	24,569	30,116	1,193	1,365	0.98	1.16
2006/07	24.80	19.60	25.13	23.93	33.39	33.47	26,195	26,870	1,098	1,183	0.80	1.10
2007/08	28.00	20.00	31.10	21.01	40.38	32.46	22,653	29,950	998	1,082	0.89	1.00
2008/09	27.30	22.40	29.97	24.50	38.23	38.16	24,466	26,200	613	979	0.88	1.15
2009/10	25.90	23.70	26.60	26.54	35.88	34.08	26,298	26,489	408	892	0.73	1.23
2010/11	26.00	20.90	26.40	22.43	37.10	29.05	25,851	29,994	436	867	0.76	1.10
2011/12	37.30	23.40	39.48	23.63	52.04	31.13	17,915	26,326	361	837	1.08	1.10
2012/13	33.02	20.57	37.82	22.88	47.57	30.76	20,827	32,716	438	1,109	1.04	1.07
2013/14	33.67	16.42	35.94	16.89	46.16	25.01	21,388	41,835	499	1,223	1.02	0.81
2014/15	42.29	15.29	47.01	15.25	60.00	22.67	17,002	41,548	376	1,137	1.34	0.73
2015/16	39.41	14.97	43.27	15.81	58.68	22.14	19,376	41,108	478	1,296	1.26	0.74
2016/17	32.45	14.29	36.89	16.65	52.82	24.41	24,470	38,118	617	1,060	1.05	0.86
2017/18	31.46	16.81	35.18	19.30	54.62	25.54	25,516	30,885	585	760	1.00	0.98
2018/19	36.80	19.83	41.57	22.90	62.97	30.61	25,553	29,156	475	688	1.25	1.18
2019/20	34.11	16.18	40.88	19.25	57.46	27.15	30,998	38,733	540	793	1.16	0.96

Table 4. Time series of negative binomial GLM estimated CPUE indices and coefficient of variation (CV) for the fish ticket based retained catch-per-pot lift for the **EAG** golden king crab stock. The GLM was fitted to the 1985/86 to 1998/99 time series of data.

Year	CPUE Index	CV
1985/86	1.63	0.05
1986/87	1.23	0.05
1987/88	0.96	0.05
1988/89	1.04	0.04
1989/90	1.08	0.03
1990/91	0.99	0.05
1991/92	0.90	0.04
1992/93	0.92	0.04
1993/94	0.91	0.05
1994/95	0.81	0.04
1995/96	0.78	0.04
1996/97	0.78	0.04
1997/98	1.05	0.05
1998/99	1.21	0.05

Table 5. The initial input number of vessel-days/trips and Stage-2 effective sample sizes iteratively estimated by the Francis method for retained, total, and groundfish discard catch size compositions of golden king crab for **model 20_1** fit to **EAG** data. NA: not available.

Year	Initial Input Retained Vessel- Days Sample Size (no)	Stage-2 Retained Effective Sample Size (no)	Initial Input Total Vessel- Days Sample Size (no)	Stage-2 Total Effective Sample Size (no)	Initial Input Groundfish Trip Sample Size (no)	Stage-2 Groundfish Effective Sample Size (no)
1985/86	57	47				
1986/87	11	9				
1987/88	61	50				
1988/89	352	288				
1989/90	792	649			9	4
1990/91	163	134	22	13	13	6
1991/92	140	115	48	28	NA	NA
1992/93	49	40	41	24	2	1
1993/94	340	279	NA	NA	2	1
1994/95	319	261	34	20	4	2
1995/96	879	720	1,117	654	5	2
1996/97	547	448	509	298	4	2
1997/98	538	441	711	416	8	4
1998/99	541	443	574	336	15	7
1999/00	463	379	607	355	14	7
2000/01	436	357	495	290	16	8
2001/02	488	400	510	298	13	6
2002/03	406	333	438	256	15	7
2003/04	405	332	416	243	17	8
2004/05	280	229	299	175	10	5
2005/06	266	218	232	136	12	6
2006/07	234	192	143	84	14	7
2007/08	199	163	134	78	17	8
2008/09	197	161	113	66	15	7
2009/10	170	139	95	56	16	8
2010/11	183	150	108	63	26	12
2011/12	160	131	107	63	13	6
2012/13	187	153	99	58	18	9
2013/14	193	158	122	71	17	8
2014/15	168	138	99	58	16	8
2015/16	190	156	125	73	10	5
2016/17	223	183	155	91	12	6
2017/18	213	175	133	78	12	6
2018/19	218	179	234	137	9	4
2019/20	208	170	230	135	8	4

Table 6. The initial input number of vessel-days/trips and Stage-2 effective sample sizes iteratively estimated by the Francis method for retained, total, and groundfish discard catch size compositions of golden king crab for **model 20_1b** fit to **EAG** data. NA: not available.

Year	Initial Input Retained Vessel- Days Sample Size (no)	Stage-2 Retained Effective Sample Size (no)	Initial Input Total Vessel- Days Sample Size (no)	Stage-2 Total Effective Sample Size (no)	Initial Input Groundfish Trip Sample Size (no)	Stage-2 Groundfish Effective Sample Size (no)
1985/86	57	47				
1986/87	11	9				
1987/88	61	50				
1988/89	352	289				
1989/90	792	650			9	4
1990/91	163	134	22	13	13	6
1991/92	140	115	48	28	NA	NA
1992/93	49	40	41	24	2	1
1993/94	340	279	NA	NA	2	1
1994/95	319	262	34	20	4	2
1995/96	879	721	1,117	650	5	2
1996/97	547	449	509	296	4	2
1997/98	538	441	711	414	8	4
1998/99	541	444	574	334	15	7
1999/00	463	380	607	353	14	7
2000/01	436	358	495	288	16	8
2001/02	488	400	510	297	13	6
2002/03	406	333	438	255	15	7
2003/04	405	332	416	242	17	8
2004/05	280	230	299	174	10	5
2005/06	266	218	232	135	12	6
2006/07	234	192	143	83	14	7
2007/08	199	163	134	78	17	8
2008/09	197	162	113	66	15	7
2009/10	170	139	95	55	16	8
2010/11	183	150	108	63	26	12
2011/12	160	131	107	62	13	6
2012/13	187	153	99	58	18	9
2013/14	193	158	122	71	17	8
2014/15	168	138	99	58	16	8
2015/16	190	156	125	73	10	5
2016/17	223	183	155	90	12	6
2017/18	213	175	133	77	12	6
2018/19	218	179	234	136	9	4
2019/20	208	171	230	134	8	4

Table 7. The initial input number of vessel-days/trips and Stage-2 effective sample sizes iteratively estimated by the Francis method for retained, total, and groundfish discard catch size compositions of golden king crab for **model 20_2** fit to **EAG** data. NA: not available.

Year	Initial Input Retained Vessel- Days Sample Size (no)	Stage-2 Retained Effective Sample Size (no)	Initial Input Total Vessel- Days Sample Size (no)	Stage-2 Total Effective Sample Size (no)	Initial Input Groundfish Trip Sample Size (no)	Stage-2 Groundfish Effective Sample Size (no)
1985/86	57	47				
1986/87	11	9				
1987/88	61	50				
1988/89	352	289				
1989/90	792	651			9	4
1990/91	163	134	22	13	13	6
1991/92	140	115	48	28	NA	NA
1992/93	49	40	41	24	2	1
1993/94	340	279	NA	NA	2	1
1994/95	319	262	34	20	4	2
1995/96	879	723	1,117	659	5	2
1996/97	547	450	509	301	4	2
1997/98	538	442	711	420	8	4
1998/99	541	445	574	339	15	7
1999/00	463	381	607	358	14	7
2000/01	436	358	495	292	16	8
2001/02	488	401	510	301	13	6
2002/03	406	334	438	259	15	7
2003/04	405	333	416	246	17	8
2004/05	280	230	299	177	10	5
2005/06	266	219	232	137	12	6
2006/07	234	192	143	84	14	7
2007/08	199	164	134	79	17	8
2008/09	197	162	113	67	15	7
2009/10	170	140	95	56	16	8
2010/11	183	150	108	64	26	12
2011/12	160	132	107	63	13	6
2012/13	187	154	99	58	18	9
2013/14	193	159	122	72	17	8
2014/15	168	138	99	58	16	8
2015/16	190	156	125	74	10	5
2016/17	223	183	155	92	12	6
2017/18	213	175	133	79	12	6
2018/19	218	179	234	138	9	4
2019/20	208	171	230	136	8	4

Table 8. Parameter estimates and coefficient of variations (CV) with the 2019 MMB (MMB estimated on 15 Feb 2020) for models 20_1, 20_1b, 20_1b Ver 2, and 20_2 for the golden king crab data from the **EAG**, 1985/86–2019/20. Recruitment and fishing mortality deviations and initial size frequency determination parameters were omitted from this list.

Parameter	Model 20_1		Model 20_1b		Model 20_1b Ver 2		Model 20_2		Limits
	Estimate	CV	Estimate	CV	Estimate	CV	Estimate	CV	
log_ω ₁ (growth incr. intercept)	2.538	0.01	2.537	0.01	2.537	0.01	2.537	0.01	1.0, 4.5
ω ₂ (growth incr. slope)	-8.282	0.21	-8.311	0.21	-8.311	0.21	-8.297	0.21	-12.0-5.0
log_a (molt prob. slope)	-2.509	0.02	-2.508	0.02	-2.508	0.02	-2.502	0.02	-4.61-1.39
log_b (molt prob. L50)	4.949	0.001	4.949	0.001	4.949	0.001	4.949	0.001	3.869,5.05
σ (growth variability std)	3.678	0.03	3.677	0.03	3.677	0.03	3.678	0.03	0.1,12.0
log_total sel deltaθ, 1985–04	3.387	0.02	3.383	0.02	3.383	0.02	3.388	0.02	0.,4.4
log_total sel deltaθ, 2005–19	2.951	0.02	2.951	0.02	2.951	0.02	2.938	0.02	0.,4.4
log_ret. sel deltaθ, 1985–19	1.868	0.02	1.868	0.02	1.868	0.02	1.869	0.02	0.,4.4
log_tot sel θ ₅₀ , 1985–04	4.835	0.002	4.834	0.002	4.834	0.002	4.836	0.002	4.0,5.0
log_tot sel θ ₅₀ , 2005–19	4.922	0.002	4.922	0.002	4.922	0.002	4.919	0.002	4.0,5.0
log_ret. sel θ ₅₀ , 1985–19	4.915	0.0003	4.915	0.0003	4.915	0.0003	4.915	0.0003	4.0,5.0
log_β _r (rec.distribution par.)	-1.079	0.17	-1.080	0.17	-1.080	0.17	-1.076	0.17	-12.0, 12.0
logq2 (catchability 1995–04)	-0.538	0.14	-0.541	0.13	-0.540	0.13	-0.541	0.13	-9.0, 2.25
logq3 (catchability 2005–19)	-0.711	0.17	-0.712	0.17	-0.712	0.17	-0.752	0.15	-9.0, 2.25
log_mean_rec (mean rec.)	0.828	0.05	0.828	0.05	0.828	0.05	0.836	0.05	0.01, 5.0
log_mean_Fpot (Pot fishery F)	-0.940	0.07	-0.943	0.07	-0.943	0.07	-0.963	0.07	-15.0, -0.01
log_mean_Fground (GF byc. F)	-9.155	0.09	-9.156	0.09	-9.156	0.09	-9.172	0.09	-15.0, -1.6
σ _e ² (observer CPUE additional var)	0.055	0.36	0.055	0.36	0.055	0.36	0.045	0.37	0.0, 0.15
σ _e ² (fishery CPUE additional var)	0.039	0.43	0.033	0.44	0.033	0.44	0.033	0.44	0.0,1.0
2019 MMB	9,765	0.22	9,762	0.22	9,775	0.22	10,099	0.21	

Table 9. Annual abundance estimates of model recruits (millions of crabs), legal male biomass (t) with coefficient of variations (CV), and mature male biomass (t) with CV for **model 20_1** for golden king crab in the **EAG**. Legal male biomass was estimated on July 1 (start of fishing year) of fishing year y . Mature male biomass for fishing year y was estimated on February 15 of year $y+1$, after the year y fishery total catch removal. Recruits estimates for 1961 to 2020 are restricted to 1985–2020. Equilibrium MMB_{eq} and $MMB_{35\%}$ are also listed.

Year	Recruits to the Model (\geq 101 mm CL)	Mature Male Biomass (\geq 111 mm CL)	CV	Legal Size Male Biomass (\geq 136 mm CL)	CV
		$MMB_{eq}=22,632$ $MMB_{35\%}=6,601$			
1985	1.68	9,486	0.04	9,723	0.06
1986	1.01	7,259	0.04	8,234	0.04
1987	4.25	6,645	0.05	6,430	0.04
1988	3.60	6,630	0.05	5,363	0.05
1989	2.02	5,771	0.06	4,793	0.07
1990	2.96	5,882	0.05	4,306	0.07
1991	3.49	5,966	0.04	4,586	0.06
1992	2.25	5,887	0.04	4,425	0.05
1993	2.15	6,044	0.03	4,452	0.05
1994	2.43	5,581	0.03	4,875	0.04
1995	2.30	5,001	0.04	4,435	0.04
1996	2.24	5,111	0.04	3,835	0.04
1997	3.00	5,363	0.05	3,969	0.04
1998	2.76	5,918	0.05	4,076	0.05
1999	2.86	6,571	0.05	4,501	0.05
2000	2.65	7,143	0.06	5,147	0.06
2001	2.00	7,456	0.06	5,746	0.06
2002	2.45	7,689	0.07	6,241	0.06
2003	2.12	7,882	0.07	6,540	0.07
2004	1.87	7,889	0.07	6,718	0.07
2005	2.76	7,902	0.07	6,830	0.07
2006	2.14	8,072	0.07	6,709	0.08
2007	2.06	8,055	0.07	6,798	0.08
2008	2.97	8,131	0.07	6,906	0.08
2009	1.93	8,314	0.06	6,837	0.08
2010	1.79	8,109	0.06	7,026	0.07
2011	2.09	7,817	0.06	7,063	0.06
2012	1.80	7,489	0.06	6,794	0.06
2013	1.55	6,963	0.06	6,465	0.06
2014	2.65	6,610	0.07	6,048	0.06
2015	3.24	6,783	0.08	5,534	0.07
2016	3.71	7,436	0.11	5,321	0.08
2017	4.97	8,770	0.14	5,670	0.11
2018	2.61	9,901	0.19	6,586	0.14
2019	2.25	9,765	0.22	7,893	0.18
2020	2.29				

Table 10. Annual abundance estimates of model recruits (millions of crabs), legal male biomass (t) with coefficient of variations (CV), and mature male biomass (t) with CV for **model 20_1b** for golden king crab in the **EAG**. Legal male biomass was estimated on July 1 (start of fishing year) of fishing year *y*. Mature male biomass for fishing year *y* was estimated on February 15 of year *y*+1, after the year *y* fishery total catch removal. Recruits estimates for 1961 to 2020 are restricted to 1985–2020. Equilibrium MMB_{eq} and $MMB_{35\%}$ are also listed.

Year	Recruits to the Model (≥ 101 mm CL)	Mature Male Biomass (≥ 111 mm CL)	CV	Legal Size Male Biomass (≥ 136 mm CL)	CV
		$MMB_{eq}=22,241$ $MMB_{35\%}=6,774$			
1985	1.71	9,454	0.04	9,671	0.06
1986	1.02	7,248	0.04	8,189	0.04
1987	4.29	6,655	0.05	6,411	0.04
1988	3.63	6,672	0.05	5,363	0.05
1989	2.02	5,830	0.06	4,820	0.07
1990	2.91	5,926	0.05	4,359	0.07
1991	3.49	5,986	0.04	4,645	0.06
1992	2.25	5,903	0.04	4,459	0.05
1993	2.16	6,057	0.03	4,471	0.05
1994	2.43	5,592	0.04	4,889	0.04
1995	2.31	5,007	0.04	4,448	0.04
1996	2.24	5,117	0.04	3,844	0.04
1997	3.01	5,368	0.05	3,976	0.04
1998	2.76	5,923	0.05	4,082	0.05
1999	2.86	6,576	0.05	4,508	0.05
2000	2.65	7,149	0.06	5,154	0.06
2001	2.00	7,461	0.06	5,753	0.06
2002	2.45	7,693	0.07	6,248	0.06
2003	2.12	7,885	0.07	6,546	0.07
2004	1.87	7,891	0.07	6,723	0.07
2005	2.77	7,904	0.07	6,833	0.07
2006	2.14	8,074	0.07	6,712	0.08
2007	2.06	8,058	0.07	6,802	0.08
2008	2.97	8,134	0.07	6,911	0.08
2009	1.93	8,318	0.06	6,842	0.08
2010	1.79	8,112	0.06	7,031	0.07
2011	2.09	7,820	0.06	7,067	0.06
2012	1.80	7,493	0.06	6,798	0.06
2013	1.55	6,967	0.06	6,470	0.06
2014	2.65	6,613	0.07	6,053	0.06
2015	3.24	6,786	0.08	5,538	0.07
2016	3.71	7,437	0.11	5,326	0.08
2017	4.96	8,770	0.14	5,674	0.11
2018	2.61	9,899	0.19	6,589	0.14
2019	2.25	9,762	0.22	7,895	0.18
2020	2.29				

Table 11. Annual abundance estimates of model recruits (millions of crabs), legal male biomass (t) with coefficient of variations (CV), and mature male biomass (t) with CV for **model 20_2** for golden king crab in the **EAG**. Legal male biomass was estimated on July 1 (start of fishing year) of fishing year y. Mature male biomass for fishing year y was estimated on February 15 of year y+1, after the year y fishery total catch removal. Recruits estimates for 1961 to 2020 are restricted to 1985–2020. Equilibrium MMB_{eq} and $MMB_{35\%}$ are also listed.

Year	Recruits to the Model (\geq 101 mm CL)	Mature Male Biomass (\geq 111 mm CL)	CV	Legal Size Male Biomass (\geq 136 mm CL)	CV
		$MMB_{eq}=23,445$ $MMB_{35\%}=6,794$			
1985	1.71	9,473	0.04	9,704	0.06
1986	1.02	7,262	0.04	8,208	0.04
1987	4.30	6,669	0.05	6,420	0.04
1988	3.62	6,685	0.05	5,370	0.05
1989	2.02	5,840	0.06	4,830	0.07
1990	2.90	5,936	0.05	4,365	0.07
1991	3.49	5,991	0.04	4,651	0.06
1992	2.22	5,899	0.04	4,463	0.05
1993	2.15	6,038	0.03	4,470	0.05
1994	2.44	5,566	0.04	4,875	0.04
1995	2.32	4,990	0.04	4,421	0.04
1996	2.26	5,114	0.04	3,819	0.04
1997	3.05	5,391	0.05	3,962	0.05
1998	2.83	5,985	0.05	4,087	0.05
1999	2.93	6,688	0.05	4,541	0.05
2000	2.72	7,314	0.06	5,229	0.06
2001	2.06	7,676	0.06	5,879	0.06
2002	2.52	7,951	0.06	6,423	0.06
2003	2.13	8,166	0.07	6,764	0.07
2004	1.87	8,160	0.07	6,977	0.07
2005	2.75	8,143	0.07	7,092	0.07
2006	2.16	8,281	0.07	6,948	0.08
2007	2.08	8,249	0.07	7,001	0.07
2008	2.98	8,313	0.07	7,085	0.07
2009	1.95	8,482	0.06	7,004	0.07
2010	1.81	8,267	0.06	7,181	0.07
2011	2.13	7,976	0.06	7,206	0.06
2012	1.82	7,658	0.06	6,932	0.06
2013	1.56	7,133	0.06	6,611	0.06
2014	2.68	6,777	0.07	6,201	0.06
2015	3.30	6,959	0.09	5,685	0.07
2016	3.82	7,648	0.11	5,473	0.09
2017	5.05	9,042	0.14	5,839	0.11
2018	2.66	10,217	0.18	6,804	0.13
2019	2.28	10,099	0.21	8,165	0.18
2020	2.31				

Table 12. Negative log-likelihood values of the fits for models 20_1 (base, last year's accepted model with additional 2019/20 data), 20_1b, 20_1b Ver 2 (21_b but mean recruitment estimation time period modified to 1987–2012), and 20_2 (observer CPUE estimated with Year an Area interaction factor) for golden king crab in the **EAG**. Likelihood components with zero entry in the entire rows are omitted. RetdcatchB= retained catch biomass.

Likelihood Component	Model 20_1	Model 20_1b	Model 20_1b Ver 2	Model 20_2
Number of free parameters	149	149	149	149
Retlencomp	-1286.4300	-1286.6600	-1286.6600	-1286.7800
Totallencomp	-1428.6400	-1427.3300	-1427.3200	-1430.6100
Observer cpue	-0.5240	-0.5376	-0.5493	-2.4792
RetdcatchB	7.7446	7.6845	7.6847	7.9245
TotalcatchB	23.3301	23.3858	23.3859	23.4631
GdiscdcatchB	0.0003	0.0003	0.0003	0.0003
Rec_dev	7.3036	7.3053	7.3061	7.3886
Pot F_dev	0.0126	0.0125	0.0125	0.0128
Gbyc_F_dev	0.0296	0.0296	0.0296	0.0296
Tag	2692.5200	2692.5100	2692.5100	2692.3100
Fishery cpue	-2.3673	-3.5143	-3.5137	-3.4738
RetcatchN	0.0054	0.0055	0.0055	0.0055
Total	12.9831	12.8964	12.8904	7.7967

Table 13. Time series of negative binomial GLM estimated CPUE indices and coefficient of variations (CV) for the fish ticket based retained catch-per-pot lift for the **WAG** golden king crab stock. The GLM was fitted to the 1985/86 to 1998/99 time series of data. GLM predictor variables were selected by R square criteria.

Year	CPUE Index	CV
1985/86	2.07	0.05
1986/87	1.59	0.04
1987/88	1.22	0.04
1988/89	1.41	0.03
1989/90	1.15	0.03
1990/91	0.87	0.03
1991/92	0.76	0.04
1992/93	0.61	0.04
1993/94	0.76	0.05
1994/95	0.83	0.04
1995/96	0.90	0.04
1996/97	0.84	0.03
1997/98	0.76	0.03
1998/99	1.06	0.03

Table 14. The initial input number of vessel-days/trips and Stage-2 effective sample sizes iteratively estimated by the Francis method for retained, total, and groundfish discard catch size compositions of golden king crab for **model 20_1** model fit to **WAG** data. NA: not available.

Year	Initial Input Retained Vessel- Days Sample Size (no)	Stage-2 Retained Effective Sample Size (no)	Initial Input Total Vessel- Days Sample Size (no)	Stage-2 Total Effective Sample Size (no)	Initial Input Groundfish Trip Sample Size (no)	Stage-2 Groundfish Effective Sample Size (no)
1985/86	45	22				
1986/87	23	11				
1987/88	8	4				
1988/89	286	139				
1989/90	513	250			7	4
1990/91	205	100	190	99	6	4
1991/92	102	50	104	54	1	1
1992/93	76	37	94	49	3	2
1993/94	378	184	62	32	NA	NA
1994/95	367	179	119	62	2	1
1995/96	705	344	907	474	5	3
1996/97	817	398	1061	554	8	5
1997/98	984	480	1116	583	6	4
1998/99	613	299	638	333	14	9
1999/00	915	446	1155	603	18	11
2000/01	1029	502	1205	629	11	7
2001/02	898	438	975	509	11	7
2002/03	628	306	675	352	16	10
2003/04	688	336	700	365	8	5
2004/05	449	219	488	255	9	6
2005/06	337	164	220	115	6	4
2006/07	337	164	321	168	14	9
2007/08	276	135	257	134	17	11
2008/09	318	155	258	135	19	12
2009/10	362	177	292	152	24	15
2010/11	328	160	222	116	13	8
2011/12	295	144	252	132	14	9
2012/13	288	140	241	126	18	11
2013/14	327	159	236	123	17	11
2014/15	305	149	219	114	18	11
2015/16	287	140	243	127	10	6
2016/17	392	191	253	132	12	8
2017/18	299	146	222	116	10	6
2018/19	328	160	318	166	5	3
2019/20	256	125	320	167	6	4

Table 15. The initial input number of vessel-days/trips and Stage-2 effective sample sizes iteratively estimated by the Francis method for retained, total, and groundfish discard catch size compositions of golden king crab for **model 20_1b** model fit to **WAG** data. NA: not available.

Year	Initial Input Retained Vessel- Days Sample Size (no)	Stage-2 Retained Effective Sample Size (no)	Initial Input Total Vessel- Days Sample Size (no)	Stage-2 Total Effective Sample Size (no)	Initial Input Groundfish Trip Sample Size (no)	Stage-2 Groundfish Effective Sample Size (no)
1985/86	45	22				
1986/87	23	11				
1987/88	8	4				
1988/89	286	142				
1989/90	513	255			7	4
1990/91	205	102	190	98	6	4
1991/92	102	51	104	54	1	1
1992/93	76	38	94	48	3	2
1993/94	378	188	62	32	NA	NA
1994/95	367	182	119	61	2	1
1995/96	705	350	907	467	5	3
1996/97	817	405	1061	546	8	5
1997/98	984	488	1116	574	6	4
1998/99	613	304	638	328	14	9
1999/00	915	454	1155	595	18	11
2000/01	1029	511	1205	620	11	7
2001/02	898	446	975	502	11	7
2002/03	628	312	675	347	16	10
2003/04	688	341	700	360	8	5
2004/05	449	223	488	251	9	6
2005/06	337	167	220	113	6	4
2006/07	337	167	321	165	14	9
2007/08	276	137	257	132	17	11
2008/09	318	158	258	133	19	12
2009/10	362	180	292	150	24	15
2010/11	328	163	222	114	13	8
2011/12	295	146	252	130	14	9
2012/13	288	143	241	124	18	11
2013/14	327	162	236	121	17	11
2014/15	305	151	219	113	18	11
2015/16	287	142	243	125	10	6
2016/17	392	195	253	130	12	8
2017/18	299	148	222	114	10	6
2018/19	328	163	318	164	5	3
2019/20	256	127	320	165	6	4

Table 16. The initial input number of vessel-days/trips and Stage-2 effective sample sizes iteratively estimated by the Francis method for retained, total, and groundfish discard catch size compositions of golden king crab for **model 20_2** model fit to **WAG** data. NA: not available.

Year	Initial Input Retained Vessel- Days Sample Size (no)	Stage-2 Retained Effective Sample Size (no)	Initial Input Total Vessel- Days Sample Size (no)	Stage-2 Total Effective Sample Size (no)	Initial Input Groundfish Trip Sample Size (no)	Stage-2 Groundfish Effective Sample Size (no)
1985/86	45	22				
1986/87	23	11				
1987/88	8	4				
1988/89	286	142				
1989/90	513	254			7	4
1990/91	205	102	190	99	6	4
1991/92	102	51	104	54	1	1
1992/93	76	38	94	49	3	2
1993/94	378	187	62	32	NA	NA
1994/95	367	182	119	62	2	1
1995/96	705	349	907	475	5	3
1996/97	817	405	1061	555	8	5
1997/98	984	488	1116	584	6	4
1998/99	613	304	638	334	14	9
1999/00	915	453	1155	605	18	11
2000/01	1029	510	1205	631	11	7
2001/02	898	445	975	510	11	7
2002/03	628	311	675	353	16	10
2003/04	688	341	700	366	8	5
2004/05	449	223	488	255	9	6
2005/06	337	167	220	115	6	4
2006/07	337	167	321	168	14	9
2007/08	276	137	257	135	17	11
2008/09	318	158	258	135	19	12
2009/10	362	179	292	153	24	15
2010/11	328	163	222	116	13	8
2011/12	295	146	252	132	14	9
2012/13	288	143	241	126	18	11
2013/14	327	162	236	124	17	11
2014/15	305	151	219	115	18	11
2015/16	287	142	243	127	10	6
2016/17	392	194	253	132	12	8
2017/18	299	148	222	116	10	6
2018/19	328	163	318	166	5	3
2019/20	256	127	320	168	6	4

Table 17. Parameter estimates and coefficient of variations (CV) with the 2019 MMB (MMB estimated on 15 Feb 2020) for models 20_1, 20_1b, 20_1b Ver 2, and 20_2 for the golden king crab data from the **WAG**, 1985/86–2019/20. Recruitment and fishing mortality deviations and initial size frequency determination parameters were omitted from this list.

Parameter	Model 20_1		Model 20_1b		Model 20_1b Ver 2		Model 20_2		Limits
	Estimate	CV	Estimate	CV	Estimate	CV	Estimate	CV	
log_ω ₁ (growth incr. intercept)	2.537	0.01	2.537	0.01	2.537	0.01	2.537	0.01	1.0, 4.5
ω ₂ (growth incr. slope)	-7.699	0.22	-7.733	0.22	-7.733	0.22	-7.717	0.22	-12.0-5.0
log_a (molt prob. slope)	-2.625	0.03	-2.626	0.03	-2.626	0.03	-2.626	0.03	-4.61-1.39
log_b (molt prob. L50)	4.947	0.001	4.947	0.001	4.947	0.001	4.947	0.001	3.869,5.05
σ (growth variability std)	3.690	0.03	3.689	0.03	3.689	0.03	3.690	0.03	0.1,12.0
log_total sel deltaθ, 1985–04	3.411	0.01	3.408	0.01	3.408	0.01	3.410	0.01	0.,4.4
log_total sel deltaθ, 2005–19	2.838	0.02	2.840	0.02	2.840	0.02	2.840	0.02	0.,4.4
log_ret. sel deltaθ, 1985–19	1.793	0.02	1.793	0.02	1.793	0.02	1.793	0.02	0.,4.4
log_tot sel θ ₅₀ , 1985–04	4.868	0.002	4.868	0.002	4.868	0.002	4.868	0.002	4.0,5.0
log_tot sel θ ₅₀ , 2005–19	4.900	0.001	4.900	0.001	4.900	0.001	4.900	0.001	4.0,5.0
log_ret. sel θ ₅₀ , 1985–19	4.916	0.0002	4.916	0.0002	4.916	0.0002	4.916	0.0002	4.0,5.0
log_β _r (rec.distribution par.)	-1.039	0.15	-1.040	0.15	-1.040	0.15	-1.037	0.15	-12.0, 12.0
logq2 (catchability 1995–04)	-0.046	1.41	-0.036	1.93	-0.036	1.93	-0.037	1.85	-9.0, 2.25
logq3 (catchability 2005–19)	-0.371	0.22	-0.372	0.22	-0.372	0.22	-0.371	0.23	-9.0, 2.25
log_mean_rec (mean rec.)	0.719	0.06	0.721	0.05	0.721	0.05	0.722	0.05	0.01, 5.0
log_mean_Fpot (Pot fishery F)	-0.691	0.09	-0.695	0.09	-0.695	0.09	-0.694	0.09	-15.0, -0.01
log_mean_Fground (GF byc. F)	-8.292	0.10	-8.294	0.10	-8.294	0.10	-8.296	0.10	-15.0, -1.6
σ _e ² (observer CPUE additional var)	0.020	0.34	0.019	0.35	0.019	0.35	0.019	0.40	0.0, 0.15
σ _e ² (fishery CPUE additional var)	0.014	0.65	0.024	0.61	0.024	0.61	0.024	0.60	0.0,1.0
2019 MMB	6,528	0.16	6,542	0.16	6,548	0.16	6,734	0.16	

Table 18. Annual abundance estimates of model recruits (millions of crabs), legal male biomass (t) with coefficient of variations (CV), and mature male biomass (t) with CV for **model 20_1** for golden king crab in the **WAG**. Legal male biomass was estimated on July 1 (start of fishing year) of fishing year *y*. Mature male biomass for fishing year *y* was estimated on February 15 of year *y*+1, after the year *y* fishery total catch removal. Recruits estimates for 1961 to 2020 are restricted to 1985–2020. Equilibrium MMB_{eq} and $MMB_{35\%}$ are also listed.

Year	Recruits to the Model (\geq 101 mm CL)	Mature Male Biomass (\geq 111 mm CL)	CV	Legal Size Male Biomass (\geq 136 mm CL)	CV
		$MMB_{eq}=17,953$ $MMB_{35\%}=5,204$			
1985	4.00	10,485	0.05	8,930	0.09
1986	3.57	8,072	0.05	8,414	0.07
1987	2.66	7,459	0.04	5,973	0.06
1988	1.76	6,376	0.04	5,631	0.04
1989	2.39	4,316	0.04	5,002	0.04
1990	1.92	3,956	0.05	3,130	0.05
1991	1.67	3,722	0.05	2,792	0.05
1992	2.10	3,895	0.04	2,692	0.05
1993	1.56	4,497	0.03	2,850	0.05
1994	1.97	3,808	0.03	3,469	0.03
1995	1.89	3,810	0.03	2,813	0.03
1996	1.71	3,821	0.04	2,762	0.03
1997	1.86	3,891	0.04	2,808	0.04
1998	1.90	4,214	0.03	2,888	0.04
1999	2.24	4,245	0.04	3,172	0.03
2000	2.50	4,394	0.04	3,114	0.04
2001	2.52	4,818	0.05	3,121	0.04
2002	2.44	5,345	0.05	3,446	0.05
2003	1.71	5,640	0.05	3,955	0.05
2004	2.23	5,715	0.06	4,421	0.05
2005	2.35	5,989	0.06	4,578	0.06
2006	2.47	6,531	0.05	4,720	0.06
2007	1.71	6,732	0.05	5,165	0.06
2008	1.51	6,563	0.05	5,483	0.05
2009	1.91	6,197	0.05	5,552	0.05
2010	1.59	5,916	0.05	5,205	0.05
2011	1.15	5,421	0.04	4,906	0.05
2012	1.84	4,823	0.05	4,564	0.05
2013	2.21	4,570	0.05	3,951	0.05
2014	1.69	4,639	0.06	3,469	0.06
2015	2.01	4,730	0.06	3,511	0.06
2016	2.14	5,101	0.07	3,635	0.07
2017	1.80	5,462	0.09	3,927	0.07
2018	3.28	5,897	0.12	4,313	0.09
2019	2.02	6,528	0.16	4,500	0.11
2020	2.05				

Table 19. Annual abundance estimates of model recruits (millions of crabs), legal male biomass (t) with coefficient of variations (CV), and mature male biomass (t) with CV for **model 20_1b** for golden king crab in the **WAG**. Legal male biomass was estimated on July 1 (start of fishing year) of fishing year *y*. Mature male biomass for fishing year *y* was estimated on February 15 of year *y*+1, after the year *y* fishery total catch removal. Recruits estimates for 1961 to 2020 are restricted to 1985–2020. Equilibrium MMB_{eq} and $MMB_{35\%}$ are also listed.

Year	Recruits to the Model (≥ 101 mm CL)	Mature Male Biomass (≥ 111 mm CL)	CV	Legal Size Male Biomass (≥ 136 mm CL)	CV
		$MMB_{eq}=18,343$ $MMB_{35\%}=5,319$			
1985	4.05	10,471	0.05	9,006	0.10
1986	3.47	8,040	0.05	8,427	0.08
1987	2.68	7,387	0.04	5,960	0.06
1988	1.86	6,326	0.04	5,580	0.05
1989	2.52	4,339	0.04	4,936	0.04
1990	1.92	4,037	0.05	3,104	0.06
1991	1.64	3,808	0.05	2,836	0.05
1992	2.02	3,950	0.04	2,775	0.05
1993	1.59	4,521	0.03	2,928	0.05
1994	1.96	3,824	0.03	3,509	0.03
1995	1.89	3,817	0.04	2,830	0.03
1996	1.71	3,827	0.04	2,771	0.04
1997	1.86	3,892	0.04	2,814	0.04
1998	1.89	4,211	0.04	2,891	0.04
1999	2.24	4,238	0.04	3,172	0.04
2000	2.49	4,384	0.04	3,111	0.04
2001	2.52	4,805	0.05	3,114	0.04
2002	2.45	5,333	0.05	3,436	0.05
2003	1.71	5,631	0.05	3,943	0.05
2004	2.23	5,712	0.06	4,411	0.05
2005	2.35	5,988	0.06	4,572	0.06
2006	2.46	6,529	0.05	4,719	0.06
2007	1.71	6,731	0.05	5,165	0.06
2008	1.51	6,562	0.05	5,482	0.06
2009	1.91	6,197	0.05	5,551	0.05
2010	1.59	5,917	0.05	5,205	0.05
2011	1.15	5,423	0.04	4,907	0.05
2012	1.84	4,824	0.05	4,566	0.05
2013	2.21	4,574	0.05	3,952	0.05
2014	1.69	4,648	0.06	3,472	0.06
2015	2.01	4,742	0.06	3,517	0.06
2016	2.14	5,113	0.07	3,646	0.07
2017	1.81	5,475	0.09	3,940	0.07
2018	3.28	5,910	0.12	4,326	0.09
2019	2.02	6,542	0.16	4,513	0.11
2020	2.06				

Table 20. Annual abundance estimates of model recruits (millions of crabs), legal male biomass (t) with coefficient of variations (CV), and mature male biomass (t) with CV for **model 20_2** for golden king crab in the **WAG**. Legal male biomass was estimated on July 1 (start of fishing year) of fishing year y. Mature male biomass for fishing year y was estimated on February 15 of year y+1, after the year y fishery total catch removal. Recruits estimates for 1961 to 2020 are restricted to 1985–2020. Equilibrium MMB_{eq} and $MMB_{35\%}$ are also listed.

Year	Recruits to the Model (≥ 101 mm CL)	Mature Male Biomass (≥ 111 mm CL)	CV	Legal Size Male Biomass (≥ 136 mm CL)	CV
		$MMB_{eq}=18,413$ $MMB_{35\%}=5,343$			
1985	4.04	10,474	0.05	9,012	0.10
1986	3.47	8,041	0.05	8,432	0.08
1987	2.68	7,389	0.04	5,961	0.06
1988	1.86	6,328	0.04	5,582	0.05
1989	2.52	4,338	0.04	4,938	0.04
1990	1.91	4,035	0.05	3,105	0.05
1991	1.64	3,805	0.05	2,835	0.05
1992	2.01	3,943	0.04	2,772	0.05
1993	1.58	4,507	0.03	2,924	0.05
1994	1.97	3,809	0.03	3,500	0.03
1995	1.89	3,807	0.04	2,816	0.03
1996	1.70	3,817	0.04	2,760	0.04
1997	1.87	3,884	0.04	2,806	0.04
1998	1.90	4,210	0.03	2,882	0.04
1999	2.24	4,240	0.04	3,168	0.03
2000	2.49	4,384	0.04	3,111	0.04
2001	2.50	4,796	0.05	3,116	0.04
2002	2.42	5,307	0.05	3,433	0.05
2003	1.70	5,589	0.05	3,929	0.05
2004	2.26	5,667	0.06	4,379	0.05
2005	2.42	5,973	0.06	4,528	0.06
2006	2.51	6,566	0.05	4,682	0.06
2007	1.69	6,794	0.05	5,168	0.06
2008	1.46	6,612	0.05	5,531	0.05
2009	1.89	6,217	0.05	5,612	0.05
2010	1.57	5,912	0.04	5,243	0.05
2011	1.14	5,397	0.04	4,917	0.05
2012	1.86	4,792	0.05	4,551	0.05
2013	2.22	4,549	0.05	3,921	0.05
2014	1.72	4,638	0.06	3,442	0.06
2015	2.07	4,765	0.07	3,497	0.06
2016	2.18	5,177	0.07	3,647	0.07
2017	1.83	5,571	0.09	3,976	0.07
2018	3.39	6,050	0.12	4,400	0.09
2019	2.03	6,734	0.16	4,616	0.11
2020	2.06				

Table 21. Negative log-likelihood values of the fits for models 20_1 (base, last year's accepted model with additional 2019/20 data), 20_1b, 20_1b Ver 2 (21_b but mean recruitment estimation time period modified to 1987–2012), and 20_2 (observer CPUE estimated with Year an Area interaction factor) for golden king crab in the **WAG**. Likelihood components with zero entry in the entire rows are omitted. RetdcatchB= retained catch biomass.

Likelihood Component	Model 20_1	Model 20_1b	Model 20_1b Ver 2	Model 20_2
Number of free parameters	149	149	149	149
Retlencomp	-1240.2800	-1244.3900	-1244.3900	-1243.7800
Totallencomp	-1564.8500	-1561.8900	-1561.8800	-1565.1200
Observer cpue	-13.0279	-13.7535	-13.7556	-11.6569
RetdcatchB	5.1206	5.2357	5.2357	5.3112
TotalcatchB	45.6044	45.7246	45.7252	45.7664
GdiscdcatchB	0.0014	0.0015	0.0015	0.0014
Rec_dev	5.0374	4.9326	4.9342	5.1016
Pot F_dev	0.0264	0.0265	0.0265	0.0266
Gbyc_F_dev	0.0384	0.0385	0.0385	0.0384
Tag	2694.2000	2694.1900	2694.1900	2694.2400
Fishery cpue	-9.3432	-5.6807	-5.6811	-5.7031
RetcatchN	0.0019	0.0019	0.0018	0.0019
Total	-77.4698	-75.5594	-75.5643	-75.7768




Figure 1. Aleutian Islands, Area O, red and golden king crab management area (from Leon *et al.* 2017).




Figure 2. Adak (Area R) and Dutch Harbor (Area O) king crab registration area and districts, 1984/85–1995/96 seasons (Leon *et al.* 2017).




Figure 3. Percent of total 1981/82–1995/96 golden king crab retained catch weight (harvest) from one-degree longitude intervals in the Aleutian Islands, with dotted line denoting the border at 171° W longitude used during the 1984/85–1995/96 seasons to divide fishery management between the Dutch Harbor Area (east of 171° W longitude) and the Adak Area (west of 171° W longitude) and solid line denoting the border at 174° W longitude used since the 1996/97 season to manage crab east and west of 174° W longitude (adapted from Figure 4-2 in Morrison *et al.* 1998).




Figure 4. Retained catch (t) of golden king crab within one-degree longitude intervals in the Aleutian Islands during the 2000/01 through 2019/20 commercial fishery seasons; solid line denotes the border at 174° W longitude that has been used since the 1996/97 season to manage Aleutian Island golden king crab as separate stocks east and west of 174° W longitude and dashed line denotes the border at 171° W longitude used during the 1984/85–1995/96 seasons to divide fishery management between the Dutch Harbor Area (east of 171° W longitude) and the Adak Area (west of 171° W longitude).




Figure 5. Average golden king crab CPUE (kg/nm²) for tows, number of tows, and average depth of tows from one-degree longitude intervals during the 2002, 2004, 2006, 2010, and 2012 NMFS Aleutian Islands bottom trawl surveys; preliminary summary of data obtained on 1 April 2013 from http://www.afsc.noaa.gov/RACE/groundfish/survey_data/default.htm.




Figure 6. Historical commercial harvest (from fish tickets; metric tons) and catch-per-unit effort (CPUE, number of crabs per pot lift) of golden king crab in the **EAG**, 1985/86–2019/20 fisheries (note: 1985 refers to the 1985/86 fishing year).




Figure 7. Historical commercial harvest (from fish tickets; metric tons) and catch-per-unit effort (CPUE, number of crabs per pot lift) of golden king crab in the **WAG**, 1985/86–2019/20 fisheries (note: 1985 refers to the 1985/86 fishing year).




Figure 8. Catch distribution by statistical area.in 2019/20.




Figure 9. Predicted (line) vs. observed (bar) retained catch relative length frequency distributions under models 20_1 (orange line), 20_1b (black line), 20_1c (dark red line), 20_2 (green line), and 20_2b (blue line) for golden king crab in the EAG, 1985/86 to 2019/20. This color scheme is used in all other figures.




Figure 10. Predicted (line) vs. observed (bar) total catch relative length frequency distributions under models 20_1, 20_1b, 20_1c, 20_2, and 20_2b for golden king crab in the EAG, 1990/91 to 2019/20.




Figure 11. Predicted (line) vs. observed (bar) groundfish discarded bycatch relative length frequency distributions under models 20_1, 20_1b, 20_1c, 20_2, and 20_2b for golden king crab in the EAG, 19989/90 to 2019/20.




Figure 12. Estimated total (black solid line) and retained selectivity (red dotted line) for pre- and post-rationalization periods under models 20_1, 20_1b, 20_1c, and 20_2 fits to golden king crab data in the **EAG**.




Figure 13. Observed (open circles) vs. predicted (solid line) tag recaptures by size bin for years 1 to 6 post tagging under model 20_1 for **EAG** golden king crab.




Figure 14. Estimated number of male recruits (crab size ≥ 101 mm CL) to the assessment model under models 20_1, 20_1b, 20_1b Ver2, 20_1c, 20_2, and 20_2b fits to **EAG** golden king crab data, 1961–2020. The numbers of recruits are standardized using $(R - \text{mean } R) / \text{mean } R$ for comparing different scenarios' results.




Figure 15. Recruit size distribution to the assessment model under models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_2, and 20_2b fits to **EAG** golden king crab data.




Figure 16. Estimated molt probability vs. carapace length of golden king crab for models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_2, and 20_2b fits to **EAG** golden king crab data.




Figure 17. Observed (open circle) vs. predicted (solid line) retained catch (top left), total catch (top right in), and groundfish bycatch (bottom left) of golden king crab for models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_2, and 20_2b fits in EAG, 1981/82–2019/20.




Figure 18. Observed (open circle) vs. predicted (solid line) retained catch of golden king crab for models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_2, and 20_2b for golden king crab fits in the **EAG**, 1981/82–1984/85. Note: Input retained catches to the model during pre-1985 fishery period were in number of crabs.




Figure 19. Bubble plot of standardized residuals of retained catch length composition for model 20_1b fit for **EAG** golden king crab, 1985/86–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 20. Bubble plot of standardized residuals of total catch length composition for model 20_1b fit for **EAG** golden king crab, 1990/91–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 21. Bubble plot of standardized residuals of retained catch length composition for model 20_2 fit for **EAG** golden king crab, 1985/86–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 22. Bubble plot of standardized residuals of total catch length composition for model 20_2 fit for **EAG** golden king crab, 1990/91–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 23. Retrospective fits of MMB by the model following removal of terminal year data under models 20_1, 20_1b, 20_1bVer2, and 20_2 for golden king crab in the EAG, 1960/61–2019/20.




Figure 24. Comparison of input CPUE indices (orange open circles with +/- 2 SE for model 20_1 and green open circles with +/- 2 SE for model 20_2) with predicted CPUE indices (colored solid lines) under models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_2, and 20_2b for **EAG** golden king crab data, 1985/86–2019/20. Model estimated additional standard error was added to each input standard error.




Figure 25. Trends in pot fishery full selection total fishing mortality of golden king crab for models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_1d, 20_2, and 20_2b fits in the **EAG** (left) and models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** (right) data, 1981/82–2019/20.




Figure 26. Trends in golden king crab mature male biomass for models 20_1, 20_1b, 20_1bVer2, 20_1c, 20_1d, 20_2, and 20_2b fits to **EAG** (left) and models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** (right) data, 1960/61–2019/20. Model 20_1bVer2 estimate has two standard error confidence limits.




Figure 27. Predicted (line) vs. observed (bar) retained catch relative length frequency distributions under models 20_1, 20_1b, and 20_2 fits to golden king crab data in the **WAG**, 1985/86 to 2019/20.




Figure 28. Predicted (line) vs. observed (bar) total catch relative length frequency distributions under models 20_1, 20_1b, and 20_2 fits to golden king crab data in the **WAG**, 1990/91 to 2019/20.




Figure 29. Predicted (line) vs. observed (bar) groundfish discarded bycatch relative length frequency distributions under models 20_1, 20_1b, and 20_2 fits to golden king crab data in the **WAG**, 1989/90 to 2019/20.




Figure 30. Estimated total (black solid line) and retained selectivity (red dotted line) for pre- and post- rationalization periods under models 20_1, 20_1b, and 20_2 fits to golden king crab data in the **WAG**.




Figure 31. Observed (open circles) vs. predicted (solid line) tag recaptures by size bin for years 1 to 6 post tagging under model 20_1 fit to **WAG** golden king crab data.




Figure 32. Estimated number of male recruits (crab size ≥ 101 mm CL) to the assessment model under models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** golden king crab data, 1961–2020. The numbers of recruits are standardized using $(R - \text{mean } R) / \text{mean } R$ for comparing different scenarios' results.




Figure 33. Recruit size distribution to the assessment model under models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** golden king crab data.




Figure 34. Estimated molt probability vs. carapace length of golden king crab for models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** golden king crab data.




Figure 35. Observed (open circle) vs. predicted (solid line) retained catch (top left), total catch (top right in), and groundfish bycatch (bottom left) of golden king crab for models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** data, 1981/82–2019/20.




Figure 36. Observed (open circle) vs. predicted (solid line) retained catch of golden king crab for models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** data, 1981/82–1984/85. Note: Input retained catches to the model during pre-1985 fishery period was in number of crabs.




Figure 37. Bubble plot of standardized residuals of retained catch length composition for model 20_1b fit to **WAG** golden king crab data, 1985/86–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 38. Bubble plot of standardized residuals of total catch length composition for model 20_1b fit to **WAG** golden king crab data, 1990/91–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 39. Bubble plot of standardized residuals of retained catch length composition for model 20_2 fit to **WAG** golden king crab data, 1985/86–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 40. Bubble plot of standardized residuals of total catch length composition for model 20_2 fit to **WAG** golden king crab data, 1990/91–2019/20. Green circles are the positive and pink circles are the negative standardized residuals. The area of the circle is the relative magnitude of the residual.




Figure 41. Retrospective fits of MMB by the model following removal of terminal year data under models 20_1, 20_1b, 20_1bVer2, and 20_2 fits for golden king crab in the **WAG**, 1960/61–2019/20.




Figure 42. Comparison of input CPUE indices (orange open circles with +/- 2 SE for model 20_1 and green open circles with +/- 2 SE for model 20_2) with model predicted CPUE indices (colored solid lines) under models 20_1, 20_1b, 20_1bVer2, and 20_2 fits to **WAG** golden king crab data, 1985/86–2019/20. Model estimated additional standard error was added to each input standard error.




Figure 43. Relationships between full fishing mortalities for the directed pot fishery and mature male biomass during 1981/82–2019/20 under models 20_1bVer2 and 20_2 fits to **EAG** and **WAG** data. F in 2019/20 (red) and 1981/82 (black) are shown in the plots.

Appendix A: Integrated model

Aleutian Islands Golden King Crab (*Lithodes aequispinus*) Stock Assessment Model Development- east of 174° W (EAG) and west of 174° W (WAG) Aleutian Island stocks

Basic population dynamics

The annual [male] abundances by size are modeled using the equation:

$$N_{t+1,j} = \sum_{i=1}^j [N_{t,i} e^{-M} - (\hat{C}_{t,i} + \hat{D}_{t,i} + \hat{Tr}_{t,i}) e^{(y_t-1)M}] X_{i,j} + R_{t+1,j} \quad (\text{A.1})$$

where $N_{t,i}$ is the number of [male] crab in length class i on 1 July (start of fishing year) of year t ; $\hat{C}_{t,i}$, $\hat{D}_{t,i}$, and $\hat{Tr}_{t,i}$ are respectively the predicted fishery retained, pot fishery discard dead, and groundfish fishery discard dead catches in length class i during year t ; $\hat{D}_{t,i}$ is estimated from the intermediate total ($\hat{T}_{t,i,temp}$) catch and the retained ($\hat{C}_{t,i}$) catch by Equation A.2c. $X_{i,j}$ is the probability of length-class i growing into length-class j during the year; y_t is elapsed time period from 1 July to the mid –point of fishing period in year t ; M is instantaneous rate of natural mortality; and $R_{t+1,j}$ recruitment to length class j in year $t+1$.

The catches are predicted using the equations

$$\hat{T}_{t,j,temp} = \frac{F_t s_{t,j}^T}{Z_{t,j}} N_{t,j} e^{-y_t M} (1 - e^{-Z_{t,j}}) \quad (\text{A.2a})$$

$$\hat{C}_{t,j} = \frac{F_t s_{t,j}^T s_{t,j}^r}{Z_{t,j}} N_{t,j} e^{-y_t M} (1 - e^{-Z_{t,j}}) \quad (\text{A.2b})$$

$$\hat{D}_{t,j} = 0.2(\hat{T}_{t,j,temp} - \hat{C}_{t,j}) \quad (\text{A.2c})$$

$$\hat{Tr}_{t,j} = 0.65 \frac{F_t^{Tr} s_j^{Tr}}{Z_{t,j}} N_{t,j} e^{-y_t M} (1 - e^{-Z_{t,j}}) \quad (\text{A.2d})$$

$$\hat{T}_{t,j} = \hat{C}_{t,j} + \hat{D}_{t,j} \quad (\text{A.2e})$$

where $Z_{t,j}$ is total fishery-related mortality on animals in length-class j during year t :

$$Z_{t,j} = F_t s_{t,j}^T s_{t,j}^r + 0.2 F_t s_{t,j}^T (1 - s_{t,j}^r) + 0.65 F_t^{Tr} s_j^{Tr} \quad (\text{A.3})$$

F_t is the full selection fishing mortality in the pot fishery, F_t^{Tr} is the full selection fishing mortality in the trawl fishery, $s_{t,j}^T$ is the total selectivity for animals in length-class j by the pot fishery during year t , s_j^{Tr} is the selectivity for animals in length-class j by the trawl fishery, $s_{t,j}^r$ is the probability

of retention for animals in length-class j by the pot fishery during year t . Pot bycatch mortality of 0.2 and groundfish bycatch mortality of 0.65 (average of trawl (0.8) and fish pot (0.5) mortality) were assumed.

Initial abundance

The initial conditions are computed as the equilibrium initial condition using the following relations:

The equilibrium stock abundance is

$$N = X.S.N + R \quad (\text{A.4})$$

The equilibrium abundance in 1960, N_{1960} , is

$$\underline{N}_{1960} = (I - XS)^{-1}R \quad (\text{A.5})$$

where X is the growth matrix, S is a matrix with diagonal elements given by e^{-M} , I is the identity matrix, and R is the product of average recruitment and relative proportion of total recruitment to each size-class.

We used the mean number of recruits from 1987 to 2012 in equation (A.5) to obtain the equilibrium solution under only natural mortality in year 1960, and then projected the equilibrium abundance under natural mortality with recruitment estimated for each year after 1960 up to 1985 with removal of retained catches during 1981/82 to 1984/85.

Growth Matrix

The growth matrix X is modeled as follows:

$$X_{i,j} = \begin{cases} 0 & \text{if } j < i \\ P_{i,j} + (1 - m_i) & \text{if } j = i \\ P_{i,j} & \text{if } j > i \end{cases} \quad (\text{A.6})$$

where:

$$P_{i,j} = m_i \begin{cases} \int_{-\infty}^{j_2 - L_i} N(x | \mu_i, \sigma^2) dx & \text{if } j = i \\ \int_{j_1 - L_i}^{j_2 - L_i} N(x | \mu_i, \sigma^2) dx & \text{if } i < j < n, \\ \int_{j_1 - L_i}^{\infty} N(x | \mu_i, \sigma^2) dx & \text{if } i = n \end{cases}$$

$$N(x | \mu_i, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\left(\frac{x - \mu_i}{\sqrt{2}\sigma}\right)^2}, \text{ and}$$

μ_i is the mean growth increment for crab in size-class i :

$$\mu_i = \omega_1 + \omega_2 * \bar{L}_i. \quad (\text{A.7})$$

ω_1 , ω_2 , and σ are estimable parameters, and j_1 and j_2 are the lower and upper limits of the receiving length-class j (in mm CL), and \bar{L}_i is the mid-point of the contributing length interval i . The quantity m_i is the molt probability for size-class i :

$$m_i = \frac{1}{1 + e^{c(\tau_i - d)}} \quad (\text{A.8})$$

where τ_i is the mid-length of the i -th length-class, c and d are parameters.

Selectivity and retention

Selectivity and retention are both assumed to be logistic functions of length. Selectivity depends on the fishing period for the pot fishery:

$$S_i = \frac{1}{1 + e^{\left[-\ln(19) \frac{\tau_i - \theta_{50}}{\theta_{95} - \theta_{50}}\right]}} \quad (\text{A.9})$$

where θ_{95} and θ_{50} are the parameters of the selectivity/ retention pattern (Mark Maunder, unpublished generic crab model). In the program, we re-parameterized the denominator ($\theta_{95} - \theta_{50}$) to $\log(\delta\theta)$ so that the difference is always positive and transformed θ_{50} to $\log(\theta_{50})$ to keep the estimate always positive.

Recruitment

Recruitment to length-class i during year t is modeled as $R_{t,i} = \bar{R}e^{\epsilon_i}\Omega_i$ where Ω_i is a normalized gamma function

$$\text{gamma}(x|\alpha_r, \beta_r) = \frac{x^{\alpha_r - 1} e^{-\frac{x}{\beta_r}}}{\beta_r^{\alpha_r} \Gamma(\alpha_r)} \quad (\text{A.10})$$

with α_r and β_r (restricted to the first five length classes).

Parameter estimation

Table A1 lists the parameters of the model indicating which are estimated and which are pre-specified. The objective function includes contributions related to the fit of the model to the available data and penalties (priors on various parameters).

Tables A2 lists parameter values (with the corresponding coefficient of variations in parentheses) used to weight the components of the objective functions for **EAG** and **WAG**.

Likelihood components

Catches

The contribution of the catch data (retained, total, and groundfish discarded) to the objective function is given by:

$$LL_r^{catch} = \lambda_r \sum_t \left\{ \ln \left(\sum_j \hat{C}_{t,j} w_j + c \right) - \ln \left(\sum_j C_{t,j} w_j + c \right) \right\}^2 \quad (\text{A.11a})$$

$$LL_T^{catch} = \lambda_T \sum_t \left\{ \ln \left(\sum_j \hat{T}_{t,j} w_j + c \right) - \ln \left(\sum_j T_{t,j} w_j + c \right) \right\}^2 \quad (\text{A.11b})$$

$$LL_{GD}^{catch} = \lambda_{GD} \sum_t \left\{ \ln \left(\sum_j \hat{Tr}_{t,j} w_j + c \right) - \ln \left(\sum_j Tr_{t,j} w_j + c \right) \right\}^2 \quad (\text{A.11c})$$

where λ_r , λ_T , and λ_{GD} are weights assigned to likelihood components for the retained, pot total, and groundfish discard catches; w_j is the average mass of a crab in length-class j ; $C_{t,j}$, $T_{t,j}$, and $Tr_{t,j}$ are, respectively, the observed numbers of crab in size class j for retained, pot total, and groundfish fishery discarded crab during year t , and c is a small constant value. We assumed $c = 0.001$.

An additional retained catch likelihood (using Equation A.11a without w) for the retained catch in number of crabs during 1981/82 to 1984/85 was also considered in all scenarios.

Catch-rate indices

The catch-rate indices are assumed to be lognormally distributed about the model prediction. Account is taken of variation in addition to that related to sampling variation:

$$LL_r^{CPUE} = \lambda_{r,CPUE} \left\{ 0.5 \sum_t \ln [2\pi(\sigma_{r,t}^2 + \sigma_e^2)] + \sum_t \frac{(\ln(CPUE_t^r + c) - \ln(\widehat{CPUE}_t^r + c))^2}{2(\sigma_{r,t}^2 + \sigma_e^2)} \right\} \quad (A.12)$$

where $CPUE_t^r$ is the standardized retained catch-rate index for year t , $\sigma_{r,t}$ is standard error of the logarithm of $CPUE_t^r$, and \widehat{CPUE}_t^r is the model-estimate of $CPUE_t^r$:

$$\widehat{CPUE}_t^r = q_k \sum_j S_j^T S_j^r (N_{t,j} - 0.5[\widehat{C}_{t,j} + \widehat{D}_{t,j} + \widehat{Tr}_{t,j}]) e^{-y_t M} \quad (A.13)$$

in which q_k is the catchability coefficient during the k -th time period (e.g., pre- and post-rationalization time periods), σ_e is the extent of over-dispersion, c is a small constant to prevent zero values (we assumed $c = 0.001$), and $\lambda_{r,CPUE}$ is the weight assigned to the catch-rate data. We used the same likelihood formula (A.12) for fish ticket and cooperative survey retained catch rate indices. However, for cooperative survey catch rate prediction we used a different catchability parameter.

Following Burnham *et al.* (1987), we computed the $\ln(CPUE)$ variance by:

$$\sigma_{r,t}^2 = \ln(1 + CV_{r,t}^2) \quad (A.14)$$

Length-composition data

The length-composition data are included in the likelihood function using the robust normal for proportions likelihood, i.e., generically:

$$LL_r^{LF} = 0.5 \sum_t \sum_j \ln(2\pi\sigma_{t,j}^2) - \sum_t \sum_j \ln \left[\exp\left(-\frac{(P_{t,j} - \widehat{P}_{t,j})^2}{2\sigma_{t,j}^2}\right) + 0.01 \right] \quad (A.15)$$

where $P_{t,j}$ is the observed proportion of crabs in length-class j in the catch during year t , $\widehat{P}_{t,j}$ is the model-estimate corresponding to $P_{t,j}$, i.e.:

$$\widehat{L}_{t,j}^r = \frac{\widehat{C}_{t,j}}{\sum_j \widehat{C}_{t,j}}$$

$$\begin{aligned}\hat{L}_{t,j}^T &= \frac{\hat{T}_{t,j}}{\sum_j^n \hat{T}_{t,j}} \\ \hat{L}_{t,j}^{GF} &= \frac{\hat{Tr}_{t,j}}{\sum_j^n \hat{Tr}_{t,j}}\end{aligned}\tag{A.16}$$

$\sigma_{t,j}^2$ is the variance of $P_{t,j}$:

$$\sigma_{t,j}^2 = \left[(1 - P_{t,j})P_{t,j} + \frac{0.1}{n} \right] / S_t\tag{A.17}$$

and S_t is the effective sample size for year t and n is the number of size classes.

Note: The likelihood calculation for retained length composition starts from length-class 6 (mid length 128 mm CL) because the length-classes 1 to 5 mostly contain zero data.

Tagging data

Let $v_{j,t,y}$ be the number of tagged male crab that were released during year t that were in size-class j when they were released and were recaptured after y years, and $\underline{\rho}_{j,t,y}$ be the vector of recaptures by size-class from the males that were released in year t that were in size-class j when they were released and were recaptured after y years. The log-likelihood corresponding to the multinomial distribution for the tagging data is then:

$$\ln L = \lambda_{y,tag} \sum_j \sum_t \sum_y \sum_i \rho_{j,t,y,i} \ln \hat{\rho}_{j,t,y,i}\tag{A.18}$$

where $\lambda_{y,tag}$ is the weight assigned to the tagging data for recapture year y , $\hat{\rho}_{j,t,y,i}$ is the proportion in size-class i of the recaptures of males that were released during year t that were in size-class j when they were released and were recaptured after y years:

$$\hat{\rho}_{j,t,y} \propto \underline{s}^T [\mathbf{X}]^y \underline{Z}^{(j)}\tag{A.19}$$

where $\underline{Z}^{(j)}$ is a vector with $v_{j,t,y}$ at element j and 0 otherwise, and \underline{s}^T is the vector of total selectivity for tagged male crab by the pot fishery. This log-likelihood function is predicated on the assumption that all recaptures are in the pot fishery and the reporting rate is independent of the size of crab.

Penalties

Penalties are imposed on the deviations of annual pot fishing mortality about mean pot fishing mortality, annual trawl fishing mortality about mean trawl fishing mortality, recruitment about mean recruitment, and the posfunction (fpen):

$$P_1 = \lambda_F \sum_t (\ln F_t - \ln \bar{F})^2\tag{A.20}$$

$$P_2 = \lambda_{F^{Tr}} \sum_t (\ln F_t^{Tr} - \ln \bar{F}^{Tr})^2$$

(A.21)

$$P_3 = \lambda_R \sum_t (\ln \varepsilon_t)^2 \quad (\text{A.22})$$

$$P_5 = \lambda_{\text{posfn}} * \text{fpen} \quad (\text{A.23})$$

Standardized Residual of Length Composition

$$\text{Std. Res}_{t,j} = \frac{P_{t,j} - \widehat{P}_{t,j}}{\sqrt{2\sigma_{t,j}^2}} \quad (\text{A.24})$$

Output Quantities

Harvest rate

Total pot fishery harvest rate:

$$E_t = \frac{\sum_{j=1}^n (\widehat{C}_{j,t} + \widehat{D}_{j,t})}{\sum_{j=1}^n N_{j,t}} \quad (\text{A.25})$$

Exploited legal male biomass at the start of year t :

$$LMB_t = \sum_{j=\text{legal size}}^n s_j^T s_j^r N_{j,t} w_j \quad (\text{A.26})$$

where w_j is the weight of an animal in length-class j .

Mature male biomass on 15 February spawning time (NPFMC 2007a, b) in the following year:

$$MMB_t = \sum_{j=\text{mature size}}^n \{N_{j,t} e^{-y'M} - (\widehat{C}_{j,t} + \widehat{D}_{j,t} + \widehat{T}_{r,j,t}) e^{(y_t - y')M}\} w_j \quad (\text{A.27})$$

where y' is the elapsed time from 1 July to 15 February in the following year.

For estimating the next year limit harvest levels from current year stock abundances, a F_{OFL} value is needed. Current crab management plan specifies five different Tier formulas for different stocks depending on the strength of information available for a stock, for computing F_{OFL} (NPFMC 2007a, b). For the golden king crab, the following Tier 3 formula is applied to compute F_{OFL} :

If,

$$MMB_{\text{current}} > MMB_{35\%}, F_{OFL} = F_{35\%}$$

If,

$$MMB_{\text{current}} \leq MMB_{35\%} \text{ and } MMB_{\text{current}} > 0.25MMB_{35\%},$$

$$F_{OFL} = F_{35\%} \frac{\left(\frac{MMB_{\text{current}}}{MMB_{35\%}} - \alpha\right)}{(1-\alpha)} \quad (\text{A.28})$$

If,

$$MMB_{\text{current}} \leq 0.25MMB_{35\%},$$

$$F_{OFL} = 0.$$

where α is a parameter, MMB_{current} is the mature male biomass in the current year and $MMB_{35\%}$ is the proxy MMB_{MSY} for Tier 3 stocks. We assumed $\alpha = 0.1$.

Because projected MMB_t (i.e., MMB_{current}) depends on the intervening retained and discard catch (i.e., MMB_t is estimated after the fishery), an iterative procedure is applied using Equations A.27 and A.28 with retained and discard catch predicted from Equations A.2b-d. The next year limit harvest catch is estimated using Equations A.2b-d with the estimated F_{OFL} value.

Table A1. Pre-specified and estimated parameters of the population dynamics model

Parameter	Number of parameters
<i>Fishing mortalities:</i>	
Pot fishery, F_t	1981–2019 (estimated)
Mean pot fishery fishing mortality, \bar{F}	1 (estimated)
Groundfish fishery, F_t^{Tr}	1989–2019 (the mean F for 1989 to 1994 was used to estimate groundfish discards back to 1981 (estimated))
Mean groundfish fishery fishing mortality, \bar{F}^{Tr}	1 (estimated)
<i>Selectivity and retention:</i>	
Pot fishery total selectivity, θ_{50}^T	2 (1981–2004; 2005+) (estimated)
Pot fishery total selectivity difference, $\Delta\theta^T$	2 (1981–2004; 2005+) (estimated)
Pot fishery retention, θ_{50}^R	1 (1981+) (estimated)
Pot fishery retention selectivity difference, $\Delta\theta^R$	1 (1981+) (estimated)
Groundfish fishery selectivity	fixed at 1 for all size-classes
<i>Growth:</i>	
Expected growth increment, ω_1, ω_2	2 (estimated)
Variability in growth increment, σ	1 (estimated)
Molt probability (size transition matrix with tag data), a	1 (estimated)
Molt probability (size transition matrix with tag data), b	1 (estimated)
Natural mortality, M	1 (pre-specified, 0.21yr^{-1})
<i>Recruitment:</i>	
Number of recruiting length-classes	5 (pre-specified)
Mean recruit length	1 (pre-specified, 110 mmCL)
Distribution to length-class, β_r	1 (estimated)
Median recruitment, \bar{R}	1 (estimated)
Recruitment deviations, \mathcal{E}_t	60 (1961–2020) (estimated)
Fishery catchability, q	2 (1985–2004; 2005+) (estimated)
Additional CPUE indices standard deviation, σ_e	1 (estimated)
Likelihood weights (coefficient of variation)	Pre-specified, varies by scenario

Table A2. Specifications for the weights with corresponding coefficient of variations* in parentheses for each model for **EAG** and **WAG**.

Weight	Model 20_1	Model 20_1b	Model 20_2
<i>Catch:</i>			
Retained catch for 1981–1984 and/or 1985–2019, λ_r	500 (0.032)	500	500
Total catch for 1990–2019, λ_T	Number of sampled pots scaled to a max 250	Number of sampled pots scaled to a max 250	Number of sampled pots scaled to a max 250
Groundfish bycatch for 1989–2019, λ_{GD}	0.2 (3.344)	0.2	0.2
<i>Catch-rate:</i>			
Observer legal size crab catch-rate for 1995–2019, $\lambda_{r,CPUE}$	1 (0.805)	1	1
Fish ticket retained crab catch-rate for 1985–1998, $\lambda_{r,CPUE}$	1 (0.805)	1	1
<i>Penalty weights:</i>			
Pot fishing mortality dev, λ_F	Initially 1000, relaxed to 0.001 at phases \geq select. phase	Initially 1000, relaxed to 0.001 at phases \geq select. phase	Initially 1000, relaxed to 0.001 at phases \geq select. phase
Groundfish fishing mortality dev, λ_{F^*}	Initially 1000, relaxed to 0.001 at phases \geq select. phase	Initially 1000, relaxed to 0.001 at phases \geq select. phase	Initially 1000, relaxed to 0.001 at phases \geq select. phase
Recruitment, λ_R	2 (0.533)	2	2
Posfunction (to keep abundance estimates always positive), λ_{posfn}	1000 (0.022)	1000	1000
Tagging likelihood	EAG individual tag returns	EAG tag data	EAG tag data

* Coefficient of Variation, $CV = \sqrt{\exp\left[\frac{1}{2w}\right] - 1}$, w = weight

Appendix B: Catch and CPUE data

The commercial catch and length frequency distribution were estimated from ADF&G landing records and dockside sampling (Bowers *et al.* 2008, 2011). The annual retained catch, total catch, and groundfish (or trawl) discarded mortality are provided in Tables 1, 2, and 2b for **EAG** and **WAG**. The weighted length frequency data were used to distribute the catch into 5-mm size intervals. The length frequency data for a year were weighted by each sampled vessel's catch as follows. The i -th length-class frequency was estimated as:

$$\sum_{j=1}^k C_j \frac{LF_{j,i}}{\sum_{i=1}^n LF_{j,i}} \quad (\text{B.1})$$

where k = number of sampled vessels in a year, $LF_{j,i}$ = number of crabs in the i -th length-class in the sample from j -th vessel, n = number of size classes, C_j = number of crabs caught by j -th vessel. Then the relative frequency for the year was calculated and applied to the annual retained catch (in number of crabs) to obtain retained catch by length-class.

The annual total catch (in number of crabs) was estimated by the observer nominal (unstandardized) total CPUE considering all vessels multiplied by the total fishing effort (number of pot lifts). The weighted length frequency of the observer samples across the fleet was estimated using Equation B.1. Observer measurement of crab ranged from 20 to 220 mm CL. To restrict the total number of crabs to the model assumed size range (101–185+ mm CL), the proportion of observer total relative length frequency corresponding to this size range was multiplied by the total catch (number of crabs). This total number of crabs was distributed into length-classes using the weighted relative length frequency. Thus, crab sizes < 101 mm CL were excluded from the model. In addition, all crab >185 mm CL were pooled into a plus length class. Note that the total crab catch by size that went into the model did not consider retained and discard components separately. However, once the model estimated the annual total catch, then retained catch was deducted from this total and multiplied by handling mortality [we used a 20% handling mortality (Siddeek *et al.* 2005) to obtain the directed fishery discarded (dead) catch].

Observer data have been collected since 1988 (Moore *et al.* 2000; Barnard *et al.* 2001; Barnard and Burt 2004; Gaeuman 2011), but data were not comprehensive in the initial years, so a shorter time series of data for the period 1990/91–2018/19 was selected for this analysis. During 1990/91–1994/95, observers were only deployed on catcher-processor vessels. During 1995/96–2004/05, observers were deployed on all fishing vessels during fishing activity. Observers have been deployed on all fishing vessels since 2005/06, but catcher-only vessels are only required to carry observers for a minimum of 50% of their fishing activity during a season; catcher-processor vessels are still required to carry observers during all fishing activity. Onboard observers sample seven pots per day (it can be different number of pots per string) and count and measure all crabs caught and categorize catch as females, sublegal males, retained legal males, and non-retained legal males in a sampled pot. Prior to the 2009/10

season, depending on season, area, and type of fishing vessel, observers were also instructed to sample additional pots in which all crab were only counted and categorized as females, sublegal males, retained legal males, and non-retained legal males, but were not measured. Annual mean nominal CPUEs of retained and total crabs were estimated considering all sampled pots within each season (Table 3). The observer CPUE data collection improved over the years and the data since 1995/96 are more reliable. Thus, for model fitting, the observer CPUE time series was restricted to 1995/96–2019/20. The 1990/91–2019/20 observer database consists of 116,508 records and that of 1995/96–2019/20 contains 112,229 records. For CPUE standardization, these data were further reduced by 5% cutoff of Soak time and 1% cutoff of Depth on both ends of the variable range to remove unreliable data or data from dysfunctional pot operations, and restricting to vessels which have made five trips per year for at least three years during 1985/86–2019/20.

Length-specific CPUE data collected by observers provides information on a wider size range of the stock than did the commercial catch length frequency data obtained from mostly legal-sized landed males.

There were significant changes in fishing practice due to changes in management regulations (e.g., since 1996/97 constant TAC and since 2005/06 crab rationalization), pot configuration (escape web on the pot door increased to 9” since 1999), and improved observer recording in Aleutian Islands golden king crab fisheries since 1998. These changes prompted us to consider two separate observer CPUE time series, 1995/96–2004/05 and 2005/06–2019/20, to estimate CPUE indices for model input.

To include a long time series of CPUE indices for stock abundance contrast, we also considered the 1985/86–1998/99 legal size standardized CPUE as a separate likelihood component in all scenarios. Because of the lack of soak time data before 1990, we estimated the CPUE index considering a limited set of explanatory variables (e.g., vessel, captain, area, month) and fitting the lognormal and negative binomial GLM models to fish ticket data (Tables 4 and 13).

When using CPUE indices in the model fit, we compared the predicted with the observed legal male CPUE in the observer CPUE likelihoods because legal male (retained plus non-retained) data are more reliable than total in the observer samples.

The CPUE standardization followed the GLM fitting procedure (Maunder and Punt 2004; Starr 2012; Siddeek et al. 2018). Following a suggestion made by the CIE reviewers (CIE, June 2018) we reduced the number of gear codes in the database after consulting with the fishing industry (Rip Carlton, Chad Hoefler, and Scott Goodman, personal communication December 2018; Table B1). Following SSC (October 2018) suggestion, we used a hybrid procedure: First, selected a scope of variables set by Akaike Information Criterion, AIC (Burnham and Anderson 2002). An increase of more than 2 units in the AIC was used to identify the variable to be included successively (stepAIC program, R Core Team 2018). Then, the model parsimony was

improved further by successively removing the term that explained the least proportion of deviance ($R^2 < 0.01$) (stepCPUE R function was used, Siddeek et al. 2018). Feenstra, et al. (unpublished 2019) used a similar hybrid approach.

Table B.1. Updated Gear code for observer data analysis. Only gear code # 5, 6, 7, 8, and 13 were considered following crab industry suggestion. Note: Identical codes were given to those gear codes with similar catchability/selectivity. X stands for the gear codes that were ignored.

Original Gear code	Pot gear description	Mark X against the code that can be ignored	Number Encountered by Observers during 1990-2016	Updated Gear Code
1	Dungeness crab pot, small & round	X	2	X
2	Pyramid pot, tunnel openings usually on sides, stackable	X	2121	X
3	Conical pot, opening at top of cone, stackable	X	2000	X
4	4' X 4' rectangular pot		60	X
5	5' X 5' rectangular pot		18032	5
6	6' X 6' rectangular pot		17508	6
7	7' X 7' rectangular pot		23806	7
8	8' X 8' rectangular pot		1936	8
9	5 1/2' X 5 1/2' rectangular pot		6934	5
10	6 1/2' X 6 1/2' rectangular pot		22085	6
11	7 1/2' X 7 1/2' rectangular pot		387	7
12	Round king crab pot, enlarged version of Dungeness crab pot		8259	X
13	10' X 10' rectangular pot		466	13
14	9' X 9' rectangular pot	X	1	X
15	8 1/2' X 8 1/2' rectangular pot	X	1	X
16	9 1/2' X 9 1/2' rectangular pot	X	Not used	X
17	8' X 9' rectangular pot	X	1	X
18	8' X 10' rectangular pot	X	1	X
19	9' X 10' rectangular pot		Not used	X
20	7' X 8' rectangular pot	X	252	X
21	Hair crab pot, longlined and small, stackable		Not used	X
22	snail pot	X	1	X

23	Dome-shaped pot, tunnel opening on top, often longlined in deep-water fisheries ADF&G shellfish research 7' X 7' X34"	X	6756	X
24	rectangular pot with 2.75" stretch mesh and no escapement rings or mesh		Research pot	X
80	Historical: Cod pot, any shape pot targeting cod, usually with tunnel fingers	X	711	X
81	Historical: Rectangular pot, unknown size, with escape rings	X	1123	X

All scenarios used CPUE indices estimated by the hybrid GLM method. Following January 2019 CPT request, we considered an Year:Area interaction factor as a special case for a CPUE standardization scenario.

Thus we estimated two sets of observer CPUE indices for model input, 20_1 (reduced number of gear codes), and 20_2 (reduced number of gear codes and Year:Area interaction).

Observer CPUE index by GLM:

a. Non-interaction GLM model:

The CPUE standardization followed the GLM fitting procedure (Maunder and Punt 2004; Starr 2012; Siddeek *et al.* 2016b). We considered the negative binomial GLM on positive and zero catches to select the explanatory variables. The response variable CPUE is the observer sample catch record for a pot haul. The negative binomial model uses the log link function for the GLM fit.

For the non-interaction model, we assumed the null model to be

$$\ln(\text{CPUE}_i) = \text{Year}_{y_i} \tag{B.2}$$

where Year is a factorial variable.

The maximum set of model terms offered to the stepwise selection procedure was:

$$\ln(\text{CPUE}_i) = \text{Year}_{y_i} + \text{ns}(\text{Soak}_{s_i}, \text{df}) + \text{Month}_{m_i} + \text{Vessel}_{v_i} + \text{Captain}_{c_i} + \text{Area}_{a_i} + \text{Gear}_{g_i} + \text{ns}(\text{Depth}_{d_i}, \text{df}), \tag{B.3}$$

where Soak is in unit of days and is numeric; Month, Area (Block) code, Vessel code, Captain code, and Gear code are factorial variables; Depth in fathom is a numeric variable; ns=cubic spline, and df = degree of freedom.

We used a log link function and a dispersion parameter (θ) in the GLM fitting process. We used the R^2 criterion for predictor variable selection (Siddeek *et al.* 2016b).

The degrees of freedom and dispersion parameters were determined by calculating AICs for a range of values and locating the best value at the minimum AIC (results are not shown but available with the first author).

Instead of using the traditional AIC ($-2\log_{\text{likelihood}}+2p$) we used the Consistent Akaike Information Criteria (CAIC) (Bozdogan 1987) $\{-2\log_{\text{likelihood}}+[\ln(n)+1]*p\}$ for variable selection by StepAIC, where n=number of observations and p= number of parameters to be estimated. The number of selected variables were further reduced for parsimony, if feasible, by the R^2 criterion using the StepCPUE function. i.e., a hybrid selection procedure (Feenstra *et al.* 2019).

Example R codes used for main effect GLM fitting are as follows:

For EAG 1995_04 CPUE indices:

```
library(MASS)
```

```
library(splines)
```

Step 1:

```
glm.object<- glm(Legals~Year,family = negative.binomial(1.38),data=datacore)
```

```
epotsampleoutAIC<-stepAIC(glm.object,scope=list(upper=  
~(Year+ns(SoakDays,df=4)+Month+Vessel+Captain+Area+Gear+ns(Depth,df=16)),lower  
=~Year),family=negative.binomial(1.38),direction='forward',trace=9,k=log(nrow(datacor  
e))+1.0)
```

Step 2:

```
glm.object<- glm(Legals~Year,family = negative.binomial(1.38),data=atacore)
```

```
epotsampleout<-  
stepCPUE(glm.object,scope=list(upper=~(Year+Gear+Captain+ns(SoakDays,df=4)+  
Month+Area),lower=~Year),family=negative.binomial(1.38),direction="forward",trace=9,  
r2.change=0.01)
```

The final main effect models for **EAG** were:

Model 20_1:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Gear} + \text{Captain} + \text{ns}(\text{Soak}, 4) + \text{Month} + \text{Block}$$

AIC=203808

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Gear} + \text{Captain} + \text{ns}(\text{Soak}, 4) + \text{Month} \quad (\text{B.4})$$

for the 1995/96–2004/05 period [$\theta=1.38$, $R^2 = 0.2205$]

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Captain} + \text{Gear} + \text{ns}(\text{Soak}, 16)$$

AIC=72738

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Captain} + \text{ns}(\text{Soak}, 16) + \text{Gear} \quad (\text{B.5})$$

for the 2005/06–2019/20 period [$\theta = 2.33$, $R^2 = 0.1125$].

The final models for **WAG** were:

Model 20_1:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Captain} + \text{ns}(\text{Soak}, 15) + \text{Gear} + \text{Area} + \text{Month} + \text{Vessel}$$

AIC=191025

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Captain} + \text{ns}(\text{Soak}, 15) + \text{Gear} \quad (\text{B.6})$$

for the 1995/96–2004/05 period [$\theta=0.97$, $R^2 = 0.1684$]

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Gear} + \text{Vessel} + \text{Month} + \text{ns}(\text{Soak}, 19)$$

AIC=110148

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Gear} + \text{ns}(\text{Soak}, 19) \quad (\text{B.7})$$

for the 2005/06–2019/20 period [$\theta = 1.13$, $R^2 = 0.0525$, Soak forced in].

b. Year:Area interaction GLM:

For year and area interaction analysis, we designed the areas in to 1 X 1 nmi grids enmeshed in 10 larger blocks as follows. The number of blocks was restricted to a few to prevent GLM fitting problems.




Figure B.1. The 1995/96 to 2019/20 observer pot samples enmeshed in 10 blocks for the Aleutian Islands golden king crab.

The blocks were determined from visually exploring each year's pot distribution locations (each year's data plots are available with the first author). The blocks contain observed patches of crab distribution during this time period.

Table B.1. Number of 1 x 1 nmi grids containing observer sample locations within each block by fishing year for the Aleutian Islands golden king crab, 1995/96–2019/20 data. Blocks 1–4 belong to **EAG** and 5 – 10 to **WAG**. Sum of ever fished number of grids for each block is listed at the bottom row.

Year	Block_1	Block_2	Block_3	Block_4	Block_5	Block_6	Block_7	Block_8	Block_9	Block_10
1995	125	529	748	379	218	373	112	722	166	122
1996	149	814	761	372	89	473	359	799	200	35
1997	116	530	755	257	202	443	104	568	274	0
1998	78	581	453	236	18	318	157	251	132	0
1999	123	593	454	231	163	476	182	627	193	145
2000	72	540	754	301	187	440	195	555	547	47
2001	123	507	507	329	45	369	288	634	256	9
2002	97	387	584	271	71	341	205	335	242	37
2003	43	492	530	299	111	347	212	465	150	61
2004	81	289	377	216	77	319	150	359	172	116
2005	0	205	221	118	8	220	83	261	54	0
2006	0	154	248	122	15	191	58	220	39	0
2007	0	111	177	110	24	228	78	173	20	0
2008	0	111	203	93	12	181	67	196	0	0
2009	0	59	146	60	6	137	95	220	25	0
2010	0	81	141	85	1	115	73	260	39	0
2011	0	126	117	33	3	83	73	266	9	0
2012	0	146	110	56	7	91	85	312	53	0
2013	2	149	129	51	12	144	105	293	86	0
2014	1	138	96	41	39	120	114	319	37	0
2015	0	135	147	61	46	163	106	280	16	48
2016	0	145	231	63	26	134	89	210	106	0

2017	0	97	170	110	11	87	79	198	118	0
2018	0	91	158	95	7	69	82	204	121	0
2019	1	112	171	101	0	0	89	316	138	0

	Block_1	Block_2	Block_3	Block_4	Block_5	Block_6	Block_7	Block_8	Block_9	Block_10
1995-2019 - Sum of 1x1 cells ever fished	375	1363	1754	907	452	1026	777	1940	998	325

We assumed the null model to be

$$\ln(\text{CPUE}_i) = \text{Year}_{y_i}:\text{Area}_{a_i} \quad (\text{B.8})$$

The maximum set of model terms offered to the stepwise selection procedure was:

$$\ln(\text{CPUE}_i) = \text{Year}_{y_i}:\text{Area}_{a_i} + \text{ns}(\text{Soak}_{s_i}, \text{df}) + \text{Month}_{m_i} + \text{Vessel}_{v_i} + \text{Captain}_{c_i} + \text{Area}_{a_i} + \text{Gear}_{g_i} + \text{ns}(\text{Depth}_{d_i}, \text{df}). \quad (\text{B.9})$$

Example R codes used for interaction effect GLM fitting are as follows:

For WAG 1995_04 CPUE indices:

```
library(MASS)
```

```
library(splines)
```

Step 1:

```
glm.object<- glm(Legals~Year:Area,family = negative.binomial(0.97),data=datacore)
```

```
wpotsampleoutAIC<-stepAIC(glm.object,scope=list(upper=  
~(Year:Area+ns(SoakDays,df=15)+Month+Vessel+Captain+Area+Gear  
ns(Depth,df=18)),lower=~Year:Area),family=  
negative.binomial(0.97),direction="forward",trace=9,k=log(nrow(datacore))+1.0)
```

Step 2:

```
glm.object<- glm(Legals~Year:Area,family = negative.binomial(0.97),data=datacore)
```

```
wpotsampleout<-stepCPUE(glm.object,scope=list(upper=  
~(Captain+ns(SoakDays,df=15)+Gear+Area+Month+Year:Area),lower=  
~Year:Area),family=  
negative.binomial(0.97),direction="forward",trace=9,r2.change=0.01)
```

The final interaction effect models for **EAG** were:

Model 20_2:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Gear} + \text{Captain} + \text{ns}(\text{Soak}, 4) + \text{Month} + \text{Year: Area}$$

AIC=203851

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Gear} + \text{Captain} + \text{ns}(\text{Soak}, 4) + \text{Year: Area} \quad (\text{B.10})$$

for the 1995/96–2004/05 period [$\theta=1.38$, $R^2 = 0.2235$]

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Vessel} + \text{Gear} + \text{ns}(\text{Soak}, 16) + \text{Year: Area}$$

AIC=72860

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Vessel} + \text{ns}(\text{Soak}, 16) + \text{Gear} + \text{Year: Area} \quad (\text{B.11})$$

for the 2005/06–2019/20 period [$\theta = 2.33$, $R^2 = 0.1238$].

The final interaction effect models for **WAG** were:

Model 20_2:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Vessel} + \text{ns}(\text{Soak}, 15) + \text{Gear} + \text{Month} + \text{Year: Area}$$

AIC=191140

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Vessel} + \text{ns}(\text{Soak}, 15) + \text{Gear} + \text{Year: Area} \quad (\text{B.12})$$

for the 1995/96–2004/05 period [$\theta=0.97$, $R^2 = 0.1721$]

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Gear} + \text{Vessel} + \text{Month} + \text{Year: Area} + ns(\text{Soak}, 19)$$

$$\text{AIC}=110438$$

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Gear} + \text{Year: Area} + ns(\text{Soak}, 19) \quad (\text{B.13})$$

for the 2005/06–2019/20 period [$\theta = 1.13$, $R^2 = 0.0708$, Soak forced in].

Steps:

1. *Block-scale analysis:*

The estimate of the CPUE index in each Year-Area (Area=Block) was first obtained:

$$\text{CPUE}_{ij} = e^{YB_{ij} + \sigma_{ij}^2/2} \quad (\text{B.14})$$

Where CPUE_{ij} is the CPUE index in the i th year and j th block, YB_{ij} is the coefficient of the i th year and j th block interaction, and σ_{ij} is the biased correction standard error for expected CPUE value.

The number of 1 x 1 nmi grids in each block can change from year to year; so, we considered using the number of grids **ever fished** in a block, N_{everj} [this is equivalent to assuming that the grids fished in any year randomly sample the stock in that block (see Campbell, 2004)].

The abundance index for j th block in i th year is

$$B_{ij} = N_{everj} \text{CPUE}_{ij} \quad (\text{B.15})$$

As you noticed in Table B.1 that there are no-observer samplings took place in certain years for a whole block. We filled the B_{ij} index gaps by filling them using a log-linear model, i.e.:

$$\hat{B}_{i,j} = e^{A_i + C_j} \quad (\text{B.16})$$

where $B_{i,j}$ is the index of biomass for year i and block j , A_i is a year factor, and C_j is a block factor, and used this model to predict the biomass index for blocks x years with no (or very limited) data.

Annual biomass index, B_i , was estimated as,

$$B_i = \sum_j B_{ij} \quad (\text{B.17})$$

The variance of the total biomass index was computed as:

$$\text{Var}(B_i) = \sum_j N_{\text{ever},j}^2 \text{var}(CPUE_{i,j}) \quad (\text{B.18})$$

where $N_{\text{ever},j}$ is the total number of 1x1 mni cells ever fished in block j , and $CPUE_{i,j}$ is the CPUE index for year i and block j .

To compare with other CPUE index estimates (Figures 24 for **EAG** and 42 for **WAG**) as well as to input into the assessment model (models 20_2 for **EAG** and **WAG**, and 20_2b for **EAG**), we rescaled the B_i indices by the geometric mean of estimated B_i values separately for the pre- and post-rationalization periods. The corresponding coefficient of variation (CV_i) of $CPUE_i$ was estimated by

$$\sqrt{\frac{\text{Var}(B_i)}{(B_i)^2}} \quad (\text{B.19})$$

Following Burnham *et al.* (1987), the variance of $\ln(CPUE_i)$ for input to assessment models were estimated by $\sigma_i^2 = \ln(1 + CV_i^2)$.

c. Commercial fishery CPUE index by non-interaction model:

We fitted separate lognormal and negative binomial GLM models for fish ticket retained CPUE time series 1985/86 – 1998/99 offering Year, Month, Vessel, Captain, and Area as explanatory variables and applying the hybrid selection method. Reduced area resolution (grouped ADF&G code- AreaGP) was used for model fitting.

The final model under lognormal error structure for **EAG** was:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Month}$$

AIC=5,805

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Month} \quad (\text{B.20})$$

for the 1985/86–1998/99 period [$R^2 = 0.3700$]

and that for **WAG** was:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Area}$$

AIC= 11,082

Final selection by stepCPUE

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel}, R^2 = 0.3679 \quad (\text{B.21})$$

The final model under negative binomial error structure for **EAG** was:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Month}$$

AIC=16,997

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Month} \quad (\text{B.22})$$

for the 1985/86–1998/99 period [$\theta=10.45, R^2 = 0.3328$]

and that for **WAG** was:

Initial selection by stepAIC:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Area}$$

AIC=31,701

Final selection by stepCPUE:

$$\ln(\text{CPUE}) = \text{Year} + \text{Vessel} + \text{Area} \quad (\text{B.23})$$

for the 1985/86–1998/99 period [$\theta=6.67, R^2 = 0.3569$]

Appendix C. Cooperative Survey

1. Brief summary of the survey method

The ADF&G and industry collaborative pot survey was initiated in 2015 in the **EAG** and continued since then. The survey was extended to **WAG** in 2018. A stratified two-stage sampling design has been implemented in a 2 X 2 nmi grids within 1000 m depth covering the entire golden king crab fishing area. The 2 x 2 nmi choice was the best compromise between scale of fishing gear, accuracy of defining habitat, and number of possible stations (Figure C1).



Figure C.1. Survey design: 2 x 2 nmi grids overlaid on observer pot sample locations (green squares) in **EAG**.

There are nearly 1100 grids in the **EAG** divided into three equal size strata for selecting random pot sampling locations (Figures C.2 and C.3).




Figure C.2. Survey design: 2 x 2 nmi grids stratified by three equal sizes for selecting random pot sampling locations in **EAG**.




Figure C.3. Random sample of 22 cells selected in each of three sub strata in **EAG** during the 2019 fishery.

Survey occurs during the first month of each fishing season with one to two ADF&G biologists onboard the fishing vessel to collect fishery and biological data. Fishing operation takes place in a randomly selected set of grids in each strata with long-line pots. The number of pots per string ranges from 30 to 40, 200 m apart, and a vessel carry on average 35 strings. Pot sizes range from 5.5 x 5.5 ft to 7 x 7 ft with large mesh sizes for retention of legal king crab. A few small mesh size research pots are also deployed for special studies. Fishing operation is not standardized for depth or soak time to allow normal fishing practices.

There are multiple pots (typically about 5 pots) sampled for each long-line string with approximately 35 crab measurement made per pot. For example, if 100 crabs are caught in a sampled pot, the biologist measures every third crab. The following snapshot of an observation record will provide details of what stock assessment data are collected.

Work on details size composition plots and CPUE by size, year, and area is not yet finished to present at this time.

fishery	year	vessel	skipper	String#	pot_size	mesh_size	bait	subsample_rate	species_code	sex	size	legal
EAG	2015	20556	Chad_Hoefer	1	5x5	king(large)	halibut	2	923	1	187	1

Pot#	date_in	time_in	depth_start	start_lat	start_lon	depth_out	end_lat	end_lon	date_out	time_out	comments	soak_time
1	8/4/2015	17:00	132	52.74133	-170.692	133	52.7515	-170.675	8/17/2015	3:00		12.41667

2. Standardization of cooperative survey CPUE by mixed random effects model:

The unique property of cooperative survey is that multiple pots from multiple strings are sampled. All sample measurements were taken in **EAG** except for 2018 and 2019, during which measurements were also taken from **WAG**. The CPT and SSC suggested to use the random effects model to standardize the survey CPUE data.

Data:

There are 27,255 records from five-year (2015–2019) cooperative surveys.

Data preparation for CPUE standardization:

- i.) Created two new columns by concatenating Vessel Code with String# as well as Pot# because String# and Pot# are not unique numbers to each vessel. The new column names were identified as VesString and VesPot. For example, a Vessel Code 20556 with a String# 3 was concatenated to be 205563 in a new column VesString, and a Vessel Code 20556 with a Pot# 5 was concatenated to be 205565 in a new column VesPot.
- ii.) Raised the Catch in each record by the Sample Rate.
- iii.) Subset the data by large mesh king crab pot (Mesh ID not equal to 2), legal size (Size > 135 mm CL), and **EAG** (EAGWAG=1). The female (Sex=2) catch without any male (Sex=1) in a crab pot was set to 0 to account for the possibility of zero catch for expected CPUE determination.
- iv.) Further subset the data by 5% to 95% trimmed Soak time and 1% to 99% trimmed Depth. This is to exclude catches from any unusual pot operations.
- v.) Summed up the catch across sizes for each Pot# and labelled it as SumCatch. Thus, each Pot# has a single catch number.

The mixed random effects model considered a random intercept procedure with the following model formulation:

$$\text{Sum Catch} = Y + \text{ns}(\text{Soak}, \text{df}=16) + \text{ns}(\text{Depth}, \text{df}=10) + (1|\text{Vessel/Pot}) + (1|\text{Block/String})$$

We used the “lme4” library in R (version 3.5.1, R Core Team, 2018) with the “glmer()” function to fit the mixed random effects model. The glmer() function allows to use any type of error model (we used the negative binomial model) to fit the data:

library(MASS)

library(splines)

library(Matrix)

library(lme4)


```
best.lmeFit <- glmer(SumCatch ~ Year + ns(SoakDays, df=16) + ns(Depth,
df=10) + (1/Vessel/VesPot) + (1/Block/VesString), family =
negative.binomial(2.33), control = glmerControl(optimizer = "bobyqa",
optCtrl = list(maxfun = 100000)), data = eSurvey15_19Subtrim)
```

where Sum Catch = observed CPUE, best.lmeFit = expected CPUE. Year, SoakDays and Depth are fixed effect variables. The fixed effect variables were selected from fit of a fixed effect model on the survey data. The dispersion parameter value for the negative binomial error model and the degrees of freedom for cubic splines were borrowed from the observer final GLM model estimate for **EAG** for the post rationalization period.

The QQ plot for the fit assured model assumptions were correct (Figure C.4).




Figure C4. Studentized residual plot for the mixed random effects model fit. The 2015–2019 **EAG** data were used.

Comparison of standardized CPUE from cooperative survey data (2015–19) for **EAG** and the corresponding years' observer CPUE indices indicated similar pattern except for 2019 (Figure C5).




Figure C5. Comparison of cooperative survey CPUE indices (green) and model 20_1 CPUE indices (red). The confidence limits are determined with $\pm 2SE$.

We standardized the yearly mean of predicted survey CPUEs for 2015–2019 by the geometric mean to obtain the CPUE indices for input to the assessment model (20_1c and 20_2b) (Table C.1).

Table C.1. The cooperative survey expected legal size male standardized (by geometric mean) CPUE indices by the mixed random effects model, standard errors (SE), and lower- and upper-95% confidence limits for assessment model input for **EAG**, 2015–2019 data.

Year	Predicted CPUE index	SE	Lower Limit	Upper Limit
2015	1.1137	0.0265	1.0562	1.1743
2016	0.9459	0.0266	0.8968	0.9976
2017	1.1075	0.0417	1.0189	1.2038
2018	1.1690	0.0365	1.0868	1.2575
2019	0.7332	0.0382	0.6793	0.7914

Appendix D: Jittering

Jittering of models 20_1b and 20_2 parameter estimates:

We followed the Stock Synthesis approach to do 100 jitter runs of models 20_1b and 20_2 parameter estimates to use as initial parameter values (as .PIN file in ADMB) to assess model stability and to determine whether a global as opposed to local minima has been reached by the search algorithm:

The *Jitter* factor of 0.3 was multiplied by a random normal deviation $rdev=N(0,1)$, to a transformed parameter value based upon the predefined parameter:

$$temp = 0.5 * rdev * Jitterfactor * \ln\left(\frac{P_{max} - P_{min} + 0.0000002}{P_{val} - P_{min} + 0.0000001} - 1\right), \quad (D.1)$$

with the final jittered initial parameter value back transformed as:

$$P_{new} = P_{min} + \frac{P_{max} - P_{min}}{1.0 + \exp(-2.0 temp)}, \quad (D.2)$$

where P_{max} and P_{min} are upper and lower bounds of parameter search space and P_{val} is the estimated parameter value before the jittering.

The jitter results are summarized for scenarios 20_1b in Tables D.1 and D.2; and 20_2 in Tables D.3 and D.4 for **EAG** and **WAG**, respectively. Almost all runs converged to the highest log likelihood values. We concluded from jitter results that optimization of 20_1b and 20_2 models achieved global minima.

Table D.1. Results from 100 jitter runs for scenario 20_1b for **EAG**. Jitter run 0 corresponds to the original optimized estimates.

Jitter Run	Objective Function	Maximum Gradient	B _{35%} (t)	OFL (t)	Current MMB (t)
0	12.9831	0.003023	6,774	2,986	8,470
1	12.8964	0.000280	6,774	2,986	8,470
2	12.8964	0.000192	6,774	2,986	8,470
3	12.8964	0.000159	6,774	2,986	8,470
4	12.8964	0.000426	6,774	2,986	8,470
5	12.8964	0.000180	6,774	2,986	8,470
6	12.8964	0.000053	6,774	2,986	8,470
7	12.8964	0.000093	6,774	2,986	8,470
8	12.8964	0.000054	6,774	2,986	8,470
9	12.8964	0.000593	6,774	2,986	8,470
10	12.8964	0.000032	6,774	2,986	8,470
11	12.8964	0.000125	6,774	2,986	8,470
12	12.8964	0.000022	6,774	2,986	8,470

13	12.8964	0.000350	6,774	2,986	8,470
14	12.8964	0.000350	6,774	2,986	8,470
15	12.8964	0.000216	6,774	2,986	8,470
16	12.8964	0.000017	6,774	2,986	8,470
17	12.8964	0.000035	6,774	2,986	8,470
18	12.8964	0.000285	6,774	2,986	8,470
19	12.8964	0.000014	6,774	2,986	8,470
20	12.8964	0.000085	6,774	2,986	8,470
21	12.8964	0.000057	6,774	2,986	8,470
22	12.8964	0.000025	6,774	2,986	8,470
23	12.8964	0.000025	6,774	2,986	8,470
24	12.8964	0.000089	6,774	2,986	8,470
25	12.8964	0.000015	6,774	2,986	8,470
26	12.8964	0.000153	6,774	2,986	8,470
27	12.8964	0.000072	6,774	2,986	8,470
28	12.8964	0.000113	6,774	2,986	8,470
29	12.8964	0.000050	6,774	2,986	8,470
30	12.8964	0.000364	6,774	2,986	8,470
31	12.8964	0.000090	6,774	2,986	8,470
32	20.9858	0.000041	7,180	3,225	8,995
33	12.8964	0.000170	6,774	2,986	8,470
34	12.8964	0.000088	6,774	2,986	8,470
35	12.8964	0.000226	6,774	2,986	8,470
36	12.8964	0.000175	6,774	2,986	8,470
37	12.8964	0.000296	6,774	2,986	8,470
38	12.8964	0.000136	6,774	2,986	8,470
39	12.8964	0.000248	6,774	2,986	8,470
40	12.8964	0.000116	6,774	2,986	8,470
41	12.8964	0.000096	6,774	2,986	8,470
42	12.8964	0.000259	6,774	2,986	8,470
43	12.8964	0.000036	6,774	2,986	8,470
44	12.8964	0.000019	6,774	2,986	8,470
45	12.8964	0.000063	6,774	2,986	8,470
46	12.8964	0.000085	6,774	2,986	8,470
47	12.8964	0.000244	6,774	2,986	8,470
48	12.8964	0.000057	6,774	2,986	8,470
49	12.8964	0.000021	6,774	2,986	8,470
50	12.8964	0.000052	6,774	2,986	8,470
51	12.8964	0.000078	6,774	2,986	8,470
52	12.8964	0.000107	6,774	2,986	8,470
53	12.8964	0.000147	6,774	2,986	8,470
54	12.8964	0.000054	6,774	2,986	8,470
55	12.8964	0.000063	6,774	2,986	8,470

56	12.8964	0.000275	6,774	2,986	8,470
57	12.8964	0.000067	6,774	2,986	8,470
58	12.8964	0.000166	6,774	2,986	8,470
59	12.8964	0.000060	6,774	2,986	8,470
60	12.8964	0.000037	6,774	2,986	8,470
61	12.8964	0.000037	6,774	2,986	8,470
62	12.8964	0.000251	6,774	2,986	8,470
63	12.8964	0.000157	6,774	2,986	8,470
64	12.8964	0.000041	6,774	2,986	8,470
65	12.8964	0.000043	6,774	2,986	8,470
66	12.8964	0.000183	6,774	2,986	8,470
67	12.8964	0.000010	6,774	2,986	8,470
68	12.8964	0.000062	6,774	2,986	8,470
69	12.8964	0.000398	6,774	2,986	8,470
70	12.8964	0.000091	6,774	2,986	8,470
71	12.8964	0.000046	6,774	2,986	8,470
72	12.8964	0.000027	6,774	2,986	8,470
73	12.8964	0.000108	6,774	2,986	8,470
74	12.8964	0.000016	6,774	2,986	8,470
75	12.8964	0.000143	6,774	2,986	8,470
76	12.8964	0.000004	6,774	2,986	8,470
77	12.8964	0.000167	6,774	2,986	8,470
78	12.8964	0.000179	6,774	2,986	8,470
79	12.8964	0.000147	6,774	2,986	8,470
80	12.8964	0.000009	6,774	2,986	8,470
81	12.8964	0.000080	6,774	2,986	8,470
82	12.8964	0.000075	6,774	2,986	8,470
83	12.8964	0.000092	6,774	2,986	8,470
84	12.8964	0.000035	6,774	2,986	8,470
85	12.8964	0.000005	6,774	2,986	8,470
86	12.8964	0.000037	6,774	2,986	8,470
87	12.8964	0.000141	6,774	2,986	8,470
88	12.8964	0.000081	6,774	2,986	8,470
89	12.8964	0.000091	6,774	2,986	8,470
90	12.8964	0.000697	6,774	2,986	8,470
91	12.8964	0.000140	6,774	2,986	8,470
92	12.8964	0.000134	6,774	2,986	8,470
93	12.8964	0.000129	6,774	2,986	8,470
94	12.8964	0.000212	6,774	2,986	8,470
95	12.8964	0.000044	6,774	2,986	8,470
96	12.8964	0.000022	6,774	2,986	8,470
97	12.8964	0.000013	6,774	2,986	8,470
98	12.8964	0.000021	6,774	2,986	8,470

99	12.8964	0.000109	6,774	2,986	8,470
100	12.8964	0.000035	6,774	2,986	8,470

Table D.2 Results from 100 jitter runs for scenario 20_1b for **WAG**. Jitter run 0 corresponds to the original optimized estimates.

Jitter Run	Objective Function	Maximum Gradient	B _{35%} (t)	OFL (t)	Current MMB (t)
0	-75.5594	0.000060	5,319	1,807	6,290
1	-79.6389	0.000115	5,815	1,911	6,641
2	-75.5594	0.000228	5,319	1,807	6,290
3	-75.5594	0.000013	5,319	1,807	6,290
4	-75.5594	0.000048	5,319	1,807	6,290
5	-75.5594	0.000220	5,319	1,807	6,290
6	-75.5594	0.000096	5,319	1,807	6,290
7	-75.5594	0.000040	5,319	1,807	6,290
8	-75.5594	0.000332	5,319	1,807	6,290
9	-75.5594	0.000051	5,319	1,807	6,290
10	-75.5594	0.000144	5,319	1,807	6,290
11	-75.5594	0.000087	5,319	1,807	6,290
12	-75.5594	0.000105	5,319	1,807	6,290
13	-75.5594	0.000085	5,319	1,807	6,290
14	NA	NA	NA	NA	NA
15	-74.3830	0.000516	5,756	1,908	6,583
16	-79.6389	0.000150	5,815	1,911	6,641
17	-75.5594	0.000280	5,319	1,807	6,290
18	-75.5594	0.000088	5,319	1,807	6,290
19	-80.1879	0.000369	5,829	1,902	6,582
20	-75.5594	0.000042	5,319	1,807	6,290
21	-80.1879	0.000046	5,829	1,902	6,582
22	-75.5594	0.000023	5,319	1,807	6,290
23	-75.5594	0.000175	5,319	1,807	6,290
24	-79.6389	0.000163	5,815	1,911	6,641
25	-79.6389	0.000008	5,815	1,911	6,641
26	-75.5594	0.000095	5,319	1,807	6,290
27	-75.5594	0.000033	5,319	1,807	6,290
28	-75.5594	0.000033	5,319	1,807	6,290
29	-75.5594	0.000047	5,319	1,807	6,290
30	-75.5594	0.000103	5,319	1,807	6,290
31	-75.5594	0.000134	5,319	1,807	6,290
32	-75.5594	0.000196	5,319	1,807	6,290
33	-75.5594	0.000051	5,319	1,807	6,290

34	-75.5594	0.000364	5,319	1,807	6,290
35	-75.5594	0.000077	5,319	1,807	6,290
36	-75.5594	0.000119	5,319	1,807	6,290
37	-75.5594	0.000082	5,319	1,807	6,290
38	-75.5594	0.000176	5,319	1,807	6,290
39	-75.5594	0.000099	5,319	1,807	6,290
40	-75.5594	0.000051	5,319	1,807	6,290
41	-75.5594	0.000030	5,319	1,807	6,290
42	-75.5594	0.000235	5,319	1,807	6,290
43	-75.5594	0.000063	5,319	1,807	6,290
44	-75.5594	0.000141	5,319	1,807	6,290
45	-75.5594	0.000102	5,319	1,807	6,290
46	-75.5594	0.000050	5,319	1,807	6,290
47	-80.6251	0.000074	6,107	1,932	6,687
48	-79.6389	0.000407	5,815	1,911	6,641
49	-75.5594	0.000018	5,319	1,807	6,290
50	-75.5594	0.000188	5,319	1,807	6,290
51	-75.5594	0.000205	5,319	1,807	6,290
52	-75.5594	0.000569	5,319	1,807	6,290
53	-75.5594	0.000083	5,319	1,807	6,290
54	-75.5594	0.000137	5,319	1,807	6,290
55	-75.5594	0.000065	5,319	1,807	6,290
56	-75.5594	0.000056	5,319	1,807	6,290
57	-75.5594	0.000131	5,319	1,807	6,290
58	-79.6389	0.000008	5,815	1,911	6,641
59	-75.5594	0.000141	5,319	1,807	6,290
60	-75.5594	0.000159	5,319	1,807	6,290
61	-75.5594	0.000098	5,319	1,807	6,290
62	-75.5594	0.000015	5,319	1,807	6,290
63	-75.5594	0.000129	5,319	1,807	6,290
64	-75.5594	0.000242	5,319	1,807	6,290
65	-75.5594	0.000073	5,319	1,807	6,290
66	-75.5594	0.000022	5,319	1,807	6,290
67	-75.5594	0.000082	5,319	1,807	6,290
68	-75.5594	0.000055	5,319	1,807	6,290
69	-75.5594	0.000105	5,319	1,807	6,290
70	-75.5594	0.000026	5,319	1,807	6,290
71	-80.1879	0.000161	5,829	1,902	6,582
72	-75.5594	0.000076	5,319	1,807	6,290
73	-75.5594	0.000212	5,319	1,807	6,290
74	-75.5594	0.000030	5,319	1,807	6,290
75	-75.5594	0.000214	5,319	1,807	6,290
76	-75.5594	0.000185	5,319	1,807	6,290

77	-75.5594	0.000134	5,319	1,807	6,290
78	-74.2426	0.000012	5,731	1,896	6,564
79	-75.5594	0.000111	5,319	1,807	6,290
80	NA	NA	NA	NA	NA
81	-79.6389	0.000396	5,815	1,911	6,641
82	-75.5594	0.000206	5,319	1,807	6,290
83	-75.5594	0.000406	5,319	1,807	6,290
84	-75.5594	0.000101	5,319	1,807	6,290
85	-75.5594	0.000078	5,319	1,807	6,290
86	-75.5594	0.000156	5,319	1,807	6,290
87	-75.5594	0.000207	5,319	1,807	6,290
88	-75.5594	0.000189	5,319	1,807	6,290
89	-75.5594	0.000088	5,319	1,807	6,290
90	-75.5594	0.000252	5,319	1,807	6,290
91	-75.5594	0.000058	5,319	1,807	6,290
92	-75.5594	0.000174	5,319	1,807	6,290
93	-80.6251	0.000245	6,107	1,932	6,687
94	-75.5594	0.000131	5,319	1,807	6,290
95	-80.1879	0.000158	5,829	1,902	6,582
96	-75.5594	0.000610	5,319	1,807	6,290
97	-75.5594	0.000052	5,319	1,807	6,290
98	-75.5594	0.000107	5,319	1,807	6,290
99	-75.5594	0.000342	5,319	1,807	6,290
100	-74.3830	0.000277	5,756	1,908	6,583

Table D.3. Results from 100 jitter runs for scenario 20_2 for **EAG**. Jitter run 0 corresponds to the original optimized estimates.

Jitter Run	Objective Function	Maximum Gradient	B _{35%} (t)	OFL (t)	Current MMB (t)
0	7.7967	0.001281	6,794	3,133	8,665
1	7.7966	0.000182	6,794	3,133	8,665
2	7.7966	0.000091	6,794	3,133	8,665
3	7.7966	0.000218	6,794	3,133	8,665
4	7.7966	0.000092	6,794	3,133	8,665
5	7.7966	0.000500	6,794	3,133	8,665
6	7.7966	0.000013	6,794	3,133	8,665
7	7.7966	0.000020	6,794	3,133	8,665
8	7.7966	0.000254	6,794	3,133	8,665
9	7.7966	0.000058	6,794	3,133	8,665
10	7.7966	0.000145	6,794	3,133	8,665
11	7.7966	0.000047	6,794	3,133	8,665

12	7.7966	0.000355	6,794	3,133	8,665
13	7.7966	0.000123	6,794	3,133	8,665
14	7.7966	0.000188	6,794	3,133	8,665
15	7.7966	0.000100	6,794	3,133	8,665
16	7.7966	0.000017	6,794	3,133	8,665
17	7.7966	0.000141	6,794	3,133	8,665
18	7.7966	0.000141	6,794	3,133	8,665
19	7.7966	0.000198	6,794	3,133	8,665
20	7.7966	0.000361	6,794	3,133	8,665
21	7.7966	0.000447	6,794	3,133	8,665
22	7.7966	0.000490	6,794	3,133	8,665
23	7.7966	0.000255	6,794	3,133	8,665
24	7.7966	0.000116	6,794	3,133	8,665
25	7.7966	0.000059	6,794	3,133	8,665
26	7.7966	0.000081	6,794	3,133	8,665
27	7.7966	0.000386	6,794	3,133	8,665
28	7.7966	0.000004	6,794	3,133	8,665
29	7.7966	0.000053	6,794	3,133	8,665
30	7.7966	0.000112	6,794	3,133	8,665
31	7.7966	0.000074	6,794	3,133	8,665
32	7.7966	0.000052	6,794	3,133	8,665
33	7.7966	0.000175	6,794	3,133	8,665
34	7.7966	0.000154	6,794	3,133	8,665
35	7.7966	0.000503	6,794	3,133	8,665
36	7.7966	0.000289	6,794	3,133	8,665
37	7.7966	0.000340	6,794	3,133	8,665
38	7.7966	0.000088	6,794	3,133	8,665
39	7.7966	0.000045	6,794	3,133	8,665
40	7.7966	0.000056	6,794	3,133	8,665
41	7.7966	0.000231	6,794	3,133	8,665
42	7.7966	0.000074	6,794	3,133	8,665
43	7.7966	0.000062	6,794	3,133	8,665
44	7.7966	0.000051	6,794	3,133	8,665
45	7.7966	0.000122	6,794	3,133	8,665
46	7.7966	0.000036	6,794	3,133	8,665
47	7.7966	0.000078	6,794	3,133	8,665
48	7.7966	0.000038	6,794	3,133	8,665
49	7.7966	0.000492	6,794	3,133	8,665
50	7.7966	0.000089	6,794	3,133	8,665
51	7.7966	0.000124	6,794	3,133	8,665
52	7.7966	0.000031	6,794	3,133	8,665
53	7.7966	0.000035	6,794	3,133	8,665
54	7.7966	0.000275	6,794	3,133	8,665

55	7.7966	0.000196	6,794	3,133	8,665
56	7.7966	0.000208	6,794	3,133	8,665
57	7.7966	0.000014	6,794	3,133	8,665
58	7.7966	0.000140	6,794	3,133	8,665
59	7.7966	0.000618	6,794	3,133	8,665
60	7.7966	0.000026	6,794	3,133	8,665
61	7.7966	0.000088	6,794	3,133	8,665
62	7.7966	0.000142	6,794	3,133	8,665
63	7.7966	0.000488	6,794	3,133	8,665
64	7.7966	0.000160	6,794	3,133	8,665
65	7.7966	0.000021	6,794	3,133	8,665
66	7.7966	0.000228	6,794	3,133	8,665
67	7.7966	0.000026	6,794	3,133	8,665
68	7.7966	0.000070	6,794	3,133	8,665
69	7.7966	0.000147	6,794	3,133	8,665
70	7.7966	0.000287	6,794	3,133	8,665
71	7.7966	0.000172	6,794	3,133	8,665
72	7.7966	0.000353	6,794	3,133	8,665
73	7.7966	0.000126	6,794	3,133	8,665
74	7.7966	0.000251	6,794	3,133	8,665
75	7.7966	0.000253	6,794	3,133	8,665
76	7.7966	0.000075	6,794	3,133	8,665
77	7.7966	0.000064	6,794	3,133	8,665
78	7.7966	0.000091	6,794	3,133	8,665
79	7.7966	0.000431	6,794	3,133	8,665
80	7.7966	0.000222	6,794	3,133	8,665
81	7.7966	0.000131	6,794	3,133	8,665
82	7.7966	0.000044	6,794	3,133	8,665
83	7.7966	0.000307	6,794	3,133	8,665
84	7.7966	0.000240	6,794	3,133	8,665
85	7.7966	0.000102	6,794	3,133	8,665
86	7.7966	0.000100	6,794	3,133	8,665
87	7.7966	0.000175	6,794	3,133	8,665
88	7.7966	0.000295	6,794	3,133	8,665
89	7.7966	0.000150	6,794	3,133	8,665
90	7.7966	0.000034	6,794	3,133	8,665
91	7.7966	0.000081	6,794	3,133	8,665
92	7.7966	0.000252	6,794	3,133	8,665
93	7.7966	0.000089	6,794	3,133	8,665
94	7.7966	0.000043	6,794	3,133	8,665
95	7.7966	0.000131	6,794	3,133	8,665
96	7.7966	0.000137	6,794	3,133	8,665
97	7.7966	0.000232	6,794	3,133	8,665

98	7.7966	0.000018	6,794	3,133	8,665
99	7.7966	0.000041	6,794	3,133	8,665
100	7.7966	0.000015	6,794	3,133	8,665

Table D.4 Results from 100 jitter runs for scenario 20_2 for **WAG**. Jitter run 0 corresponds to the original optimized estimates.

Jitter Run	Objective Function	Maximum Gradient	B _{35%} (t)	OFL (t)	Current MMB (t)
0	-75.7768	0.000171	5,343	1,860	6,441
1	-75.7768	0.000073	5,343	1,860	6,441
2	-75.7768	0.000131	5,343	1,860	6,441
3	-75.7768	0.000048	5,343	1,860	6,441
4	-75.7768	0.000052	5,343	1,860	6,441
5	-79.5165	0.000122	5,869	1,960	6,750
6	-75.7768	0.000375	5,343	1,860	6,441
7	-75.7768	0.000126	5,343	1,860	6,441
8	-75.7768	0.000262	5,343	1,860	6,441
9	-75.7768	0.000084	5,343	1,860	6,441
10	-75.7768	0.000134	5,343	1,860	6,441
11	-75.7768	0.000099	5,343	1,860	6,441
12	-75.7768	0.000227	5,343	1,860	6,441
13	-75.7768	0.000240	5,343	1,860	6,441
14	-75.7768	0.000447	5,343	1,860	6,441
15	-75.7768	0.000158	5,343	1,860	6,441
16	-75.7768	0.000018	5,343	1,860	6,441
17	-75.7768	0.000124	5,343	1,860	6,441
18	-79.5165	0.000134	5,869	1,960	6,750
19	-74.0867	0.000154	5,769	1,947	6,722
20	-75.7768	0.000029	5,343	1,860	6,441
21	-75.7768	0.000010	5,343	1,860	6,441
22	-75.7768	0.000387	5,343	1,860	6,441
23	-75.7768	0.000218	5,343	1,860	6,441
24	-75.7768	0.000004	5,343	1,860	6,441
25	-75.7768	0.000158	5,343	1,860	6,441
26	-75.7768	0.000566	5,343	1,860	6,441
27	-75.7768	0.000050	5,343	1,860	6,441
28	-75.7768	0.000042	5,343	1,860	6,441
29	-75.7768	0.000084	5,343	1,860	6,441
30	-74.0867	0.000038	5,769	1,947	6,722
31	-75.7768	0.000010	5,343	1,860	6,441
32	-75.7768	0.000093	5,343	1,860	6,441

33	-75.7768	0.000116	5,343	1,860	6,441
34	-75.7768	0.000037	5,343	1,860	6,441
35	-75.7768	0.000126	5,343	1,860	6,441
36	-75.7768	0.000079	5,343	1,860	6,441
37	-75.7768	0.000473	5,343	1,860	6,441
38	-75.7768	0.000459	5,343	1,860	6,441
39	-75.7768	0.000122	5,343	1,860	6,441
40	-75.7768	0.000020	5,343	1,860	6,441
41	-75.7768	0.000124	5,343	1,860	6,441
42	-74.0867	0.000081	5,769	1,947	6,722
43	-75.7768	0.000153	5,343	1,860	6,441
44	-75.7768	0.000287	5,343	1,860	6,441
45	-75.7768	0.000651	5,343	1,860	6,441
46	-75.7768	0.000007	5,343	1,860	6,441
47	-75.7768	0.000247	5,343	1,860	6,441
48	-75.7768	0.000093	5,343	1,860	6,441
49	-75.7768	0.000243	5,343	1,860	6,441
50	-75.7768	0.000183	5,343	1,860	6,441
51	-75.7768	0.000168	5,343	1,860	6,441
52	-75.7768	0.000131	5,343	1,860	6,441
53	-75.7768	0.000080	5,343	1,860	6,441
54	-75.7768	0.000042	5,343	1,860	6,441
55	-75.7768	0.000153	5,343	1,860	6,441
56	-75.7768	0.000297	5,343	1,860	6,441
57	-75.7768	0.000080	5,343	1,860	6,441
58	-75.7768	0.000051	5,343	1,860	6,441
59	-75.7768	0.000013	5,343	1,860	6,441
60	-75.7768	0.000077	5,343	1,860	6,441
61	-75.7768	0.000029	5,343	1,860	6,441
62	-75.7768	0.000050	5,343	1,860	6,441
63	-79.5165	0.000169	5,869	1,960	6,750
64	-75.7768	0.000058	5,343	1,860	6,441
65	-79.0546	0.000104	5,848	1,969	6,810
66	-75.7768	0.000048	5,343	1,860	6,441
67	-75.7768	0.000021	5,343	1,860	6,441
68	-75.7768	0.000060	5,343	1,860	6,441
69	-75.7768	0.000040	5,343	1,860	6,441
70	-75.7768	0.000063	5,343	1,860	6,441
71	-75.7768	0.000527	5,343	1,860	6,441
72	-75.7768	0.000149	5,343	1,860	6,441
73	-75.7768	0.000291	5,343	1,860	6,441
74	-75.7768	0.000058	5,343	1,860	6,441
75	-75.7768	0.000077	5,343	1,860	6,441

76	-75.7768	0.000045	5,343	1,860	6,441
77	-75.7768	0.000059	5,343	1,860	6,441
78	-75.7768	0.000016	5,343	1,860	6,441
79	-75.7768	0.000107	5,343	1,860	6,441
80	-75.7768	0.000178	5,343	1,860	6,441
81	-75.7768	0.000459	5,343	1,860	6,441
82	-75.7768	0.000148	5,343	1,860	6,441
83	-75.7768	0.000505	5,343	1,860	6,441
84	-75.7768	0.000115	5,343	1,860	6,441
85	-75.7768	0.000315	5,343	1,860	6,441
86	-79.5165	0.000168	5,869	1,960	6,750
87	-79.0546	0.000066	5,848	1,969	6,810
88	-75.7768	0.000018	5,343	1,860	6,441
89	-75.7768	0.000086	5,343	1,860	6,441
90	-75.7768	0.000123	5,343	1,860	6,441
91	-75.7768	0.000034	5,343	1,860	6,441
92	-75.7768	0.000392	5,343	1,860	6,441
93	-75.7768	0.000543	5,343	1,860	6,441
94	-75.7768	0.000036	5,343	1,860	6,441
95	-75.7768	0.000102	5,343	1,860	6,441
96	-75.7768	0.000085	5,343	1,860	6,441
97	NA	NA	NA	NA	NA
98	-75.7768	0.000140	5,343	1,860	6,441
99	-75.7768	0.000038	5,343	1,860	6,441
100	-75.7768	0.000357	5,343	1,860	6,441
