Gulf of Alaska 2018

Ecosystem Status Report


Subsurface Temperature Anomaly (°C) 0-100m Depth October 2018 NCEP GODAS 1981-2010 Climatology 60N 50N 30N 1 20N 10N 10S 205 140E 160E 180 160W 140W 120W 100W 80W -2 -1.5-0.50.5 1.5 -.32

Stephani Zador and Ellen Yasumiishi Alaska Fisheries Science Center GOA Groundfish Plan Team November 13, 2018

Complete Recap of 2017

Complete recap of 2017

- Temperatures moderated relative to heatwave years 2014–2016; Blob dissipated
- 2. Reduced transport in the Alaska Current and low eddy activity
- 3. Abundant, but small, zooplankton taxa and few euphausiids




2017 Zooplankton

Batten, Hopcroft, Ressler, Fergusson

- Continuous Plankton Recorder
- Copepods small during heatwave
- Switched to larger in 2017
- Mesozooplankton biomass high
- Seward Line copepods abundant, but euphausiids low
- Euphausiids also low in acoustic survey
- Lipid content low Icy Strait

2017 Zooplankton


Campbell et al



Complete recap of 2017

- Temperatures moderated relative to heatwave years 2014–2016; Blob dissipated
- 2. Reduced transport into the Alaska Current and low eddy activity
- 3. Abundant, but small, zooplankton taxa and few euphausiids
- 4. Few cod larvae, but above-average abundance of pollock larvae. Wide distribution, unlike in 2013. Late summer age-0 abundant around Shelikof and Kodiak.


2017 Larval and age-0 groundfish




Larvae

EcoFOCI, EMA, RECA

- Mostly average abundance during larval surveys except few cod
- Positive signs for 2017 pollock
 - Above average larval densities
 - Widely distributed age-0s
 - Energy density high EGOA
 - Seen again as age-1s during winter Shelikof surveys






Complete recap of 2017

- Temperatures moderated relative to heatwave years 2014–2016; Blob dissipated
- 2. Reduced transport into the Alaska Current and low eddy activity
- 3. Abundant, but small, zooplankton taxa and few euphausiids
- 4. Few cod larvae, but above-average abundance of pollock larvae. Wide distribution, unlike in 2013. Late summer age-0 abundant around Shelikof and Kodiak.
- All groundfish but cod had poor body condition. Record low for arrowtooth (apex) and northern rockfish (pelagic)
- 6. Salmon marine survival low
- 7. Seabirds and marine mammals poor reproduction overall
- 8. Seabird bycatch up but other non-target catch low (jellyfish, inverts)


Current Conditions Western GOA

West of Cape Suckling 144°W


2018 climate




Mid-Oct IRI/CPC Model-Based Probabilistic ENSO Forecasts




• PDO remains neutral



Another heatwave!



GOA Region 📕 WGOA 📕 EGOA 📒 SEAK Inside



59.5°N

58.5°N

58°N

2000

2005

2010

Latitude N°65

May temperation of the second second

2015

2018 oceanography


Watson, Danielson and Hopcroft

Satellite-derived sea surface temperature


May temperature averaged over top 100m depth at the Seward Line

2018 zooplankton

Hopcroft and Coyle




- Seward Line sampling during May
- 0.5 mm mesh
- Averaged over top 100m
- Copepods remained abundant in 2018
- Euphausiids increased in 2018





2018 forage fish

Hatch et al, Pegau et al


- Sampled from seabirds provisioning chicks at MIddleton Island
- Time series also available for herring, salmon
- PWS herring mile-days of milt continued decline to lowest on record









Standarized Anomoly Value





2018 groundfish Worton

- Catches remain below the long term mean for ATF, pollock (inshore), flathead sole, cod, skates, halibut
- Overall catch increased from 2017 due to pollock and ATF



Renner et al

FISH & WILDLIFE SERVICE	AL EXPENSION OF MARKEN	Alaska Maritime National Wildlife Refuge												
		2018 9	Seabird	l Repo	rt Card									
	SYSTEM	1	1	S.	7	R				6.		X	S CAR	
Region	Annual monitoring site	Red-faced cormorants	Glaucous- winged gulls	Common murres	Thick- billed murres	Horned puffins	Tufted puffins	Red- legged kittiwakes	Black- legged kittiwakes	Northern fulmar	Fork-tailed storm- petrels	Leach's storm- petrels	Parakeet auklets	Least auklets
Chukchi Sea	Cape Lisburne			(XXX) XXX XXX	The second secon									
Bering Sea	St. Paul			\odot				TT X X		The second secon				
Berin	St. George	$\overline{}$		\odot				TA X X		The second secon				
Aleutian Islands	Buldir		$\overline{}$	The second secon		$\overline{}$	JS X X	$\overline{}$	$\overline{}$					\odot
Aleutiar	Aiktak			(The second seco	The second secon									
Alaska Penin.	Chowiet	\bigcirc	$\overline{}$											
Gulf of Alaska	East Amatuli						\bigcirc				\bigcirc			
Gulf of	St. Lazaria											\bigcirc		
\bigcirc		<pre></pre>	ily fish eater	rs 🗧		803	()		Partie Partie	~		Primarily zoo	oplankton ea	ters >>>
White eg	Eggs represent overall productivity relative to the long-term average. White eggs indicate productivity derived from monitoring data; colored eggs indicate productivity based on anecdotal observations.													

Few murres at cliffs, but those there did well reproductively

In general, piscivorous birds didn't do well during the heatwave, but planktivorous birds did (except Chowiet this year)

Renner et al

FISH & WILDLIFE SERVICE	A COMPANY OF MARKEN	Alaska Maritime National Wildlife Refuge												
		2018 9	Seabird	l Repo	rt Card									
	SYSTEM	1	2	S.	7	R		R		6		X	S CAR	
Region	Annual monitoring site	Red-faced cormorants	Glaucous- winged gulls	Common murres	Thick- billed murres	Horned puffins	Tufted puffins	Red- legged kittiwakes	Black- legged kittiwakes	Northern fulmar	Fork-tailed storm- petrels	Leach's storm- petrels	Parakeet auklets	Least auklets
Chukchi Sea	Cape Lisburne			(J' X X Y)	The second				\bigcirc					
Bering Sea	St. Paul			\odot				TT X X X		The second secon				
Berin	St. George	$\overline{}$		\odot				TT X X		Ly x x x				
Aleutian Islands	Buldir		$\overline{}$	JS X X		$\overline{}$	JS X X	$\overline{}$	$\overline{}$		$\overline{}$			\odot
Aleutiar	Aiktak			The second secon	The second secon	(\mathbf{i})	(ô ô)							
Alaska Penin.	Chowiet	\bigcirc	$\overline{}$											
Gulf of Alaska	East Amatuli						$\overline{\odot}$				\bigcirc			
Gulf of	St. Lazaria											\bigcirc		
\bigcirc		<pre></pre>	rily fish eater	rs ←	+		()		Prese			Primarily zoc	plankton ea	ters >>>
White eg	Eggs represent overall productivity relative to the long-term average. White eggs indicate productivity derived from monitoring data; colored eggs indicate productivity based on anecdotal observations.													

Few murres at cliffs, but those there did well reproductively

In general, piscivorous birds didn't do well during the heatwave, but planktivorous birds did (except Chowiet this year)

Renner et al

FISH & WILDLIFE SERVICE	Attentions of whether I have near hour Wattonal Wattonal SEFTER	Alaska Maritime National Wildlife Refuge												
		2018 9	Seabird	l Repo	rt Card									
	<u>SYSTEM</u>	1	1	54	2	R		R		6		X	S CAR	
Region	Annual monitoring site	Red-faced cormorants	Glaucous- winged gulls	Common murres	Thick- billed murres	Horned puffins	Tufted puffins	Red- legged kittiwakes	Black- legged kittiwakes	Northern fulmar	Fork-tailed storm- petrels	Leach's storm- petrels	Parakeet auklets	Least auklets
Chukchi Sea	Cape Lisburne			TAS X X	The xx				\bigcirc					
Bering Sea	St. Paul			$\overline{}$				The second secon		The second secon				
Berin	St. George	$\overline{}$		\odot				The second secon		The second secon				
Aleutian Islands	Buldir		$\overline{}$	JS X X		$\overline{}$	JS X X	$\overline{}$	$\overline{}$		$\overline{}$			\odot
Aleutiar	Aiktak			The second secon	The second secon									
Alaska Penin.	Chowiet	\bigcirc	$\overline{}$											
Gulf of Alaska	East Amatuli						\bigcirc				\odot			
Gulf of	St. Lazaria											\bigcirc		
\bigcirc		<pre>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>></pre>	ily fish eater	rs ←		30	()		Partie .			Primarily zoo	oplankton ea	ters >>>
White e	Eggs represent overall productivity relative to the long-term average. White eggs indicate productivity derived from monitoring data; colored eggs indicate productivity based on anecdotal observations. Way above average!													

Few murres at cliffs, but those there did well reproductively

In general, piscivorous birds didn't do well during the heatwave, but planktivorous birds did (except Chowiet this year)

2018 humpback whales

Moran and Straley

Fall survey of humpback whales in Prince William Sound

Month/year	Counts of whales	Nautical miles surveyed	Encounter rate Whale/NM			
Sep 2007*	24	370	0.06			
Sep 2008	71	412	0.17			
Oct 2011	62	441	0.14			
Sep 2012	81	444	0.18			
Sep 2013	113	355	0.32			
Sep 2014	181	427	0.42			
Sep 2017	12	543	0.02			
Sep 2018	17	541	0.03			


* didn't survey hotspot at Montague Entrance

- Some with poor body condition in 2018 but lesser extent than in 2016-2017
- PWS whale decline mirrors herring decline



Current Conditions Eastern GOA


GOA Region 📕 WGOA 📕 EGOA 📒 SEAK



2018 oceanography Watson, Fergusson


- Satellite derived data indicate cooler winter & spring sea temperatures in the EGOA.
- Icy Strait 20-m integrated SST (May-Aug.) was average (9.2 °C) for the 22 year time series.

Icy Strait 1997-2018



2018 zooplankton Fergusson, Fugate


- Increase in zooplankton densities, but
- Decrease in large copepods
- Increase in lipid content



- Auke Creek weir, 1980–2018
- Low marine survival of coho (piscivores) and pink (planktivores) salmon, 2017 & 2018 out migrants
- Record low returns of adult pink salmon in 2018


2018 salmon

Vulstek, Russell, Yasumiishi



Smolt Year

- lcy Strait, 1997–2018
- Low CPUE of juvenile salmon sampled at sea
- Small juveniles in length and weight



1997

1999

2001 2003

2005

2007

2009 2011 2013


2015 2017

2018 salmon

Watson, Fergusson

CPUE


Renner et al



Piscivorous and planktivorous seabirds did well at St Lazaria (Sitka)

- Low crude birth rate (1/99)
- Low calf #
- Improved adult body condition

Humpback whales in Glacier Bay, northern southeast Alaska



Year

2018 marine mammals


Gabriele, Neilson, Moran

Predictions and Forecasts

2019 Sea Surface Temperature Forecasts Bond

Feb-Apr 2019

NMME Forecast of SST Anom IC=201811 for Lead 3 2019FMA



- Aleutian low deeper than normal in late winter of 2018/19 = warm weather for Alaska enhanced by warm waters.
- No typical PDO pattern in N Pacific in early 2019 (warm everywhere).

Summary

2018

- The GOA is currently in a heatwave (21 Oct 2018)
- Warm SST forecasted across North Pacific (Feb-Mar-Apr 2019)
- In general, during the 2014–2016 heatwave, copepods were small but abundant, but forage fish were absent
- In 2018, mixed zooplankton signals in the western GOA, but good forage fish signals
- Seabirds and marine mammals continue to show lagged negative effects of the heatwave


Contributors

Mayumi Arimitsu, Steve Barbeaux, Sonia Batten, James Bodkin, Nick Bond, Rob Campbell, Heather Coletti, Ken Coyle, Seth Danielson, Thomas Dean, Alison Deary, Sherri Dressel, Brie Drummond, Anne Marie Eich, Emily Fergusson, Benjamin Fissel, Shannon Fitzgerald, Corey Fugate, Chris Gabriele, Tom Gelatt, Bradley P. Harris, Scott Hatch, Stormy Haught, Kyle Hebert, Ron Heintz, Paul K. Hershberger, Russ Hopcroft, Katrin Iken, Tayler Jarvis, Stephen Kasperski, Arthur Kettle, Brenda Konar, Carol Ladd, Geoffrey Lang, Anna Lavoie, Jean Lee, Daniel Lew, Mandy Lindeberg, Caitlin McKinstry, Kathryn Miers, Jennifer Mondragon, Daniel Monson, John Moran, Franz Mueter, Janet Neilson, John Olson, W. Scott Pegau, John Piatt, Heather Renner, Lauren Rogers, Nora Rojek, Joshua R. Russell, Sioned E. Sitkiewicz, T. Scott Smeltz, Kim Sparks, William Stockhausen, Janice Straley, Rob Suryan, Kathryn Sweeney, Marysia Szymkowiak, John Trochta, Scott Vulstek, Jordan Watson, Sarah R. Webster, George A. Whitehouse, Sarah Wise, Nathan Wolf, Carrie Worton, Ellen Yasumiishi, and Stephani Zador

Thank you!

2018 salmon

Fergusson, Watson



Sablefish Matrix? score 2

Environmental/ecosystem considerations

a. Adverse trends in environmental/ecosystem indicators


b. Ecosystem model results

c. Decreases in ecosystem productivity d. Decreases in prey abundance or availability

e. Increases in predator abundance

2017 Steller sea lions


Sweeney and Gelatt



• Stabilized in WGOA

• Oscillating near carrying capacity in EGOA?

GULF NP



SEAK NP Region

