

# Saint Matthew Island Blue King Crab DRAFT Stock Assessment

## May 2022

Katie Palof<sup>1</sup>, Jie Zheng<sup>1</sup>, Jim Ianelli<sup>2</sup>

<sup>1</sup>Alaska Department of Fish and Game, katie.palof@alaska.gov and jie.zheng@alaska.gov

<sup>2</sup>NOAA, jim.ianelli@noaa.gov

May 2022

## Executive Summary

1. **Stock:** Blue king crab, *Paralithodes platypus*, Saint Matthew Island (SMBKC), Alaska.
2. **Catches:** Peak historical harvest was 4,288 t (9.454 million pounds) in 1983/84<sup>1</sup>. The fishery was closed for 10 years after the stock was declared overfished in 1999. Fishing resumed in 2009/10 with a fishery-reported retained catch of 209 t (0.461 million pounds), less than half the 529.3 t (1.167 million pound) TAC. Following three more years of modest harvests supported by a fishery catch per unit effort (CPUE) of around 10 crab per pot lift, the fishery was again closed in 2013/14 due to declining trawl-survey estimates of abundance and concerns about the health of the stock. The directed fishery resumed again in 2014/15 with a TAC of 300 t (0.655 million pounds), but the fishery performance was relatively poor with a retained catch of 140 t (0.309 million pounds). The retained catch in 2015/16 was even lower at 48 t (0.105 million pounds) and the fishery has remained closed since 2016/17.
3. **Stock biomass:** The 1978-2021 NMFS trawl survey mean biomass is 5,605 t with the 2021 value being the 9th lowest (1,929 t; the seventh lowest since 2000). This 2021 biomass of  $\geq 90$  mm carapace length (CL) male crab is 35% of the long term mean at 4.25 million pounds (with a CV of 43%), and an 39% decrease from the 2019 biomass (due to the cancellation of the 2020 survey estimates are compared to 2019 as the last survey year). The most recent 3-year average of the NMFS survey is 41% of the mean value, indicating a decline in biomass compared to historical survey estimates, notably in 2010 and 2011 that were over four times the current average. Additionally, the 2021 value decreased from the 2019 value, falling closer to the two previous years (1,929 t compared to 1,731 t in 2018 and 1,794 t in 2017). The ADFG pot survey last occurred in 2018, when the relative biomass index was the lowest in the time series (12% of the mean from the 11 surveys conducted since 1995). The assessment model estimates temper this increase and suggest that the stock (in survey biomass units) is presently at about 32% of the long term model-predicted survey biomass average, similar to the last three years. The trend from these values suggests a steady state in the last few years, which does not fit the 2019 observed survey data point well.
4. **Recruitment:** Recruitment is based on estimated number of male crab within the 90-104 mm CL size class in each year. The 2021 trawl-survey area-swept estimate of 0.423 million male SMBKC in this size class is the 15th lowest in the 43 years since 1978 and follows two of the lowest previously observed values in 2017 and 2018. The recent six-year (2015 - 2021) average recruitment is only 43% of the long-term mean. In the pot-survey, the abundance of this size group in 2017 was also the second-lowest in the time series (22% of the mean for the available pot-survey data) whereas in 2018 the value was the lowest observed at only 10% of the mean value.

---

<sup>1</sup>1983/84 refers to a fishing year that extends from 1 July 1983 to 30 June 1984.

5. **Management performance:** In this assessment, estimated total male catch is the sum of fishery-reported retained catch, estimated male discard mortality in the directed fishery, and estimated male bycatch mortality in the groundfish fisheries. Based on the reference model for SMBKC, the estimate for mature male biomass was below the minimum stock-size threshold (MSST) in 2018/19 and is in an “overfished” condition, despite a directed fishery closure since the 2016/17 season (and hence overfishing has not occurred) (Tables 1, 3, and 4). Computations which indicate the relative impact of fishing (i.e., the “dynamic  $B_0$ ”) suggests, that the current spawning stock biomass has been reduced to 61% of what it would have been in the absence of fishing, assuming the same level of recruitment as estimated.

Table 1: Status and catch specifications (1000 t) for the base model.

Year	Biomass		TAC	Retained	Total	OFL	ABC
	MSST	( $MMB_{\text{mat}})$		catch	male catch		
2016/17	1.97	2.23	0.00	0.00	0.001	0.14	0.11
2017/18	1.85	2.05	0.00	0.00	0.003	0.12	0.10
2018/19	1.74	1.15	0.00	0.00	0.001	0.04	0.03
2019/20	1.67	1.06	0.00	0.00	0.001	0.04	0.03
2020/21	1.65	1.14	0.00	0.00	0.001	0.05	0.04
2021/22		1.12				0.05	0.04

Table 2: Status and catch specifications (million pounds) for the base model.

Year	Biomass		TAC	Retained	Total	OFL	ABC
	MSST	( $MMB_{\text{mat}})$		catch	male catch		
2016/17	4.3	4.91	0.000	0.000	0.002	0.31	0.25
2017/18	4.1	2.85	0.000	0.000	0.007	0.27	0.22
2018/19	3.84	2.54	0.000	0.000	0.002	0.08	0.07
2019/20	3.68	2.34	0.000	0.000	0.002	0.096	0.08
2020/21	3.64	2.52	0.000	0.000	0.002	0.112	0.08
2021/22		2.46				0.109	0.08

6. **Basis for the OFL:** Estimated mature-male biomass (MMB) on 15 February is used as the measure of biomass for this Tier 4 stock, with males measuring  $\geq 105$  mm CL considered mature. The  $B_{MSY}$  proxy is obtained by averaging estimated MMB over a specific reference period, and current CPT/SSC guidance recommends using the full assessment time frame (1978 - 2019) as the default reference period.

Table 3: Basis for the OFL (1000 t) from the base model.

Year	Tier	Biomass					Basis for $B_{MSY}$	Natural mortality
		$B_{MSY}$	( $MMB_{\text{mat}})$	$B/B_{MSY}$	$F_{OFL}$	$\gamma$		
2016/17	4b	3.67	2.23	0.61	0.09	1	1978-2016	0.18
2017/18	4b	3.86	2.05	0.53	0.08	1	1978-2017	0.18
2018/19	4b	3.7	1.15	0.35	0.043	1	1978-2017	0.18
2019/20	4b	3.48	1.06	0.31	0.042	1	1978-2018	0.18
2020/21	4b	3.34	1.14	0.32	0.047	1	1978-2019	0.18
2021/22	4b	3.3	1.12	0.34	0.048	1	1978-2019	0.18

## A. Summary of Major Changes

### Changes in Management of the Fishery

There are no new changes in management of the fishery.

### Changes to the Input Data

Data used in this assessment have been updated to include the most recently available fishery data. This assessment includes one new survey data points from the 2021 NMFS trawl-survey, which is included here since this assessment is now on a biennial cycle. The triennial ADF&G pot surveys were last conducted in 2018, and are back on a semi-triennial cycle, with the next survey planned for 2022. There is new size composition data from the trawl survey. The assessment was updated with 2010-2020 groundfish trawl and fixed gear bycatch estimates based on NMFS Alaska Regional Office (AKRO) data. The directed fishery has been closed since 2016/17, so no recent fishery data are available.

### Changes in Assessment Methodology

This assessment uses the General Model for Alaska Crab Stocks (GMACS) framework. The model is configured to track three stages of length categories and was first presented in May 2011 by W.Gaeuman, ADF&G, and accepted by the CPT in May 2012. A difference from the original approach and that used here is that natural and fishing mortalities are continuous within 5 discrete time blocks within a year (using the appropriate catch equation rather than assuming an applied pulse removal). The time blocks within a year in GMACS are controlled by changing the proportion of natural mortality that is applied each block. Diagnostic output includes estimates of the “dynamic  $B_0$ ” which simply computes the ratio of the estimated spawning biomass relative to the spawning biomass that would have occurred had there been no historical fishing mortality. Details of this implementation and other model details are provided in Appendix A.

### Changes in Assessment Results

Both surveys indicate a decline over the past few years. The “reference” model is that which was selected for use in 2020, the year of the last full assessment. The base model presented here is the reference model with updated groundfish bycatch data for the 2020/21 crab season and updated survey data - biomass and size composition - from the 2021 NMFS trawl survey (model 16.0 2022).

In addition to the reference/base model other models are presented here for sensitivity of life history constants. Here two models are presented that raise the natural mortality from the accepted fixed value of 0.18 for king crab. Model 22.0a has a  $M = 0.21$  and Model 22.0b has a  $M = 0.26$ . These values were taken from methods of Then et al. (2015) using a maximum age of 25 or 30 years. Assuming a maximum age of less than 25 years produced a high estimate of natural mortality using Then et al. methods.

## B. Responses to SSC and CPT

### CPT and SSC Comments on Assessments in General

*Comment: Regarding general code development, the SSC and CPT outstanding requests continue to be as follows:*

1. *Continued exploration of data weighting (Francis and other approaches) and evaluation of models with and without the 1998 natural mortality spike. The authors are encouraged to bring other models forward for CPT and SSC consideration*

We continued with the iterative re-weighting for composition data (Table 17). We did not address models without the natural mortality spike. These have been considered previously.

2. *Retrospective analyses*

These were provided in the Sept. 2020 SAFE document and will be provided again in the Sept. 2022 final SAFE.

Comment: *Explore potential explanations for the discrepancy in the time trends of the two types of survey data, including movement hypotheses using spatial models (not necessarily VAST)*

Exploration of the spatial extent and density differences between the surveys was done on all male crab included in the model (Appendix C). The authors plan to use this and further analyses to better characterize catchability/availability for the pot survey.

Comment: *Explore VAST estimates compared to design based, and ones that combine the two surveys*

Progress is underway to refine the SMBKC VAST estimates using preliminary code that incorporates the island effect. Jon Richar (NMFS) is working on these estimates. At the time of this final SAFE there are no additional improvements to this data set and therefore the VAST model is not presented as a model option. Future work on VAST models for this stock includes VAST data output for the NMFS trawl survey incorporating the island effect and VAST output using both survey data sets together.

Comment: *Random walk or exploration of catchability*

The initial model of time blocks for Q did not show much potential for this in May 2020, therefore time blocks were not a focus for May 2022. More coding work is needed to make a true random walk for catchability in GMACS and this will be added to model development.

Comment: *Consider increasing the number of size bins so that cohorts might be more easily tracked and growth better estimated*

A full review of the research and literature for blue king crab is underway but no changes to the assessment model size bins were considered in this review due to the lack of a concrete basis for these changes and concern over sample size reductions with increased size bins.

Comment: *Explore the assumed and estimated life history parameters (e.g., natural mortality, growth, and maturity) to ensure the best available science is being used to assess this stock.*

Specific research on St. Matthew blue king crab life history parameters is not available and therefore these are borrowed from other stocks/species. At this time only sensitivities of the model to increased natural mortality (M) were looked at here (Models 22.0a and 22.0b). Sensitivities to the model assumptions on growth and maturity will be explored at a later date.

## C. Introduction

### Scientific Name

The blue king crab is a lithodid crab, *Paralithodes platypus* (Brant 1850).

### Distribution

Blue king crab are sporadically distributed throughout the North Pacific Ocean from Hokkaido, Japan, to southeastern Alaska (Figure 1). In the eastern Bering Sea small populations are distributed around

St. Matthew Island, the Pribilof Islands, St. Lawrence Island, and Nunivak Island. Isolated populations also exist in some other cold water areas of the Gulf of Alaska (NPFMC 1998). The St. Matthew Island Section for blue king crab is within Area Q2 (Figure 2), which is the Northern District of the Bering Sea king crab registration area and includes the waters north of Cape Newenham (58°39' N. lat.) and south of Cape Romanzof (61°49' N. lat.).

## Stock Structure

The Alaska Department of Fish and Game (ADF&G) Gene Conservation Laboratory, has detected regional population differences between blue king crab collected from St. Matthew Island and the Pribilof Islands<sup>2</sup>. The NMFS tag-return data from studies on blue king crab in the Pribilof Islands and St. Matthew Island support the idea that legal-sized males do not migrate between the two areas (Otto and Cummiskey 1990). St. Matthew Island blue king crab tend to be smaller than their Pribilof conspecifics, and the two stocks are managed separately.

## Life History

Like the red king crab, *Paralithodes camtschaticus*, the blue king crab is considered a shallow water species by comparison with other lithodids such as golden king crab, *Lithodes aequispinus*, and the scarlet king crab, *Lithodes couesi* (Donaldson and Byersdorfer 2005). Adult male blue king crab are found at an average depth of 70 m (NPFMC 1998). The reproductive cycle appears to be annual for the first two reproductive cycles and biennial thereafter (Jensen and Armstrong 1989), and mature crab seasonally migrate inshore where they molt and mate. Unlike red king crab, juvenile blue king crab do not form pods, but instead rely on cryptic coloration for protection from predators and require suitable habitat such as cobble and shell hash. Somerton and MacIntosh (1983) estimated SMBKC male size at sexual maturity to be 77 mm carapace length (CL). Paul et al. (1991) found that spermatophores were present in the vas deferens of 50% of the St. Matthew Island blue king crab males examined with sizes of 40-49 mm CL and in 100% of the males at least 100 mm CL. Spermatophore diameter also increased with increasing CL with an asymptote at ~ 100 mm CL. It was noted, however, that although spermatophore presence indicates physiological sexual maturity, it may not be an indicator of functional sexual maturity. For purposes of management of the St. Matthew Island blue king crab fishery, the State of Alaska uses 105 mm CL to define the lower size bound of functionally mature males (Pengilly and Schmidt 1995). Otto and Cummiskey (1990) report an average growth increment of 14.1 mm CL for adult SMBKC males.

## Management History

The SMBKC fishery developed subsequent to baseline ecological studies associated with oil exploration (Otto 1990). Ten U.S. vessels harvested 545 t (1.202 million pounds) in 1977, and harvests peaked in 1983 when 164 vessels landed 4,288 t (9.454 million pounds) (Fitch et al. 2012; Table 7).

The fishing seasons were generally short, often lasting only a few days. The fishery was declared overfished and closed in 1999 when the stock biomass estimate was below the minimum stock-size threshold (MSST) of 4,990 t (11.0 million pounds) as defined by the Fishery Management Plan (FMP) for the Bering Sea/Aleutian Islands King and Tanner crabs (NPFMC 1999). Zheng and Kruse (2002) hypothesized a high level of SMBKC natural mortality from 1998 to 1999 as an explanation for the low catch per unit effort (CPUE) in the 1998/99 commercial fishery and the low numbers across all male crab size groups caught in the annual NMFS eastern Bering Sea trawl survey from 1999 to 2005 (see survey data in next section). In November 2000, Amendment 15 to the FMP for Bering Sea/Aleutian Islands king and Tanner crabs was approved to implement a rebuilding plan for the SMBKC stock (NPFMC 2000). The rebuilding plan included a State of Alaska regulatory harvest strategy (5 AAC 34.917), area closures, and gear modifications. In addition,

---

<sup>2</sup>NOAA grant Bering Sea Crab Research II, NA16FN2621, 1997.

commercial crab fisheries near St. Matthew Island were scheduled in fall and early winter to reduce the potential for bycatch mortality of vulnerable molting and mating crab.

NMFS declared the stock rebuilt on 21 September 2009, and the fishery was reopened after a 10-year closure on 15 October 2009 with a TAC of 529 t (1.167 million pounds), closing again by regulation on 1 February 2010. Seven participating vessels landed a catch of 209 t (0.461 million pounds) with a reported effort of 10,697 pot lifts and an estimated CPUE of 9.9 retained individual crab per pot lift. The fishery remained open the next three years with modest harvests and similar CPUE, but large declines in the NMFS trawl-survey estimate of stock abundance raised concerns about the health of the stock. This prompted ADF&G to close the fishery again for the 2013/14 season. The fishery was reopened for the 2014/15 season with a low TAC of 297 t (0.655 million pounds) and in 2015/16 the TAC was further reduced to 186 t (0.411 million pounds) then completely closed the 2016/17 season.

Although historical observer data are limited due to low sampling effort, bycatch of female and sublegal male crab from the directed blue king crab fishery off St. Matthew Island was relatively high historically, with estimated total bycatch in terms of number of crab captured sometimes more than twice as high as the catch of legal crab (Moore et al. 2000; ADF&G Crab Observer Database). Pot-lift sampling by ADF&G crab observers (Gaeuman 2013; ADF&G Crab Observer Database) indicates similar bycatch rates of discarded male crab since the reopening of the fishery (Table 5), with total male discard mortality in the 2012/13 directed fishery estimated at about 12% (88 t or 0.193 million pounds) of the reported retained catch weight, assuming 20% handling mortality.

These data suggest a reduction in the bycatch of females, which may be attributable to the later timing of the contemporary fishery and the more offshore distribution of fishery effort since reopening in 2009/10<sup>3</sup>. Some bycatch of discarded blue king crab has also been observed historically in the eastern Bering Sea snow crab fishery, but in recent years it has generally been negligible. The St. Matthew Island golden king crab fishery, the third commercial crab fishery to have taken place in the area, typically occurred in areas with depths exceeding blue king crab distribution. The NMFS observer data suggest that variable, but mostly limited, SMBKC bycatch has also occurred in the eastern Bering Sea groundfish fisheries (Table 6).

## D. Data

### Summary of New Information

Data used in this assessment were updated to include the most recently available fishery and survey estimates. Since this stock is now on a biennial assessment cycle, the new data for these models includes updated bycatch estimates and 2021 NMFS trawl survey biomass and size composition data. The assessment uses updated 1993-2020 groundfish and fixed gear bycatch estimates based on NMFS AKRO data. The directed fishery has been closed since the 2016/17 season, and therefore no directed fishery catch data are available. The data used in each of the new models is shown in Figure 3.

### Major Data Sources

Major data sources used in this assessment include annual directed-fishery retained-catch statistics from fish tickets (1978/79-1998/99, 2009/10-2012/13, and 2014/15-2015/16; Table 7); results from the annual NMFS eastern Bering Sea trawl survey (1978-2021; Table 8); results from the ADF&G SMBKC pot survey (every third year during 1995-2013, then every year during 2015-2018; Table 9); mean somatic mass given length category by year (Table 10); size-frequency information from ADF&G crab-observer pot-lift sampling (1990/91-1998/99, 2009/10-2012/13, and 2014/15-2016/17; Table 5); and the NMFS groundfish-observer bycatch biomass estimates (1992/93-2020/21; Table 6).

---

<sup>3</sup>D. Pengilly, ADF&G, pers. comm.

Figure 4 maps stations from which SMBKC trawl-survey and pot-survey data were obtained. Further information concerning the NMFS trawl survey as it relates to commercial crab species is available in Daly et al. (2014); see Gish et al. (2012) for a description of ADF&G SMBKC pot-survey methods. It should be noted that the two surveys cover different geographic regions and that each has in some years encountered proportionally large numbers of male blue king crab in areas not covered by the other survey (Figure 5). Crab-observer sampling protocols are detailed in the crab-observer training manual (ADF&G 2013). Groundfish SMBKC bycatch data come from the NMFS Regional office and have been compiled to coincide with the SMBKC management area.

## Other Data Sources

The growth transition matrix used is based on Otto and Cummiskey (1990), as in the past. Other relevant data sources, including assumed population and fishery parameters, are presented in Appendix A, which also provides a detailed description of the model configuration used for this assessment.

## E. Analytic Approach

### History of Modeling Approaches for this Stock

A four-stage catch-survey-analysis (CSA) assessment model was used before 2011 to estimate abundance and biomass and prescribe fishery quotas for the SMBKC stock. The four-stage CSA is similar to a full length-based analysis, the major difference being coarser length groups, which are more suited to a small stock with consistently low survey catches. In this approach, the abundance of male crab with a CL  $\geq 90$  mm is modeled in terms of four crab stages: stage 1: 90-104 mm CL; stage 2: 105-119 mm CL; stage 3: newshell 120-133 mm CL; and stage 4: oldshell  $\geq 120$  mm CL and newshell  $\geq 134$  mm CL. Motivation for these stage definitions comes from the fact that for management of the SMBKC stock, male crab measuring  $\geq 105$  mm CL are considered mature, whereas 120 mm CL is considered a proxy for the legal size of 5.5 in carapace width, including spines. Additional motivation for these stage definitions comes from an estimated average growth increment of about 14 mm per molt for SMBKC (Otto and Cummiskey 1990).

Concerns about the pre-2011 assessment model led to the CPT and SSC recommendations that included development of an alternative model with provisional assessment based on survey biomass or some other index of abundance. An alternative 3-stage model was proposed to the CPT in May 2011, but a survey-based approach was requested for the Fall 2011 assessment. In May 2012 the CPT approved a slightly revised and better documented version of the alternative model for assessment. Subsequently, the model developed and used since 2012 was a variant of the previous four-stage SMBKC CSA model and similar in complexity to that described by Collie et al. (2005). Like the earlier model, it considered only male crab  $\geq 90$  mm in CL, but combined stages 3 and 4 of the earlier model, resulting in three stages (male size classes) defined by CL measurements of (1) 90-104 mm, (2) 105-119 mm, and (3) 120 mm+ (i.e., 120 mm and above). This consolidation was driven by concern about the accuracy and consistency of shell-condition information, which had been used in distinguishing stages 3 and 4 of the earlier model.

In 2016 the accepted SMBKC assessment model made use of the modeling framework GMACS encompassing a three-stage model structure (Webber et al. 2016). In that assessment, an effort was made to match the 2015 SMBKC stock assessment model to bridge a framework which provided greater flexibility and opportunity to evaluate model assumptions more fully.

### Assessment Methodology

This assessment model again uses the modeling framework GMACS and is detailed in Appendix A. An updated version of GMACS (version 2.01.F, 2022-04-16) was used.

## Model Selection and Evaluation

The base model is presented here as the last accepted model in 2020, and an updated model to include data from the 2021 crab year (fishery, survey, etc.). In addition, three other models explore the model sensitivity to natural mortality estimates. In summary, the following lists the models presented and the naming convention used:

1. **16.0 - 2020 Model:** 2020 accepted model, fixed  $M = 0.18$  all years except 1998 time block where  $M$  is estimated
2. **16.0 - 2022 base model:** updated with 2020/21 groundfish bycatch & 2021 NMFS trawl data
3. **22.0a - 2022 base model with fixed  $M = 0.21$ :** natural mortality is increased to 0.21 instead of 0.18
4. **22.0b - 2022 base model with fixed  $M = 0.26$ :** natural mortality is increased to 0.26 instead of 0.18
5. **22.0c - 2022 base model, without  $M$  time blocks:** natural mortality is fixed for all years at 0.18, no time blocks

Note the change in naming convention (per SSC comments). The base model is model 16.0 since that was the year of model development and acceptance.

## Results

### a. Sensitivity to new data

The last accepted model for SMBKC was in Sept. 2020, new data added to the current 2022 model here includes both groundfish bycatch data for the 2020/21 crab season and NMFS trawl survey data - biomass and length compositions - from the 2021 summer survey. Additionally, the groundfish bycatch data was updated for past years due to some changes in the weights used to estimate crab bycatch in the groundfish fisheries (per. comm. NMFS AKRO). The 2022 reference model is compared here to the 2020 accepted model (both model 16.0), which is shown in Figures 6 and 7 with recruitment and spawning biomass shown in Figures 8 and 9, respectively. The 2020 accepted model and the 2022 base model have identical fits to the survey data, as well as nearly identical estimates of SSB and recruitment. This is expected since there is only one new survey data point between these models. As has been noted in the past, the reference model (16.0) still does not capture the recent survey declines in the ADF&G pot survey, or fit post 2005 trawl survey data points well.

### b. Effective sample sizes and weighting factors

Observed and estimated effective sample sizes are compared in Table 11. Data weighting factors, standard deviation of normalized residuals (SDNRs), and median absolute residual (MAR) are presented in Table 17. Currently the SDNR and MAR are not outputting correctly for the survey data in GMACS. This is on the list to address at the next modeling workshop. In Sept. 2019 the SDNR for the trawl survey was acceptable at 1.66 in the base model. Francis (2011) weighting was applied in 2017 but given the relatively few size bins in this assessment, this application was suspended for this assessment.

In Sept. 2019 the SDNRs for the pot surveys showed a similar pattern in each of the scenarios, but are much higher suggesting an inconsistency between the pot survey data and the model structure and other data components. Rather than re-weighting, we chose to retain the values as specified, noting that down-weighting these data would effectively exclude the signal from this series. The MAR values for the trawl and pot surveys showed the same pattern among each of the scenarios as the SDNR. The MAR values for the trawl survey and pot survey size compositions were adequate, ranging from 0.61 to 0.69 for the reference case. The SDNRs for the directed pot fishery and other size compositions were similar to previous estimates.



### c. Parameter estimates

Model parameter estimates for each of the GMACS scenarios are summarized in Tables 12, 13, 14 and 14. These parameter estimates are compared in Table 16. Negative log-likelihood values and management measures for each of the model configurations are compared in Tables 4 and 18.

There are differences in parameter estimates among models as reflected in the log-likelihood components and the management quantities. The most drastic of these differences comes in Model 22.0c which has a fixed natural mortality for the entire time series (Table 16). Removing the period of higher natural mortality in 1998 forces the model to accommodate the large drop in survey biomass by increases in recruitment and overall changes in mature male biomass in the late 90s.

Selectivity estimates for the directed fishery show some variability between the base model and model 22.0c, these differences accommodate the changes to the survey biomass estimates without increased natural mortality in 1998 (Figure 10). Estimated recruitment changes in magnitude with range of natural mortality levels in the models, with the largest differences being in the early part of the time series. Estimates of recruitment post-2000 are similar in all model runs, with the recent 10 years being the most similar regardless of the natural mortality assumed (Figure 11). Estimated mature male biomass (MMB) on 15 February is similar among the models (Figure 12). Only model 22.0c (fixed  $M = 0.18$  with no time blocks) has a different trajectory for MMB during the late 90s leading up to the large decline in 1998. Fits to the NMFS trawl survey data are similar among the models, once again with only model 22.0c having a lower trend in the 90s to fit the large drop in survey biomass from 1998 to 1999.

Estimated natural mortality in each year ( $M_t$ ) is presented in Figure 13, showing the mortality event in the late 90s for all models except model 22.0c. Estimates of fishing mortality, from the base model (16.0), are shown to assist with the rebuilding and reference point time frame discussions (Figure 26). Fishing mortality can not be ruled out as being an influential factor in the current low stock status.

### d. Evaluation of the fit to the data.

The base model (model 16.0) fit to total male ( $\geq 90$  mm CL) NMFS trawl survey biomass tends to miss the recent peak around 2010 and fits recent survey data points on the lower end of their error bars (Figures 14). These fits are most likely being pulled down by the recent decline in the ADF&G pot survey data points, since the **no pot** model that was run in the 2020 final SAFE captures more of the error bars for these data points when the NMFS trawl survey data is the only abundance index in the model. However, this model, similar to the additional CV models presenting in May 2020, tend to overfit the recent trawl survey data points (Figure 14).

The base model fit to the pot survey CPUE is similar to past reference models, fitting the overall trends in the data but not capturing some of the high and low points (Figure 15).

For the trawl survey the standardized residuals for all model scenarios have a positive residual pattern in the last 15 years, continually under predicting the observed data points (Figure 16). The standardized residuals for the ADF&G pot survey have similar patterns to past reference model iterations (Figure 17).

Fits to the size compositions for trawl survey, pot survey, and commercial observer data are reasonable but miss the largest size category in some years (Figures 18, 19, and 20) for most of the scenarios. Size composition fits in 1998 and 1999 for model 22.0c, the model that removed the 1998 time block increase in  $M$ , are very poor in these years for all three data sets. Representative residual plots of the composition data generally have a similar fit to the three composition data sources for the model scenarios with decreased fit for model 22.0c (Figures 21, 22, 23, and 24). The model fits to different types of retained and discarded catch values performed as expected given the assumed levels of uncertainty on the input data (Figure 25).

### e. Retrospective and historical analyses

This is the fourth year GMACS has been used for this stock. As such, retrospective patterns and historical analyses of GMACS assessments are limited. However, completion of a retrospective analysis, for the base model, will be completed in Sept.

### f. Uncertainty and sensitivity analyses.

Estimated standard deviations of parameters and selected management measures for the models are summarized for each individual model in Tables 12, 13, 14, and compiled in Table 16. Model estimates of mature male biomass and OFL in 2022 are presented in Section F.

### g. Comparison of alternative model scenarios.

The model scenarios presented here explore the sensitivity of the base model (16.0) to assumptions of level of natural mortality for this stock. With the exception of model 22.0c, the rest of the model suite produces a similar mature male biomass trajectory, with similar terminal year estimates of MMB (Figure 12). Removing the mortality time block in 1998 (model 22.0c) produces a similar mature male biomass trend for the last 20 years, however it allows other parameters in the model (such as recruitment and length composition fit) to change to accommodate the large drop in biomass from 1998 to 1999. As the length composition fits illustrate these changes are not supported by the data.

In summary, the suite of model scenarios were provided to explore the sensitivity of this model. Currently, the base model is still the most appropriate model for setting reference points and model specifications. Research on alternative model specifications that may address the disparities between the trawl and pot survey data are ongoing, and a preliminary spatial overview is presented in Appendix C. Additionally, the overfished status of this stock lends itself to maintaining the status quo base model until an appropriate resolution is found to deal with the trawl and pot survey data fit issues. The recommended model for fall 2022 would be the base model (16.0) to maintain consistency for this stock during the rebuilding time frame.

## F. Calculation of the OFL and ABC

The overfishing level (OFL) is the total catch associated with the  $F_{OFL}$  fishing mortality. The SMBKC stock is currently managed as Tier 4, and only a Tier 4 analysis is presented here. Thus, given stock estimates or suitable proxy values of  $B_{MSY}$  and  $F_{MSY}$ , along with two additional parameters  $\alpha$  and  $\beta$ ,  $F_{OFL}$  is determined by the control rule

$$F_{OFL} = \begin{cases} F_{MSY}, & \text{when } B/B_{MSY} > 1 \\ F_{MSY} \frac{(B/B_{MSY} - \alpha)}{(1 - \alpha)}, & \text{when } \beta < B/B_{MSY} \leq 1 \end{cases} \quad (1)$$
$$F_{OFL} < F_{MSY} \text{ with directed fishery } F = 0 \text{ when } B/B_{MSY} \leq \beta$$

where  $B$  is quantified as mature-male biomass (MMB) at mating with time of mating assigned a nominal date of 15 February. Note that as  $B$  itself is a function of the fishing mortality  $F_{OFL}$  (therefore numerical approximation of  $F_{OFL}$  is required). As implemented for this assessment, all calculations proceed according to the model equations given in Appendix A.  $F_{OFL}$  is taken to be full-selection fishing mortality in the directed pot fishery and groundfish trawl and fixed-gear fishing mortalities set at their geometric mean values over years for which there are data-based estimates of bycatch-mortality biomass.

The currently recommended Tier 4 convention is to use the full assessment period, currently 1978 - 2020, to define a  $B_{MSY}$  proxy in terms of average estimated MMB and to set  $\gamma = 1.0$  with assumed stock natural mortality  $M = 0.18 \text{ yr}^{-1}$  in setting the  $F_{MSY}$  proxy value  $\gamma M$ . The parameters  $\alpha$  and  $\beta$  are assigned

their default values  $\alpha = 0.10$  and  $\beta = 0.25$ . The  $F_{OFL}$ , OFL, ABC, and MMB in 2019 for all scenarios are summarized in Table 4. The currently recommended ABC is 75% of the OFL (ABC buffer = 25%).

Table 4: Comparisons of management measures for the model scenarios. Biomass and OFL are in tons.

Component	Ref	M0.21	M0.26	Notimeblock
MMB <sub>2021</sub>	1144.575	1077.861	1047.205	1032.327
$B_{MSY}$	3298.391	3333.337	3531.318	3005.061
$MMB/B_{MSY}$	0.339	0.308	0.288	0.336
$F_{OFL}$	0.048	0.049	0.000	0.047
OFL <sub>2021</sub>	49.323	45.887	1.207	44.021
ABC <sub>2021</sub>	36.992	34.415	0.905	33.016

## G. Rebuilding Analysis

This stock was declared overfished in fall of 2018 and a rebuilding plan went before the Council for final review in June 2020. The most updated rebuilding plan can be found on the NPFMC website for the June 2020 meeting. Progress towards rebuilding will be updated in the Sept. 2022 SAFE report.

## H. Data Gaps and Research Priorities

The following topics have been listed as areas where more research on SMBKC is needed:

1. Growth increments and molting probabilities as a function of size.
2. Trawl survey catchability and selectivities.
3. Pot survey catchability and selectivities.
4. Temporal changes in spatial distributions near the island.
5. Natural mortality.

## I. Projections and outlook

The outlook for recruitment is pessimistic and the abundance relative to the proxy  $B_{MSY}$  is low. The NMFS survey results in 2019 noted ocean conditions warmer than normal with an absence of a “cold pool” in the region. This could have detrimental effects on the SMBKC stock and should be carefully monitored. Relative to the impact of historical fishing, we again conducted a “dynamic- $B_0$ ” analysis. This procedure simply projects the population based on estimated recruitment but removes the effect of fishing. For the reference case, this suggests that the impact of fishing has reduced the stock to about 61% of what it would have been in the absence of fishing (Figure 27, supporting the hypothesis that fishing pressure is not the sole contributor to the decline of this stock in recent years. The other non-fishing contributors to the observed depleted stock trend (ignoring stock-recruit relationship) may reflect variable survival rates due to environmental conditions and also range shifts.

## J. Acknowledgements

We thank the Crab Plan Team and ADF&G staff for reviewing an earlier draft of this report and Andre Punt for his continued input into refinements to the GMACS model code, specifically the projections module.

## K. References

- Alaska Department of Fish and Game (ADF&G). 2013. Crab observer training and deployment manual. Alaska Department of Fish and Game Shellfish Observer Program, Dutch Harbor. Unpublished.
- Collie, J.S., A.K. Delong, and G.H. Kruse. 2005. Three-stage catch-survey analysis applied to blue king crabs. Pages 683-714 [In] Fisheries assessment and management in data-limited situations. University of Alaska Fairbanks, Alaska Sea Grant Report 05-02, Fairbanks.
- Daly, B., R. Foy, and C. Armistead. 2014. The 2013 eastern Bering Sea continental shelf bottom trawl survey: results for commercial crab species. NOAA Technical Memorandum 295, NMFS-AFSC.
- Donaldson, W.E., and S.C. Byersdorfer. 2005. Biological field techniques for lithodid crabs. University of Alaska Fairbanks, Alaska Sea Grant Report 05-03, Fairbanks.
- Fitch, H., M. Deiman, J. Shaishnikoff, and K. Herring. 2012. Annual management report for the commercial and subsistence shellfish fisheries of the Bering Sea, 2010/11. Pages 75-1776 [In] Fitch, H., M. Schwenzfeier, B. Baechler, T. Hartill, M. Salmon, and M. Deiman, E.
- Evans, E. Henry, L. Wald, J. Shaishnikoff, K. Herring, and J. Wilson. 2012. Annual management report for the commercial and subsistence shellfish fisheries of the Aleutian Islands, Bering Sea and the Westward Region's Shellfish Observer Program, 2010/11. Alaska Department of Fish and Game, Fishery Management Report No. 12-22, Anchorage.
- Fournier, D.A., H.J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J. Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim. Methods Softw.* 27:233-249.
- Francis, R.I.C.C. 2011. Data weighting in statistical fisheries stock assessment models. *Can. J. Fish. Aquat. Sci.* 68: 1124-1138.
- Gaeuman, W.B. 2013. Summary of the 2012/13 mandatory crab observer program database for the Bering Sea/Aleutian Islands commercial crab fisheries. Alaska Department of Fish and Game, Fishery Data Series No. 13-54, Anchorage.
- Gish, R.K., V.A. Vanek, and D. Pengilly. 2012. Results of the 2010 triennial St. Matthew Island blue king crab pot survey and 2010/11 tagging study. Alaska Department of Fish and Game, Fishery Management Report No. 12-24, Anchorage.
- Ianelli, J., D. Webber, and J. Zheng, 2017. Stock assessment of Saint Matthews Island blue king crab. North Pacific Fishery Management Council. Anchorage AK.
- Ianelli, J., and J. Zheng, 2018. Stock assessment of Saint Matthews Island blue king crab. North Pacific Fishery Management Council. Anchorage AK.
- Jensen, G.C., and D.A. Armstrong. 1989. Biennial reproductive cycle of blue king crab, *Paralithodes platypus*, at the Pribilof Islands, Alaska and comparison to a congener, *P. camtschatica*. *Can. J. Fish. Aquat. Sci.* 46: 932-940.
- Moore, H., L.C. Byrne, and D. Connolly. 2000. Alaska Department of Fish and Game summary of the 1998 mandatory shellfish observer program database. Alaska Dept. Fish and Game, Commercial Fisheries Division, Reg. Inf. Rep. 4J00-21, Kodiak.
- North Pacific Fishery Management Council (NPFMC). 1998. Fishery Management Plan for Bering Sea/Aleutian Islands king and Tanner crabs. North Pacific Fishery Management Council, Anchorage.
- North Pacific Fishery Management Council (NPFMC). 1999. Environmental assessment/regulatory impact review/initial regulatory flexibility analysis for Amendment 11 to the Fishery Management Plan for Bering Sea/Aleutian Islands king and Tanner crabs. North Pacific Fishery Management Council, Anchorage.
- North Pacific Fishery Management Council (NPFMC). 2000. Environmental assessment/regulatory impact review/initial regulatory flexibility analysis for proposed Amendment 15 to the Fishery Management Plan for

king and Tanner crab fisheries in the Bering Sea/Aleutian Islands and regulatory amendment to the Fishery Management Plan for the groundfish fishery of the Bering Sea and Aleutian Islands area: A rebuilding plan for the St. Matthew blue king crab stock. North Pacific Fishery Management Council, Anchorage. Draft report.

North Pacific Fishery Management Council (NPFMC). 2007. Public Review Draft: Environmental assessment for proposed Amendment 24 to the Fishery Management Plan for Bering Sea and Aleutian Islands king and Tanner crabs to revise overfishing definitions. 14 November 2007. North Pacific Fishery Management Council, Anchorage.

Otto, R.S. 1990. An overview of eastern Bering Sea king and Tanner crab fisheries. Pages 9-26 [In] Proceedings of the international symposium on king and Tanner crabs. University of Alaska Fairbanks, Alaska Sea Grant Program Report 90-4, Fairbanks.

Otto, R.S., and P.A. Cummiskey. 1990. Growth of adult male blue king crab (*Paralithodes platypus*). Pages 245-258 [In] Proceedings of the international symposium on king and Tanner crabs. University of Alaska Fairbanks, Alaska Sea Grant Report 90-4, Fairbanks.

Palof, K.J., J. Zheng, and Ianelli, J., 2019. Stock assessment of Saint Matthews Island blue king crab. North Pacific Fishery Management Council. Anchorage AK.

Paul, J.M., A. J. Paul, R.S. Otto, and R.A. MacIntosh. 1991. Spermatophore presence in relation to carapace length for eastern Bering Sea blue king crab (*Paralithodes platypus*, Brandt, 1850) and red king crab (*P. camtschaticus*, Tilesius, 1815). J. Shellfish Res. 10: 157-163.

Pengilly, D. and D. Schmidt. 1995. Harvest Strategy for Kodiak and Bristol Bay red king crab and St. Matthew Island and Pribilof blue king crab. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Special Publication Number 7, Juneau.

Somerton, D.A., and R.A. MacIntosh. 1983. The size at sexual maturity of blue king crab, *Paralithodes platypus*, in Alaska. Fishery Bulletin 81: 621-828.

Then, A. Y., Hoenig, J. M., Hall, N. G., and Hewitt, D. A. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. ICES Journal of Marine Science, 72: 82-92.

Thorson, J.T., and L.A.K. Barnett. 2017. Comparing estimates of abundance trends and distribution shifts using single- and multispecies models of fishes and biogenic habitat. ICES Journal of Marine Science 75:1311-1321.

Thorson, J.T., J.N. Ianelli, E. Larsen, L. Ries, M.D. Scheuerell, C. Szuwalski, and E. Zipkin. 2016. Joint dynamic species distribution models: a tool for community ordination and spatiotemporal monitoring. Glob. Ecol. Biogeogr. 25(9): 1144-1158. doi:10.1111/geb.12464.

Thorson, J.T., Scheuerell, M.D., Shelton, A.O., See, K.E., Skaug, H.J., and Kristensen, K. 2015. Spatial factor analysis: a new tool for estimating joint species distributions and correlations in species range. Methods Ecol. Evol. 6(6): 627-637. doi:10.1111/2041-210X.12359.

Webber, D., J. Zheng, and J. Ianelli, 2016. Stock assessment of Saint Matthews Island Blue King Crab. North Pacific Fishery Management Council. Anchorage AK.

Zheng, J. 2005. A review of natural mortality estimation for crab stocks: data-limited for every stock? Pages 595-612 [In] Fisheries Assessment and Management in Data-Limited Situations. University of Alaska Fairbanks, Alaska Sea Grant Program Report 05-02, Fairbanks.

Zheng, J., and G.H. Kruse. 2002. Assessment and management of crab stocks under uncertainty of massive die-offs and rapid changes in survey catchability. Pages 367-384 [In] A.J. Paul, E.G. Dawe, R. Elner, G.S. Jamieson, G.H. Kruse, R.S. Otto, B. Sainte-Marie, T.C. Shirley, and D. Woodby (eds.). Crabs in Cold Water Regions: Biology, Management, and Economics. University of Alaska Fairbanks, Alaska Sea Grant Report 02-01, Fairbanks.

Zheng, J., M.C. Murphy, and G.H. Kruse. 1997. Application of catch-survey analysis to blue king crab stocks near Pribilof and St. Matthew Islands. Alaska Fish. Res. Bull. 4:62-74.

## Tables

Table 5: Observed proportion of crab by size class during the ADF&G crab observer pot-lift sampling.  
Source: ADF&G Crab Observer Database.

Year	Total pot lifts	Pot lifts sampled	Number of crab (90 mm+ CL)	Stage 1	Stage 2	Stage 3
1990/91	26,264	10	150	0.113	0.393	0.493
1991/92	37,104	125	3,393	0.133	0.177	0.690
1992/93	56,630	71	1,606	0.191	0.268	0.542
1993/94	58,647	84	2,241	0.281	0.210	0.510
1994/95	60,860	203	4,735	0.294	0.271	0.434
1995/96	48,560	47	663	0.148	0.212	0.640
1996/97	91,085	96	489	0.160	0.223	0.618
1997/98	81,117	133	3,195	0.182	0.205	0.613
1998/99	91,826	135	1,322	0.193	0.216	0.591
1999/00 - 2008/09			FISHERY CLOSED			
2009/10	10,484	989	19,802	0.141	0.324	0.535
2010/11	29,356	2,419	45,466	0.131	0.315	0.553
2011/12	48,554	3,359	58,666	0.131	0.305	0.564
2012/13	37,065	2,841	57,298	0.141	0.318	0.541
2013/14			FISHERY CLOSED			
2014/15	10,133	895	9,906	0.094	0.228	0.679
2015/16	5,475	419	3,248	0.115	0.252	0.633
2016/17 - 2020/21			FISHERY CLOSED			

Table 6: Groundfish SMBKC male bycatch biomass (t) estimates. Trawl includes pelagic trawl and non-pelagic trawl types. Source: J. Zheng, ADF&G, and author estimates based on data from R. Foy, NMFS. Estimates used after 2008/09 are from NMFS Alaska Regional Office.

Year	Trawl bycatch	Fixed gear bycatch
1978	0.000	0.000
1979	0.000	0.000
1980	0.000	0.000
1981	0.000	0.000
1982	0.000	0.000
1983	0.000	0.000
1984	0.000	0.000
1985	0.000	0.000
1986	0.000	0.000
1987	0.000	0.000
1988	0.000	0.000
1989	0.000	0.000
1990	0.000	0.000
1991	3.538	0.045
1992	1.996	2.268
1993	1.542	0.500
1994	0.318	0.091
1995	0.635	0.136
1996	0.500	0.045
1997	0.500	0.181
1998	0.500	0.907
1999	0.500	1.361
2000	0.500	0.500
2001	0.500	0.862
2002	0.726	0.408
2003	0.998	1.134
2004	0.091	0.635
2005	0.500	0.590
2006	2.812	1.451
2007	0.045	69.717
2008	0.272	6.622
2009	0.638	7.522
2010	0.360	9.564
2011	0.170	0.796
2012	0.011	0.739
2013	0.163	0.341
2014	0.010	0.490
2015	0.010	0.711
2016	0.229	1.630
2017	0.048	5.935
2018	0.001	1.224
2019	0.030	1.124
2020	0.001	0.671

Table 7: Fishery characteristics and update. Columns include the 1978/79 to 2015/16 directed St. Matthew Island blue king crab pot fishery. The Guideline Harvest Level (GHL) and Total Allowable Catch (TAC) are in millions of pounds. Harvest includes deadloss. Catch per unit effort (CPUE) in this table is simply the harvest number / pot lifts. The average weight is the harvest weight / harvest number in pounds. The average CL is the average of retained crab in mm from dockside sampling of delivered crab. Source: Fitch et al 2012; ADF&G Dutch Harbor staff, pers. comm. Note that management (GHL) units are in pounds, for conserving space, conversion to tons is ommitted.

Year	Dates	GHL/TAC	Harvest		Pot lifts	CPUE	avg wt	avg CL
			Crab	Pounds				
1978/79	07/15 - 09/03		436,126	1,984,251	43,754	10	4.5	132.2
1979/80	07/15 - 08/24		52,966	210,819	9,877	5	4.0	128.8
1980/81	07/15 - 09/03		CONFIDENTIAL					
1981/82	07/15 - 08/21		1,045,619	4,627,761	58,550	18	4.4	NA
1982/83	08/01 - 08/16		1,935,886	8,844,789	165,618	12	4.6	135.1
1983/84	08/20 - 09/06	8.0	1,931,990	9,454,323	133,944	14	4.9	137.2
1984/85	09/01 - 09/08	2.0-4.0	841,017	3,764,592	73,320	11	4.5	135.5
1985/86	09/01 - 09/06	0.9-1.9	436,021	2,175,087	46,988	9	5.0	139.0
1986/87	09/01 - 09/06	0.2-0.5	219,548	1,003,162	22,073	10	4.6	134.3
1987/88	09/01 - 09/05	0.6-1.3	227,447	1,039,779	28,230	8	4.6	134.1
1988/89	09/01 - 09/05	0.7-1.5	280,401	1,236,462	21,678	13	4.4	133.3
1989/90	09/01 - 09/04	1.7	247,641	1,166,258	30,803	8	4.7	134.6
1990/91	09/01 - 09/07	1.9	391,405	1,725,349	26,264	15	4.4	134.3
1991/92	09/16 - 09/20	3.2	726,519	3,372,066	37,104	20	4.6	134.1
1992/93	09/04 - 09/07	3.1	545,222	2,475,916	56,630	10	4.5	134.1
1993/94	09/15 - 09/21	4.4	630,353	3,003,089	58,647	11	4.8	135.4
1994/95	09/15 - 09/22	3.0	827,015	3,764,262	60,860	14	4.9	133.3
1995/96	09/15 - 09/20	2.4	666,905	3,166,093	48,560	14	4.7	135.0
1996/97	09/15 - 09/23	4.3	660,665	3,078,959	91,085	7	4.7	134.6
1997/98	09/15 - 09/22	5.0	939,822	4,649,660	81,117	12	4.9	139.5
1998/99	09/15 - 09/26	4.0	635,370	2,968,573	91,826	7	4.7	135.8
1999/00 - 2008/09			FISHERY CLOSED					
2009/10	10/15 - 02/01	1.17	103,376	460,859	10,697	10	4.5	134.9
2010/11	10/15 - 02/01	1.60	298,669	1,263,982	29,344	10	4.2	129.3
2011/12	10/15 - 02/01	2.54	437,862	1,881,322	48,554	9	4.3	130.0
2012/13	10/15 - 02/01	1.63	379,386	1,616,054	37,065	10	4.3	129.8
2013/14			FISHERY CLOSED					
2014/15	10/15 - 02/05	0.66	69,109	308,582	10,133	7	4.5	132.3
2015/16	10/19 - 11/28	0.41	24,076	105,010	5,475	4	4.4	132.6
2016/17			FISHERY CLOSED					
2017/18			FISHERY CLOSED					
2018/19			FISHERY CLOSED					
2019/20			FISHERY CLOSED					
2020/21			FISHERY CLOSED					



Table 8: NMFS EBS trawl-survey area-swept estimates of male crab abundance ( $10^6$  crab) and male ( $\geq 90$  mm CL) biomass ( $10^6$  lbs). Total number of captured male crab  $\geq 90$  mm CL is also given. Source: J.Richar, NMFS. The "+" refer to plus group.

Year	Abundance					Biomass		Number of crabs
	Stage-1 (90-104 mm)	Stage-2 (105-119 mm)	Stage-3 (120+ mm)	Total	CV	Total (90+ mm CL)	CV	
1978	2.213	1.991	1.521	5.726	0.411	15.064	0.394	157
1979	3.061	2.281	1.808	7.150	0.472	17.615	0.463	178
1980	2.856	2.563	2.541	7.959	0.572	22.017	0.507	185
1981	0.483	1.213	2.263	3.960	0.368	14.443	0.402	140
1982	1.669	2.431	5.884	9.984	0.401	35.763	0.344	271
1983	1.061	1.651	3.345	6.057	0.332	21.240	0.298	231
1984	0.435	0.497	1.452	2.383	0.175	8.976	0.179	105
1985	0.379	0.376	1.117	1.872	0.216	6.858	0.210	93
1986	0.203	0.447	0.374	1.025	0.428	3.124	0.388	46
1987	0.325	0.631	0.715	1.671	0.302	5.024	0.291	71
1988	0.410	0.816	0.957	2.183	0.285	6.963	0.252	81
1989	2.169	1.154	1.786	5.109	0.314	13.974	0.271	208
1990	1.053	1.031	2.338	4.422	0.302	14.837	0.274	170
1991	1.147	1.665	2.233	5.046	0.259	15.318	0.248	197
1992	1.074	1.382	2.291	4.746	0.206	15.638	0.201	220
1993	1.521	1.828	3.276	6.626	0.185	21.051	0.169	324
1994	0.883	1.298	2.257	4.438	0.187	14.416	0.176	211
1995	1.025	1.188	1.741	3.953	0.187	12.574	0.178	178
1996	1.238	1.891	3.064	6.193	0.263	20.746	0.241	285
1997	1.165	2.228	3.789	7.182	0.367	24.084	0.337	296
1998	0.660	1.661	2.849	5.170	0.373	17.586	0.355	243
1998	0.223	0.222	0.558	1.003	0.192	3.515	0.182	52
2000	0.282	0.285	0.740	1.307	0.303	4.623	0.310	61
2001	0.419	0.502	0.938	1.859	0.243	6.242	0.245	91
2002	0.111	0.230	0.640	0.981	0.311	3.820	0.320	38
2003	0.449	0.280	0.465	1.194	0.399	3.454	0.336	65
2004	0.247	0.184	0.562	0.993	0.369	3.360	0.305	48
2005	0.319	0.310	0.501	1.130	0.403	3.620	0.371	42
2006	0.917	0.642	1.240	2.798	0.339	8.585	0.334	126
2007	2.518	2.020	1.193	5.730	0.420	14.266	0.385	250
2008	1.352	0.801	1.457	3.609	0.289	10.261	0.284	167
2009	1.573	2.161	1.410	5.144	0.263	13.892	0.256	251
2010	3.937	3.253	2.458	9.648	0.544	24.539	0.466	388
2011	1.800	3.255	3.207	8.263	0.587	24.099	0.558	318
2012	0.705	1.970	1.808	4.483	0.361	13.669	0.339	193
2013	0.335	0.452	0.807	1.593	0.215	5.043	0.217	74
2014	0.723	1.627	1.809	4.160	0.503	13.292	0.449	181
2015	0.992	1.269	1.979	4.240	0.774	12.958	0.770	153
2016	0.535	0.660	1.178	2.373	0.447	7.685	0.393	108
2017	0.091	0.323	0.663	1.077	0.657	3.955	0.600	42
2018	0.154	0.232	0.660	1.047	0.298	3.816	0.281	62
2019	0.403	0.482	1.170	2.056	0.352	6.990	0.337	105
2021	0.423	0.168	0.682	1.273	0.496	4.253	0.427	59

Table 9: Size-class and total CPUE (90+ mm CL) with estimated CV and total number of captured crab (90+ mm CL) from the 96 common stations surveyed during the ADF&G SMBKC pot surveys. Source: ADF&G.

Year	Stage-1 (90-104 mm)	Stage-2 (105-119 mm)	Stage-3 (120+ mm)	Total CPUE	CV	Number of crabs
1995	1.919	3.198	6.922	12.042	0.13	4624
1998	0.964	2.763	8.804	12.531	0.06	4812
2001	1.266	1.737	5.487	8.477	0.08	3255
2004	0.112	0.414	1.141	1.667	0.15	640
2007	1.086	2.721	4.836	8.643	0.09	3319
2010	1.326	3.276	5.607	10.209	0.13	3920
2013	0.878	1.398	3.367	5.643	0.19	2167
2015	0.198	0.682	1.924	2.805	0.18	1077
2016	0.198	0.456	1.724	2.378	0.19	777
2017	0.177	0.429	1.083	1.689	0.25	643
2018	0.076	0.161	0.508	0.745	0.14	286

Table 10: Mean weight (kg) by stage in used in all of the models (provided as a vector of weights at length each year to GMACS).

Year	Stage-1	Stage-2	Stage-3
1978	0.7	1.2	1.9
1979	0.7	1.2	1.7
1980	0.7	1.2	1.9
1981	0.7	1.2	1.9
1982	0.7	1.2	1.9
1983	0.7	1.2	2.1
1984	0.7	1.2	1.9
1985	0.7	1.2	2.1
1986	0.7	1.2	1.9
1987	0.7	1.2	1.9
1988	0.7	1.2	1.9
1989	0.7	1.2	2.0
1990	0.7	1.2	1.9
1991	0.7	1.2	2.0
1992	0.7	1.2	1.9
1993	0.7	1.2	2.0
1994	0.7	1.2	1.9
1995	0.7	1.2	2.0
1996	0.7	1.2	2.0
1997	0.7	1.2	2.1
1998	0.7	1.2	2.0
1999	0.7	1.2	1.9
2000	0.7	1.2	1.9
2001	0.7	1.2	1.9
2002	0.7	1.2	1.9
2003	0.7	1.2	1.9
2004	0.7	1.2	1.9
2005	0.7	1.2	1.9
2006	0.7	1.2	1.9
2007	0.7	1.2	1.9
2008	0.7	1.2	1.9
2009	0.7	1.2	1.9
2010	0.7	1.2	1.8
2011	0.7	1.2	1.8
2012	0.7	1.2	1.8
2013	0.7	1.2	1.9
2014	0.7	1.2	1.9
2015	0.7	1.2	1.9
2016	0.7	1.2	1.9
2017	0.7	1.2	1.9
2018	0.7	1.2	1.9
2019	0.7	1.2	1.9
2020	0.7	1.2	1.9

Table 11: Observed and input sample sizes for observer data from the directed pot fishery, the NMFS trawl survey, and the ADF&G pot survey.

Year	Number measured			Input sample sizes		
	Observer pot	NMFS trawl	ADF&G pot	Observer pot	NMFS trawl	ADF&G pot
1978		157			50	
1979		178			50	
1980		185			50	
1981		140			50	
1982		271			50	
1983		231			50	
1984		105			50	
1985		93			46.5	
1986		46			23	
1987		71			35.5	
1988		81			40.5	
1989		208			50	
1990	150	170		15	50	
1991	3393	197		25	50	
1992	1606	220		25	50	
1993	2241	324		25	50	
1994	4735	211		25	50	
1995	663	178	4624	25	50	100
1996	489	285		25	50	
1997	3195	296		25	50	
1998	1323	243	4812	25	50	100
1999		52			26	
2000		61			30.5	
2001		91	3255		45.5	100
2002		38			19	
2003		65			32.5	
2004		48	640		24	100
2005		42			21	
2006		126			50	
2007		250	3319		50	100
2008		167			50	
2009	19802	251		50	50	
2010	45466	388	3920	50	50	100
2011	58667	318		50	50	
2012	57282	193		50	50	
2013		74	2167		37	100
2014	9906	181		50	50	
2015	3248	153	1077	50	50	100
2016		108	777		50	100
2017		42	643		21	100
2018		62	286		31	100
2019		105			50	
2020						
2021		59			50	

Table 12: Model parameter estimates, selected derived quantities, and their standard deviations (SD) for the base (16.0) model 2022.

Parameter	Estimate	SD
Natural mortality deviation in 1998/99 ( $\delta_{1998}^M$ )	1.581	0.137
$\log(\bar{R})$	13.868	0.196
$\log(n_1^0)$	14.954	0.174
$\log(n_2^0)$	14.511	0.211
$\log(n_3^0)$	14.327	0.207
$q_{pot}$	3.786	0.246
$\log(\bar{F}^{df})$	-2.130	0.052
$\log(\bar{F}^{tb})$	-9.636	0.073
$\log(\bar{F}^{fb})$	-8.082	0.073
log Stage-1 directed pot selectivity 1978-2008	-0.920	0.180
log Stage-2 directed pot selectivity 1978-2008	-0.561	0.132
log Stage-1 directed pot selectivity 2009-2017	-0.542	0.163
log Stage-2 directed pot selectivity 2009-2017	-0.000	0.000
log Stage-1 NMFS trawl selectivity	-0.316	0.066
log Stage-2 NMFS trawl selectivity	-0.000	0.000
log Stage-1 ADF&G pot selectivity	-0.720	0.125
log Stage-2 ADF&G pot selectivity	-0.000	0.000
$F_{OFL}$	0.048	0.007
OFL	49.323	11.990

Table 13: Model parameter estimates, selected derived quantities, and their standard deviations (SD) for Model 22.0a with increased M=0.21).

Parameter	Estimate	SD
Natural mortality deviation in 1998/99 ( $\delta_{1998}^M$ )	1.372	0.147
$\log(\bar{R})$	13.995	0.197
$\log(n_1^0)$	15.027	0.176
$\log(n_2^0)$	14.544	0.212
$\log(n_3^0)$	14.366	0.209
$q_{pot}$	3.635	0.258
$\log(\bar{F}^{df})$	-2.112	0.057
$\log(\bar{F}^{tb})$	-9.685	0.073
$\log(\bar{F}^{fb})$	-8.131	0.073
log Stage-1 directed pot selectivity 1978-2008	-1.013	0.180
log Stage-2 directed pot selectivity 1978-2008	-0.610	0.132
log Stage-1 directed pot selectivity 2009-2017	-0.658	0.169
log Stage-2 directed pot selectivity 2009-2017	-0.000	0.000
log Stage-1 NMFS trawl selectivity	-0.360	0.068
log Stage-2 NMFS trawl selectivity	-0.000	0.000
log Stage-1 ADF&G pot selectivity	-0.840	0.127
log Stage-2 ADF&G pot selectivity	-0.018	0.076
$F_{OFL}$	0.049	0.008
OFL	45.887	11.475

Table 14: Model parameter estimates, selected derived quantities, and their standard deviations (SD) for Model 22.0b increased  $M = 0.26$ .

Parameter	Estimate	SD
Natural mortality deviation in 1998/99 ( $\delta_{1998}^M$ )	1.205	0.149
$\log(\bar{R})$	14.224	0.198
$\log(n_1^0)$	15.239	0.178
$\log(n_2^0)$	14.623	0.220
$\log(n_3^0)$	14.358	0.236
$q_{pot}$	3.456	0.251
$\log(\bar{F}^{df})$	-2.145	0.054
$\log(\bar{F}^{tb})$	-9.769	0.075
$\log(\bar{F}^{fb})$	-8.216	0.075
log Stage-1 directed pot selectivity 1978-2008	-1.234	0.183
log Stage-2 directed pot selectivity 1978-2008	-0.754	0.130
log Stage-1 directed pot selectivity 2009-2017	-0.853	0.175
log Stage-2 directed pot selectivity 2009-2017	-0.000	0.000
log Stage-1 NMFS trawl selectivity	-0.592	0.076
log Stage-2 NMFS trawl selectivity	-0.094	0.066
log Stage-1 ADF&G pot selectivity	-1.050	0.130
log Stage-2 ADF&G pot selectivity	-0.155	0.076
$F_{OFL}$	0.000	0.000
OFL	1.207	0.252

Table 15: Model parameter estimates, selected derived quantities, and their standard deviations (SD) for Model 22.0c constant  $M = 0.18$  without 1998 time block.

Parameter	Estimate	SD
Natural mortality deviation in 1998/99 ( $\delta_{1998}^M$ )		
$\log(\bar{R})$	13.768	0.196
$\log(n_1^0)$	14.950	0.176
$\log(n_2^0)$	14.568	0.209
$\log(n_3^0)$	14.366	0.209
$q_{pot}$	3.925	0.254
$\log(\bar{F}^{df})$	-1.947	0.051
$\log(\bar{F}^{tb})$	-9.541	0.069
$\log(\bar{F}^{fb})$	-7.987	0.069
log Stage-1 directed pot selectivity 1978-2008	-0.926	0.180
log Stage-2 directed pot selectivity 1978-2008	-0.632	0.148
log Stage-1 directed pot selectivity 2009-2017	-0.558	0.165
log Stage-2 directed pot selectivity 2009-2017	-0.000	0.000
log Stage-1 NMFS trawl selectivity	-0.242	0.066
log Stage-2 NMFS trawl selectivity	-0.000	0.000
log Stage-1 ADF&G pot selectivity	-0.453	0.122
log Stage-2 ADF&G pot selectivity	-0.000	0.000
$F_{OFL}$	0.047	0.000
OFL	44.021	0.252

Table 16: Comparisons of parameter estimates for the model scenarios.

Parameter	Ref	M0.21	M0.26	Mall
$\log(\bar{F}^{\text{df}})$	-2.130	-2.112	-2.145	-1.947
$\log(\bar{F}^{\text{fb}})$	-8.082	-8.131	-8.216	-7.987
$\log(\bar{F}^{\text{tb}})$	-9.636	-9.685	-9.769	-9.541
$\log(\bar{R})$	13.868	13.995	14.224	13.768
$\log(n_1^0)$	14.954	15.027	15.239	14.950
$\log(n_2^0)$	14.511	14.544	14.623	14.568
$\log(n_3^0)$	14.327	14.366	14.358	14.366
$F_{\text{OFL}}$	0.048	0.049	0.000	0.047
$q_{\text{pot}}$	3.786	3.635	3.456	3.925
log Stage-1 ADF&G pot selectivity	-0.720	-0.840	-1.050	-0.453
log Stage-1 directed pot selectivity 1978-2008	-0.920	-1.013	-1.234	-0.926
log Stage-1 directed pot selectivity 2009-2017	-0.542	-0.658	-0.853	-0.558
log Stage-1 NMFS trawl selectivity	-0.316	-0.360	-0.592	-0.242
log Stage-2 ADF&G pot selectivity	-0.000	-0.018	-0.155	-0.000
log Stage-2 directed pot selectivity 1978-2008	-0.561	-0.610	-0.754	-0.632
log Stage-2 directed pot selectivity 2009-2017	-0.000	-0.000	-0.000	-0.000
log Stage-2 NMFS trawl selectivity	-0.000	-0.000	-0.094	-0.000
Natural mortality deviation in 1998/99 ( $\delta_{1998}^M$ )	1.581	1.372	1.205	-
OFL	49.323	45.887	1.207	44.021

Table 17: Comparisons of data weights, SDNR and MAR (standard deviation of normalized residuals and median absolute residual) values for the model scenarios.

Component	Ref	M0.21	M0.26	Notimeblock
NMFS trawl survey weight	1.00	1.00	1.00	1.00
ADF&G pot survey weight	1.00	1.00	1.00	1.00
Directed pot LF weight	1.00	1.00	1.00	1.00
NMFS trawl survey LF weight	1.00	1.00	1.00	1.00
ADF&G pot survey LF weight	1.00	1.00	1.00	1.00
SDNR NMFS trawl survey	0.00	0.00	0.00	0.00
SDNR ADF&G pot survey	0.00	0.00	0.00	0.00
SDNR directed pot LF	0.67	0.66	0.67	1.40
SDNR NMFS trawl survey LF	1.29	1.33	1.24	1.77
SDNR ADF&G pot survey LF	0.95	0.92	0.92	3.75
MAR NMFS trawl survey	0.00	0.00	0.00	0.00
MAR ADF&G pot survey	0.00	0.00	0.00	0.00
MAR directed pot LF	0.46	0.51	0.54	0.39
MAR NMFS trawl survey LF	0.61	0.57	0.66	0.73
MAR ADF&G pot survey LF	0.69	0.62	0.66	0.75

Table 18: Comparisons of negative log-likelihood values for the selected model scenarios. It is important to note that comparisons among models may be limited since the number of parameters between models changes (e.g., **nopot** model).

Component	Ref	M0.21	M0.26	Notimeblock
Pot Retained Catch	-68.46	-68.88	-69.26	-68.63
Pot Discarded Catch	6.67	5.62	4.27	12.05
Trawl bycatch Discarded Catch	-8.26	-8.26	-8.26	-8.26
Fixed bycatch Discarded Catch	-8.23	-8.24	-8.25	-8.23
NMFS Trawl Survey	6.56	5.43	0.35	15.12
ADF&G Pot Survey CPUE	85.15	77.52	68.75	73.22
Directed Pot LF	-104.67	-104.81	-104.64	-96.15
NMFS Trawl LF	-260.01	-259.01	-261.38	-237.57
ADF&G Pot LF	-91.22	-91.91	-91.92	-84.16
Recruitment deviations	61.86	61.96	61.54	63.16
F penalty	9.66	9.66	9.66	9.66
M penalty	6.46	6.45	6.45	0.00
Prior	13.71	13.71	13.71	13.02
Total	-350.76	-360.75	-378.98	-316.77
Total estimated parameters	150.00	150.00	150.00	149.00



Table 19: Population abundances ( $n$ ) by crab stage in numbers of crab at the time of the survey and mature male biomass (MMB) in tons on 15 February for the **model configuration used in 2019**.

Year	$n_1$	$n_2$	$n_3$	MMB	CV MMB
1978	3122227	2005666	1668140	4567	0.177
1979	4229414	2371064	2292298	6467	0.123
1980	3795952	3180118	3465983	10172	0.084
1981	1440711	3205532	4831471	10632	0.063
1982	1604844	1833818	4862354	7555	0.072
1983	801695	1446966	3446924	4508	0.100
1984	656553	860353	1971017	3008	0.124
1985	921365	623196	1399968	2646	0.144
1986	1357544	704853	1182651	2593	0.140
1987	1322467	985840	1274668	3066	0.128
1988	1234967	1057915	1480517	3352	0.126
1989	2908115	1031051	1633815	3840	0.121
1990	1861819	1964265	1937494	4980	0.092
1991	1913082	1671231	2424141	5005	0.094
1992	2093140	1584760	2385980	5178	0.085
1993	2367043	1671825	2496114	5434	0.077
1994	1613228	1846492	2576446	5216	0.070
1995	1743790	1470990	2479775	5105	0.072
1996	1748347	1434367	2378653	4815	0.074
1997	908710	1423495	2279695	4184	0.093
1998	605756	936324	1856918	2763	0.109
1999	372308	311921	716235	1690	0.102
2000	411246	314513	790708	1833	0.083
2001	375417	337554	858113	1984	0.076
2002	131246	324815	922531	2090	0.070
2003	294929	181983	946573	1974	0.071
2004	188200	227591	910048	1958	0.071
2005	477608	182179	892733	1874	0.071
2006	711699	331441	882927	2022	0.072
2007	413273	513299	971534	2362	0.068
2008	853429	399027	1094554	2494	0.059
2009	693645	616062	1195711	2539	0.053
2010	608835	588523	1272530	2156	0.055
2011	465219	516479	1120792	1562	0.068
2012	228870	399616	802718	1008	0.106
2013	251329	233118	512385	1165	0.096
2014	207389	219847	569439	1095	0.103
2015	163672	186544	539866	1076	0.105
2016	169765	153478	537101	1122	0.102
2017	131810	147097	541787	1122	0.100
2018	141800	123251	537994	1091	0.100
2019	250784	121250	524312	1065	0.101
2020	550402	182481	522840	1086	0.103

Table 20: Population abundances ( $n$ ) by crab stage in numbers of crab at the time of the survey (1 July, season 1) and mature male biomass (MMB) in tons on 15 February for the 2020 base model.

Year	$n_1$	$n_2$	$n_3$	MMB	CV MMB
1978	3122937	2004519	1667550	4565	0.177
1979	4231872	2371147	2291325	6466	0.123
1980	3797136	3181545	3465448	10173	0.084
1981	1436971	3206679	4831857	10634	0.063
1982	1605397	1832149	4862920	7554	0.072
1983	799872	1446836	3446713	4507	0.099
1984	655623	859410	1970674	3006	0.124
1985	920947	622428	1399170	2644	0.144
1986	1356725	704394	1181578	2590	0.140
1987	1321951	985240	1273467	3063	0.129
1988	1234325	1057437	1479166	3349	0.126
1989	2910079	1030542	1632387	3837	0.121
1990	1862455	1965213	1936234	4979	0.093
1991	1914728	1671906	2423614	5005	0.094
1992	2095878	1585932	2386042	5180	0.085
1993	2371914	1673774	2497020	5438	0.077
1994	1616875	1849901	2578653	5224	0.070
1995	1750933	1474195	2483692	5116	0.072
1996	1756356	1439471	2384211	4831	0.074
1997	914406	1429722	2287656	4208	0.093
1998	610090	941635	1867261	2784	0.108
1999	373770	312892	719259	1697	0.101
2000	413487	315665	793863	1840	0.083
2001	377241	339208	861544	1993	0.075
2002	131183	326398	926405	2099	0.070
2003	295947	182471	950601	1982	0.071
2004	188966	228330	913756	1965	0.071
2005	479186	182859	896277	1881	0.071
2006	715147	332562	886380	2030	0.072
2007	414453	515628	975313	2372	0.068
2008	855895	400471	1098998	2504	0.059
2009	696265	617941	1200387	2549	0.053
2010	612147	590641	1277639	2167	0.055
2011	469143	519068	1126466	1575	0.068
2012	231482	402707	809190	1022	0.105
2013	254986	235691	519700	1181	0.094
2014	210664	222775	577197	1112	0.101
2015	166431	189375	548146	1093	0.102
2016	173025	155983	545710	1141	0.100
2017	134486	149776	550552	1141	0.098
2018	145340	125652	546903	1109	0.097
2019	249238	124051	533288	1084	0.098
2020	183324	182525	531584	1145	0.109
2021	451372	164485	553444	1117	0.112

## Figures




Figure 1: Distribution of blue king crab (*Paralithodes platypus*) in the Gulf of Alaska, Bering Sea, and Aleutian Islands waters (shown in blue).




Figure 2: Blue king crab Registration Area Q (Bering Sea)




Figure 3: Data extent for the SMBKC assessment.




Figure 4: Trawl and pot-survey stations used in the SMBKC stock assessment.




Figure 5: Catches (in numbers) of male blue king crab > 90mm CL from the 2012-2021 NMFS trawl-survey at the 56 stations used to assess the SMBKC stock.




Figure 6: Fits to NMFS area-swept trawl estimates of total (> 90mm) male survey biomass for the base model only (16.0 ref for 2022 and 16.0 2020 accepted model). Error bars are plus and minus 2 standard deviations.






Figure 7: Comparisons of fits to CPUE from the ADFG pot surveys for model 16.0 the reference model in 2020 and 2022. Error bars are plus and minus 2 standard deviations.




Figure 8: Estimated recruitment 1979-2020 comparing ref model (16.0) for 2020 and 2022. The solid horizontal lines in the background represent the estimate of the average recruitment parameter ( $\bar{R}$ ) in each model scenario.




Figure 9: Sensitivity of new data in 2022 on estimated mature male biomass (MMB); 1978-2021.




Figure 10: Comparisons of the estimated stage-1 and stage-2 selectivities for the different model scenarios (the stage-3 selectivities are all fixed at 1). Estimated selectivities are shown for the directed pot fishery, the trawl bycatch fishery, the fixed bycatch fishery, the NMFS trawl survey, and the ADFG pot survey. Two selectivity periods are estimated in the directed pot fishery, from 1978-2008 and 2009-2021.

## Recruitment model scenarios




Figure 11: Estimated recruitment 1979-2020 comparing model alternatives. The solid horizontal lines in the background represent the estimate of the average recruitment parameter ( $\bar{R}$ ) in each model scenario. Note the high uncertainty in recruitment in the terminal year.




Figure 12: Comparisons of estimated mature male biomass (MMB) time series on 15 February during 1978-2021 for each of the model scenarios.




Figure 13: Time-varying natural mortality ( $M_t$ ). Estimated pulse period occurs in 1998/99 (i.e.  $M_{1998}$ ).




Figure 14: Comparisons of area-swept estimates of total (90+ mm CL) male survey biomass (tons) and model predictions for the model scenarios. The error bars are plus and minus 2 standard deviations.






Figure 15: Comparisons of total (90+ mm CL) male pot survey CPUEs and model predictions for the model scenarios. The error bars are plus and minus 2 standard deviations.




Figure 16: Standardized residuals for area-swept estimates of total male survey biomass for the model scenarios.




Figure 17: Standardized residuals for total male pot survey CPUEs for each of the GMACS model scenarios.




Figure 18: Observed and model estimated size-frequencies of SMBKC by year retained in the directed pot fishery for the model scenarios.




Figure 19: Observed and model estimated size-frequencies of discarded male SMBKC by year in the NMFS trawl survey for the model scenarios.




Figure 20: Observed and model estimated size-frequencies of discarded SMBKC by year in the ADFG pot survey for the model scenarios.




Figure 21: Bubble plots of residuals by stage and year for the all the size composition data sets (ADFG pot survey, NMFS trawl survey, and the directed pot fishery) for SMBKC in the 'base' model (16.0).




Figure 22: Bubble plots of residuals by stage and year for the all the size composition data sets (NMFS trawl survey, and the directed pot fishery) for SMBKC in the M 0.21 model (22.0a).




Figure 23: Bubble plots of residuals by stage and year for the all the size composition data sets (NMFS trawl survey, and the directed pot fishery) for SMBKC in the M 0.26 model (22.0b).






Figure 24: Bubble plots of residuals by stage and year for the all the size composition data sets (NMFS trawl survey, and the directed pot fishery) for SMBKC in model (22.0c) which removes the increased M time block in 1998 and has a fixed M of 0.18.




Figure 25: Comparison of observed and model predicted retained catch and bycatches in each of the GMACS models. Note that difference in units between each of the panels, some panels are expressed in numbers of crab, some as biomass (tons).




Figure 26: Fishing mortality estimates from the base model (16.0) for directed and bycatch fleets




Figure 27: Comparison of mature male biomass relative to the dynamic B zero value, (15 February, 1978-2019) for each of the model scenarios.

# Appendix A: SMBKC Model Description

## 1. Introduction

The GMACS model has been specified to account only for male crab  $\geq 90$  mm in carapace length (CL). These are partitioned into three stages (size- classes) determined by CL measurements of (1) 90-104 mm, (2) 105-119 mm, and (3) 120+ mm. For management of the St. Matthew Island blue king crab (SMBKC) fishery, 120 mm CL is used as the proxy value for the legal measurement of 5.5 inch carapace width (CW), whereas 105 mm CL is the management proxy for mature-male size (state regulation *5 AAC 34.917 (d)*). Accordingly, within the model only stage-3 crab are retained in the directed fishery, and stage-2 and stage-3 crab together comprise the collection of mature males. Some justification for the 105 mm value is presented in Pengilly and Schmidt (1995), who used it in developing the current regulatory SMBKC harvest strategy. The term “recruit” here designates recruits to the model, i.e., annual new stage-1 crab, rather than recruits to the fishery. The following description of model structure reflects the GMACS base model configuration.

## 2. Model Population Dynamics

Within the model, the beginning of the crab year is assumed contemporaneous with the NMFS trawl survey, nominally assigned a date of 1 July. Although the timing of the fishery is different each year, MMB is estimated at 15 February, which is the reference date for calculation of federal management biomass quantities. To accommodate this, each model year is split into 5 seasons ( $t$ ) and a proportion of the natural mortality ( $\tau_t$ ), scaled relative to the portions of the year, is applied in each of these seasons where  $\sum_{t=1}^{t=5} \tau_t = 1$ . Each model year consists of the following processes with time-breaks denoted here by “Seasons.” However, it is important to note that actual seasons are survey-to-fishery, fishery-to Feb 15, and Feb 15 to July 1. The following breakdown accounts for events and fishing mortality treatments:

1. Season 1 (survey period)
  - Beginning of the SMBKC fishing year (1 July)
  - $\tau_1 = 0$
  - Surveys
2. Season 2 (natural mortality until pulse fishery)
  - $\tau_2$  ranges from 0.05 to 0.44 depending on the time of year the fishery begins each year (i.e., a higher value indicates the fishery begins later in the year; see Table reftab:smbkc-fishery)
3. Season 3 (pulse fishery)
  - $\tau_3 = 0$
  - fishing mortality applied
4. Season 4 (natural mortality until spawning)
  - $\tau_4 = 0.63 - \sum_{i=1}^{i=4} \tau_i$
  - Calculate MMB (15 February)
5. Season 5 (natural mortality and somatic growth through to June 30th)
  - $\tau_5 = 0.37$
  - Growth and molting
  - Recruitment (all to stage-1)

The proportion of natural mortality ( $\tau_t$ ) applied during each season in the model is provided in Table 21. The beginning of the year (1 July) to the date that MMB is measured (15 February) is 63% of the year. Therefore 63% of the natural mortality must be applied before the MMB is calculated. Because the timing of the fishery is different each year,  $\tau_2$  varies and thus  $\tau_4$  varies also.

With boldface lower-case letters indicating vector quantities we designate the vector of stage abundances during season  $t$  and year  $y$  as

$$\mathbf{n}_{t,y} = n_{l,t,y} = [n_{1,t,y}, n_{2,t,y}, n_{3,t,y}]^\top. \quad (2)$$

The number of new crab, or recruits, of each stage entering the model each season  $t$  and year  $y$  is represented as the vector  $\mathbf{r}_{t,y}$ . The SMBKC formulation of GMACS specifies recruitment to stage-1 only during season  $t = 5$ , thus the recruitment size distribution is

$$\phi_l = [1, 0, 0]^\top, \quad (3)$$

and the recruitment is

$$\mathbf{r}_{t,y} = \begin{cases} 0 & \text{for } t < 5 \\ \bar{R}\phi_l\delta_y^R & \text{for } t = 5. \end{cases} \quad (4)$$

where  $\bar{R}$  is the average annual recruitment and  $\delta_y^R$  are the recruitment deviations each year  $y$

$$\delta_y^R \sim \mathcal{N}(0, \sigma_R^2). \quad (5)$$

Using boldface upper-case letters to indicate a matrix, we describe the size transition matrix  $\mathbf{G}$  as

$$\mathbf{G} = \begin{bmatrix} 1 - \pi_{12} - \pi_{13} & \pi_{12} & \pi_{13} \\ 0 & 1 - \pi_{23} & \pi_{23} \\ 0 & 0 & 1 \end{bmatrix}, \quad (6)$$

with  $\pi_{jk}$  equal to the proportion of stage- $j$  crab that molt and grow into stage- $k$  within a season or year.

The natural mortality each season  $t$  and year  $y$  is

$$M_{t,y} = \bar{M}\tau_t + \delta_y^M \text{ where } \delta_y^M \sim \mathcal{N}(0, \sigma_M^2) \quad (7)$$

Fishing mortality by year  $y$  and season  $t$  is denoted  $F_{t,y}$  and calculated as

$$F_{t,y} = F_{t,y}^{\text{df}} + F_{t,y}^{\text{tb}} + F_{t,y}^{\text{fb}} \quad (8)$$

where  $F_{t,y}^{\text{df}}$  is the fishing mortality associated with the directed fishery,  $F_{t,y}^{\text{tb}}$  is the fishing mortality associated with the trawl bycatch fishery,  $F_{t,y}^{\text{fb}}$  is the fishing mortality associated with the fixed bycatch fishery. Each of these are derived as

$$\begin{aligned} F_{t,y}^{\text{df}} &= \bar{F}^{\text{df}} + \delta_{t,y}^{\text{df}} & \text{where } \delta_{t,y}^{\text{df}} &\sim \mathcal{N}(0, \sigma_{\text{df}}^2), \\ F_{t,y}^{\text{tb}} &= \bar{F}^{\text{tb}} + \delta_{t,y}^{\text{tb}} & \text{where } \delta_{t,y}^{\text{tb}} &\sim \mathcal{N}(0, \sigma_{\text{tb}}^2), \\ F_{t,y}^{\text{fb}} &= \bar{F}^{\text{fb}} + \delta_{t,y}^{\text{fb}} & \text{where } \delta_{t,y}^{\text{fb}} &\sim \mathcal{N}(0, \sigma_{\text{fb}}^2), \end{aligned} \quad (9)$$

where  $\delta_{t,y}^{\text{df}}$ ,  $\delta_{t,y}^{\text{tb}}$ , and  $\delta_{t,y}^{\text{fb}}$  are the fishing mortality deviations for each of the fisheries, each season  $t$  during each year  $y$ ,  $\bar{F}^{\text{df}}$ ,  $\bar{F}^{\text{tb}}$ , and  $\bar{F}^{\text{fb}}$  are the average fishing mortalities for each fishery. The total mortality  $Z_{l,t,y}$  represents the combination of natural mortality  $M_{t,y}$  and fishing mortality  $F_{t,y}$  during season  $t$  and year  $y$

$$\mathbf{Z}_{t,y} = Z_{l,t,y} = M_{t,y} + F_{t,y}. \quad (10)$$

The survival matrix  $\mathbf{S}_{t,y}$  during season  $t$  and year  $y$  is

$$\mathbf{S}_{t,y} = \begin{bmatrix} 1 - e^{-Z_{1,t,y}} & 0 & 0 \\ 0 & 1 - e^{-Z_{2,t,y}} & 0 \\ 0 & 0 & 1 - e^{-Z_{3,t,y}} \end{bmatrix}. \quad (11)$$

The basic population dynamics underlying GMACS can thus be described as

$$\begin{aligned} \mathbf{n}_{t+1,y} &= \mathbf{S}_{t,y}\mathbf{n}_{t,y}, & \text{if } t < 5 \\ \mathbf{n}_{t,y+1} &= \mathbf{G}\mathbf{S}_{t,y}\mathbf{n}_{t,y} + \mathbf{r}_{t,y} & \text{if } t = 5. \end{aligned} \quad (12)$$

### 3. Model Data

Data inputs used in model estimation are listed in Table 22.

### 4. Model Parameters

Table 23 lists fixed (externally determined) parameters used in model computations. In all scenarios, the stage-transition matrix is

$$\mathbf{G} = \begin{bmatrix} 0.2 & 0.7 & 0.1 \\ 0 & 0.4 & 0.6 \\ 0 & 0 & 1 \end{bmatrix} \quad (13)$$

which is the combination of the growth matrix and molting probabilities.

Estimated parameters are listed in Table 24 and include an estimated natural mortality deviation parameter in 1998/99 ( $\delta_{1998}^M$ ) assuming an anomalous mortality event in that year, as hypothesized by Zheng and Kruse (2002), with natural mortality otherwise fixed at  $0.18 \text{ yr}^{-1}$ .

### 5. Model Objective Function and Weighting Scheme

The objective function consists of the sum of several “negative log-likelihood” terms characterizing the hypothesized error structure of the principal data inputs (Table 18). A lognormal distribution is assumed to characterize the catch data and is modelled as

$$\sigma_{t,y}^{\text{catch}} = \sqrt{\log \left( 1 + \left( CV_{t,y}^{\text{catch}} \right)^2 \right)} \quad (14)$$

$$\delta_{t,y}^{\text{catch}} = \mathcal{N} \left( 0, \left( \sigma_{t,y}^{\text{catch}} \right)^2 \right) \quad (15)$$

where  $\delta_{t,y}^{\text{catch}}$  is the residual catch. The relative abundance data is also assumed to be lognormally distributed

$$\sigma_{t,y}^{\text{I}} = \frac{1}{\lambda} \sqrt{\log \left( 1 + \left( CV_{t,y}^{\text{I}} \right)^2 \right)} \quad (16)$$

$$\delta_{t,y}^{\text{I}} = \log \left( I^{\text{obs}} / I^{\text{pred}} \right) / \sigma_{t,y}^{\text{I}} + 0.5 \sigma_{t,y}^{\text{I}} \quad (17)$$

and the likelihood is

$$\sum \log \left( \delta_{t,y}^{\text{I}} \right) + \sum 0.5 \left( \sigma_{t,y}^{\text{I}} \right)^2 \quad (18)$$

GMACS calculates standard deviation of the normalised residual (SDNR) values and median of the absolute residual (MAR) values for all abundance indices and size compositions to help the user come up with reasonable likelihood weights. For an abundance data set to be well fitted, the SDNR should not be much greater than 1 (a value much less than 1, which means that the data set is fitted better than was expected, is not a cause for concern). What is meant by “much greater than 1” depends on  $m$  (the number of years in the data set). Francis (2011) suggests upper limits of 1.54, 1.37, and 1.26 for  $m = 5, 10,$  and  $20,$  respectively. Although an SDNR not much greater than 1 is a necessary condition for a good fit, it is not sufficient. It is important to plot the observed and expected abundances to ensure that the fit is good.

GMACS also calculates Francis weights for each of the size composition data sets supplied (Francis 2011). If the user wishes to use the Francis iterative re-weighting method, first the weights applied to the abundance indices should be adjusted by trial and error until the SDNR (and/or MAR) are adequate. Then the Francis weights supplied by GMACS should be used as the new likelihood weights for each of the size composition data sets the next time the model is run. The user can then iteratively adjust the abundance index and size composition weights until adequate SDNR (and/or MAR) values are achieved, given the Francis weights.

## 6. Estimation

The model was implemented using the software AD Model Builder (Fournier et al. 2012), with parameter estimation by minimization of the model objective function using automatic differentiation. Parameter estimates and standard deviations provided in this document are AD Model Builder reported values assuming maximum likelihood theory asymptotics.



Table 21: Proportion of the natural mortality ( $\tau_t$ ) that is applied during each season ( $t$ ) in the model.

Year	Season 1	Season 2	Season 3	Season 4	Season 5
1978	0.00	0.07	0.00	0.56	0.37
1979	0.00	0.06	0.00	0.57	0.37
1980	0.00	0.07	0.00	0.56	0.37
1981	0.00	0.05	0.00	0.58	0.37
1982	0.00	0.07	0.00	0.56	0.37
1983	0.00	0.12	0.00	0.51	0.37
1984	0.00	0.10	0.00	0.53	0.37
1985	0.00	0.14	0.00	0.49	0.37
1986	0.00	0.14	0.00	0.49	0.37
1987	0.00	0.14	0.00	0.49	0.37
1988	0.00	0.14	0.00	0.49	0.37
1989	0.00	0.14	0.00	0.49	0.37
1990	0.00	0.14	0.00	0.49	0.37
1991	0.00	0.18	0.00	0.45	0.37
1992	0.00	0.14	0.00	0.49	0.37
1993	0.00	0.18	0.00	0.45	0.37
1994	0.00	0.18	0.00	0.45	0.37
1995	0.00	0.18	0.00	0.45	0.37
1996	0.00	0.18	0.00	0.45	0.37
1997	0.00	0.18	0.00	0.45	0.37
1998	0.00	0.18	0.00	0.45	0.37
1999	0.00	0.18	0.00	0.45	0.37
2000	0.00	0.18	0.00	0.45	0.37
2001	0.00	0.18	0.00	0.45	0.37
2002	0.00	0.18	0.00	0.45	0.37
2003	0.00	0.18	0.00	0.45	0.37
2004	0.00	0.18	0.00	0.45	0.37
2005	0.00	0.18	0.00	0.45	0.37
2006	0.00	0.18	0.00	0.45	0.37
2007	0.00	0.18	0.00	0.45	0.37
2008	0.00	0.18	0.00	0.45	0.37
2009	0.00	0.44	0.00	0.19	0.37
2010	0.00	0.44	0.00	0.19	0.37
2011	0.00	0.44	0.00	0.19	0.37
2012	0.00	0.44	0.00	0.19	0.37
2013	0.00	0.44	0.00	0.19	0.37
2014	0.00	0.44	0.00	0.19	0.37
2015	0.00	0.44	0.00	0.19	0.37
2016	0.00	0.44	0.00	0.19	0.37
2017	0.00	0.44	0.00	0.19	0.37
2018	0.00	0.44	0.00	0.19	0.37
2019	0.00	0.44	0.00	0.19	0.37
2020	0.00	0.44	0.00	0.19	0.37

Table 22: Data inputs used in model estimation.

Data	Years	Source
Directed pot-fishery retained-catch number (not biomass)	1978/79 - 1998/99 2009/10 - 2015/16	Fish tickets (fishery closed 1999/00 - 2008/09 and 2016/17 - 2018/19)
Groundfish trawl bycatch biomass	1992/93 - 2018/19	NMFS groundfish observer program
Groundfish fixed-gear bycatch biomass	1992/93 - 2018/19	NMFS groundfish observer program
NMFS trawl-survey biomass index (area-swept estimate) and CV	1978-2019	NMFS EBS trawl survey
ADF&G pot-survey abundance index (CPUE) and CV	1995-2018	ADF&G SMBKC pot survey
NMFS trawl-survey stage proportions and total number of measured crab	1978-2019	NMFS EBS trawl survey
ADF&G pot-survey stage proportions and total number of measured crab	1995-2018	ADF&G SMBKC pot survey
Directed pot-fishery stage proportions and total number of measured crab	1990/91 - 1998/99 2009/10 - 2015/16	ADF&G crab observer program (fishery closed 1999/00 - 2008/09 and 2016/17 - 2018/19)

Table 23: Fixed model parameters for all scenarios.

Parameter	Symbol	Value	Source/rationale
Trawl-survey catchability	$q$	1.0	Default
Natural mortality	$M$	0.18 yr <sup>-1</sup>	NPFMC (2007)
Size transition matrix	$\mathbf{G}$	Equation 13	Otto and Cummiskey (1990)
Stage-1 and stage-2 mean weights	$w_1, w_2$	0.7, 1.2 kg	Length-weight equation (B. Foy, NMFS) applied to stage midpoints
Stage-3 mean weight	$w_{3,y}$	Depends on year	Fishery reported average retained weight from fish tickets, or its average, and mean weights of legal males
Recruitment SD	$\sigma_R$	1.2	High value
Natural mortality SD	$\sigma_M$	10.0	High value (basically free parameter)
Directed fishery handling mortality		0.2	2010 Crab SAFE
Groundfish trawl handling mortality		0.8	2010 Crab SAFE
Groundfish fixed-gear handling mortality		0.5	2010 Crab SAFE

Table 24: The lower bound (LB), upper bound (UB), initial value, prior, and estimation phase for each estimated model parameter.

Parameter	LB	Initial value	UB	Prior	Phase
Average recruitment $\log(\bar{R})$	-7	10.0	20	Uniform(-7,20)	1
Stage-1 initial numbers $\log(n_1^0)$	5	14.5	20	Uniform(5,20)	1
Stage-2 initial numbers $\log(n_2^0)$	5	14.0	20	Uniform(5,20)	1
Stage-3 initial numbers $\log(n_3^0)$	5	13.5	20	Uniform(5,20)	1
ADF&G pot survey catchability $q$	0	3.0	5	Uniform(0,5)	1
Stage-1 directed fishery selectivity 1978-2008	0	0.4	1	Uniform(0,1)	3
Stage-2 directed fishery selectivity 1978-2008	0	0.7	1	Uniform(0,1)	3
Stage-1 directed fishery selectivity 2009-2017	0	0.4	1	Uniform(0,1)	3
Stage-2 directed fishery selectivity 2009-2017	0	0.7	1	Uniform(0,1)	3
Stage-1 NMFS trawl survey selectivity	0	0.4	1	Uniform(0,1)	4
Stage-2 NMFS trawl survey selectivity	0	0.7	1	Uniform(0,1)	4
Stage-1 ADF&G pot survey selectivity	0	0.4	1	Uniform(0,1)	4
Stage-2 ADF&G pot survey selectivity	0	0.7	1	Uniform(0,1)	4
Natural mortality deviation during 1998 $\delta_{1998}^M$	-3	0.0	3	Normal(0, $\sigma_M^2$ )	4
Recruitment deviations $\delta_y^R$	-7	0.0	7	Normal(0, $\sigma_R^2$ )	3
Average directed fishery fishing mortality $\bar{F}^{\text{df}}$	-	0.2	-	-	1
Average trawl bycatch fishing mortality $\bar{F}^{\text{tb}}$	-	0.001	-	-	1
Average fixed gear bycatch fishing mortality $\bar{F}^{\text{fb}}$	-	0.001	-	-	1



```

0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370
0.000 0.440 0.000 0.190 0.370 # (updated)
0.000 0.440 0.000 0.190 0.370 # (updated 4-14-22)
#0 0.0025 0 0.6245 0.373
# Fishing fleet names (delimited with spaces no spaces in names)
Pot_Fishery Trawl_Bycatch Fixed_bycatch
# Survey names (delimited with spaces no spaces in names)
NMFS_Trawl ADFG_Pot
# Are the fleets instantaneous (0) or continuous (1)
1 1 1 1 1
# Number of catch data frames
4
# Number of rows in each data frame
27 18 30 30 # (updated - all should increase 1 if value for current year NO placeholder for direct fishery if closed)
## CATCH DATA
## Type of catch: 1 = retained, 2 = discard
## Units of catch: 1 = biomass, 2 = numbers
## for SMBKC Units are in number of crab for landed & 1000 kg for discards.
## Male Retained
# year seas fleet sex obs cv type units mult effort discard_mortality
1978 3 1 1 436126 0.03 1 2 1 0 0.2
1979 3 1 1 52966 0.03 1 2 1 0 0.2
1980 3 1 1 33162 0.03 1 2 1 0 0.2
1981 3 1 1 1045619 0.03 1 2 1 0 0.2
1982 3 1 1 1935886 0.03 1 2 1 0 0.2
1983 3 1 1 1931990 0.03 1 2 1 0 0.2
1984 3 1 1 841017 0.03 1 2 1 0 0.2
1985 3 1 1 436021 0.03 1 2 1 0 0.2
1986 3 1 1 219548 0.03 1 2 1 0 0.2
1987 3 1 1 227447 0.03 1 2 1 0 0.2
1988 3 1 1 280401 0.03 1 2 1 0 0.2
1989 3 1 1 247641 0.03 1 2 1 0 0.2
1990 3 1 1 391405 0.03 1 2 1 0 0.2
1991 3 1 1 726519 0.03 1 2 1 0 0.2
1992 3 1 1 545222 0.03 1 2 1 0 0.2
1993 3 1 1 630353 0.03 1 2 1 0 0.2
1994 3 1 1 827015 0.03 1 2 1 0 0.2
1995 3 1 1 666905 0.03 1 2 1 0 0.2
1996 3 1 1 660665 0.03 1 2 1 0 0.2
1997 3 1 1 939822 0.03 1 2 1 0 0.2
1998 3 1 1 635370 0.03 1 2 1 0 0.2
2009 3 1 1 103376 0.03 1 2 1 0 0.2
2010 3 1 1 298669 0.03 1 2 1 0 0.2
2011 3 1 1 437862 0.03 1 2 1 0 0.2
2012 3 1 1 379386 0.03 1 2 1 0 0.2
2014 3 1 1 69109 0.03 1 2 1 0 0.2
2015 3 1 1 24407 0.03 1 2 1 0 0.2
#2016 3 1 1 10.000 0.03 1 2 1 0 0.2
#2017 3 1 1 10.000 0.03 1 2 1 0 0.2
#2018 3 1 1 10.000 0.03 1 2 1 0 0.2 # placeholder no fishery
# Male discards Pot fishery
1990 3 1 1 254.9787861 0.6 2 1 1 0 0.2
1991 3 1 1 531.4483252 0.6 2 1 1 0 0.2
1992 3 1 1 1050.387026 0.6 2 1 1 0 0.2
1993 3 1 1 951.4626128 0.6 2 1 1 0 0.2
1994 3 1 1 1210.764588 0.6 2 1 1 0 0.2
1995 3 1 1 363.112032 0.6 2 1 1 0 0.2
1996 3 1 1 528.5244687 0.6 2 1 1 0 0.2
1997 3 1 1 1382.825328 0.6 2 1 1 0 0.2
1998 3 1 1 781.1032977 0.6 2 1 1 0 0.2
2009 3 1 1 123.3712279 0.2 2 1 1 0 0.2
2010 3 1 1 304.6562225 0.2 2 1 1 0 0.2
2011 3 1 1 481.3572126 0.2 2 1 1 0 0.2
2012 3 1 1 437.3360731 0.2 2 1 1 0 0.2
2014 3 1 1 45.4839749 0.2 2 1 1 0 0.2
2015 3 1 1 21.19378597 0.2 2 1 1 0 0.2

```

2016	3	1	1	0.021193786	0.2	2	1	1	0	0.2
2017	3	1	1	0.021193786	0.2	2	1	1	0	0.2
2018	3	1	1	0.214868020	0.2	2	1	1	0	0.2 # (updated)
#2019	3	1	1	0.0	0.2	2	1	1	0	0.2
#2020	3	1	1	0.0	0.2	2	1	1	0	0.2
#2021	3	1	1	0.0	0.2	2	1	1	0	0.2
# Trawl fishery discards										
1991	2	2	1	3.538	0.31	2	1	1	0	0.8
1992	2	2	1	1.996	0.31	2	1	1	0	0.8
1993	2	2	1	1.542	0.31	2	1	1	0	0.8
1994	2	2	1	0.318	0.31	2	1	1	0	0.8
1995	2	2	1	0.635	0.31	2	1	1	0	0.8
1996	2	2	1	0.500	0.31	2	1	1	0	0.8
1997	2	2	1	0.500	0.31	2	1	1	0	0.8
1998	2	2	1	0.500	0.31	2	1	1	0	0.8
1999	2	2	1	0.500	0.31	2	1	1	0	0.8
2000	2	2	1	0.500	0.31	2	1	1	0	0.8
2001	2	2	1	0.500	0.31	2	1	1	0	0.8
2002	2	2	1	0.726	0.31	2	1	1	0	0.8
2003	2	2	1	0.998	0.31	2	1	1	0	0.8
2004	2	2	1	0.091	0.31	2	1	1	0	0.8
2005	2	2	1	0.500	0.31	2	1	1	0	0.8
2006	2	2	1	2.812	0.31	2	1	1	0	0.8
2007	2	2	1	0.045	0.31	2	1	1	0	0.8
2008	2	2	1	0.272	0.31	2	1	1	0	0.8
2009	2	2	1	0.638	0.31	2	1	1	0	0.8
2010	2	2	1	0.360	0.31	2	1	1	0	0.8
2011	2	2	1	0.170	0.31	2	1	1	0	0.8
2012	2	2	1	0.011	0.31	2	1	1	0	0.8
2013	2	2	1	0.163	0.31	2	1	1	0	0.8
2014	2	2	1	0.010	0.31	2	1	1	0	0.8
2015	2	2	1	0.010	0.31	2	1	1	0	0.8
2016	2	2	1	0.229	0.31	2	1	1	0	0.8
2017	2	2	1	0.048	0.31	2	1	1	0	0.8 # updated in 2020 was 0.052, now 0.48?
2018	2	2	1	0.001	0.31	2	1	1	0	0.8 # (data is 0 but small value for placeholder)
2019	2	2	1	0.030	0.31	2	1	1	0	0.8 # (updated )
2020	2	2	1	0.001	0.31	2	1	1	0	0.8 # (4-14-22)
# Fixed fishery discards										
1991	2	3	1	0.045	0.31	2	1	1	0	0.5
1992	2	3	1	2.268	0.31	2	1	1	0	0.5
1993	2	3	1	0.500	0.31	2	1	1	0	0.5
1994	2	3	1	0.091	0.31	2	1	1	0	0.5
1995	2	3	1	0.136	0.31	2	1	1	0	0.5
1996	2	3	1	0.045	0.31	2	1	1	0	0.5
1997	2	3	1	0.181	0.31	2	1	1	0	0.5
1998	2	3	1	0.907	0.31	2	1	1	0	0.5
1999	2	3	1	1.361	0.31	2	1	1	0	0.5
2000	2	3	1	0.500	0.31	2	1	1	0	0.5
2001	2	3	1	0.862	0.31	2	1	1	0	0.5
2002	2	3	1	0.408	0.31	2	1	1	0	0.5
2003	2	3	1	1.134	0.31	2	1	1	0	0.5
2004	2	3	1	0.635	0.31	2	1	1	0	0.5
2005	2	3	1	0.590	0.31	2	1	1	0	0.5
2006	2	3	1	1.451	0.31	2	1	1	0	0.5
2007	2	3	1	69.717	0.31	2	1	1	0	0.5
2008	2	3	1	6.622	0.31	2	1	1	0	0.5
2009	2	3	1	7.522	0.31	2	1	1	0	0.5
2010	2	3	1	9.564	0.31	2	1	1	0	0.5
2011	2	3	1	0.796	0.31	2	1	1	0	0.5
2012	2	3	1	0.739	0.31	2	1	1	0	0.5
2013	2	3	1	0.341	0.31	2	1	1	0	0.5
2014	2	3	1	0.490	0.31	2	1	1	0	0.5
2015	2	3	1	0.711	0.31	2	1	1	0	0.5
2016	2	3	1	1.630	0.31	2	1	1	0	0.5 # updated from 1.632
2017	2	3	1	5.935	0.31	2	1	1	0	0.5 # updates was 6.032
2018	2	3	1	1.224	0.31	2	1	1	0	0.5 # updated was 1.281
2019	2	3	1	1.124	0.31	2	1	1	0	0.5 # (updated - bycatch_groundfish.R)
2020	2	3	1	0.671	0.31	2	1	1	0	0.5 # (4-14-22 - bycatch_groundfish.R)
## RELATIVE ABUNDANCE DATA										
## Units of abundance: 1 = biomass, 2 = numbers										
## for SMBKC pot survey Units are in crabs for Abundance.										
## Number of relative abundance indicies										

```

2
# Index Type (1=Selectivity; 2=retention)
# AEPAP
1 1
## Number of rows in each index, need to update when survey data is added
54
# Survey data (abundance indices, units are mt for trawl survey and crab/potlift for pot survey)
# Index, Year, Seas, Fleet, Sex, Maturity, Abundance, CV abundance units timing
1 1978 1 4 1 0 6832.819 0.394 1 0
1 1979 1 4 1 0 7989.881 0.463 1 0
1 1980 1 4 1 0 9986.83 0.507 1 0
1 1981 1 4 1 0 6551.132 0.402 1 0
1 1982 1 4 1 0 16221.933 0.344 1 0
1 1983 1 4 1 0 9634.25 0.298 1 0
1 1984 1 4 1 0 4071.218 0.179 1 0
1 1985 1 4 1 0 3110.541 0.21 1 0
1 1986 1 4 1 0 1416.849 0.388 1 0
1 1987 1 4 1 0 2278.917 0.291 1 0
1 1988 1 4 1 0 3158.169 0.252 1 0
1 1989 1 4 1 0 6338.622 0.271 1 0
1 1990 1 4 1 0 6730.13 0.274 1 0
1 1991 1 4 1 0 6948.184 0.248 1 0
1 1992 1 4 1 0 7093.272 0.201 1 0
1 1993 1 4 1 0 9548.459 0.169 1 0
1 1994 1 4 1 0 6539.133 0.176 1 0
1 1995 1 4 1 0 5703.591 0.178 1 0
1 1996 1 4 1 0 9410.403 0.241 1 0
1 1997 1 4 1 0 10924.107 0.337 1 0
1 1998 1 4 1 0 7976.839 0.355 1 0
1 1999 1 4 1 0 1594.546 0.182 1 0
1 2000 1 4 1 0 2096.795 0.31 1 0
1 2001 1 4 1 0 2831.44 0.245 1 0
1 2002 1 4 1 0 1732.599 0.32 1 0
1 2003 1 4 1 0 1566.675 0.336 1 0
1 2004 1 4 1 0 1523.869 0.305 1 0
1 2005 1 4 1 0 1642.017 0.371 1 0
1 2006 1 4 1 0 3893.875 0.334 1 0
1 2007 1 4 1 0 6470.773 0.385 1 0
1 2008 1 4 1 0 4654.473 0.284 1 0
1 2009 1 4 1 0 6301.47 0.256 1 0
1 2010 1 4 1 0 11130.898 0.466 1 0
1 2011 1 4 1 0 10931.232 0.558 1 0
1 2012 1 4 1 0 6200.219 0.339 1 0
1 2013 1 4 1 0 2287.557 0.217 1 0
1 2014 1 4 1 0 6029.22 0.449 1 0
1 2015 1 4 1 0 5877.433 0.77 1 0
1 2016 1 4 1 0 3485.909 0.393 1 0
1 2017 1 4 1 0 1793.76 0.599 1 0
1 2018 1 4 1 0 1730.742 0.281 1 0
1 2019 1 4 1 0 3170.467 0.337 1 0 # (updated - EBSsurvey_analysis.R)
1 2021 1 4 1 0 1929.298 0.427 1 0 # updated 4-14-22
2 1995 1 5 1 0 12042 0.13 2 0
2 1998 1 5 1 0 12531 0.06 2 0
2 2001 1 5 1 0 8477 0.08 2 0
2 2004 1 5 1 0 1667 0.15 2 0
2 2007 1 5 1 0 8643 0.09 2 0
2 2010 1 5 1 0 10209 0.13 2 0
2 2013 1 5 1 0 5643 0.19 2 0
2 2015 1 5 1 0 2805 0.18 2 0
2 2016 1 5 1 0 2378 0.186 2 0
2 2017 1 5 1 0 1689 0.25 2 0
2 2018 1 5 1 0 745 0.14 2 0 # no smbk pot survey in 2019, 2020, 2021
## Number of length frequency matrices
3
## Number of rows in each matrix
15 43 11 # (updated 4-14-22)
## Number of bins in each matrix (columns of size data)
3 3 3
## SIZE COMPOSITION DATA FOR ALL FLEETS
## SIZE COMP LEGEND
## Sex: 1 = male, 2 = female, 0 = both sexes combined
## Type of composition: 1 = retained, 2 = discard, 0 = total composition

```

```

## Maturity state: 1 = immature, 2 = mature, 0 = both states combined
## Shell condition: 1 = new shell, 2 = old shell, 0 = both shell types combined
##length proportions of pot discarded males
##Year, Seas, Fleet, Sex, Type, Shell, Maturity, Nsamp, DataVec
1990 3 1 1 0 0 0 15 0.1133 0.3933 0.4933
1991 3 1 1 0 0 0 25 0.1329 0.1768 0.6902
1992 3 1 1 0 0 0 25 0.1905 0.2677 0.5417
1993 3 1 1 0 0 0 25 0.2807 0.2097 0.5096
1994 3 1 1 0 0 0 25 0.2942 0.2714 0.4344
1995 3 1 1 0 0 0 25 0.1478 0.2127 0.6395
1996 3 1 1 0 0 0 25 0.1595 0.2229 0.6176
1997 3 1 1 0 0 0 25 0.1818 0.2053 0.6128
1998 3 1 1 0 0 0 25 0.1927 0.2162 0.5911
2009 3 1 1 0 0 0 50 0.1413 0.3235 0.5352
2010 3 1 1 0 0 0 50 0.1314 0.3152 0.5534
2011 3 1 1 0 0 0 50 0.1314 0.3051 0.5636
2012 3 1 1 0 0 0 50 0.1417 0.3178 0.5406
2014 3 1 1 0 0 0 50 0.0939 0.2275 0.6786
2015 3 1 1 0 0 0 50 0.1148 0.2518 0.6333 #no fishery so not updated
##length proportions of trawl survey males
##Year, Seas, Fleet, Sex, Type, Shell, Maturity, Nsamp, DataVec
1978 1 4 1 0 0 0 50 0.3865 0.3478 0.2657
1979 1 4 1 0 0 0 50 0.4281 0.3190 0.2529
1980 1 4 1 0 0 0 50 0.3588 0.3220 0.3192
1981 1 4 1 0 0 0 50 0.1219 0.3065 0.5716
1982 1 4 1 0 0 0 50 0.1671 0.2435 0.5893
1983 1 4 1 0 0 0 50 0.1752 0.2726 0.5522
1984 1 4 1 0 0 0 50 0.1823 0.2085 0.6092
1985 1 4 1 0 0 0 46.5 0.2023 0.2010 0.5967
1986 1 4 1 0 0 0 23 0.1984 0.4364 0.3652
1987 1 4 1 0 0 0 35.5 0.1944 0.3779 0.4277
1988 1 4 1 0 0 0 40.5 0.1879 0.3737 0.4384
1989 1 4 1 0 0 0 50 0.4246 0.2259 0.3496
1990 1 4 1 0 0 0 50 0.2380 0.2332 0.5288
1991 1 4 1 0 0 0 50 0.2274 0.3300 0.4426
1992 1 4 1 0 0 0 50 0.2263 0.2911 0.4826
1993 1 4 1 0 0 0 50 0.2296 0.2759 0.4945
1994 1 4 1 0 0 0 50 0.1989 0.2926 0.5085
1995 1 4 1 0 0 0 50 0.2593 0.3005 0.4403
1996 1 4 1 0 0 0 50 0.1998 0.3054 0.4948
1997 1 4 1 0 0 0 50 0.1622 0.3102 0.5275
1998 1 4 1 0 0 0 50 0.1276 0.3212 0.5511
1999 1 4 1 0 0 0 26 0.2224 0.2214 0.5562
2000 1 4 1 0 0 0 30.5 0.2154 0.2180 0.5665
2001 1 4 1 0 0 0 45.5 0.2253 0.2699 0.5048
2002 1 4 1 0 0 0 19 0.1127 0.2346 0.6527
2003 1 4 1 0 0 0 32.5 0.3762 0.2345 0.3893
2004 1 4 1 0 0 0 24 0.2488 0.1848 0.5663
2005 1 4 1 0 0 0 21 0.2825 0.2744 0.4431
2006 1 4 1 0 0 0 50 0.3276 0.2293 0.4431
2007 1 4 1 0 0 0 50 0.4394 0.3525 0.2081
2008 1 4 1 0 0 0 50 0.3745 0.2219 0.4036
2009 1 4 1 0 0 0 50 0.3057 0.4202 0.2741
2010 1 4 1 0 0 0 50 0.4081 0.3371 0.2548
2011 1 4 1 0 0 0 50 0.2179 0.3940 0.3881
2012 1 4 1 0 0 0 50 0.1573 0.4393 0.4034
2013 1 4 1 0 0 0 37 0.2100 0.2834 0.5065
2014 1 4 1 0 0 0 50 0.1738 0.3912 0.4350
2015 1 4 1 0 0 0 50 0.2340 0.2994 0.4666
2016 1 4 1 0 0 0 50 0.2255 0.2780 0.4965
2017 1 4 1 0 0 0 21 0.0849 0.2994 0.6157
2018 1 4 1 0 0 0 31 0.1475 0.2219 0.6306
2019 1 4 1 0 0 0 50 0.1961 0.2346 0.5692 # no survey so not updated
2021 1 4 1 0 0 0 50 0.3323 0.1320 0.5357 # updated 4-14-22
##length proportions of pot survey
##Year, Seas, Fleet, Sex, Type, Shell, Maturity, Nsamp, DataVec
1995 1 5 1 0 0 0 100 0.1594 0.2656 0.5751
1998 1 5 1 0 0 0 100 0.0769 0.2205 0.7026
2001 1 5 1 0 0 0 100 0.1493 0.2049 0.6457
2004 1 5 1 0 0 0 100 0.0672 0.2484 0.6845
2007 1 5 1 0 0 0 100 0.1257 0.3148 0.5595
2010 1 5 1 0 0 0 100 0.1299 0.3209 0.5492

```









```

-4 21 1 0 595 1 999 0 1 999 -3 1978 2021 # update end yr
# Gear-5
-5 22 1 0 595 1 999 0 1 999 -3 1978 2021 # update end yr

# Number of asymptotic parameters
1
# Fleet Sex Year ival lb ub phz
1 1 1978 0.000001 0 1 -3

## ===== ##
## PRIORS FOR CATCHABILITY
## If a uniform prior is selected for a parameter then the lb and ub are used (p1 ##
## and p2 are ignored). ival must be > 0 ##
## only allowed to use uniform or lognormal prior ##
## if analytic q estimation step is chosen, turn off estimating q by changing the estimation phase to be -ve ##
## LEGEND ##
## prior: 0 = uniform, 1 = normal, 2 = lognormal, 3 = beta, 4 = gamma ##
## ===== ##
## LAMBDA: Arbitrary relative weights for each series, 0 = do not fit.
## SURVEYS/INDICES ONLY
## Analytic (0 = not analytically solved q, use uniform or lognormal prior; 1 = analytic) ##
## Lambda = multiplier for input CV, Emphasis = multiplier for likelihood ##
## ival lb ub phz prior p1 p2 Analytic? LAMBDA Emphasis
1.0 0.5 1.2 -4 0 0 9.0 0 1 1 # NMFS trawl
0.003 0 5 3 0 0 9.0 0 1 1 # ADF&G pot
## ===== ##
## if uniform prior is specified then use lb and ub rather than p1 and p2
## ===== ##
## ADDITIONAL CV FOR SURVEYS/INDICES
## If a uniform prior is selected for a parameter then the lb and ub are used (p1 ##
## and p2 are ignored). ival must be > 0 ##
## LEGEND ##
## prior: 0 = uniform, 1 = normal, 2 = lognormal, 3 = beta, 4 = gamma ##
## ===== ##
## ival lb ub phz prior p1 p2
0.0000001 0.000000001 10.0 -4 4 1.0 100 # NMFS (PHASE -4)
0.00000001 0.000000001 10.0 -4 4 1.0 100 # ADF&G
## ===== ##
### Pointers to how the additional CVs are used (0 ignore; >0 link to one of the paramters)
0 0
## ===== ##
## PENALTIES FOR AVERAGE FISHING MORTALITY RATE FOR EACH GEAR
## ===== ##
## Mean_F Female Offset STD_PHZ1 STD_PHZ2 PHZ_M PHZ_F Fbar_l Fbar_h Fdev_L Fdev_h Foff_l Foff_h # Pot
0.2 0.0 3.0 50.0 1 -1 -12 4 -10 10 -10 10 # Pot
0.0001 0.0 4.0 50.0 1 -1 -12 4 -10 10 -10 10 # Trawl
0.0001 0.0 4.0 50.0 1 -1 -12 4 -10 10 -10 10 # Fixed
0.0 0.0 2.00 20.00 -1 -1 -12 4 -10 10 -10 10 # NMFS
0.00 0.0 2.00 20.00 -1 -1 -12 4 -10 10 -10 10 # ADF&G
## ===== ##

## ===== ##
## OPTIONS FOR SIZE COMPOSTION DATA (COLUMN FOR EACH MATRIX)
## ===== ##
## LIKELIHOOD OPTIONS
## -1) Multinomial with estimated/fixed sample size
## -2) Robust approximation to multinomial
## -3) logistic normal (NIY)
## -4) multivariate-t (NIY)
## -5) Dirichlet
## AUTOTAIL COMPRESSION
## pmin is the cumulative proportion used in tail compression.
## ===== ##
# 1 1 1 # Type of likelihood
2 2 2 # Type of likelihood
# 5 5 5 # Type of likelihood
0 0 0 # Auto tail compression (pmin)
1 1 1 # Initial value for effective sample size multiplier
-4 -4 -4 # Phz for estimating effective sample size (if appl.)
1 2 3 # Composition aggregator
1 2 2 # set to 2 for survey-like predictions; 1 for catch like predictions #AEP
1 1 1 # LAMBDA

```



```

## EMPHASIS FACTORS (CATCH)
## ===== ##
#Ret_POT Disc_POT Disc_trawl Disc_fixed
      1      1      1      1

## EMPHASIS FACTORS (Priors) by fleet: Fdev_total, Fdov_total, Fdev_year, Fdov_year
1 0 0.000 0 # Pot fishery
1 0 0.000 0 # Trawl bycatch
1 0 0.000 0 # fixed gear bycatch
1 0 0.000 0 # NMFS survey
1 0 0.000 0 # ADF&G survey

## ===== ##
## EMPHASIS FACTORS (Priors)
## ===== ##
# Log_fdevs   meanF      Mdevs  Rec_devs  Initial_devs  Fst_dif_dev  Mean_sex-Ratio   Molt_prob  Free selectivity  Init_n_at_len  Fvecs  Fdovss (!!!N
      10000      0      1.0      1      0      0      1      0      0      0      0      0
## EOF
9999

```