

Analysis of past BSAI and GOA Groundfish Plan Team recommendations to set ABC below the maximum permissible level

Grant G. Thompson

Resource Ecology and Fisheries Management Division
Alaska Fisheries Science Center
National Marine Fisheries Service
National Oceanic and Atmospheric Administration
7600 Sand Point Way NE., Seattle, WA 98115-6349

Introduction

The minutes of the February 2018 SSC meeting state, “The SSC recommends *identification of clear and transparent rules* for defining the specific criteria to be used when adjusting the recommended ABC” (emphasis added). This paper describes an attempt to develop a starting point for satisfying the SSC’s request by reverse-engineering a set of rules from previous ABC recommendations.

Methods

During the 15 years spanning the period 2003-2017, the BSAI and GOA Groundfish Plan Teams (“Teams”) recommended setting ABC below the maximum permissible level (“maxABC”) in a total of 76 instances, for an average of 5.07 instances per year (Figure 1). In all but 4 instances the Teams listed at least one reason for their recommendations, with an average of 2.68 reasons per instance. After omitting some non-essential verbiage, a total of 122 unique reasons were listed over the course of the 15 years (some reasons were listed more than once).

To make the analysis more tractable, the 122 unique reasons were condensed into a set of 66 initial paraphrased reasons (“Round 1” variables, Table 1). This resulted in a 76×66 (instance \times reason) matrix \mathbf{X} , where each element is either 0 or 1 depending on whether the given reason (column) was listed by the Teams for the given ABC reduction (row). The product $\mathbf{X}'\mathbf{X}$ turned out to be singular, due in part to confounding of several of the Round 1 variables. Specifically, the members of 8 subsets of Round 1 variables never occurred in the matrix apart from the other members of the respective subset. This was considered problematic insofar as it rendered ordinary least-squares regression impossible.

Each of the 8 confounded subsets of Round 1 variables was therefore consolidated into a single “Round 2” variable (Table 2; note that all Round 1 variables not listed in this table were retained for Round 2). This resulted in a new \mathbf{X} matrix of dimension 76×52 (22 Round 1 variables were included in the 8 confounded subsets, resulting in $66 - (22 - 8) = 52$ variables in Round 2). However, the product $\mathbf{X}'\mathbf{X}$ was still singular, with 10 of the 52 eigenvalues being zero. The zero eigenvalues were associated with 20 of the Round 2 variables.

Each of the 20 Round 2 variables associated with the zero eigenvalues was therefore combined with one Round 2 variable that was *not* associated with the zero eigenvalues, giving the list of “Round 3” variables shown in Table 3 (note that all Round 2 variables not listed in this table were retained for Round 3). The choices regarding which variables to combine were subjective, with a goal of minimizing the degree of controversy resulting from those choices. This resulted in another new \mathbf{X} matrix of dimension 76×32 (52 Round 1 variables – 20 Round 1 variables associated with the zero eigenvalues = 32 variables in Round 3). The product $\mathbf{X}'\mathbf{X}$ was now non-singular.

The set of reductions in ABC, expressed as proportions of maxABC (“y”), were logit-transformed for $i=1,2,\dots,76$ as follows:

$$\text{logit}_i y_i = \ln \left(\frac{1 + y_i}{1 - y_i} \right).$$

Least-squares parameter estimates were then calculated as follows:

$$\beta_1 = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\text{logit}_i \mathbf{y}.$$

Some of the resulting β_1 values were negative, implying that, if any subset of the corresponding variables were the only ones that were to apply in a particular situation, the ABC would have to be adjusted upward, rather than downward, from maxABC. Therefore, the β_1 estimates were used as initial values for β_2 in the following constrained optimization problem, which ensures that adjustments relative to maxABC are always in the downward direction (note that the least-squares reduction estimates have undergone a logistic back-transformation):

$$\text{minimize } \sum_{i=1}^{76} \left(y_i - \left(2(1 + \exp(-\mathbf{x}\beta_2)_i) \right)^{-1} - 1 \right)^2 \text{ subject to } \beta_{2i} \geq 0 \text{ for all } i = 1, 2, \dots, 76.$$

Results

Figure 2 compares the β_1 and β_2 estimates for the 32 Round 3 variables, sorted in increasing order of β_2 . Note that 8 of the β_1 estimates are less than zero. The corresponding β_2 estimates for the first 7 of these are zero, and the eighth (variable 9 in Figure 2) is positive. Back-transforming the least-squares model (β_1) to the natural scale gives an R^2 of 0.826, while the constrained model (β_2 , which has already been back-transformed) gives an R^2 of 0.824. The β_2 values, sorted in increasing order, are shown in Table 4 along with the implied univariate ABC reduction for each (i.e., the ABC reduction that would result if no other variable was applicable).

The ABC reductions estimated by the two models are compared against the observed ABC reductions in Figure 3. The errors (absolute value) from both models were less than 0.2 in all but six instances. The following five instances of errors greater than 0.2 were common to both models:

Stock	Year
EBS pollock	2011
EBS pollock	2014
AI pollock	2008
BSAI Greenland turbot	2004
BSAI Greenland turbot	2005

The sixth instance of an error greater than 0.2 occurred with respect to 2014 EBS Pacific cod for the unconstrained model (β_1) and 2003 BSAI Greenland turbot for the constrained model (β_2).

Discussion

First, some caveats/clarifications:

- Some of the steps involved in this analysis were necessarily subjective, and it is likely that other analysts would have arrived at a somewhat different final model (just like in a stock assessment).
- The analysis is necessarily one-sided in that the Teams did not check consistently to determine which, if any, of the variables listed in Table 1 were applicable to each stock/complex in each year; rather, the reasons for reducing ABC were typically articulated after the assessment author or the Team became concerned that a reduction for a particular stock/complex in a particular year might be warranted, without consistent regard for whether those same reasons, or reasons articulated for the same stock or other stocks in previous years, might apply elsewhere.
- A zero coefficient (β_2) does not necessarily imply that the corresponding variable, considered in isolation, is unimportant; only that it does not provide any explanatory power over and above those of the other variables with which it has been paired in previous applications.

The multivariate logistic nature of the final model developed here (β_2) is appealing in that the functional form is fairly simple, all coefficients are positive (meaning that if only a single variable applies in a particular situation, the resulting reduction is guaranteed to be positive), and the resulting reductions will always be between 0 and 1 (this would not necessarily be the case if a linear, rather than logistic, form were used).

However, this analysis is intended as a starting point only. If the basic approach is adopted, it is expected that the Teams will want to both add to, and subtract from, the list of variables in Table 4. For example:

- “None” is not a particularly compelling reason to reduce ABC below maxABC.
- Variables such as “SSC’s rule for Bogoslof pollock ABC” are likely of historical interest only.
- To satisfy the SSC’s request for “identification of clear and transparent rules,” it may be necessary to frame each of the variables in objectively quantifiable terms (e.g., “most recent survey CV is in excess of 50%,” may be preferable to “large survey CV”).
- Potential candidate variables suggested by the SSC (February 2018) include:
 - Stock assessment uncertainty relative to levels upon which the Tier system was constructed.
 - Atypical data availability or usage (e.g., reliance on only catch-per-unit-effort vs. a survey index).
 - Ecosystem considerations.

Not only is the list of variables subject to modification, but it is expected that the Teams will want to modify the corresponding coefficients as well. When setting a coefficient value, it may be desirable to reverse-engineer it from a specified univariate reduction through the logit transform described above. For example, to achieve a 10% univariate reduction for some particular variable, the corresponding coefficient would be set equal to $\ln((1+0.1)/(1-0.1)) = \ln(1.1/0.9) \approx 0.201$.

Once the set of variables to be included and the values of the corresponding coefficients β have been determined, the proportional reduction in any particular situation (i.e., combination of stock/complex and year) can be computed as

$$reduction = 2 \left(1 + \exp \left(- \sum_{var=1}^{nvar} x_{var} \beta_{var} \right) \right)^{-1} - 1 ,$$

where $nvar$ represents the number of variables included in the decision process, and \mathbf{x} is a vector with elements equal to 0 or 1 depending on whether the variable applies in the particular situation.

Tables

Table 1 (page 1 of 2). Round 1 paraphrased reasons (variables) for setting $ABC < maxABC$.

Round 1 variables	N Assessments, in alphabetical order
Catching maxABC would require large increase in effort	1 2017 EBS pollock
Change in biomass estimates between assessments	1 2008 BSAI Greenland turbot
Dome-shaped selectivity may overestimate abundance	1 2003 BSAI Pacific cod
Dominance of single year class	7 2010-2014 EBS pollock, 2003-2004 GOA Atka mackerel
Dorn's buffer	13 2003-2015 GOA pollock
Estimated selectivity may not capture year class targeting	1 2011 EBS pollock
Few older fish	4 2017 BSAI sablefish, 2016-2017 EBS pollock, 2017 GOA sablefish
Habitat-specific residency	12 2003-2013 and 2015 GOA demersal shelf rockfish
Hedge against future regime shift	1 2010 EBS pollock
Important role in ecosystem as prey	2 2016-2017 EBS pollock
Increased patchiness of distribution	1 2016 EBS pollock
Increasing catch on a declining stock	2 2003 BSAI sablefish, 2014 BSAI skates
Increasing predation pressure	1 2006 EBS pollock
Keep spawning exploitation rate below all-time high	1 2007 EBS pollock
Large survey CV	2 2016-2017 Bogoslof pollock
Late maturation	12 2003-2013 and 2015 GOA demersal shelf rockfish
Late-breaking analysis (not included in assessment)	1 2014 GOA Pacific cod
Longevity	12 2003-2013 and 2015 GOA demersal shelf rockfish
Long-term biomass decline	2 2003 and 2005 BSAI Greenland turbot
Long-term poor recruitment	3 2008 AI pollock, 2003 BSAI Greenland turbot, 2004 GOA pollock
Low biomass	5 2008 AI pollock, 2017 BSAI sablefish, 2011 and 2013 EBS pollock, 2017 GOA sablefish
Low roe recovery rates	2 2013 and 2016 EBS pollock
Maintain fishery CPUE	3 2011, 2014, and 2015 EBS pollock
maxABC would be a dramatic increase over historic catch	4 2008 AI pollock, 2004 and 2006 BSAI Greenland turbot, 2008 BSAI Pacific cod
Multispecies model estimates higher biomass reference points	2 2016-2017 EBS pollock
No history of rebuilding from current level when catch=maxABC	1 2007 EBS pollock
No new survey	1 2006 BSAI Greenland turbot
No reliable biomass estimate	2 2003-2004 GOA Atka mackerel
None	4 2014 and GOA demersal shelf rockfish, 2003-2004 GOA other slope rockfish
Possibility of localized depletion	2 2003-2004 GOA Atka mackerel
Potential overharvest in distant past	2 2003-2004 GOA Atka mackerel
Precedent	3 2017 GOA demersal shelf rockfish, 2003-2004 GOA shorttraker/rougheye rockfish
Preferred model does not account for structural uncertainty	1 2016 EBS pollock

Table 1 (page 2 of 2). Round 1 paraphrased reasons for setting $ABC < maxABC$.

Round 1 variables	N Assessments, in alphabetical order
Prevent $B < B_{20\%}$	1 2017 GOA Pacific cod
Projected biomass decline	4 2017 EBS pollock, 2006 GOA Pacific cod, 2004 GOA pollock, 2003 GOA sablefish
Projected poor recruitment	2 2016-2017 EBS pollock
Protect against disproportionate harvest within complex	1 2004 GOA shortraker/rougheye rockfish
Recent bird die-offs	3 2017 EBS Pacific cod, 2016-2017 EBS pollock
Recent poor recruitment	3 2017 EBS Pacific cod, 2007 EBS pollock, 2006 GOA Pacific cod
Recent year class estimated to be extremely large	2 2017 BSAI sablefish, 2017 GOA sablefish
Reduce ABC to account for State GHLL	11 2003-2013 GOA pollock
Reduce bycatch	2 2014-2015 EBS pollock
Reduced condition factor	2 2017 EBS Pacific cod, 2013 EBS pollock
Reduced prey availability	2 2006 and 2016 EBS pollock
Reliable minimum biomass estimate available for Tier 6 stock	1 2010 GOA sharks
Retrospective bias	6 2014-2015 EBS Pacific cod, 2007 and 2011-2012 EBS pollock, 2014 GOA Pacific cod
Sedentary life history	12 2003-2013 and 2015 GOA demersal shelf rockfish
Short-term biomass decline	3 2003 BSAI Pacific cod, 2017 EBS Pacific cod, 2011 EBS pollock
Significant probability of exceeding FMSY	1 2012 EBS pollock
SSC's rule for Bogoslof pollock ABC	4 2007-2010 Bogoslof pollock
Survey index lower than expected	2 2006 EBS pollock, 2004 GOA pollock
Trend of increasing F on oldest ages	1 2013 EBS pollock
Uncertain biomass estimates	2 2013 and 2015 GOA demersal shelf rockfish
Uncertain recruitment estimates	4 2004 BSAI Greenland turbot, 2017 BSAI sablefish, 2011 EBS pollock, 2017 GOA sablefish
Uncertain stock structure	2 2011 and 2017 EBS pollock
Uncertain stock trend	1 2008 BSAI Greenland turbot
Uncertainty surrounding M and Q	2 2004 AI pollock, 2003 GOA Pacific cod
Uncertainty surrounding Q	1 2015 EBS Pacific cod
Vessels are needing to travel farther	1 2006 EBS pollock
Whale depredation	4 2016-2017 BSAI sablefish, 2016-2017 GOA sablefish
Yield variability	2 2004 BSAI Pacific cod, 2004 GOA Pacific cod

Table 2. Mapping confounded Round 1 variables into Round 2 variables. (Round 1 variables not shown here were retained in Round 2).

Round 1 variables	Round 2 variables
Habitat-specific residency	Some concerns unique to most GOA demersal shelf rockfish assessments
Late maturation	Some concerns unique to most GOA demersal shelf rockfish assessments
Longevity	Some concerns unique to most GOA demersal shelf rockfish assessments
Sedentary life history	Some concerns unique to most GOA demersal shelf rockfish assessments
No reliable biomass estimate	Some concerns unique to the 2003 and 2004 GOA Atka mackerel assessments
Possibility of localized depletion	Some concerns unique to the 2003 and 2004 GOA Atka mackerel assessments
Potential overharvest in distant past	Some concerns unique to the 2003 and 2004 GOA Atka mackerel assessments
Increasing predation pressure	Some concerns unique to the 2006 EBS pollock assessment
Vessels are needing to travel farther	Some concerns unique to the 2006 EBS pollock assessment
Keep spawning exploitation rate below all-time high	Some concerns unique to the 2007 EBS pollock assessment
No history of rebuilding from current level when catch=maxABC	Some concerns unique to the 2007 EBS pollock assessment
No history of strong year classes emerging from current stock level	Some concerns unique to the 2007 EBS pollock assessment
Change in biomass estimates between assessments	Some concerns unique to the 2008 BSAI Greenland turbot assessment
Uncertain stock trend	Some concerns unique to the 2008 BSAI Greenland turbot assessment
Important role in ecosystem as prey	Some concerns unique to the 2016 and 2017 EBS pollock assessments
Multispecies model estimates higher biomass reference points	Some concerns unique to the 2016 and 2017 EBS pollock assessments
Projected poor recruitment	Some concerns unique to the 2016 and 2017 EBS pollock assessments
Increased patchiness of distribution	Some concerns unique to the 2016 EBS pollock assessment
Preferred model does not account for structural uncertainty	Some concerns unique to the 2016 EBS pollock assessment
Recent bird die-offs	Some concerns unique to the 2017 EBS Pacific cod assessment
Recent high age 1 mortality from multispecies model	Some concerns unique to the 2017 EBS Pacific cod assessment
Recent low crab abundance	Some concerns unique to the 2017 EBS Pacific cod assessment

Table 4. Constrained regression coefficients (beta) with corresponding univariate ABC reductions.

Round 3 variables	beta	univariate reduction
Age structure concerns	0.00	0.00
Protect against disproportionate harvest within complex	0.00	0.00
Recent poor recruitment	0.00	0.00
Retrospective bias	0.00	0.00
Survey index lower than expected	0.00	0.00
Uncertain biomass estimates	0.00	0.00
Whale depredation	0.00	0.00
Reduce ABC to account for State GHL	0.08	0.04
Dramatic increase in catch or increasing catch on a declining stock	0.11	0.06
Prevent B<B20%	0.14	0.07
Projected biomass decline	0.21	0.10
Long-term poor recruitment	0.22	0.11
Dorn's buffer	0.23	0.11
None	0.24	0.12
Late-breaking analysis (not included in assessment)	0.25	0.12
Yield variability	0.31	0.15
Precedent	0.33	0.16
Some concerns unique to various EBS pollock and Pacific cod assessments	0.37	0.18
Uncertain recruitment estimates, including recent large estimates	0.40	0.20
Model uncertainty	0.43	0.21
Some concerns unique to most GOA demersal shelf rockfish assessments	0.47	0.23
Uncertainty surrounding Q	0.47	0.23
Uncertainty surrounding M and Q	0.60	0.29
Some concerns unique to the 2008 BSAI Greenland turbot assessment	0.80	0.38
Low or long-term declining biomass	0.82	0.39
Significant probability of exceeding FMSY	0.86	0.40
Hedge against future regime shift	0.87	0.41
Reliable minimum biomass estimate available for Tier 6 stock	0.89	0.42
Large survey CV	0.91	0.43
No new survey	1.81	0.72
Some concerns unique to the 2003 and 2004 GOA Atka mackerel assessments	2.69	0.87
SSC's rule for Bogoslof pollock ABC	2.99	0.90

Figures




Figure 1. Frequency of reductions, binned in 10% increments.




Figure 2. Comparison of constrained and unconstrained coefficients (sorted in order of the former).




Figure 3. Estimated (constrained and unconstrained) versus observed ABC reductions.