

Table 8. Recommended Pacific halibut discard mortality rates (DMRs), expressed as percentages, for 2016-2018 CDQ and non-CDQ groundfish fisheries off Alaska. Values noted with an asterisk (*) indicate the use of the Pacific cod target DMR for that fishery, as data for 2005-2014 were unavailable. Underlined DMR values from Tables 4, 5 and 7 not used in calculating these recommendations.

Non-CDQ

Bering Sea/Aleutians			Gulf of Alaska		
Gear/Target	Used in 2013-2015	2016-2018 Recommendation	Gear/Target	Used in 2013-2015	2016-2018 Recommendation
<i>Trawl</i>			<i>Trawl</i>		
Atka mack	77	80	Bottom poll	60	58
Bottom poll	77	81	Pacific cod	62	62
Pacific cod	71	66	Dpwtr flats	43	62*
Other Flats	71	63	Shallwtr flats	67	67
Rockfish	79	81	Rockfish	66	65
Flathead sole	73	76	Flathead sole	65	67
Midwtr poll	88	88	Midwtr poll	71	72
Rock sole	85	86	Sablefish	71	71
Sablefish	75	66*	Arr. fldr	73	76
Turbot	64	82	Rex sole	69	72
Arr. fldr	76	84			
YF sole	83	84			
<i>Pot</i>			<i>Pot</i>		
Pacific cod	8	9	Pacific cod	17	15
<i>Longline</i>			<i>Longline</i>		
Pacific cod	9	9	Pacific cod	11	10
Rockfish	4	9*	Rockfish	9	10*
Turbot	13	14			

Bering Sea/Aleutians CDQ

Gear/Target	Used in 2013-2015	2016-2018 Recommendation
<i>Trawl</i>		
Atka mackerel	86	81
Bottom pollock	83	86
Pacific cod	90	90
Rockfish	80	80
Flathead sole	79	79
Midwtr pollock	90	90
Rock sole	88	88
Turbot	89	89
Yellowfin sole	86	85
<i>Pot</i>		
Pacific cod	-	1
Sablefish	34	39
<i>Longline</i>		
Pacific cod	10	10
Turbot	4	10*