


NOAA
FISHERIES
AFSC, Seattle

CEATTLE: Climate enhanced Age-based model with Temperature specific Trophic linkages and Energetics

Kirstin Holsman
Jim Ianelli , Kerim Aydin,
André Punt, Elizabeth Moffitt

Sept. 2016, NPMFC PT



kirstin.holsman@noaa.gov

Multi-species models for EBFM


- Quantify **relative effects** of climate variability, trophic interactions, and fisheries on species productivity
- **Non-stationary mortality**, B_0 , and MSY
- Can identify **indirect effects** on other species and fisheries
- **Quantify trade-offs among fisheries**

Holsman et al. in press. Deep Sea Res II

Photo: Mark Holsman

2016 multispecies assessment

- Appendix to 2016 pollock assessment as alternative model
- Provides comparative M , Rec , and SSB
- Captures climate and trophic impacts on BRPs



Outline:

- CEATTLE Overview
- 2016 Assessment
- Other applications



CEATTLE overview



Photo: Mark Holsman

CEATTLE methods references



ELSEVIER

Contents lists available at [ScienceDirect](#)

Deep-Sea Research II

journal homepage: www.elsevier.com/locate/dsr2



A comparison of fisheries biological reference points estimated from temperature-specific multi-species and single-species climate-enhanced stock assessment models

Kirstin K. Holsman ^{a,*}, James Ianelli ^a, Kerim Aydin ^a, André E. Punt ^b, Elizabeth A. Moffitt ^{b,1}

^a Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, 7600 Sand Point Way N.E., Building 4, Seattle, Washington 98115, USA

^b University of Washington School of Aquatic and Fisheries Sciences, 1122 NE Boat St., Seattle, WA 98105, USA

ARTICLE INFO

ABSTRACT

Multi-species statistical catch at age models (MSCAA) can quantify interacting effects of climate and fisheries harvest on species populations, and evaluate management trade-offs for fisheries that target


Holsman, KK, J Ianelli, K Aydin, AE Punt, EA Moffitt (2015). Comparative biological reference points estimated from temperature-specific multispecies and single species stock assessment models. Deep Sea Res II. doi:10.1016/j.dsr2.2015.08.001.

Moffitt, E, AE Punt, KK Holsman, KY Aydin, JN Ianelli, I Ortiz (2015). Moving towards Ecosystem Based Fisheries Management: options for parameterizing multi-species harvest control rules. Deep Sea Res II. doi:10.1016/j.dsr2.2015.08.002




NOAA FISHERIES

CEATTLE Multi-species model



Walleye pollock
(*Gadus chalcogrammus*)




Arrowtooth flounder
(*Atheresthes stomias*)



Pacific cod
(*Gadus macrocephalus*)

Eastern Bering Sea, Alaska, USA



$W @ \text{Age} \sim f(\text{Temperature})$
 $\text{Pred/prey} \sim f(\text{Temperature})$

Climate-Enhanced, Age-based model with Temperature-specific Trophic Linkages and Energetics



NOAA FISHERIES

Table 1: Model equations

Definition	Equation		
Recruitment	$N_{i1,y} = R_{i,y} = R_{0,i} e^{\tau_{i,y}}$	$\tau_{i,y} \sim N(0, \sigma^2)$	T1.1
Initial abundance	$N_{ij,1} = \begin{cases} R_{0,i} e^{(-j M1_{ij})} N_{0,ij} & y=1 \quad 1 < j \leq A_i \\ R_{0,i} e^{(-j M1_{i,A_i})} N_{0,i,A_i} / (1 - e^{(-j M1_{i,A_i})}) & y=1 \quad j > A_i \end{cases}$		T1.2
Numbers at age	$N_{i,j+1,y+1} = N_{ij,y} e^{-Z_{ij,y}} \quad 1 \leq y \leq n_y \quad 1 \leq j < A_i$ $N_{i,A_i,y+1} = N_{i,A_i-1,y} e^{-Z_{i,A_i-1,y}} + N_{i,A_i,y} e^{-Z_{i,A_i,y}} \quad 1 \leq y \leq n_y \quad j > A_i$		T1.3
Catch	$C_{ij,y} = \frac{F_{ij,y}}{Z_{ij,y}} (1 - e^{-Z_{ij,y}}) N_{ij,y}$		T1.4
Total yield (kg)	$Y_{i,y} = \sum_j^{A_i} \left(\frac{F_{ij,y}}{Z_{ij,y}} (1 - e^{-Z_{ij,y}}) N_{ij,y} W_{ij,y} \right)$		T1.5
	$B_{ij,y} = N_{ij,y} W_{ij,y}$		
	$F_{ij,y} = F_{0,i} e^{\tau_{i,y}} S_{ij}^s$	$\varepsilon_{i,y} \sim N(0, \sigma_{F,i}^2)$	
	$W_{ij,y} = W_{\infty,ij} \left(1 - e^{(-K_i(1-d_{i,y})(j-t_{0,i}))} \right)^{\frac{1}{1-d_{i,y}}}$		T1.10b
	$d_{i,y} = e^{(\alpha_{d,i,y} + \alpha_{0,d,i} + \beta_{d,i} T_y)}$		T1.10c
	$W_{\infty,ij} = \left(\frac{H_i}{K_i} \right)^{1/(1-d_{i,y})}$		T1.11
BT survey biomass (kg)	$\hat{\rho}_{i,y}^s = \sum_j^{A_i} \left(N_{ij,y} e^{-0.5 Z_{ij,y}} W_{ij,y} S_{ij}^s \right)$		T1.12
EIT survey			T1.13
Fishery			T1.14
BT survey			T1.15
EIT survey			T1.16
BT selectivity	$S_{ij}^s = \frac{1}{1 + e^{(-b_{i,j}^s \cdot j - a_{i,j}^s)}}$		T1.17
Fishery selectivity	$S_{ij}^f = \begin{cases} e^{\eta_{ij}} & j \leq A_{\eta,i} \\ e^{\eta_{i,A_{\eta,i}}} & j > A_{\eta,i} \end{cases}$	$\eta_{ij} \sim N(0, \sigma_{\eta,i}^2)$	T1.18
Proportion females	$\omega_{ij} = \frac{e^{-j M_{fem}}}{e^{-j M_{fem}} + e^{-j M_{male}}}$		T1.19
Proportion of mature females	$\rho_{ij} = \omega_{ij} \phi_{ij}$		T1.20
Weight at age (kg)	$W_{ij,y} = W_{ij,y}^{fem} \omega_{ij} + (1 - \omega_{ij}) W_{ij,y}^{male}$		T1.21
Residual natural mortality	$M1_{ij} = M_{ij}^{fem} \omega_{ij} + (1 - \omega_{ij}) M_{ij}^{male}$		

Residual
Natural Mortality

Predation
Natural Mortality

$$Z_{ij,y} = M1_{ij} + M2_{ij,y} + F_{ij,y}$$



Table 2: Pred. Mortality (M2)

Definition	Equation
Predation mortality	$M2_{ij,y} = \sum_{pa} \left(\frac{N_{pa,y} \delta_{pa,y} \bar{S}_{paij}}{\sum_{ij} (\bar{S}_{paij} B_{ij,y}) + B_p^{other} (1 - \sum_{ij} (\bar{S}_{paij}))} \right) \quad T2.1$
Predator-prey suitability	Age-specific prey selectivity
Mean gravimetric diet proportion	$\sum_{ij} \dots \quad T2.3$
Individual specific ratio (kg kg ⁻¹ yr ⁻¹)	Size-specific annual ration
Temperature scaling algorithm	$f(T) = V^X (X(1-V)) \quad T2.5$
	Temperature specific
	$X = \left(Z^2 \left(1 + (1 + 40/Y)^{0.5} \right)^2 \right) / 400 \quad T2.5b$
	$Z = \ln(Q_p^c) (T_p^{cm} - T_p^{co}) \quad T2.5c$
	$Y = \ln(Q_p^c) (T_p^{cm} - T_p^{co} + 2) \quad T2.5d$



Temperature specific VonB


$$W_{ij,y} = W_{\infty,iy} \left(1 - e^{(-K_i(1-d_{i,y})(j-t_{0,i}))} \right)^{\frac{1}{1-d_{i,y}}}$$

$$d_{i,y} = e^{(\alpha_{d,i,y} + \alpha_{0d,i} + \beta_{d,i} T_y)}$$

$$W_{\infty,iy} = \left(\frac{H_i}{K_i} \right)^{1/(1-d_{i,y})}$$


Bottom Temp

Fit outside of the model presently (expect for projections)



Predator- Prey Overlap


Set to 1.0 in this assessment




Stabeno et al. (2013) A comparison of the physics of the northern and southern shelves of the eastern Bering Sea and some implications for the ecosystem. *Deep-Sea Res II* 65-7014-30.




CEATTLE




CEATTLE




CEATTLE



CEATTLE



CEATTLE



CEATTLE: Options

- Trophic Interactions (on/off)
- Recruitment under projections (MCMC on/off):
 - Mean RS (Ricker or BevHolt), linear, linear+ B_{y-1}
 - Above + covariates (cold pool, BT, etc.)
- Harvest scenarios (0, mean historical F or C, set F or C, F profile, and HCRs):
 - ABC proxy
 - SPR
 - Aggregate MSY
 - MEY (in development)

2016 Assessment



Photo: Mark Holsman

Changes from Holsman et al. 2016

- P. cod fish comp data based on lengths
- Bottom Temp: BTS & updated through 2015
- Projected Bottom Temp: constant avg (BT)
- Recruitment in projections is Ricker
- 2 harvest scenarios are presented here:
 1. F 40% of unfished biomass (for all three species simultaneously)
 2. aggregate multi-species MSY
- Survey and catch data updated from 2012



2016 multispecies data

Source	Type	Years
Fishery	Catch biomass	1979-2015
Fishery	Catch age composition	1979-2014 (plk &atf); 1979-2015 (P. cod)
EBS bottom trawl	Area-swept abundance (numbers) index by age	1982-2015
EBS bottom trawl	Gravimetric length-based diet data	1982-2105
Acoustic trawl	Population abundance (numbers) index by age	1979,1982,1985,1988, 1991, 1994, 1996-1997, 1999-2000, 2002 , 2004, 2006-2010

Bottom Temperature (BT Survey)

2016
4.5 deg




Table 4: Objective functions

Description	Equation	Data source	
Data components			
BT survey biomass	$\sum_i \sum_y \frac{[\ln(\hat{\beta}_{i,y}^s) - \ln(\beta_{i,y}^s)]^2}{2\sigma_{s,i}^2}$	NFMS annual EBS BT survey (1979–2012)	T4.1
BT survey age composition	$-\sum_i n_i \sum_y \sum_j (O_{ij,y}^s + v) \ln(\hat{O}_{ij,y}^s + v)$	NFMS annual EBS BT survey (1979–2012)	T4.2
EIT survey biomass	$\sum_y \frac{[\ln(\hat{\beta}_y^{eit}) - \ln(\beta_y^{eit})]^2}{2\sigma_{eit}^2}, \sigma_{eit} = 0.2$	Pollock acoustic trawl survey (1979–2012)	T4.3
EIT age composition	$-n \sum_y \sum_j (O_{1j,y}^{eit} + v) \ln(\hat{O}_{1j,y}^{eit} + v)$	Pollock acoustic trawl survey (1979–2012)	T4.4
Total catch	$\sum_i \sum_y \frac{[\ln(C_{i,y}^*) - \ln(C_{i,y}^*)]^2}{2\sigma_c^2}, \sigma_c = 0.05$	Fishery observer data (1979–2012)	T4.5
Fishery age composition	$-\sum_i n_i \sum_y \sum_j (O_{ij,y}^f + v) \ln(\hat{O}_{ij,y}^f + v)$	Fishery observer data (1979–2012)	T4.6
Penalties			
Fishery selectivity	$\sum_i \sum_j^{A_i-1} \chi \cdot \left[\ln\left(\frac{\eta_{ij}^f}{\eta_{ij+1}^f}\right) - \ln\left(\frac{\eta_{ij+1}^f}{\eta_{ij+2}^f}\right) \right]^2, \chi = \begin{cases} 20, & \text{if } \eta_{ij}^f > \eta_{ji+1}^f \\ 0, & \text{if } \eta_{ij}^f \leq \eta_{ji+1}^f \end{cases}$		T4.7
Priors			
	$\sum_i \sum_y (\tau_{i,y})^2$		T4.8
	$\sum_i \sum_y (N_{0,ij})^2$		T4.9
	$\sum_i \sum_y (\varepsilon_{i,y})^2$		T4.10

$v=0.001$.

Description	Equation
Data components	
BT survey biomass	$\sum_i \sum_y \frac{[\ln(\beta_{i,y}^s) - \ln(\hat{\beta}_{i,y}^s)]^2}{2\sigma_{s,i}^2}$
BT survey age composition	$-\sum_i n_i \sum_y \sum_j (O_{ij,y}^s + v) \ln(\hat{O}_{ij,y}^s + v)$
EIT survey biomass	$\sum_y \frac{[\ln(\beta_y^{eit}) - \ln(\hat{\beta}_y^{eit})]^2}{2\sigma_{eit}^2}, \sigma_{eit} = 0.2$
EIT age composition	$-n \sum_y \sum_j (O_{1j,y}^{eit} + v) \ln(\hat{O}_{1j,y}^{eit} + v)$
Total catch	$\sum_i \sum_y \frac{[\ln(C_{i,y}^*) - \ln(\hat{C}_{i,y}^*)]^2}{2\sigma_c^2}, \sigma_c = 0.05$
Fishery age composition	$-\sum_i n_i \sum_y \sum_j (O_{ij,y}^f + v) \ln(\hat{O}_{ij,y}^f + v)$
Penalties	
Fishery selectivity	$\sum_i \sum_j^{A_i-1} \chi \cdot \left[\ln\left(\frac{\eta_{ij}^f}{\eta_{ij+1}^f}\right) - \ln\left(\frac{\eta_{ij+1}^f}{\eta_{ij+2}^f}\right) \right]^2, \chi = \begin{cases} 20, & \text{if } \eta_{ij}^f > \eta_{ji+1}^f \\ 0, & \text{if } \eta_{ij}^f \leq \eta_{ij+1}^f \end{cases}$
Priors	
	$\sum_i \sum_y (\tau_{i,y})^2$
	$\sum_i \sum_y (N_{0,ij})^2$
	$\sum_i \sum_y (\varepsilon_{i,y})^2$

$v = 0.001.$

Definition	Equation		
Recruitment	$N_{i1,y} = R_{i,y} = R_{0,i} e^{\epsilon_{i,y}}$	$\epsilon_{i,y} \sim N(0, \sigma^2)$	T1.1
Initial abundance	$N_{ij,1} = \begin{cases} R_{0,i} e^{(-j M1_{ij})} N_{0,ij} & y = 1 \quad 1 < j \leq A_i \\ R_{0,i} e^{(-j M1_{i,A_i})} N_{0,i,A_i} / (1 - e^{(-j M1_{i,A_i})}) & y = 1 \quad j > A_i \end{cases}$		T1.2
Numbers at age	$N_{i,j+1,y+1} = N_{ij,y} e^{-Z_{ij,y}} \quad 1 \leq y \leq n_y \quad 1 \leq j < A_i$ $N_{i,A_i,y+1} = N_{i,A_i-1,y} e^{-Z_{i,A_i-1,y}} + N_{i,A_i,y} e^{-Z_{i,A_i,y}} \quad 1 \leq y \leq n_y \quad j > A_i$		T1.3
Catch	$C_{ij,y} = \frac{F_{ij,y}}{Z_{ij,y}} (1 - e^{-Z_{ij,y}}) N_{ij,y}$		T1.4
Total yield (kg)	$Y_{i,y} = \sum_j^{A_i} \left(\frac{F_{ij,y}}{Z_{ij,y}} (1 - e^{-Z_{ij,y}}) N_{ij,y} W_{ij,y} \right)$		T1.5
Biomass at age (kg)	$B_{ij,y} = N_{ij,y} W_{ij,y}$		T1.6
Spawning biomass at age (kg)	$SSB_{ij,y} = B_{ij,y} \rho_{ij}$		T1.7
Total mortality at age	$Z_{ij,y} = M1_{ij} + M2_{ij,y} + F_{ij,y}$		T1.8
Fishing mortality at age	$F_{ij,y} = F_{0,i} e^{\epsilon_{i,y}} s_{ij}^f$	$\epsilon_{i,y} \sim N(0, \sigma_{F,i}^2)$	T1.9
Weight at age (kg)	$W_{ij,y} = W_{\infty,iy} \left(1 - e^{(-K_i(1-d_{i,y})(j-t_{0i}))} \right)^{\frac{1}{1-d_{i,y}}}$		T1.10a
	$d_{i,y} = e^{(\alpha_{d,iy} + \alpha_{0d,i} + \beta_{d,i} T_y)}$		T1.10b
	$W_{\infty,iy} = \left(\frac{H_i}{K_i} \right)^{1/(1-d_{i,y})}$		T1.10c
BT survey biomass (kg)	$\hat{\beta}_{i,y}^s = \sum_j^{A_i} \left(N_{ij,y} e^{-0.5 Z_{ij,y}} W_{ij,y} s_{ij}^s \right)$		T1.11
EIT survey biomass (kg)	$\hat{\beta}_y^{eit} = \sum_j^{A_i} \left(N_{1j,y} e^{0.5(-Z_{1j,y})} W_{1j,y} s_{1j}^{eit} q_{1j}^{eit} \right)$	(pollock only)	T1.12
Fishery age composition	$\hat{O}_{ij,y}^f = \frac{C_{ij,y}}{\sum_j C_{ij,y}}$		T1.13
BT survey age composition	$\hat{O}_{ij,y}^s = \frac{N_{ij,y} e^{0.5(-Z_{ij,y})} s_{ij}^s}{\sum_j \left(N_{ij,y} e^{0.5(-Z_{ij,y})} s_{ij}^s \right)}$		T1.14
EIT survey age composition	$\hat{O}_{1j,y}^{eit} = \frac{N_{1j,y} e^{0.5(-Z_{1j,y})} s_{1j}^{eit} q_{1j}^{eit}}{\sum_j \left(N_{1j,y} e^{0.5(-Z_{1j,y})} s_{1j}^{eit} q_{1j}^{eit} \right)}$	(pollock only)	T1.15
BT selectivity	$s_{ij}^s = \frac{1}{1 + e^{(-b_i^s \cdot j - a_i^s)}}$		T1.16
Fishery selectivity	$s_{ij}^f = \begin{cases} e^{\eta_{ij}} & j \leq A_{\eta,i} \\ e^{\eta_{i,A_i}} & j > A_{\eta,i} \end{cases}$	$\eta_{ij} \sim N(0, \sigma_{f,i}^2)$	T1.17
Proportion females	$\omega_{ij} = \frac{e^{-j M_{fem}}}{e^{-j M_{fem}} + e^{-j M_{male}}}$		T1.18
Proportion of mature females	$\rho_{ij} = \omega_{ij} \phi_{ij}$		T1.19
Weight at age (kg)	$W_{ij,y} = W_{ij,y}^{fem} \omega_{ij} + (1 - \omega_{ij}) W_{ij,y}^{male}$		T1.20
Residual natural mortality	$M1_{ij} =: M_{ij}^{fem} \omega_{ij} + (\dots \omega_{ij}) \hat{M}_{ij}^{male}$		T1.21

Definition	Equation	
Recruitment	$N_{i1,y} = R_{i,y} = R_{0,i} e^{\tau_{i,y}}$	$\tau_{i,y} \sim N(0, \sigma^2)$
Initial abundance	$N_{ij,1} = \begin{cases} R_{0,i} e^{(-j M1_{ij})} N_{0,ij} \\ R_{0,i} e^{(-j M1_{iA_i})} N_{0,iA_i} / \left(1 - e^{(-j M1_{iA_i})}\right) \end{cases}$	$y = 1 \quad 1 < j \leq A_i$ $y = 1 \quad j > A_i$
Numbers at age	$N_{i,j+1,y+1} = N_{ij,y} e^{-Z_{ij,y}} \quad 1 \leq y \leq n_y \quad 1 \leq j < A_i$ $N_{iA_i,y+1} = N_{iA_i-1,y} e^{-Z_{iA_i-1,y}} + N_{iA_i,y} e^{-Z_{iA_i,y}} \quad 1 \leq y \leq n_y \quad j > A_i$	
Catch	$C_{ij,y} = \frac{F_{ij,y}}{Z_{ij,y}} (1 - e^{-Z_{ij,y}}) N_{ij,y}$	
Total yield (kg)	$Y_{i,y} = \sum_j^{A_i} \left(\frac{F_{ij,y}}{Z_{ij,y}} (1 - e^{-Z_{ij,y}}) N_{ij,y} W_{ij,y} \right)$	
Biomass at age (kg)	$B_{ij,y} = N_{ij,y} W_{ij,y}$	
Spawning biomass at age (kg)	$SSB_{ij,y} = B_{ij,y} \rho_{ij}$	
Total mortality at age	$Z_{ij,y} = M1_{ij} + M2_{ij,y} + F_{ij,y}$	
Fishing mortality at age	$F_{ij,y} = F_{0,i} e^{\varepsilon_{i,y}} S_{ij}^f$	$\varepsilon_{i,y} \sim N(0, \sigma_{F,i}^2)$
Weight at age (kg)	$W_{ij,y} = W_{\infty,iy} \left(1 - e^{(-K_i(1-d_{iy})(j-t_{0,i}))}\right)^{\frac{1}{1-d_{iy}}}$ $d_{i,y} = e^{(\alpha_{d,iy} + \alpha_{0d,i} + \beta_{d,i} T_y)}$ $W_{\infty,iy} = \left(\frac{H_i}{K_i}\right)^{1/(1-d_{iy})}$	
BT survey biomass (kg)	$\hat{\beta}_{i,y}^S = \sum_j^{A_i} \left(N_{ij,y} e^{-0.5 Z_{ij,y}} W_{ij,y} S_{ij}^S \right)$	

BT survey biomass (kg)

$$\hat{\beta}_{i,y}^s = \sum_j^{A_i} \left(N_{ij,y} e^{-0.5 Z_{ij,y}} W_{ij,y} S_{ij}^s \right)$$

EIT survey biomass (kg)

$$\hat{\beta}_y^{eit} = \sum_j^{A_1} \left(N_{1j,y} e^{0.5(-Z_{1j,y})} W_{1j,y} S_{1j}^{eit} q_{1j}^{eit} \right)$$

(pollock only)

Fishery age composition

$$\hat{O}_{ij,y}^f = \frac{C_{ij,y}}{\sum_j C_{ij,y}}$$

BT survey age composition

$$\hat{O}_{ij,y}^s = \frac{N_{ij,y} e^{0.5(-Z_{ij,y})} S_{ij}^s}{\sum_j \left(N_{ij,y} e^{0.5(-Z_{ij,y})} S_{ij}^s \right)}$$

EIT survey age composition

$$\hat{O}_{1j,y}^{eit} = \frac{N_{1j,y} e^{0.5(-Z_{1j,y})} S_{1j}^{eit} q_{1j}^{eit}}{\sum_j \left(N_{1j,y} e^{0.5(-Z_{1j,y})} S_{1j}^{eit} q_{1j}^{eit} \right)}$$

(pollock only)

BT selectivity

$$S_{ij}^s = \frac{1}{1 + e^{(-b_i^s \cdot j - a_i^s)}}$$

Fishery selectivity

$$S_{ij}^f = \begin{cases} e^{\eta_{ij}} & j \leq A_{\eta,i} \\ e^{\eta_{i,A_{\eta,i}}} & j > A_{\eta,i} \end{cases}$$

$\eta_{ij} \sim N(0, \sigma_{f,i}^2)$

Proportion females

$$\omega_{ij} = \frac{e^{-j M_{fem}}}{e^{-j M_{fem}} + e^{-j M_{male}}}$$

Proportion of mature females

$$\rho_{ij} = \omega_{ij} \phi_{ij}$$

Weight at age (kg)

$$W_{ij,y} = W_{ij,y}^{fem} \omega_{ij} + (1 - \omega_{ij}) W_{ij,y}^{male}$$

Residual natural mortality

$$M_{1j} = M_{1j}^{fem} \omega_{ij} + (1 - \omega_{ij}) M_{1j}^{male}$$

Table 8. Proportion mature and residual natural mortality for each species in the model.

Age	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Proportion mature																					
Walleye pollock	0.00	0.01	0.29	0.64	0.84	0.90	0.95	0.96	0.97	1.00	1.00	1.00									
Pacific cod	0.00	0.02	0.06	0.14	0.30	0.53	0.75	0.89	0.95	0.98	0.99	1.00									
Arrowtooth flounder	0.00	0.00	0.01	0.02	0.06	0.16	0.34	0.59	0.80	0.92	0.97	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Residual mortality (M1)																					
Walleye pollock	0.52	0.52	0.45	0.26	0.26	0.27	0.28	0.28	0.27	0.29	0.30	0.30									
Pacific cod	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37									
Arrowtooth flounder	0.27	0.26	0.26	0.25	0.25	0.24	0.24	0.23	0.23	0.23	0.22	0.22	0.22	0.22	0.21	0.21	0.21	0.21	0.21	0.21	0.21

Definition**Equation**

Predation mortality

$$M2_{ij,y} = \sum_{pa} \left(\frac{N_{pa,y} \delta_{pa,y} \bar{S}_{paij}}{\sum_{ij} (\bar{S}_{paij} B_{ij,y}) + B_p^{other} (1 - \sum_{ij} (\bar{S}_{paij}))} \right) \quad T2.1$$

Predator-prey suitability

$$\bar{S}_{paij} = \frac{1}{n_y} \sum_y \left(\frac{\frac{\bar{U}_{paij}}{B_{ij,y}}}{\sum_{ij} \left(\frac{\bar{U}_{paij}}{B_{ij,y}} \right) + \frac{1 + \sum_{ij} \bar{U}_{paij}}{B_p^{other}}} \right) \quad T2.2$$

Mean gravimetric diet
proportion

$$\bar{U}_{paij} = \frac{\sum_y U_{paij,y}}{n_y} \quad T2.3$$

Individual specific ration
(kg kg⁻¹ yr⁻¹)

$$\delta_{pa,y} = \hat{\varphi}_p \alpha_\delta W_{pa,y}^{(1+\beta_\delta)} f(T_y)_p \quad T2.4$$

Temperature scaling
algorithm

$$f(T_y)_p = V^X e^{X(1-V)} \quad T2.5$$


$$V = (T_p^{cm} - T_y) / (T_p^{cm} - T_p^{co}) \quad T2.5a$$

$$X = \left(Z^2 \left(1 + (1 + 40/Y)^{0.5} \right)^2 \right) / 400 \quad T2.5b$$


$$Z = \ln(Q_p^c) (T_p^{cm} - T_p^{co}) \quad T2.5c$$

$$Y = \ln(Q_p^c) (T_p^{cm} - T_p^{co} + 2) \quad T2.5d$$


CEATTLE 2016: Catch



CEATTLE 2016: Survey Biomass






Biomass






Understand mechanisms of change

Age 1 Mortality




Recruitment




Understand mechanisms of change

Age 1 Mortality


Recruitment




Understand mechanisms of change



Fishing mortality



Selectivity




CEATTLE 2016: Pollock age classes


years	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12+
1979	1.732	1.739	1.022	0.583	0.552	0.354	0.206	0.181	0.205	0.197	0.171	0.231
1980	0.933	3.884	2.283	0.988	0.498	0.356	0.181	0.095	0.088	0.098	0.091	0.177
1981	1.385	1.650	5.164	2.249	0.857	0.325	0.183	0.084	0.047	0.042	0.045	0.118
1982	0.955	1.914	2.056	4.925	1.992	0.610	0.192	0.099	0.046	0.025	0.022	0.080
1983	2.710	1.000	2.295	2.004	4.785	1.669	0.446	0.131	0.067	0.031	0.016	0.062
1984	0.993	3.639	1.299	2.457	2.203	4.698	1.466	0.369	0.108	0.055	0.025	0.058
1985	2.352	0.945	4.202	1.205	2.301	1.856	3.539	1.028	0.254	0.072	0.036	0.051
1986	1.246	2.378	1.091	3.914	1.146	1.958	1.413	2.520	0.722	0.175	0.048	0.055
1987	0.850	1.264	2.955	1.104	4.053	1.073	1.648	1.119	1.968	0.555	0.131	0.075
1988	0.508	0.652	1.402	2.730	1.063	3.592	0.865	1.252	0.830	1.437	0.394	0.143
1989	0.943	0.441	0.758	1.361	2.764	0.981	2.985	0.679	0.963	0.632	1.067	0.396
1990	2.375	0.704	0.404	0.571	1.053	1.928	0.607	1.715	0.379	0.525	0.335	0.759
1991	1.285	3.376	0.936	0.444	0.637	1.037	1.674	0.496	1.410	0.309	0.421	0.836
1992	0.866	1.549	4.165	0.916	0.435	0.544	0.778	1.175	0.346	0.969	0.206	0.792
1993	1.685	1.131	2.026	4.246	0.903	0.363	0.389	0.518	0.787	0.228	0.621	0.597
1994	0.722	2.344	1.432	1.961	3.970	0.708	0.245	0.244	0.325	0.486	0.136	0.701
1995	0.642	0.772	2.919	1.402	1.882	3.247	0.503	0.163	0.162	0.212	0.305	0.487
1996	1.265	0.670	1.009	3.083	1.486	1.733	2.638	0.384	0.124	0.121	0.154	0.552
1997	1.666	1.428	0.810	0.970	2.975	1.254	1.296	1.842	0.265	0.084	0.079	0.429
1998	1.016	2.186	1.853	0.841	1.018	2.753	1.036	1.005	1.416	0.200	0.061	0.346
1999	0.751	0.928	2.429	1.619	0.734	0.777	1.860	0.652	0.620	0.853	0.117	0.217
2000	1.155	0.912	1.197	2.505	1.688	0.672	0.635	1.429	0.497	0.465	0.621	0.231
2001	1.690	1.387	1.142	1.192	2.503	1.467	0.517	0.458	1.022	0.349	0.316	0.578
2002	1.130	2.216	1.825	1.195	1.241	2.243	1.154	0.381	0.337	0.743	0.246	0.600
2003	0.970	1.433	2.917	1.899	1.229	1.093	1.723	0.829	0.274	0.239	0.509	0.547
2004	0.470	1.017	1.759	2.840	1.816	1.002	0.773	1.134	0.545	0.176	0.149	0.635
2005	0.336	0.453	1.147	1.581	2.509	1.365	0.651	0.465	0.679	0.319	0.100	0.410
2006	0.743	0.325	0.520	1.075	1.483	2.019	0.949	0.421	0.302	0.433	0.198	0.293
2007	1.516	0.863	0.404	0.525	1.090	1.288	1.513	0.663	0.297	0.210	0.293	0.318
2008	0.666	2.198	1.088	0.399	0.513	0.904	0.920	1.005	0.443	0.195	0.134	0.377
2009	2.421	0.897	2.934	1.151	0.419	0.462	0.708	0.674	0.739	0.322	0.137	0.338
2010	1.375	4.306	1.234	3.156	1.225	0.385	0.372	0.535	0.507	0.548	0.230	0.321
2011	0.978	1.751	5.858	1.319	3.369	1.137	0.316	0.287	0.408	0.382	0.398	0.384
2012	0.607	1.023	1.937	4.935	1.073	2.320	0.678	0.174	0.155	0.215	0.192	0.378
2013	1.739	0.701	1.266	1.924	4.915	0.927	1.768	0.484	0.123	0.108	0.145	0.364
2014	1.461	2.087	0.866	1.257	1.931	4.335	0.727	1.301	0.352	0.088	0.075	0.331
2015	0.781	1.377	2.535	0.851	1.260	1.710	3.421	0.538	0.952	0.253	0.061	0.262
2016	0.504	0.596	1.623	2.464	0.845	1.106	1.332	2.496	0.389	0.675	0.174	0.206



CEATTLE 2016: Fish age / length comp



CEATTLE 2016: Srvy age / length comp



Projections for BRPs




Photo: Mark Holsman

CEATTLE results



Summary of assessment results for 2016:

Quantity	Walleye pollock		Pacific cod		Arrowtooth flounder	
	SSM	MSM	SSM	MSM	SSM	MSM
2016 M (natural mortality age 1)	0.518	2.291	0.367	0.979	0.269	0.852
2016 Average 3+ M (across ages)	0.297	0.315	0.370	0.370	0.226	0.229
2016 total (age 3+) biomass (t)	12,765,196	11,310,126	1,364,563	1,335,013	500,469	493,279
2016 SSB (female spawning biomass; t)	4,973,790	4,429,230	241,188	236,601	362,375	358,115
*Projected SSB_0 (t)	5,037,200	3,016,240	419,961	382,908	500,716	456,072
*Projected $SSB_{40\%}$ (t)	2,014,460	1,206,530	168,253	153,163	200,284	182,426
**Projected SSB_{mMSY}	2,482,140	2,781,050	146,696	141,982	2,229	6,717
ABC ₂₁₀₀ (t)	1,884,950	1,683,130	172,056	167,148	34,880	36,005
** $mMSY_{2100}$ (t)	1,817,130	2,363,310	170,896	168,683	1,095	3,282
$F_{40\%}$	0.577	1.155	0.372	0.404	0.115	0.130
F_{mMSY}	0.405	0.518	0.434	0.447	0.322	0.317

* SSB is based on the projected SSB at 2100 (equilibrium).



** $mMSY$ is aggregate multi-species yield

CEATTLE Recruitment: Pollock





Spawning biomass (million t)

CEATTLE Recruitment: P. cod




Spawning biomass (million t)

CEATTLE Recruitment: Arrowtooth




Spawning biomass (million t)


ABC multi-species proxy approach (Moffitt et al. in press)



ABC multi-species proxy approach (Moffitt et al. in press)



ABC multi-species proxy approach (Moffitt et al. in press)






Single-species


Harvest

Multi-species

Harvest scenario 1



Harvest scenario 1





Holsman et al. in press



NOAA FISHERIES

Single-species Multi-species




Differences between Harvest Scenarios Is greatest for prey spp


Holsman et al. in press




Multispecies control rules (Holsman et al. in press)




Projected SSB






Projected Recruitment (age 1)



Projected Yield



max F_{ABC}



NOAA FISHERIES

CEATTLE Applications

Photo: Mark Holsman

Ianelli et al. in press: Blended Forecasts

Deep-Sea Research II ■ (■■■■) ■■■-■■■



Contents lists available at [ScienceDirect](#)

Deep-Sea Research II

journal homepage: www.elsevier.com/locate/dsr2



Multi-model inference for incorporating trophic and climate uncertainty into stock assessments

James Ianelli^{a,*}, Kirstin K. Holsman^b, André E. Punt^c, Kerim Aydin^a

^a Alaska Fisheries Science Center NOAA Fisheries, 7600 Sand Point Way N.E., Building 4, Seattle, WA 98115, USA

^b University of Washington JISAO/Alaska Fisheries Science Center NOAA Fisheries, 7600 Sand Point Way N.E., Building 4, Seattle, WA 98115, USA

^c University of Washington School of Aquatic and Fisheries Sciences, 1122 NE Boat St., Seattle, WA 98105, USA

ARTICLE INFO

Keywords:

Model averaging
Model ensemble
Multi-species model



ABSTRACT

Ecosystem-based fisheries management (EBFM) approaches allow a broader and more extensive consideration of objectives than is typically possible with conventional single-species approaches. Ecosystem linkages may include trophic interactions and climate change effects on productivity for the relevant species within the system. Presently, models are evolving to include a comprehensive set of fishery and ecosystem information to address these broader management considerations. The increased


CEATTLE: Single-species

Mean R


Temp → W@A



Blended results (all three models)



Management Strategy Evaluations



ACLIM: Alaska Climate-change Integrated Modeling project

NOAA NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE

Alaska CLIMATE Project

Anne Hollowed (AFSC, SSMA/REFM)
Kirstin Holsman (AFSC, REEM/REFM)
Alan Haynie (AFSC ESSR/REFM)
Stephen Kasperski (AFSC ESSR/REFM)
Jim Ianello (AFSC, SSMA/REFM)
Kerim Aydin (AFSC, REEM/REFM)
Trond Kristiansen (IMR, Norway)
Al Hermann (UW JISAO/PMEL)
Wei Cheng (UW JISAO/PMEL)
André Punt (UW SAFS)

FATE: Fisheries & the Environment
SAAM: Stock Assessment Analytical Methods
S&T: Climate Regimes & Ecosystem Productivity



IPCC Scenarios (x3)

- AR4 A1B
- ARS RCP6.0
- ARS RCP8.5

Global Climate Models (x 11)

- ECHO-G (AR4 A1B)
- MIROC3.2 med res. (AR4 A1B)
- CGCM3-147 (AR4 A1B)
- CCSM4-NCAR-PO (ARS RCP 6.0 & 8.5)
- MIROCESM-C-PO (ARS RCP 6.0 & 8.5)
- GFDL-ESM2M*-PO (ARS RCP 6.0 & 8.5)
- GFDL-ESM2M*-PON (ARS RCP 6.0 & 8.5)


Future Climate Scenarios




Climate-enhanced Biological Models



Fishing Scenarios




Carbon Emission Scenarios



ACLIM

*The Alaska Climate change
Integrated Modeling project*




Holsman et al. in prep

Climate-specific Harvest &
Population Projections




NOAA FISHERIES


CEATTLE: Options




CEATTLE: Options



CEATTLE: Options





Slide courtesy of J. Duffy-Anderson



Thanks!

Photo: Mark Holsman

NOAA IEA program
FATE 13-07
SAAM 02

"Behind these numbers lies, of course, an infinity of movements and of destinies."

– von Bertalanffy 1938

...and of people!

Jim Ianelli, K. Aydin, Ingrid Spies, Grant Thompson

Stephanie Zador, Paul Spence, A. Hobday, A. Hollowed,, Isaac Kaplan, Troy Buckley, Matt Baker, Buck Stockhausen, P. Sean McDonald


Mark Holsman (Brother; photos)



END

























AFSC Ecosystem Based Fisheries Management





















Alaska Fisheries Science Center
NOAA Fisheries


Fisheries
Management

Levels	Scientific Advice	Management Framework
EBM Ecosystem Based Management	 Fisheries  Development  Energy  Eco Tourism  Oil & Gas	
	 Conservation  Shipping  Sanctuaries  Aquaculture  Etc	
EBFM Ecosystem Based Fisheries Management	 Fisheries  Climate  Habitat  Predator	
EAFM Ecosystem Approach to Fisheries Management	 Fisheries  Climate  Habitat  Predator	
SSFM Single Species	 Fisheries	



Levels	Scientific Advice	Management Framework
EBM Ecosystem Based Management	 Fisheries  Development  Energy  Eco Tourism  Oil & Gas	
	 Conservation  Shipping  Sanctuaries  Aquaculture  Etc	
EBFM Ecosystem Based Fisheries Management	 Fisheries  Climate  Habitat  Predator	
EAFM Ecosystem Approach to Fisheries Management	<div style="background-color: orange; color: white; padding: 10px; text-align: center;"> <p>Ecosystem Considerations Report Socio-Econ Report Ecosystem / ecological context</p> </div>	
SSFM Single Species		








When to use multi-species models:

- When there is sufficient data/information
- When expert advice is not enough
- Relationships are non-linear and complex
- Species are fully exploited
- Pressures/impacts are high, efficiency is needed
- Scope for fishery response to management is high
- When management is ready for tradeoffs analysis






Intergovernmental Panel on Climate Change (IPCC) 5th Assessment Report



"We figured it out!"

+ 1.8 °C



"Business as usual"

+4 °C



Table 4

Components of the likelihood function for each species *i* of age *j* in year *y*. EBS: Eastern Bering Sea. See [Tables 2–4](#) for parameter definitions.

Description	Equation	Data source	
Data components			
BT survey biomass	$\sum_i \sum_y \frac{[\ln(\beta_{i,y}^s) - \ln(\hat{\beta}_{i,y}^s)]^2}{2\sigma_{si}^2}$	NFMS annual EBS BT survey (1979–2012)	T4.1
BT survey age composition	$-\sum_i n_i \sum_y \sum_j (O_{ij,y}^s + v) \ln(\hat{O}_{ij,y}^s + v)$	NFMS annual EBS BT survey (1979–2012)	T4.2
EIT survey biomass	$\sum_y \frac{[\ln(\beta_y^{eit}) - \ln(\hat{\beta}_y^{eit})]^2}{2\sigma_{eit}^2}, \sigma_{eit} = 0.2$	Pollock acoustic trawl survey (1979–2012)	T4.3
EIT age composition	$-n \sum_y \sum_j (O_{1j,y}^{eit} + v) \ln(\hat{O}_{1j,y}^{eit} + v)$	Pollock acoustic trawl survey (1979–2012)	T4.4
Total catch	$\sum_i \sum_y \frac{[\ln(C_{i,y}^*) - \ln(\hat{C}_{i,y}^*)]^2}{2\sigma_c^2}, \sigma_c = 0.05$	Fishery observer data (1979–2012)	T4.5
Fishery age composition	$-\sum_i n_i \sum_y \sum_j (O_{ij,y}^f + v) \ln(\hat{O}_{ij,y}^f + v)$	Fishery observer data (1979–2012)	T4.6
Penalties			
Fishery selectivity	$\sum_i \sum_j A_i^{-1} \chi \cdot \left[\ln\left(\frac{\eta_{ij}^f}{\eta_{ij+1}^f}\right) - \ln\left(\frac{\eta_{ij+1}^f}{\eta_{ij+2}^f}\right) \right]^2, \chi = \begin{cases} 20, & \text{if } \eta_{ij}^f > \eta_{ij+1}^f \\ 0, & \text{if } \eta_{ij}^f \leq \eta_{ij+1}^f \end{cases}$		T4.7
Priors			
	$\sum_i \sum_y (\tau_{i,y})^2$		T4.8
	$\sum_i \sum_y (N_{0,i})^2$		T4.9
	$\sum_i \sum_y (\varepsilon_{i,y})^2$		T4.10

$v=0.001$.



Walleye pollock





Harvest scenario 1

Harvest scenario 2





Harvest scenario 3

Harvest scenario 4





Spawning biomass (million t)



Pacific cod



Arrowtooth flounder




Years




NOAA FISHERIES


Evaluate projection sensitivity

Temp effect \ll F effect




Holsman et al. in press. Deep Sea Res II






Get constant future
F rate that results in
40% unfished
biomass in year
2100







Find F rate that results in 40% unfished biomass in year 2100




Take yield in 2100 and plug it back into model to derive FABC




NOAA FISHERIES








max F_{ABC}




max F_{ABC}




NOAA FISHERIES

SSB at max ABC



Max ABC



Recruitment at max ABC

