

Public Testimony Sign-Up Sheet


Agenda Item D-1(d) BSAI Specs

NAME (PLEASE PRINT)		AFFILIATION
1	✓ AVE WOOD	US Seafoods
2	✓ John Gauvin	H + G workgroup
3	✓ Mike Szymanski / Bill McGill	FCA-97870 - Am 80 Limited Access ^{GROUP}
4	✓ ED RICHARDSON	POLLOCK CONSERVATION CONF.
5	✓ LORI Swanson	GFP
6	✓ GILES	Capt. O'Hara Corp.
7	✓ BRETT FRANK	LCF
8	✓ Bubba Cook	WWF
9	✓ Jen Warrick	Ocean
10	✓ Mike Hyde / Morgan Crow	American Seafoods / CVRF
11	✓ GLEN REED	PSPA
12	✓ GEORGE PLETNIKOF	GREEN PEACE
13	✓ DAVE FRASER	
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		

NOTE to persons providing oral or written testimony to the Council: Section 307(1)(I) of the Magnuson-Stevens Fishery Conservation and Management Act prohibits any person "to knowingly and willfully submit to a Council, the Secretary, or the Governor of a State false information (including, but not limited to, false information regarding the capacity and extent to which a United State fish processor, on an annual basis, will process a portion of the optimum yield of a fishery that will be harvested by fishing vessels of the United States) regarding any matter that the Council, Secretary, or Governor is considering in the course of carrying out this Act.

MEMORANDUM

TO: Council, AP and SSC

FROM: Chris Oliver 
Executive Director

DATE: November 27, 2007

SUBJECT: Bering Sea/Aleutians Islands SAFE Report and 2008/2009 harvest specifications

ESTIMATED TIME 8 HOURS (all D-1 items)
--

ACTION REQUIRED

Final action to approve the 2007 BSAI Stock Assessment and Fishery Evaluation (SAFE) report and final BSAI groundfish harvest specifications for 2008 and 2009:

1. Acceptable Biological Catch (ABC) and annual Total Allowable Catch (TAC)
2. Prohibited Species Catch Limits and seasonal apportionments of Pacific halibut, red king crab, Tanner crab, opilio crab, and herring to target fishery categories

BACKGROUND

At this meeting, the Council is scheduled to make final recommendations on groundfish and PSC specifications to manage the 2008 and 2009 Bering Sea/Aleutian Islands (BSAI) groundfish fisheries.

BSAI SAFE Report Since 2005, the Council has recommended ABCs and TACs for the next two fishing years and allows the preparation of updated assessments for species whose assessments are dependent largely on data from the EBS slope survey and the Aleutian Islands shelf survey. These surveys are conducted only in even-numbered years; therefore, the BSAI SAFE report does not contain new assessments for five rockfish categories.

The BSAI Groundfish Plan Team met in Seattle on November 13-17, 2007, to prepare the BSAI Groundfish SAFE report. The SAFE report forms the basis for BSAI groundfish harvest specifications for the 2008 and 2009 fishing years. The introduction to the BSAI SAFE report was mailed to the Council and Advisory Panel in late November 2007. The full report was mailed to the SSC.

The BSAI Groundfish Plan Team final recommendations for 2008 and 2009 are under Item D-1(d)(1). In September, preliminary projections of ABC and OFL were made on the basis of last year's stock assessments (Item D-1(d)(2)). In this SAFE report, the Plan Team has revised most of those projections. Such revisions are typically due to the development of new models; collection of new catch, survey, age composition, or size composition data; or use of new methodology for recommending ABCs. The SSC and AP recommendations will be provided to the Council during the meeting.

ABCs, TACs, and Apportionments The BSAI Groundfish Plan Team recommended OFLs and ABCs for 2008 and 2009. The sums of the recommended ABCs for 2008 and 2009 are 2,440,000 t and 2,560,000 t, respectively. They are approximately 236,000 t and 118,000 t below the sum of the 2007 ABCs. However, these values still exceed the 2 million t cap set by the Council as a conservation measure in setting TACs. Overall, the status of the stocks continues to appear favorable, although many stocks are

declining due to poor recruitment in recent years. The total biomass of 16.6 million t for 2008 declined by 300,000 t from 2007.

Overall groundfish exploitable biomass is high but declining, especially for pollock and Pacific cod. The bottom trawl survey biomass estimate for pollock in 2007 was 4.3 million t, only 87% of the long-term mean of the bottom-trawl survey. The 2007 echo-integration (EIT) survey biomass estimate was 1.88 million t, only 55% of the long-term mean for this survey. Both surveys indicate that the 2006 year class is strong and that the 2005 year class is now apparently below average. The biomass estimate from the 2007 bottom trawl survey for Pacific cod of 424,000 t is down about 18% from the 2006 estimate, and is the all-time low. Plan Team ABC recommendations are trending down for gadoids, but generally up for flatfishes. The abundances of AI pollock, sablefish, all rockfishes, all flatfishes, and Atka mackerel are projected to be above target stock size. The abundances of EBS pollock and Pacific cod are projected to be below target stock size.

The 2004 Consolidated Appropriations Act requires the Council to allocate pollock TAC to the Aleut Corporation for a directed pollock fishery in the Aleutian Islands. Starting in 2005, the Council has recommended a separate Total Allowable Catch (TAC) level of 19,000 t for the AI fishery. A mandatory 10% CDQ allocation (1,900 t) and an incidental catch allowance (ICA) of 1,600 t to cover bycatch of pollock in other AI fisheries are deducted from the TAC. The result is a directed pollock fishery allocation for the Aleut Corporation of 15,100 t. The Council has notified its intent to examine the ICA amount in recommending future AI pollock TACs.

Adopt prohibited species catch limits for Pacific halibut, crab, and herring

Beginning in 2008, the head and gut trawl catcher/processor sector, which targets flatfish, Pacific cod, and Atka mackerel, will be allocated groundfish TACs and PSCs among members of the "Amendment 80" sector that joined a cooperative. Regulations now require that crab and halibut trawl PSC be apportioned between the BSAI trawl limited access and Amendment 80 sectors after subtraction of prohibited species quota (PSQ) reserves, as presented in Table 7a for proposed 2008 and 2009 PSCs under Item D-1(d)(3). Crab and halibut trawl PSC assigned to the Amendment 80 sector is then sub-allocated to Amendment 80 cooperatives as PSC cooperative quota (CQ) and to the Amendment 80 limited access fishery as presented in Tables 7d and 7e. PSC CQ assigned to Amendment 80 cooperatives is not allocated to specific fishery categories. Regulations require the apportionment of each trawl PSC limit not assigned to Amendment 80 cooperatives be assigned into PSC bycatch allowances for seven specified fishery categories.

The Council may revise the proposed 2008 and 2009 fishery category allocations for the BSAI trawl limited access and the Amendment 80 limited access sectors as shown in Tables 7b, 7c, and 7e. Specifications for PSCs as shown in Tables 7a and 7d are fixed.

Halibut Trawl Fisheries: A 3,675 t limit on halibut mortality has been established for trawl gear. This limit can be apportioned to the trawl fishery categories as shown in the adjacent box.

Halibut Fixed Gear Fisheries: A 900 t non-trawl gear halibut mortality limit can be apportioned to the fishery categories listed in the adjacent box. Beginning in 2008, Amendment 85 divides the halibut PSC limit for the hook-and-line Pacific cod fishery

Categories used for prohibited species catch

Trawl fisheries

1. Greenland turbot, arrowtooth flounder and sablefish
2. rock sole, flathead sole, and "other flatfish"
3. yellowfin sole
4. rockfish
5. Pacific cod
6. pollock, Atka mackerel and "other species"

Non-trawl fisheries

1. Pacific cod
2. other non-trawl (longline sablefish and rockfish, and jig gear)
3. groundfish pot (exempt in recent years)

between the hook-and-line CP and CV sectors (CVs ≥ 60 ft (18.3 m) LOA and CVs < 60 ft (18.3 m) LOA combined). The Council can provide varying amounts of halibut PSC by season to each sector, tailoring PSC limits to suit the needs and timing of each sector.

Crab: Since 1997, prescribed bottom trawl fisheries in specific areas are closed when PSC limits of *C. bairdi* Tanner crab, *C. opilio* crab, and red king crab are taken. A stair step procedure for determining PSC limits for red king crab taken in Zone 1 trawl fisheries based on abundance of Bristol Bay red king crab has been in place. Based on the 2007 estimate of effective spawning biomass of 73 million pounds, the PSC limit for 2008 is 197,000 red king crabs. Up to 25% of the red king crab PSC limit can be used in the 56° - 56°10'N strip of the Red King Crab Savings Area. The red king crab cap has generally been allocated among the pollock/mackerel/other species, Pacific cod, rock sole, and yellowfin sole fisheries.

Species	Zone	Crab Abundance	PSC Limit
Red King Crab	Zone 1	< threshold or 14.5 million lb effective spawning biomass (ESB)	33,000
		> threshold, but < 55 million lb of ESB	97,000
		> 55 million lb of ESB	197,000
Tanner Crab	Zone 1	0-150 million crabs	0.5% of abundance
		150-270 million crabs	750,000
		270-400 million crabs	850,000
		> 400 million crabs	1,000,000
Tanner Crab	Zone 2	0-175 million crabs	1.2% of abundance
		175-290 million crabs	2,100,000
		290-400 million crabs	2,550,000
		> 400 million crabs	3,000,000

PSC limits for *bairdi* in Zones 1 and 2 have been based on total abundance of *bairdi* crab as indicated by the NMFS trawl survey. Based on 2007 abundance (787 million crab), and an additional reduction implemented in 1999, the PSC limit in 2008 for *C. bairdi* will be 980,000 (1,000,000 minus 20,000) *bairdi* crab in Zone 1 and 2,970,000 (3,000,000 minus 30,000) crab in Zone 2.

Since 1998, PSC limits for snow crab (*C. opilio*) are based on total abundance of *opilio* crab as indicated by the NMFS standard trawl survey. The snow crab PSC cap is set at 0.1133% of its abundance index, with a minimum PSC of 4.5 million snow crab and a maximum of 13 million snow crab. This number was further reduced by 150,000 crab in 1999. The 2007 survey estimate of 3.33 billion crabs results in a 2008 *opilio* crab PSC limit of 3,775,156 crabs, if left unadjusted. However, the crab FMP mandates a minimum of 4,350,000 snow crab. Snow crab taken within the "Snow Crab Bycatch Limitation Zone" accrues toward the PSC limits established for individual trawl fisheries.

Herring: In 1991, an overall herring PSC bycatch cap of 1 percent of the EBS biomass of herring was implemented. This cap is apportioned to the seven PSC fishery categories. Annual herring assessments indicate there will be very little change in the Bering Sea herring PSC limit for 2008. The herring biomass estimate for spring 2007 for the eastern Bering Sea was 178,652 t. The corresponding herring PSC limit for 2007 at 1% of this amount was be 1,787 t. ADF&G will provide the 2008 herring biomass estimate at the meeting.

Seasonal apportionment of bycatch limits The Council may also seasonally apportion the bycatch allowances. Regulations require that seasonal apportionments of bycatch allowances be based on information listed in the adjacent box.

Halibut discard mortality rates Halibut bycatch mortality rates for the 2007-2009 open access fisheries were adopted by the Council in October 2006. In October 2007, the Council adopted International Pacific Halibut Commission staff recommendations for DMRs for the 2008 BSAI CDQ fisheries (shown in the summary table below). Rates for CDQ fisheries will likely be set on a 3-year cycle when the next cycle commences for the non-CDQ fisheries in 2009 for 2010-2012.

- Factors to be considered for seasonal apportionments of bycatch allowances.**
1. Seasonal distribution of prohibited species;
 2. Seasonal distribution of target groundfish species relative to prohibited species distribution;
 3. Expected prohibited species bycatch needs on a seasonal basis relevant to change in prohibited species biomass and expected catches of target groundfish species;
 4. Expected variations in bycatch rates throughout the fishing year;
 5. Expected changes in directed groundfish fishing seasons;

Minutes from the BSAI Groundfish Plan Team meeting will be distributed at the meeting.

CDQ Fisheries	
Gear/Target	Recommended DMR
<i>Trawl</i>	
Atka mackerel	85
Bottom pollock	80
Rockfish	82
Flathead sole	87
Pelagic pollock	90
Rock sole	86
Yellowfin sole	86
<i>Pot</i>	
Sablefish	34
<i>Longline</i>	
Pacific cod	10
Turbot	4

Table 1. B Sea Aleutian Islands Groundfish Plan Team OFL, ABC, and TAC Recommendations for the 2008-2009 Fisheries (revised 11/29/07)

Species	Area	2007				2008			2009		
		OFL	ABC	TAC	Catch	OFL	ABC	TAC	OFL	ABC	TAC
Pollock	EBS	1,640,000	1,394,000	1,394,000	1,350,000	1,440,000	1,000,000		1,320,000	1,000,000	
	Aleutian Islands	54,500	44,500	19,000	2,488	34,000	28,200		26,100	22,700	
	Bogoslof District	48,000	5,220	10	0	58,400	7,970		58,400	7,970	
Pacific cod	BSAI	207,000	176,000	171,000	172,655	176,000	150,000		190,000	162,000	
Sablefish	BS	3,520	2,980	2,980	1,090	3,380	2,860		2,910	2,610	
	AI	3,320	2,810	2,810	1,080	2,890	2,440		2,510	2,230	
Yellowfin sole	BSAI	240,000	225,000	136,000	119,332	265,000	248,000		296,000	276,000	
Greenland turbot	Total	15,600	2,440	2,440	1,946	15,600	2,540		16,000	2,540	
	BS		1,680	1,680	1,435		1,750			1,750	
	AI		760	760	511		790			790	
Arrowtooth flounder	BSAI	193,000	158,000	20,000	11,700	297,000	244,000		300,000	246,000	
Northern rock sole	BSAI	200,000	198,000	55,000	37,013	304,000	301,000		379,000	375,000	
Flathead sole	BSAI	95,300	79,200	30,000	19,500	86,000	71,700		83,700	69,700	
Alaska plaice	BSAI	241,000	190,000	25,000	19,411	248,000	194,000		277,000	217,000	
Other flatfish	BSAI	28,500	21,400	10,000	5,840	28,800	21,600		28,800	21,600	
Pacific Ocean perch	BSAI	26,100	21,900	19,900	17,800	25,700	21,700		25,400	21,300	
	BS		4,160	2,160	811		4,200			4,100	
	AI total		17,740	17,740	16,960		17,500			17,200	
	WAI		7,720	7,720	7,421		7,610			7,490	
	CAI		5,050	5,050	4,423		4,990			4,900	
	EAI		4,970	4,970	5,116		4,900			4,810	
Northern rockfish	BSAI	9,750	8,190	8,190	3,940	9,740	8,180		9,680	8,130	
Shortraker	BSAI	564	424	424	318	564	424		564	424	
Rougheye	BSAI	269	202	202	163	269	202		269	202	
Other rockfish	BSAI	1,330	999	999	635	1,330	999		1,290	968	
	BS		414	414	205		414			414	
	AI		585	585	430		585			554	
Atka mackerel	Total	86,900	74,000	63,000	56,620	71,400	60,700		50,600	47,500	
	WAI		20,600	9,600			16,900			13,200	
	CAI		29,600	29,600			24,300			19,000	
	EAI/BS		23,800	23,800			19,500			15,300	
Squid	BSAI	2,620	1,970	1,970	1,190	2,620	1,970		2,620	1,970	
Other species	BSAI	91,700	68,800	37,400	26,500	91,200	71,800		91,200	71,800	
	Sharks					617	463		617	463	
	Skates					37,200	31,300		36,800	30,900	
	Sculpins					53,100	39,800		53,100	39,800	
	Octopus					324	243		324	243	
Total	BSAI	3,188,973	2,676,035	2,000,000	1,849,221	3,161,893	2,440,285		3,162,043	2,557,644	

**2007 catch is through October 27, 2007 (includes CDQ and state water harvests).

Council Proposed BSAI Specifications

Rollover of 2008 TACs for 2009

Species	Area	2007				2008			2009		
		OFL	ABC	TAC	Catch	OFL	ABC	TAC	OFL	ABC	TAC
Pollock	EBS	1,640,000	1,394,000	1,394,000	1,168,092	1,431,000	1,318,000	1,318,000	1,431,000	1,318,000	1,318,000
	AI	54,500	44,500	19,000	2,394	50,300	41,000	19,000	50,300	41,000	19,000
	Bogoslof	48,000	5,220	10	0	48,000	5,220	10	48,000	5,220	10
Pacific cod	BSAI	207,000	176,000	170,720	148,349	154,000	131,000	127,070	154,000	131,000	127,070
Sablefish	BS	3,520	2,980	2,980	793	3,290	2,970	2,970	3,290	2,970	2,970
	AI	3,320	2,810	2,810	915	3,100	2,800	2,800	3,100	2,800	2,800
Yellowfin sole	BSAI	240,000	225,000	136,000	116,103	261,000	245,000	150,000	261,000	245,000	150,000
Greenland turbot	Total	15,600	2,440	2,440	1,716	16,000	2,490	2,490	16,000	2,490	2,490
	BS		1,680	1,680	1,307		1,720	1,720		1,720	1,720
	AI		760	760	409		770	770		770	770
Arrowtooth flounder	BSAI	193,000	158,000	20,000	9,441	208,000	171,000	30,000	208,000	171,000	30,000
Northern rock sole	BSAI	200,000	198,000	55,000	36,648	271,000	268,000	75,000	271,000	268,000	75,000
Flathead sole	BSAI	95,300	79,200	30,000	17,685	92,800	77,200	45,000	92,800	77,200	45,000
Alaska plaice	BSAI	241,000	190,000	25,000	19,176	252,000	199,000	60,000	252,000	199,000	60,000
Other flatfish	BSAI	28,500	21,400	10,000	5,470	28,500	21,400	21,400	28,500	21,400	21,400
Pacific Ocean perch	BSAI	26,100	21,900	19,900	16,166	25,600	21,600	21,600	25,600	21,600	21,600
	BS		4,160	2,160	596		4,080	4,080		4,080	4,080
	AI total		17,740	17,740	15,570		17,520	17,520		17,520	17,520
	WAI		7,720	7,720	7,063		7,620	7,620		7,620	7,620
	CAI		5,050	5,050	3,640		5,000	5,000		5,000	5,000
	EAI		4,970	4,970	4,867		4,900	4,900		4,900	4,900
	BSAI	9,750	8,190	8,190	1,335	9,700	8,150	8,150	9,700	8,150	8,150
Northern rockfish	BSAI	564	424	424	324	564	424	424	564	424	424
Shortraker	BSAI	269	202	202	151	269	202	202	269	202	202
Rougheye	BSAI	1,330	999	999	480	1,330	999	999	1,330	999	999
Other rockfish	BS		414	414	157		414	414		414	414
	AI		585	585	323		585	585		585	585
	Total	86,900	74,000	63,000	27,904	64,200	54,900	54,900	64,200	54,900	54,900
Atka mackerel	WAI		20,600	9,600	484		15,300	15,300		15,300	15,300
	CAI		29,600	29,600	8,030		22,000	22,000		22,000	22,000
	EAI/BS		23,800	23,800	19,390		17,600	17,600		17,600	17,600
Squid	BSAI	2,620	1,970	1,970	921	2,620	1,970	1,970	2,620	1,970	1,970
Other species	BSAI	91,700	68,800	37,355	22,582	91,700	68,800	58,015	91,700	68,800	58,015
Total	BSAI	3,188,973	2,676,035	2,000,000	1,596,645	3,014,973	2,642,125	2,000,000	3,014,973	2,642,125	2,000,000

Sources: 2007 and 2008 OFLs, ABCs, and TACs from the specifications adopted by the Council in 12-07; 2009 OFLs and ABCs equal to 2008; 2007 catches through 9-8-07 from AKR Catch Accounting .

TABLE 7a-PROPOSED 2008 AND 2009 APPORTIONMENT OF PROHIBITED SPECIES CATCH ALLOWANCES TO NON-TRAWL GEAR, THE CDQ PROGRAM, AMENDMENT 80, AND THE BSAI TRAWL LIMITED ACCESS SECTORS

PSC species	Total non-trawl PSC	Non-trawl PSC remaining after CDQ PSQ ²	Total trawl PSC	Trawl PSC remaining after CDQ PSQ ²	CDQ PSQ reserve ²	Amendment 80 sector		BSAI trawl limited access fishery
						2008	2009	
Halibut mortality (mt) BSAI	900	832	3,675	3,400	343	2,525	2,475	875
Herring (mt) BSAI	n/a	n/a	1,787	n/a	n/a	n/a	n/a	n/a
Red king crab (animals) Zone 1 ¹	n/a	n/a	197,000	175,921	21,079	109,915	104,427	53,797
<u>C. opilio</u> (animals) COBLZ ¹	n/a	n/a	4,350,000	3,884,550	465,450	2,386,668	2,267,412	1,248,494
<u>C. bairdi</u> crab (animals) Zone 1 ¹	n/a	n/a	980,000	875,140	104,860	460,674	437,658	411,228
<u>C. bairdi</u> crab (animals) Zone 2 ¹	n/a	n/a	2,970,000	2,652,210	317,790	784,789	745,536	1,241,500

¹ Refer to 50 CFR § 679.2 for definitions of areas.

² Sections 679.21(e)(3)(i) and (e)(4)(i)(A) allocate 276 mt of the trawl halibut mortality and 7.5 percent, or 67 mt, of the non-trawl halibut mortality limit as the PSQ reserve for use by the groundfish CDQ program. The PSQ reserve for crab species is 10.7 percent of each crab PSC limit.

TABLE 7b—PROPOSED 2008 AND 2009 HERRING AND RED KING CRAB SAVINGS SUBAREA PROHIBITED SPECIES CATCH ALLOWANCES FOR ALL TRAWL SECTORS

Trawl gear	Herring (mt) BSAI	Red king crab (animals) Zone 1
Yellowfin sole	153	n/a
Rock sole/flathead sole/other flatfish ¹	27	n/a
Turbot/arrowtooth/sablefish ²	12	n/a
Rockfish	n/a	n/a
July 1 - December 31	10	n/a
Pacific cod	27	n/a
Midwater trawl pollock	1,364	n/a
Pollock/Atka mackerel/other species ³	194	n/a
Red king crab savings subarea	n/a	n/a
Non-pelagic trawl gear ⁴	n/a	49,250
Total trawl PSC	1,787	197,000

¹ "Other flatfish" for PSC monitoring includes all flatfish species, except for halibut (a prohibited species), flathead sole, Greenland turbot, rock sole, yellowfin sole, and arrowtooth flounder.

² Greenland turbot, arrowtooth flounder, and sablefish fishery category.

³ Pollock other than pelagic trawl pollock, Atka mackerel, and "other species" fishery category.

⁴ In October 2007 the Council recommended that the red king crab bycatch limit for non-pelagic trawl fisheries within the RKCSS be limited to 25 percent of the red king crab PSC allowance (see § 679.21(e)(3)(ii)(B)(2)).

TABLE 7c-PROPOSED 2008 AND 2009 PROHIBITED SPECIES BYCATCH ALLOWANCES FOR THE BSAI TRAWL LIMITED ACCESS SECTOR AND NON-TRAWL FISHERIES

BSAI trawl limited access fisheries	Prohibited species and zone				
	Halibut mortality (mt) BSAI	Red king crab (animals) Zone 1 ¹	<i>C. opilio</i> (animals) COBLZ ¹	<i>C. bairdi</i> (animals)	
				Zone 1 ¹	Zone 2 ¹
Yellowfin sole	145	29,938	1,170,367	259,003	1,036,505
Rock sole/flathead sole/other flatfish ²	0	0	0	0	0
Turbot/arrowtooth/sablefish ³	0	0	0	0	0
Rockfish	n/a	n/a	n/a	n/a	n/a
June 1 - December 31	3	n/a	2,000	n/a	1,000
Pacific cod	577	23,499	45,677	139,138	188,058
Pollock/Atka mackerel/other species ⁴	150	360	30,451	13,087	15,937
Total BSAI trawl limited access PSC	875	53,797	1,248,494	411,228	1,241,500
Non-trawl fisheries	Catcher processor	Catcher vessel			
Pacific cod-Total	760	15			
January 1-June 10	314	10			
June 10-August 15	0	0			
August 15-December 31	446	5			
Other non-trawl-Total		58			
May 1-December 31		58			
Groundfish pot and jig		exempt			
Sablefish hook-and-line		exempt			
Total non trawl PSC		833			

¹ Refer to § 679.2 for definitions of areas.

² "Other flatfish" for PSC monitoring includes all flatfish species, except for halibut (a prohibited species), flathead sole, Greenland turbot, rock sole, yellowfin sole, and arrowtooth flounder.

³ Greenland turbot, arrowtooth flounder, and sablefish fishery category.

TABLE 7d-PROPOSED 2008 AND 2009 PROHIBITED SPECIES BYCATCH ALLOWANCES FOR THE BSAI AMENDMENT 80 COOPERATIVES

Year	Prohibited species and zone				
	Halibut mortality (mt) BSAI	Red king crab (animals) Zone 1 ¹	<u>C. opilio</u> (animals) COBLZ ¹	<u>C. bairdi</u> (animals)	
				Zone 1 ¹	Zone 2 ¹
2008	1,837	78,631	1,632,432	340,520	580,311
2009	1,801	74,704	1,550,864	323,507	551,286

¹ Refer to § 679.2 for definitions of areas.

TABLE 7e-PROPOSED 2008 AND 2009 PROHIBITED SPECIES BYCATCH ALLOWANCES FOR THE BSAI AMENDMENT 80 LIMITED ACCESS FISHERIES

Amendment 80 trawl limited access fisheries	Prohibited species and zone									
	Halibut mortality (mt) BSAI		Red king crab (animals) Zone 1 ¹		C. opilio (animals) COBLZ ¹		C. bairdi (animals) Zone 1 ¹		Zone 2 ¹	
	2008	2009	2008	2009	2008	2009	2008	2009	2008	2009
Yellowfin sole	190	186	5,810	5,520	580,761	551,742	45,178	42,921	133,115	126,457
Jan 20 - Apr 1	63	62	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Apr 1 - May 21	39	38	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
May 21 - Jul 1	10	10	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Jul 1 - Dec 31	77	75	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Rock sole/other flat/flathead sole ²	168	164	20,844	19,803	120,677	114,647	48,422	46,003	44,372	42,152
Jan 20 - Apr 1	101	99	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Apr 1 - Jul 1	33	33	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
July 1 - Dec 31	34	33	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Turbot/arrowtooth/sablefish ³	n/a	0	n/a	n/a	7,542	7,165	n/a	n/a	n/a	n/a
Rockfish	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Jul 1 - Dec 31	14	14	n/a	n/a	7,542	7,165	n/a	n/a	818	777
Pacific cod	270	265	4,560	4,332	22,627	21,496	24,271	23,058	24,129	22,922
Pollock/Atka mackerel/other ⁴	47	46	69	66	15,085	14,331	2,283	2,169	2,045	1,943
Total Amendment 80 trawl limited access PSC	688	674	31,284	29,722	754,235	716,548	120,154	114,151	204,477	194,250

¹ Refer to § 679.2 for definitions of areas.

² "Other flatfish" for PSC monitoring includes all flatfish species, except for halibut (a prohibited species), flathead sole, Greenland turbot, rock sole, yellowfin sole, and arrowtooth flounder.

³ Greenland turbot, arrowtooth flounder, and sablefish fishery category.

⁴ Pollock other than pelagic trawl pollock, Atka mackerel, and "other species" fishery category.

FRED A. YECK, President
(541) 867-3911

AGENDA D-1(d)
DECEMBER 2007
Supplemental

F/V Seadawn Fisheries, Inc.

P.O. Box 352 • Newport, Oregon 97365
Fax (541) 867-3913



F/V Seadawn

November 17, 2007

North Pacific Fishery Management Council
605 W. 4th Avenue, Suite 306
Anchorage, Alaska 99501

RE: BERING SEA POLLOCK TAC - Agenda Item D-1(e)

Dear Chairman Olson and Council Members:

I feel compelled to write because of my concern with regard to the status of the Bering Sea Pollock stocks and the need for immediate action to reduce the TAC to protect that stock for the long term.

I am the managing owner of a 124 foot AFA catcher vessel which delivers inshore to Unisea. My family has owned this vessel since 1987 and we have been engaged in the Bering Sea Pollock fishery continuously since that time to the present. During the past four seasons we have seen the pollock available for harvest steadily decline, especially during the Pollock B Season, to the extent we could no longer obtain consistent loads in the traditional fishing areas within 250 miles of Dutch Harbor during the later half of B Season. However, up until last year we were able to find adequate stocks 400-500 miles north of Dutch Harbor but last season, even up against the Russian border, 500 plus miles north of Dutch catch rates became very slow. The pollock stocks have been showing the stress of being hammered by the entire pollock fleet in this northern area.

Traditionally, our trips were 2-3 days for consistent full loads. Now 5-6 days have become the norm and partial loads have also become almost normal. In the past several seasons fishing improved in October in our more traditional areas within 250 miles of Dutch Harbor, but this year fishing remained very difficult.

In past years it was relatively easy for the entire industry to harvest its quota well prior to the end of the season. As the record will now reflect, a substantial portion of the pollock quota remains unharvested.

These factors are reflective of the fact that the pollock resource was simply not adequate to support last year's TAC. The pollock resource available for harvest that we are seeing is the lowest in 20 years and it is disheartening to those who are used to a robust stock.

We are very concerned about the future of this resource and believe a very substantial reduction in TAC is necessary for its protection.

I am pleased to note and it gives me some reassurance in the system, that the Council's Plan Team has recommended a TAC not to exceed 1,000,000 metric tons. This recommendation is consistent with what the vessels are seeing on the grounds. I believe 1,000,000 metric tons should be considered the high end for 2008 Bering Sea TAC.

We have been involved in the pollock fishery for the past 20 years, my son is currently one of the Captains of our vessel and it is our hope that this fishery could last for the next 20 years and more. I am old enough to have witnessed first hand the over fishing that occurred under the watch of NMFS and the PFMFC on the West Coast 25-30 years ago which resulted in the collapse of the West Coast traditional groundfish fishery. I do not want to see that happen in the Bering Sea pollock fishery but action by the Council and NMFS at this time is imperative.

The NPFMC prides itself on claiming to have the best managed fisheries in the nation. That is relatively easy when quotas have been at all time highs. The test of our management is upon you now with the declining pollock stock. I trust the Council and NMFS will have the courage to prove in fact it can manage the resource for the long term and ignore those who will be pressing for short term profits.

The TAC must be reduced by at least 25-30% and if conditions do not improve, even further in years to come.

Respectfully submitted,



Fred A. Yeck
President

cc: Roy Hyder



125 Christensen Dr., Suite 2
Anchorage, AK 99501

Tel.: 907-277-8234

Fax: 907-272-6519

November 28, 2007

Mr. Eric Olsen, Chair
North Pacific Fisheries Management Council
605 W. 4th Avenue, Suite 306
Anchorage, AK 99501

NOV 29 2007
11:30 AM

Re: Agenda Item D-4, Groundfish Management

Dear Mr. Olsen and Council Members:

Greenpeace, a global environmental organization with nearly three million members and a long presence in Alaska, urges the Council to take a more precautionary, ecosystem-based approach to groundfish management. Together with nearly 7,000 others who have written you this week, we call on you to set catch limits that seek at a minimum to double in size the pollock population and maintain this higher level over time.

Please note that there was considerable concern expressed by Plan Team members about the health of Bering Sea pollock stocks. The Plan Team ultimately recommended an Allowable Biological Catch of 1,000,000 tons. Their recommendation was set lower than the maximum allowable Tier 1B allotment to reflect concerns about poor recruitment and the fact that the stock is at an all-time low. Several Plan Team members recommended an ABC of 555,000 tons, feeling that Bering Sea pollock would now be more appropriately managed under Tier 3.

This Council has long prided itself for taking what it views to be precautionary measures. There are numerous reasons for taking a precautionary approach to management of the Bering Sea pollock fishery at this time:

- Five consecutive years of below average recruitment
- Stock levels at all-time low
- No closures in place to protect pollock Essential Fish Habitat
- Similar management approach failed to protect pollock stocks in Shelikof Strait, Bogoslof, or the Aleutian Islands.
- Unprecedented spawning exploitation rates
- Heavy fishing pressure on Bering Sea spawning aggregation
- Increasing Catch per Unit Effort is exacerbating already significant bycatch concerns
- Concerns from numerous native communities about localized depletion, habitat destruction, bycatch, and food web impacts of trawl fisheries
- Continued damage to and bycatch of protected corals in known coral habitats by "pelagic" and bottom trawls

- Increasing uncertainty associated with effects of climate change
- Unprecedented levels of concern from long-time pollock fleet captains
- Increasingly clear evidence that fisheries are jeopardizing the recovery of endangered Steller sea lions and causing the depletion of northern fur seals.
- Percent catch inside the Steller sea lion Conservation Area during the pollock A season reached 57% in 2006, the highest level since 1998.
- Several species of pelagic sea birds are in decline

Setting the ABC at such a high level is not justifiable in these circumstances. Setting a TAC at a level approaching the ABC is likely to create significant problems for pollock stocks as well as for pollock predators, including endangered Steller sea lions and depleted northern fur seals. By continuing to ignore the many warning signs, the Council is putting the region at risk of a fishery collapse that would have devastating impacts on the economy, on native and coastal communities, and the entire food web.


The tools to rebuild pollock stocks and provide the basis for strong sustainable fisheries as well as healthy communities and ecosystems are already known to you. Here are a few examples of the steps that should be taken:

- Set catch limits that seek at a minimum to double in size the pollock population and maintain this higher level over time
- Maintain a buffer of at least 20% between the ABC and the TAC
- Establish a hard bycatch cap for king salmon and other vulnerable species
- Establish a network of marine reserves that include portions of pollock EFH

Failure to reduce bycatch to the extent practicable or to avoid jeopardizing the survival and recovery of endangered Steller sea lions unnecessarily opens up the Council to litigation and harms the fisheries and fishermen of Alaska.

Greenpeace will continue to work with the Council to assist efforts to manage Alaska fisheries in a more precautionary, ecosystem-based manner. We hope that our experience with fisheries in other parts of the world might be of use in helping the Council prevent the types of disastrous outcomes that have taken place elsewhere.

Sincerely,


George Pletnikoff
Alaska Oceans Campaigner



175 South Franklin Street, Suite 418 +1.907.586.4050
Juneau, AK 99801 USA www.oceana.org

November 19, 2007

Mr. Eric Olson, Chair
North Pacific Fishery Management Council
605 W. Fourth Avenue, Suite 306
Anchorage, AK 99501-2252

Dr. Jim Balsiger, Regional Administrator
NOAA Fisheries, Alaska Region
709 West Ninth Street
Juneau, AK 99802-1668

RE: 2008/2009 TAC setting

Dear Chairman Olson and Dr. Balsiger:

Each year, the National Marine Fisheries Service (Fisheries Service) authorizes the catch of billions of wild fish from the Bering Sea, Aleutian Islands, and Gulf of Alaska. This action, made within the complex catch strategy of the North Pacific Fishery Management Council (Council) process, has both known and unknown ecosystem consequences.

The Council has considerable flexibility when recommending fishing quotas each year. The upper optimum yield cap is just that, a cap. It is neither a goal nor a target, and should not be treated as something to "fill" each year. Rather, optimum yield was set as a range of values to allow the Council to provide a coarse check on fishery removals across varying ecological and ecosystem conditions. The Bering Sea/Aleutian Islands (BSAI) optimum yield cap is not 2 million metric tons, but is supposed to be an amount within a range between 1.4 and 2 million metric tons. Likewise, the optimum yield cap for the Gulf of Alaska (GOA) is to be set within a range of 116 to 800 thousand metric tons. Given prevailing ecological and ecosystem conditions and the implication of fishery removals, the Council and agency must carefully consider what to allow to be removed from the ecosystem in 2008 and 2009.

Bering Sea Pollock

The rapidly declining pollock stock in the eastern Bering Sea is of grave concern. Pollock play a central role in the North Pacific ecosystem and are responsible for much of the structure and function of the Bering Sea food web. Pollock, from juvenile to adult, function as both prey and predator for many species. Pollock are critical prey for endangered Steller sea lions, fur seals, and many other species. As predators, pollock may play a role regulating arrowtooth flounder populations.

The commercial fishery has rapidly driven the pollock stock to very low levels, and the decline likely has been hastened by increased fishing pressure on the spawning stock. In 2006 and 2007, the fishery removed more than 25% of the spawning aged female pollock¹. This level of sustained fishing pressure on the spawning stock was unprecedented, and the stock likely cannot withstand it on a sustained basis. The spawning biomass of pollock is now estimated at only 21

to 28%¹ of what there would be if commercial harvests had not occurred. In other words, without commercial fishing, the Bering Sea would have 72 to 79% greater spawning biomass of pollock than we have today.

Given the rapidly declining stock and increasing fishing pressure on the spawning biomass, the Plan Team recommended a maximum Acceptable Biological Catch (ABC) for pollock of 1 million metric tons, which is slightly less than the maximum permitted under the Tier 1 rules. We do not support such a large pollock harvest in 2008 and 2009. Several Plan Team members and a shore-based pollock harvester also questioned the high harvest rates associated with Tier 1 classification of pollock. Considerable uncertainty surrounds the reference points used to determine Tier status and harvest controls. Being on the right side of a theoretical line doesn't eliminate the potential for ecological collapse. In recent years we've seen the longest stretch of poor pollock recruitment on record. While a potentially abundant 2006 year class was seen in the surveys this year, there is no guarantee that this will be sufficient to rebuild the pollock stock. Indeed, the stock assessment authors caution; *"the high degree of uncertainty in the magnitude of these year classes cannot be overemphasized, particularly as they extend to estimates of future stock size."*² A pollock harvester recently testified at a Plan Team meeting that *"it felt like we were catching every last pollock in the Bering Sea this year."*

Scientists agree that global warming effects will be felt in the Bering Sea, and that we are currently seeing signals of these effects. While it is difficult to predict the ecological trajectory, there is general agreement that we have greater uncertainty in fishery projections and thus will have to manage in a more precautionary fashion³.

For these reasons, we urge the SSC and the Council to adopt the SAFE author's Tier 3b recommendations for EBS pollock in the table below:

Tier	Year	Max ABC	OFL
3b	2008	555 thousand mt	677 thousand mt
3b	2009	650 thousand mt	794 thousand mt

Flatfish

We are also concerned about the impacts of a substantially larger flatfish harvest in 2008 and 2009. Flatfish fisheries may undergo a significant increase and redistribution of effort if a substantially larger quota is granted. While flatfish harvests have been large in the past, particularly the early 1960's when yellowfin sole were overexploited by foreign trawlers, the flatfish fishery in more recent history has been much smaller.

The flatfish fisheries use bottom trawls, and the impacts of bottom trawls on habitat are well documented⁴. Any increase in the use of bottom trawls and any significant increase or

¹ EBS Walleye Pollock 2007 SAFE

² EBS Walleye Pollock 2007 SAFE

³ Testimony of Dr. Gordon Kruse before the U.S. Senate hearing on the effects of climate change on living marine resources, May 2007

⁴ NAS. 2002. Effects of trawling and dredging on seafloor habitat

redistribution of fishing effort requires a credible analysis of habitat impacts. Existing analyses are not sufficient for this purpose.

The Essential Fish Habitat (EFH) EIS cannot be used to predict the impacts of a large increase in bottom trawl effort. The EFH EIS reached a conclusion regarding commercial fishing impacts on habitat based on an analysis of fishing effort from 1998 to 2002. This 5-year period ostensibly was selected to represent the current level of fishing effects. Regardless of the accuracy of that assumption or the EFH EIS's conclusions, the projected 2008 yellowfin sole quota is almost double the average catch used to determine trawling impacts in the EFH EIS. The EFH analysis therefore cannot be used to address impacts to seafloor habitat because it underestimates future effort, area swept, and total habitat damaged by trawls.

In addition, the Alaska Groundfish Harvest Specifications Final EIS (NMFS, 2007) did not address the effects of significant increases or redistribution of fishing effort. As we stated in our October 2006 comment letter on the Alaska Groundfish Harvest Specifications Draft EIS, "The DEIS references the flawed conclusion reached in the EFH EIS that no effects of fishing on EFH are more than minimal, a conclusion with which we fundamentally disagree." See Oceana et al. comments on Essential Fish Habitat Draft Environmental Impact Statement. The EIS must address the effects of different harvest strategies on habitat in a credible manner.

Since neither the EFH EIS nor the Alaska Groundfish Harvest Specifications EIS addresses the effects of increasing or redistributing bottom trawl effort, you must undertake a new, credible analysis of habitat and bycatch impacts before raising flatfish quotas.

Gulf of Alaska Pollock

We are disturbed by recent increases in the amount of bycatch in the Gulf of Alaska pollock fishery from 2005 to 2006. Shortraker and roughey rockfish bycatch more than doubled to 94.7 mt. Northern rockfish bycatch also rose by more than one order of magnitude from 0.8 mt to 14.5 mt. Pelagic shelf rockfish complex bycatch rose from 2mt to 9 mt. Pacific cod bycatch doubled to 707 mt; flathead sole bycatch more than tripled to 593 mt; big and longnose skate bycatch increased 5-fold to 36 mt; rex sole increased 7-fold to 154 mt; and miscellaneous flatfish increased 100-fold to 4389 mt. The estimated bycatch of prohibited Bairdi tanner crab rose exponentially from 6 crab to over 86 thousand crab. This startling rise in the amount and composition of bycatch also clearly indicates a change in the way pollock are harvested in the Gulf of Alaska. The increase in bycatch of bottom dwelling fish and crab requires substantial analysis not only focused on bycatch, but also focused on habitat impacts.

We appreciate the opportunity to highlight some of the shared concerns that will aid you in making the best decisions for setting harvest quotas for the Bering Sea, Aleutian Islands, and Gulf of Alaska in 2008 and 2009.

Sincerely,



Jon Warrenchuk
Ocean Scientist, Oceana

November 27, 2007

North Pacific Fishery Management Council
605 W. 4th Avenue, Suite 306
Anchorage, AK 99501

Dear North Pacific Fisheries Management Council,

As concerned citizens and Greenpeace supporters, we are urging you to implement an ecosystem-based approach to fisheries management that accounts for the needs for all species that depend on pollock as their prey.

At a minimum, catch limits should seek to double in size the pollock population and maintain this higher level over time.

Many of the fish, seabirds, marine mammals and invertebrates living in the North Pacific and the Bering Sea rely on pollock as a source of food. Without enough prey they are starving and populations are dwindling. The entire structure of the ecosystem is being fundamentally changed. Marine reserves and more precautionary management strategies are needed now to turn the tide.

Please don't sit around while animals starve—reduce pollock catch limits and keep more forage fish in the seas for marine life.

Sincerely,

Copy of names (approximately 6,747) attached to letter are at Secretary's desk for review.

BSAI Groundfish Plan Team
AFSC- Seattle, WA
November 13, 2007

Loh-Lee Low (AFSC), Chair	Brenda Norcross (UAF)
Grant Thompson (AFSC), Special Envoy to the SSC	Kathy Kuletz (USFWS)
Jane DiCosimo (NPFMC), Coordinator	Kerim Aydin (AFSC)
Dave Carlile (ADF&G)	Brenda Norcross (IPHC)
Andy Smoker (AKRO)	Lowell Fritz (NMML)
Dan Lew (AFSC)	Theresa Tsou (WDFW)

The BSAI Groundfish Plan Team convened on Tuesday, November 13, 2007, at 1:00 pm. Mike Sigler participated by phone during pollock and Pacific cod discussions. Twenty five members of the public and ten agency staff attended parts of the meeting.

Pollock Taina Honkalehto presented 2007 summer *EBS Echo-Integration (EIT) pollock survey* results. She reported that 2007 was another cold year, as was 2006. Juvenile pollock concentrations were found in the extreme northwestern part of the survey area, and also the central survey area roughly 60 nmi west of the St. Paul Island. Pollock EIT biomass was distributed 13% east of 170 degrees (Pribilof Islands), 82% were NW of 170 degrees, and 5% were in the Russian waters that were surveyed (part of the Navarin Basin). Age 1 pollock were numerically dominant in the abundance estimates whereas older pollock comprised the bulk of the biomass estimate. The off-bottom component of the pollock stock (between 0.5-3 m off bottom) has consistently averaged about 20% of the water column abundance from 0.5 m to the surface.

The EIT survey group continues to evaluate the use of frequencies other than 38 kHz to help identify the non-pollock backscatter component. Inter-vessel comparisons between the *Miller Freeman* and *Oscar Dyson* are planned in winter of 2008. The Team noted its comments from September 2007 regarding the inter-vessel comparison. The Team discussed the variability between the two vessels regarding survey design and results. MACE staff reported that to date, the differences between the two vessels were small, with no significant differences in backscatter in a study conducted during the acoustic survey in the Eastern Bering Sea in 2006.

Jim Ianelli presented a summary of the *EBS pollock stock assessment*. Abundance of EBS walleye pollock has declined due to below-average recruitment from the 2001-2005 year classes. This represents the longest period of observed consecutive below-average recruitments. Spawning biomass has apparently declined from about 4% above B_{MSY} in 2007 to 28% below B_{MSY} in 2008. Although preliminary indications are that the 2006 year class is well above average, spawning biomass is unlikely to exceed the benchmark until 2010. The age 3+ biomass for 2007 is estimated to be the lowest since 1980.

The Team concurred with the SSC determination that EBS walleye pollock qualify for Tier 1 management. Projected spawning biomass for 2008 is 1.38 million t, dropping it to sub-tier "b" of Tier 1. The maximum permissible ABC harvest rate was based on the harmonic mean of the ratio of yield over fishable biomass and is 0.341 compared to last year's value of 0.243 (which used age 3+ biomass). This value is adjusted by a factor of 72% corresponding to the projected spawning biomass in 2008. The resulting maximum permissible ABC of 1.17 million t is considerably higher than the Tier 3b estimate of 555,000 t (which is adjusted downwards from the $F_{40\%}$ by stock size relative to $B_{40\%}$).

The 2006 and 2007 survey estimates of abundance were both below average levels. The 2007 surveys indicated high levels of abundance for one year old pollock. The new data in this assessment showed lower than expected 7 year old fish. This contrasts with other year classes, e.g., the 1992 year class which increased in expected abundance at later ages. The EIT and bottom trawl survey abundance at age estimates appear to be consistent with other fishery observations. A figure showing the absolute

abundance at length of pollock in the BT survey shows that the mode of 2 year old fish is rare and that 1 year old fish are more common.

Bottom trawl survey-only cohort mortality patterns show higher catches in the last several years. This is consistent with model results in that the total mortality is higher in recent years.

The model evaluations showed that adding data reduces uncertainty, and that the ABC appears to be unaffected by this. The biggest change to the ABC is a result of changes in mean weight. The reduction in ABC (compared to last year's model) is partly due to the lower biomass estimate relative to B_{MSY} (being below B_{MSY} changes the rate proportionately). It was clarified that changing from the 3+ biomass to "fishable" biomass will yield the same average catch. The motivation for the change was due to the added inter-annual variability induced by having (highly uncertain) recruitment at age 3 impact the biomass for the calculation. The underlying continuous form for F_{MSY} is unchanged. The fishery has shifted to younger pollock in recent years based on the locations of the operations. This affects future yield calculations since selectivity shifts towards younger ages and the main tenet for management is conservation of spawning biomass.

A question from the public regarding stock-recruitment stationarity was posed. The author responded that the model demonstrates lots of variability; the estimates change with new data and model configurations but that the changes have been relatively minor over time. However, if pollock productivity changes, it may be appropriate to shift the period over which the stock-recruitment relationship is estimated and revise estimates accordingly.

ABC considerations The following highlights some points presented in the assessment that the Team considered in its discussions of whether the ABC should be set to the Tier 1b maximum permissible level:

- 1) The sum of the survey catchabilities for the accepted model is over 1.5, indicating that there is considerable overlap between the availability of pollock between the two surveys (or some other mechanism such as temporary immigration of pollock into the EBS region from elsewhere).
- 2) The stock-recruitment relationship continues to be constrained within the model which causes the harvest rate to be more conservative.
- 3) 2007 weights-at-age appear to be closer to mean values and are above the lower levels observed in the 2006 fishery.
- 4) In the two surveys conducted in 2007 signs of 1-year old abundance were above average.
- 5) The precautionary MSY harvest rate has been adjusted downward to nearly 72% to account for the stock being below the B_{msy} level.
- 6) In the 1998 assessment, the outlook for 1999 was fairly pessimistic (although the age 3+ biomass was about 1 million t higher than is presently estimated for 2008). In hindsight, the perception of relatively poor stock conditions at the time changed. For example, at the time, the 1992 year class was estimated to be about average whereas now it appears to be more than twice the average and represents the third highest year class.
- 7) The stock has been at low levels in the past (but this appears to be the lowest since before 1980).
- 8) The Tier 3 ABC levels are substantially lower than the Tier 1 values due to different assumptions about reference biomass levels (hence a larger adjustment).
- 9) Future selectivity patterns are unpredictable given fish distribution.
- 10) If the 2005 year class is in fact below average, then there will have been 5 year classes in sequence that have been below average, an apparently unique event for this stock.
- 11) Spatially equitable catch rates by the fishery may continue to be impacted (lower catch rates overall are to be expected, and this can manifest as spatial differences in pollock availability).
- 12) Absorbing some of the anticipated "adjustment" from the ABC control rule will likely reduce the inter-annual variability ABC recommendations.

Arguments in support of Tier 3 (or at least the ABC at the maximum permissible level associated with Tier 3) include:

- 1) Stock is at a very low level
- 2) The stock has produced good year classes near B_{40} and hence is preferred over B_{MSY} as a target.
- 3) Size at age has declined in recent years (thought 2007 indicate about average) and experience at these low biomass levels has been limited
- 4) The stock/recruitment curve has an unusual shape
- 5) Recent period of 5 below average recruitments
- 6) Model shortcomings as noted by the lead author (e.g., retrospective patterns in the data).
- 7) Ecosystem considerations suggest that we should not rely on the stock/recruitment as under Tier 1.

Public comments also recommended results from tier 1b, tier 3, and lower than model results. Others commented that the Team should not solicit ABC recommendations from the public.

The Plan Team ultimately supported the authors' recommendation of a 1 million t ABC for 2008, but the decision was not unanimous. Some supported lower values of 976,000 t or 555,000 t as additional conservation for the ecosystem. Two issues were noted (1) Tier 3 may be more reflective of the condition of the stock and (2) recent poor recruitments may suggest a regime shift. The team discounted the regime shift argument as more data would be needed to support that a new regime exists for pollock. The team noted that the F value corresponding to an ABC of 976,000 has not been calculated.

The team recommended setting the 2009 ABC at 1 million t also, but this could be revised lower if next year's assessment does not confirm the current estimated strength of the 2006 year class.

Steve Barbeau presented the Aleutian Islands pollock assessment. Model 2B is similar to the model accepted by the SSC last year and is recommended by the authors again this year. Age 2+ biomass appears to have increased from 1999 to 2004, after which it has been stable. Spawning biomass appears to have been increasing slowly since 1999. The 2000 year class is estimated to have been well above average (third largest in the time series), and preliminary indications are that the 2005 and 2006 year classes may be slightly above average.

The Plan Team concurs with the SSC determination that this stock qualifies for Tier 3a management. And concurs with the authors' recommendation to set 2008 ABC at the maximum permissible value

No changes in the analytic approach were made in the Bogoslof pollock assessment. Survey biomass estimates since 2000 have all been lower than prior to 2000. The team concurs with management under Tier 5 and past practices for determining ABC.

Pacific cod Difficulties encountered with the 2006 assessment resulted in a thorough review of the 2006 assessment model in April 2007 during a public workshop. Many suggestions for changes and refinements of the analytical approaches were made. The assessment considered four models. Model 1 fixes M at a value of 0.34 based on life history theory. Model 2 fixes M at the traditional value of 0.37. Model 3 estimates M internally. Model 4 differs from Model 1 in several respects to respond to public comments on the use and fitting of the data. The Team observed the following regarding the assessment.

- Estimated selectivity curves appear reasonable, except that IPHC longline survey selectivity curve is peaked and thus not differentiable (increasing the number of parameters from 4 to 6 parameters still doesn't overcome stated problems with estimating selectivity in stock synthesis).
- Model 4 ignores age data. It doesn't make sense to discard data with information on relative cohort strength, even if the fit is "not very good". Length data is not as informative as age data because there is not a one-to-one relationship between length and age, except for youngest ages.

Long-term goal of assessment has been to obtain then add age data to the assessment (~10-year effort).

- Model 1, 2, 3 and 4 biomass (Fig. 2.3) and recruitment (Fig. 2.2) trends are similar. These models differ primarily in M estimate (except that Model 4 has other structural differences). This implies that data are sufficient to estimate trends, but that scale is sensitive to value of M .
- Estimated biomass dictated by choice of M (scales biomass, compare Models 1, 2 and 3). The M value of 0.34 in Model 1 is based on a derivation from standard functional relationships for fish life history (Jensen 1996); Jensen's relationship compares favorably with data and parameter estimates in other studies (e.g., Beverton 1963). Given Jensen's relationship, the M of 0.34 follows from an age at 50% maturity of 4.9 years (Stark 2007). However, it is possible that age at maturity is 3.9 years if age readers are adding a year when reading otoliths. Age at maturity values corresponding to the M values in the four models are 4.45 years ($M = 0.37$), 7.5 ($M = 0.22$), 3.58 ($M = 0.46$), and 4.9 ($M = 0.34$). However otolith reading is carefully done and best estimate is 4.9 years; in the absence of contradictory information, should go with 4.9 years.
- Modeling approach for selectivity: Asymptotic selection stipulated for some fisheries; survey selectivity allows dome-shape (except in Model 4), as well as time-variance in ascending limb.
- Recommendation for future assessments: Use 3-parameter exponential logistic (Thompson 1994) to represent selectivity.
- Abundance indices received no likelihood weight (computed "analytically"), except for trawl survey which received a weight of "1".
- Model 3 fits data the best, but model selection ranks M estimate (0.22) as "too low".
- Because the three parameters in the Schnute growth model used in SS2 are confounded in nature, it would be worth exploring a model with fixed L_{inf} at an externally estimated value and let the model estimate L_1 and K because that selectivity has more impacts on the observed sizes of young fish than those of older fish, especially when an asymptotic selectivity curve is used. Model estimated growth curves should be lower than the observed for younger age. Figure 2 from the September assessment may be a concern.

The Team appreciated the history of assessment methodology changes provided in the assessment. The Team noted that the model may be more advanced than the data can fit. The Team noted that it is very difficult to track population changes by fishery, area, and size groups. But that the Team concluded that the real debate is in choosing the correct estimate of M , and not with the models, as unexplained mortality of cod is huge. The Team recommended reducing the number of parameters in the models.

ABC considerations

- Bottom trawl survey abundance dropped two years in a row; currently at historic minimum
- Three to five below average year classes estimated, depending on the model.
- All models show decreasing abundance.
- Consistency of model estimates of biomass and recruitment trends and continued decrease of trawl survey biomass estimate implies ABC should also decrease. Author recommendation to decrease ABC to 150,000 matches decrease in trawl survey biomass.
- Low stock abundance (29% of unfished female spawning biomass) at least for next two years; It's below B_{40} but above B_{20}
- Five years of weak year classes, assuming Model 1.
- 2006 has the second highest year class that may affect the ABC as soon as 2009; sign of good recruitment is usually verified in the future for cod, compared with pollock which often tends to disappear

The Team concurred that this stock qualifies for management under sub-tier "b" of Tier 3 because the projected biomass for 2008 (398,000 t) is below $B_{40\%}$. The Team recommended setting the 2008 ABC at the maximum permissible value.

The Pacific cod chapter included a discussion paper on "results from ecosystem models on the role of Pacific cod in the eastern Bering Sea and Aleutian Islands ecosystems" as well as an extended discussion section on "ecosystem considerations." While these discussions provide useful clarifying information about the ecological dynamics of Pacific cod and its fisheries with respect to the rest of the ecosystem, there was no special identification of ecosystem features that would require adjustments to the estimated ABCs or their attendant reference points.

Sarah Gaichas briefly summarized a slide from the ecosystems considerations section. According to information collected on summer trawl surveys, Pacific cod eat pollock (25% of diet) and assorted invertebrates, but generally do not eat arrowtooth flounder (there was a question whether cod ate arrowtooth in the EBS). This diet information is not informative to the Team's discussion of model preference, although accounting of mortality sources may be. In the ecosystem model, up to half of cod mortality is from unaccounted sources. Actual observed consumption of cod by predators is relatively low, which is typical of top predators. Sources of mortality are assumed to balance production in the ecosystem model; therefore, a large amount of "unaccounted mortality" may indicate several things. First, the mortality may be actual mortality coming from sources other than predation and fishing, so it is unaccounted in the ecosystem model. Second, the unaccounted mortality may not be mortality at all, but may represent one or more of the following processes. Cod could be leaving the ecosystem, so they are not dead but effectively unavailable. Alternatively, mortality may be less than production, which suggests that the stock is increasing (this does not appear to be the case for EBS cod). Finally, the estimate of production may be too high, inflating the "unaccounted" mortality. Because production is based on natural mortality estimates used in the cod stock assessment, and cod are not increasing, the high proportion of unaccounted mortality suggests that the value of M used in the stock assessment model is not too low. A higher value of M (such as that estimated in Model 4) would exacerbate the problem, leading to an even higher proportion of unaccounted mortality.

The Team encouraged the Council's consideration of apportioning the ABC by area because of recognized differences between the BS and AI ecosystems. From a modeling perspective, the Team can justify separate ecosystem management for Pacific cod. Some avenue of exploration from a biological perspective is also justified. The Council adopted the AI Fishery Ecosystem Plan as a planning document that identified cod as an important component of the AI ecosystem. The Team has provided apportionment percentages of 84% EBS and 16% AI for several years.

Public comments noted that age at maturity may be about a year younger than the current method used and that it is difficult for the public to be fully engaged when assessment information is provided late in the process.

Sablefish The joint team discussion on sablefish can be found in the joint team minutes.

Atka mackerel Sandra Lowe reported that there were no changes in the current model. The Team accepted the author's recommendations.

Greenland turbot Jim Ianelli summarized the Greenland turbot assessment. He provided a simplified Tier 5 approach to contrast with the Stock Synthesis 2 model. The Team agreed that this stock qualified for management under Tier 3a. Although favorable recruitment appears to have occurred in recent years, the Team agreed with the author to recommend less than the maximum permissible ABC to reflect recent fishing mortalities, as has been done for a number of years.

The Team noted that the slope survey was canceled in 2006 due to lack of funding. The Team recommended that the AFSC conduct the EBS slope survey in 2008. Such information is necessary to determine whether abundance is sufficient for the Team to recommend an ABC set to the maximum permissible level in the future. The slope survey is recognized as the only one that covers the prime habitat for Greenland turbot.

Comments from the Center for Independent Experts review were noted for all flatfish assessments and several pertained to turbot. He intends to consider including information if available from Greenland turbot in the Russian zone. He will also participate in a workshop in February 2008 for a management strategy evaluation of Atlantic stock for the European Union; this stock is fished at ten times the US rate. A recent paper by Cooper (2007) published a lower M (0.112 instead of .18) for this stock. There is new information that they are longer lived. They are not found where the survey is and appear to disappear at larger size.

Yellowfin Sole Tom Wilderbuer presented all but one of the flatfish assessments. The Team concurred with the SSC's 2006 decision that yellowfin sole qualify for management under Tier 1. The Team noted that Tier 1 is about matching the stock/recruitment curve and determining where F_{MSY} is on that curve. The Team noted that there is one selectivity curve for all years, compared with pollock that has a selectivity curve for each year. Above average recruitment from the 1995 and 1999 year-classes is expected to maintain the abundance of yellowfin sole at a level above B_{40} in the near future.

Loh-lee Low observed that the Chinese market is transitioning from a consumer market to processing, also. Tom Wilderbuer noted that Amendment 80 will change how the fishery is prosecuted, in that the cooperatives will manage halibut bycatch which constrains harvest. Theresa Tsou noted that the model tends to predict more younger fish than is observed. This also occurs for northern rock sole in past years. A better fit occurs in recent years.

The catchability of yellow fin sole appears to be related to water temperature. This relationship was examined for arrowtooth flounder. However, at this time, the authors are uncertain as to how to interpret their findings. The Team suggested investigating the relationship between bottom temperature and q for all flatfish

Northern Rock Sole This stock is also now managed under tier 1. M and q were estimated within the model yielded $M=0.15$ and $q=1.87$. This was very different from the previous estimates of $M=0.16$ and $q=1.52$. While experiments indicate herding, this estimate of q means that 40% of fish are herded into the net. The authors do not think this is a realistic estimate of herding. It is most likely biased because of the skewed age composition from low recruits in 1990s. Therefore, the q estimated in past assessments was used ($M=0.15$, $q=1.5$). The rock sole stock is expected to remain stable through 2009. However, good recruitment in 2001 through 2004 should increase the stock biomass at the beginning of the next decade.

The SSC requested continued management strategy evaluation (MSE) for Tier 1 management. The authors intend to continue this exploration for climate and productivity changes and will examine a split-sex model for the fall 2008 assessment.

Arrowtooth flounder The stock assessment was expanded to include the 10 Aleutian Islands surveys and the survey size composition for arrowtooth flounder. The BSAI stock is therefore redefined with 73% of the stock on the Bering Sea shelf with the AI data added, compared with 87% when only the shelf and slope were considered. The AI component is 17% with the remaining 10% from the Bering Sea slope. While the ABC appears to have increased, it is an artifact of adding Aleutian Islands data. Structural problems with this model must be resolved before arrowtooth flounder can be changed to Tier 1 status. As there is little to no fishery for arrowtooth flounder the model is mostly driven by the survey data.

The long-term trend of increasing arrowtooth flounder biomass in the EBS is expected to continue as recruitment has also been increasing for the last 10 years. Arrowtooth flounder in the AI leveled off in 1990s, but has been steadily increasing since then. The ABC recommendation continues to include Kamchatka flounder. Though Kamchatka flounder can now be identified and separated from arrowtooth flounder on AFSC surveys, it is not distinguished in the fishery.

More female arrowtooth flounder are caught than males during the fisheries and the surveys. It is hypothesized that males are not missed by the survey, but that they experience higher mortality than the females and do not behave differently than females. Allowing the maximum selectivity for shelf males to be a bit lower than 1.0 leaves room for some males to migrate down slope or be missed in survey. The age data from the GOA indicates males reach 17 yrs, whereas many females have been aged as high as 25 yrs. Therefore, natural mortality of male arrowtooth flounder must be increased, as opposed to decreasing gear selectivity at older ages in the male population.

Alaska plaice The female spawning biomass and total biomass trends were slowly decreasing from 1985 to 2005 but have since been increasing and is expected to continue. The shelf survey biomass has been fairly steady since the mid 1980s and the stock is very lightly exploited. Above-average recruitment year classes since 1998, with an exceptionally strong year class in 2003, will support the increasing trend in this stock. The authors extended the range of the length bins to mimic the dynamics of larger fish, at the request of the SSC. The authors will implement a split-sex model next year.

Other flatfish complex Since 2002, the SSC has classified "other flatfish" as a Tier 5 species complex. No progress was made on the assessment of bottom temperature effect on catchability of other flatfish species. There was discussion regarding separating Bering Flounder from flathead sole in the combined chapter and adding Bering Flounder to the Other Flatfish chapter since there are now able to be identified in both surveys and fisheries.

Flathead sole Buck Stockhausen presented the flathead sole stock assessment. This stock qualifies for management under Tier 3. More research is needed in the Chukchi and Beaufort Seas to better characterize the geographical distribution of the flathead sole and Bering flounder stocks. In response to SSC and Plan Team comments, the author examined the distribution of Bering flounder with respect to the fishery. The northerly distribution of the species did not seem to overlap the spatial distribution of the fishery, although mismatch in seasonal timing of the survey versus the fishery means that this is not conclusive. The Team recommended placing Bering flounder in the "other flatfish" category in future assessments, but that the flathead model may always carry some Bering flounder with it into the future (due to species misidentification early in the catch and survey time series). The Team was concerned that Bering flounder maybe harvested at too high a rate. NMFS catch accounting started a species code in 2007. Historical catches could be identified for this species.

A public comment was posed regarding the control rule for this stock and the author's recommendation that ABC be set at the maximum allowable ABC for a Tier 3a stock, given the recent declining trend in stock abundance and moderate recruitment levels. Current biomass is high relative to the mean but has dropped from the 1990s. High recruitment occurred in the 1970s and 1980s, then decreased in the 1990s and currently remains around the mean. The control rule puts this stock at twice the value of B_{MSY} and B_{40} , and it is lightly exploited. The author responded that since stock biomass is high and the TAC is typically set much lower than the ABC, there is less concern that recommending the maxABC for 2008-9 will lead to overfishing or an overfished condition in the near term. Declining recruitment will be more of a concern in the future.

Rockfishes Paul Spencer reported on the off-year assessments for the rockfish assemblages. There were no changes to the models this year and he summarized plans for 2008, in response to SSC comments and the 2006 CIE review.

Other Species Olav Ormseth presented the squid assessment. The Team discussed whether this group may more appropriately reside in the forage fish category. He reviewed squid life history and indicated that the predominant species in this group (*Berryteuthis magister*) may experience multiple cohorts within a single year. The assessment includes more complete ecological, biological, and survey data (although survey biomass estimates are unreliable). The Tier 6 specifications did not change from last year.

Olav Ormseth presented the 2007 skate assessment that estimated the Alaska skate component of the assemblage within Tier 3 and applied Tier 5 to the remaining species. The authors recommended that if Tier 3 is not applied to Alaska skates this year, that a value of $M = 0.1$ should be used in the Tier 5 calculations, as it has been in past years. The Team encouraged the authors to address a discrepancy between increasing survey biomass estimate in recent years relative to the model projections of declining biomass, and discussed the idea that this discrepancy may be due to temporal changes in length composition. Since 2000, there has been a decrease in the proportion of skates in the smallest length bins. This may reflect poor recruitment in recent years which results in model estimation of declining biomass.

Jon Heifetz summarized the shark assessment. The rationale for the Team's recommendations for applying tier 5 in 2006 was that the calculation of biomass was a minimal estimate and would be better than applying tier 6 based on catch history. The assessment could be viewed more as a repository of information, rather than an assessment. The assessment included a modified tier 6 and tier 5 calculations but the latter relies on an unreliable estimate of M . The Plan Team concurred with the SSC's conclusion in 2006 that survey information was not reliable enough to promote the Tier 5 assessment. However, applying the modified Tier 6 would not be constraining on directed fisheries if this group were managed separately from the assemblage under a proposed plan amendment. This could result in a conservation issue since very limited information is available for management.

Rebecca Reuter summarized the sculpin assessment. She reported significant strides for age/growth information from North Pacific Research Board funds (for estimating M for four species) and species identification in that 2008 will be the first year that observers will be asked to identify sculpins to species. In contrast to the Tier 5 assessment in 2006, where a single M (0.19) and two biomass estimates (one for the EBS and one for the AI) were used, the 2007 assessment applies distinct M estimates from the literature and species or species group biomass estimates for the five most predominate sculpin species/species groups. The Team noted that lower estimates of M are not reasonable to assume and it was faced with two bad choices: last year's approach or justifying using new estimates of M .

The assessment authors noted that species composition differs among the BS shelf, BS slope, and the AI. The Team noted that the flatfish fisheries are the principal known remover of sculpins. Approximately 51% of sculpin catch in the flatfish bottom trawl fisheries is from the yellowfin sole fishery; 24% from the flathead sole fishery, and 24% in the rock sole fishery.

Liz Conners presented the octopus assessment. The team concurred with the author's conclusions that 1) the octopus biomass and mortality estimates are not reliable and therefore do not support a Tier 5 assessment and 2) the Tier 6 estimate based on average catch likely underestimates the population and would unnecessarily constrain fisheries, primarily the pot fishery for Pacific cod, if octopus were managed separately from the 'other species' category. The author recommended a revised Tier 6 based on recent maximum catch. Ongoing research includes species identification discard mortality estimates. The Plan Team recommended Tier 6. The Team noted that octopus biomass estimates from food web models are orders of magnitude bigger than survey biomass estimates. Therefore Tier 6 management would be

unnecessarily constraining, particularly to the Pacific cod pot fishery, if managed separately. Three years of special project data could generate pot cod survival rates (believed to be high).

The Team noted that results from a cooperative research project collected from the plants showed that larger octopus are Giant Pacific octopus *Enteroctopus dolphineii*. It may be possible in the future to separate adults of this species from smaller species based on size.


Ecosystem Issues The BSAI plan team discussed the development of a set of indicators to summarize the trend in the 'state' of the ecosystem. Time series of indices such as average trophic level of the catch, fish community diversity as evidenced in bottom trawl surveys, estimates of annual surplus production, production at upper trophic levels, and other summary indicators could be considered so that a 'movie' of ecosystem status could be produced. The ecosystems considerations authors stated that this year they have begun to develop such a set as part of moving the Ecosystem Assessment into part of an Integrated Ecosystem Assessment (IEA), but that finalizing the set of indicators would be an iterative process including input from the Council and multiple users. This was discussed with reference to a suggestion by the SSC and Council in August 2007 that the carrying capacity for Steller sea lions may now (since the early 1990s) be lower than it was in the past (1960s and 1970s), while at the same time, carrying capacity for their prey may have increased or remained stable (as evidenced by surveys, stock assessments, and fishery catches). NMFS-Protected Resources will evaluate all factors that affect the recovery of Steller sea lions, including if and why carrying capacity has changed, as it finishes the Steller sea lion recovery plan and the status quo biological opinion on the effects of the BSAI-GOA groundfish fisheries. A set of ecosystem indicators that may reflect a time series of carrying capacity for upper trophic levels would assist NMFS-Protected Resources in this process.

Adjourn The team completed its assessment reviews at 10 am on Friday; the teams worked on summaries for the introduction. The teams adjourned at approximately 4:00 pm.

APPENDIX A

**STOCK ASSESSMENT AND FISHERY EVALUATION REPORT
FOR THE GROUND FISH RESOURCES
OF THE BERING SEA/ALEUTIAN ISLANDS REGIONS**

Compiled by

The Plan Team**for the Groundfish Fisheries of the Bering Sea and Aleutian Islands**

With Contributions by

K. Aydin, S. Barbeaux, D. Clausen, M.E. Connors, D. Courtney, J. DiCosimo, M. Dorn, J. Fujioka, S. Gaichas, K. Goldman, D. Hanselman, J. Heifetz, G. Hoff, T. Honkalehto, J. Ianelli, E. Jorgenson, S. Kotwicki, R. Lauth, S. Lowe, C. Lunsford, B. Matta, D. Nichol, O.A. Ormseth, R. Reuter, C.J. Rodgveller, P. Spencer, I. Spies, W. Stockhausen, T. TenBrink, G. Thompson, C. Tribuzio, T. Wilderbuer, M. Wilkins, G. Williams, and N. Williamson

November 2007

**North Pacific Fishery Management Council
605 West 4th Ave., Suite 306
Anchorage, AK 99501**

This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by the National Marine Fisheries Service and should not be construed to represent any agency determination or policy.

Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Region

Table of Contents

Introduction.....	3
Background Information.....	3
Overview of “Stock Assessment” Section	9
Appendix A: Pacific Halibut Community Development Quota Discard Mortality Rates	40
 Stock Assessment Section	
1. Eastern Bering Sea Walleye pollock.....	41
1a. Aleutian Islands Walleye pollock	139
1b. Bogoslof Pollock.....	197
2. Pacific cod.....	209
3. Sablefish.....	329
4. Yellowfin sole.....	447
5. Greenland turbot	513
6. Arrowtooth flounder	563
7. Northern rock sole.....	627
8. Flathead sole	687
9. Alaska Plaice.....	755
10. Other flatfish	799
11. Pacific ocean perch	811
12. Northern Rockfish.....	815
13. Shortraker/Rougheye rockfish	817
14. Other rockfish	821
15. Atka mackerel	825
16. Skates	909
17. Squids.....	931
18. Sharks.....	1033
19. Octopus	1081
20. Sculpins.....	1117
 Ecosystem Considerations	 Bound Separately
Economic Status of Groundfish Fisheries off Alaska	Bound Separately

Summary

By

The Plan Team for the Groundfish Fisheries of the Bering Sea and Aleutian Islands

Introduction

The *National Standard Guidelines for Fishery Management Plans* published by the National Marine Fisheries Service (NMFS) require that a stock assessment and fishery evaluation (SAFE) report be prepared and reviewed annually for each fishery management plan (FMP). The SAFE report summarizes the best available scientific information concerning the past, present, and possible future condition of the stocks, marine ecosystems, and fisheries that are managed under Federal regulation. It provides information to the Councils for determining annual harvest levels from each stock, documenting significant trends or changes in the resource, marine ecosystems, and fishery over time, and assessing the relative success of existing state and Federal fishery management programs. For the FMP for the Groundfish Fishery of the Bering Sea and Aleutian Islands (BSAI) Area, the SAFE report is published in three sections: a "Stock Assessment" section, which comprises the bulk of this document, and "Economic Status of Groundfish Fisheries off Alaska" and "Ecosystem Considerations" sections, which are bound separately.

The BSAI Groundfish FMP requires that a draft of the SAFE report be produced each year in time for the December meeting of the North Pacific Fishery Management Council. Each stock or stock complex is represented in the SAFE report by a chapter containing the latest stock assessment. New or revised stock assessment models are generally previewed at the September Plan Team meeting, and considered again by the Plan Team at its November meeting for recommending final specifications for the following two fishing years.

This Stock Assessment section of the SAFE report for the BSAI groundfish fisheries is compiled by the BSAI Groundfish Plan Team from chapters contributed by scientists at NMFS Alaska Fisheries Science Center (AFSC). These chapters include a recommendation by the author(s) for overfishing level (OFL) and acceptable biological catch (ABC) for each stock and stock complex managed under the FMP. This introductory section includes the recommendations of the Plan Team (Table 1). The ABC recommendations are reviewed by the Scientific and Statistical Committee (SSC), which may confirm the Plan Team recommendations. The Plan Team and SSC recommendations, together with social and economic factors, are considered by the Council in determining total allowable catches (TACs) and other measures used to manage the fisheries. Neither the author(s), Plan Team, nor SSC recommends TACs.

Members of the BSAI Plan Team who compiled this SAFE report were Loh-lee Low (chair), Jane DiCosimo (BSAI FMP coordinator), Kerim Aydin, David Carlile, Lowell Fritz, Steven Hare, Kathy Kuletz, Dan Lew, Brenda Norcross, Mike Sigler, Andrew Smoker, Grant Thompson, and Theresa Tsou.

Background Information

Management Areas and Species

The BSAI management area lies within the 200-mile U.S. Exclusive Economic Zone (EEZ) of the US (Figure 1). International North Pacific Fisheries Commission (INPFC) statistical areas 1 and 2 comprise the EBS. The Aleutian Islands (AI) region is INPFC area 5.




Figure 1. Bering Sea and Aleutian Islands statistical and reporting Areas

Five categories of finfishes and invertebrates have been designated for management purposes (see below). They are prohibited species (species that must be returned to the sea when caught), target species (species for which an individual TAC is established), other species (species for which an aggregate TAC is established), forage fish (species for which targeted harvest is prohibited, with a maximum of 2 percent retainable bycatch allowed), and non-specified species (all species not included in one of the other categories). This SAFE report describes the status of the stocks in the target species and "other species" categories only. The finfish species categories, other than non-specified species, are listed below:

Prohibited Species	Target Species	Other Species	Forage Fish
Salmon	Walleye pollock	Sculpins	Eulachon
Pacific halibut	Pacific cod	Sharks	Capelin
Pacific herring	Sablefish	Skates	Sandlance
Steelhead trout	Yellowfin sole		Myctophids
	Greenland turbot		Bathylagids
	Arrowtooth flounder		Sandfish
	Northern rock sole		Pholids
	Flathead Sole		Stichaeids
	Alaska plaice		Gonostomatids
	Other flatfish		
	Pacific Ocean perch		
	Northern rockfish		
	Shortraker rockfish		
	Rougheye rockfish		
	Other rockfish		
	Atka mackerel		

The invertebrate species categories, other than non-specified species, are listed below:

Prohibited Species	Target Species	Other Species	Forage Fish
King crab		Squid	
Tanner crab		Octopus	

Historical Catch Statistics

Catch statistics since 1954 are shown for the Eastern Bering Sea (EBS) subarea in Table 2. The initial target species was yellowfin sole. During the early period of these fisheries, total catches of groundfish peaked at 674,000 t in 1961. Following a decline in abundance of yellowfin sole, other species (principally walleye pollock) were targeted, and total catches peaked at 2.2 million t in 1972. Walleye pollock is now the principal fishery, with recent catches approximately 1.4-1.5 million t due to years of high recruitment. After the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA) was adopted in 1976, catch restrictions and other management measures were placed on the fishery and total groundfish catches have since varied from one to two million t. In 2005, Congress implemented a statutory cap on TACs for BSAI groundfish of 2 million t. Catches generally total about 10 percent below the cap.

Catches in the Aleutian Islands (AI) subarea have always been much smaller than in the EBS. Target species have also differed (Table 3). Pacific Ocean perch (POP) was the initial target species. As POP abundance declined, the fishery diversified to other species. During the early years of exploitation, total AI groundfish catches peaked at 112,000 t in 1965. Atka mackerel is the largest fishery (53,700 t in 2007) in the AI, followed by Pacific cod (34,000 t in 2007). Recent catches have been about 100,000 t annually,

after peaking at 191,000 t in 1996. Total 2007 AI catches were 117,000 t, reflecting a 10,000 t increase in Pacific cod catch in 2007.

Total 2007 BSAI catches were 1.85 million t. Total catches since 1954 for the EBS and AI combined are in Table 4.

Recent Total Allowable Catches

Amendment 1 to the BSAI Groundfish FMP provides the framework to manage the groundfish resources as a complex. Maximum sustainable yield (MSY) for the BSAI groundfish complex was originally estimated at 1.8 to 2.4 million t. The optimum yield (OY) range was set at 85% of the MSY range, or 1.4 to 2.0 million t. The sum of the TACs equals OY for the complex, which is constrained by the 2.0 million t cap.

Establishment of the Western Alaska Community Development Quota (CDQ) Program annual groundfish reserves is concurrent with the annual BSAI groundfish harvest specifications. Once annual BSAI groundfish TACs are established, certain species categories are allocated to the CDQ Program. This includes 10 percent of the BS and AI pollock TACs, 20 percent of the fixed gear sablefish TAC, and 7.5 percent of the sablefish trawl gear allocation. It also receives 10.7 percent of the TACs (up from 7.5 percent prior to 2008) for Pacific cod, yellowfin sole, rock sole, flathead sole, Atka mackerel, AI Pacific ocean perch, arrowtooth flounder, and BS Greenland turbot. The program also receives allocations of prohibited species quotas.

For the non-specified reserve, both the trawl and non-trawl fisheries are prosecuted under a single TAC. The TAC specifications for the primary allocated species, and PSC specifications, are recommended by the Council at its December meeting. A portion of the TAC limits for sablefish in the trawl gear category, arrowtooth flounder, BS Greenland turbot, and those species not otherwise allocated to the CDQ Program are placed in a reserve. Apportionments to this reserve range from 4.3 percent to 15 percent of applicable TAC limits. The reserve is used for (1) correction of operational problems in the fishing fleets, (2) to promote full and efficient use of groundfish resources, (3) adjustments of species TACs according to changing conditions of stocks during fishing year, and (4) apportionments. The initial TAC (ITAC) for each species is the remainder of the TAC after the subtraction of these reserves.

New Data Summary

Since the SAFE Report for 2007 was issued (NPFMC 2006), the following new information has been incorporated into the stock assessments:

- 1) Eastern Bering Sea Walleye pollock: 1) updated 2006 and 2007 catch data; 2) an age-length transition matrix was estimated to use the current-year fishery length frequency data; 3) the ability to split the age-1 values from the rest of the age compositions for the EIT survey was added so that the interaction between age-1 variability and other age groups was reduced; 4) total age 2 and older numbers-at-age from the surveys were used to tune the model; and 5) the Tier 1 ABC estimation method now uses "fishable" biomass defined as the biomass of pollock available to the fishery as modified by the selectivity-at-age estimates.
- 1a) Aleutian Islands Walleye pollock: 1) catch from 2006 and 2007; and 2) age data from the 2006 AI bottom trawl survey.
- 1b) Bogoslof pollock: 1) updated 2006 and 2007 catch data; and 2) echo-integration trawl survey data.
- 2) Pacific cod: 1) updated 2006 and 2007 catch data; 2) commercial fishery size composition data were recompiled for 1990-2007; 3) size composition data from the 2007 EBS shelf bottom trawl survey; 4) biomass estimate from the 2007 EBS shelf bottom trawl survey; 5) the numeric abundance estimates from the 1979-2007 EBS shelf bottom trawl surveys; 6) age composition data from the 1995 and 2006 EBS shelf bottom trawl surveys; 7) seasonal catch per unit effort (CPUE)

- data for the trawl, longline, and pot fisheries from 1991-2007; 8) catch rates from the 1998-2007 International Pacific Halibut Commission (IPHC) longline surveys; 9) size composition data from the 2007 IPHC longline survey.
- 3) Sablefish: 1) updated 2006 and 2007 catch data; 2) relative abundance and length data from the 2007 longline survey; 3) relative abundance and length data from the 2006 longline and trawl fisheries; 4) age data from the 2006 longline survey and longline fishery; 5) relative abundance and length data from 2007 GOA bottom trawl survey; and 6) older growth data (1981-1993) were updated, and new growth data were added (1996-2004) in the form of new age-length conversion matrices.
 - 4) Yellowfin sole: 1) updated 2006 and 2007 catch data; 2) 2006 fishery age composition; 3) 2006 survey age composition; 4) 2007 trawl survey biomass point estimate and standard error; 5) Estimate of the discarded and retained portions of the 2006 catch; and 5) update of weight at age using biological data through 2006.
 - 5) Greenland turbot: 1) updated 2006 and 2007 catch data; 2) EBS shelf survey 2007 biomass and length composition estimates; and 3) updated aggregated longline survey data index for the EBS and AI regions.
 - 6) Arrowtooth flounder: 1) updated 2006 and 2007 catch data; 2) 2007 shelf survey size composition; 3) 2007 shelf survey biomass point-estimates and standard errors; 4) estimate of retained and discarded portion of the 2006 catch; and 5) include the 10 AI surveys and survey size compositions.
 - 7) Northern rock sole: 1) updated 2006 and 2007 catch data; 2) 2006 fishery age composition; 3) 2006 survey age composition; 4) 2007 trawl survey biomass point estimate and standard error; and 5) Estimate of retained and discarded portions of the 2006 catch.
 - 8) Flathead sole: 1) 2006 catch data; 2) 2007 fishery length compositions and 1990-2006 data were recalculated; 3) 2004 and 2005 fishery age compositions from previous years were recalculated; 4) estimated survey biomass from the 2007 EBS Trawl Survey; 5) sex-specific length compositions from the 2007 EBS Trawl Survey and previous data were recalculated; 6) sex-specific age compositions from the 2006 EBS Trawl Survey and previous data were recalculated; and 7) mean bottom temperature from the 2007 EBS trawl survey.
 - 9) Alaska plaice: 1) 2006 catch data; 2) 2007 trawl survey biomass estimate and standard error, and 2007 survey length composition; 3) 2006 survey age composition; and 4) length bins and the transition matrix were extended from 45 to 60 cm.
 - 10) Other flatfish: 1) 2007 catch; and 2) 2007 EBS trawl survey biomass estimates.
 - 11) Pacific ocean perch: updated 2006 and 2007 catch data.
 - 12) Northern rockfish: updated 2006 and 2007 catch data.
 - 13) Shortraker and roughey rockfish: updated 2006 and 2007 catch data.
 - 14) Other rockfish: updated 2006 and 2007 catch data.
 - 15) Atka mackerel: 1) 2006 and 2007 catch data; 2) 2006 fishery age composition data; 3) age data from the 2006 Aleutian Islands bottom trawl survey; 4) 2006 fishery and survey weight-at-age values; 5) updated population weight-at-age values; and 6) the years used to compute an average selectivity vector for projections was updated from 2001-2005 to 2002-2006.
 - 16) Squids: updated 2006 and 2007 data.
 - 17) Skates: 1) updated 2006 and 2007 data; 2) biomass estimates from the 2007 EBS shelf survey; 3) independent estimates of survey selectivity and catchability; 4) length frequencies from survey data; 5) mean length at age from survey and fishery collections; and 6) natural mortality, growth, and maturity parameters.
 - 18) Sharks: 1) updated 2006 and 2007 catch data; and 2) bottom trawl survey biomass estimates for the 2007 EBS shelf.

- 19) Octopuses: 1) results of a cooperative research project that verified that all octopus observed in plant deliveries was confirmed as giant Pacific octopus; 2) results of observer program special project that noted that the observed 2007 sex ratios were different between octopus observed on vessels and those in plant deliveries; and 3) biomass estimates from summer 2007 Bering Sea Shelf Survey.
- 20) Sculpins: 1) updated 2006 and 2007 catch data; 2) total catch by target fishery and gear type for 2006; 3) biomass estimates from the 2007 Bering Sea Shelf Survey; and 4) new estimates of natural mortality for four species

Biological Reference Points

A number of biological reference points are used in this SAFE report. Among these are the fishing mortality rate (F) and stock biomass level (B) associated with MSY (F_{MSY} and B_{MSY} , respectively). Fishing mortality rates reduce the level of spawning biomass per recruit to some percentage P of the pristine level ($F_{P\%}$). The fishing mortality rate used to compute ABC is designated F_{ABC} , and the fishing mortality rate used to compute the overfishing level (OFL) is designated F_{OFL} .

Definition of Acceptable Biological Catch and the Overfishing Level

Amendment 56 to the BSAI Groundfish FMP, which was implemented in 1999, defines ABC and OFL for the BSAI groundfish fisheries. The new definitions are shown below, where the fishing mortality rate is denoted F , stock biomass (or spawning stock biomass, as appropriate) is denoted B , and the F and B levels corresponding to MSY are denoted F_{MSY} and B_{MSY} respectively.

Acceptable Biological Catch is a preliminary description of the acceptable harvest (or range of harvests) for a given stock or stock complex. Its derivation focuses on the status and dynamics of the stock, environmental conditions, other ecological factors, and prevailing technological characteristics of the fishery. The fishing mortality rate used to calculate ABC is capped as described under “overfishing” below.

Overfishing is defined as any amount of fishing in excess of a prescribed maximum allowable rate. This maximum allowable rate is prescribed through a set of six tiers which are listed below in descending order of preference, corresponding to descending order of information availability. The SSC will have final authority for determining whether a given item of information is reliable for the purpose of this definition, and may use either objective or subjective criteria in making such determinations. For tier (1), a pdf refers to a probability density function. For tiers (1-2), if a reliable pdf of B_{MSY} is available, the preferred point estimate of B_{MSY} is the geometric mean of its pdf. For tiers (1-5), if a reliable pdf of B is available, the preferred point estimate is the geometric mean of its pdf. For tiers (1-3), the coefficient α is set at a default value of 0.05, with the understanding that the SSC may establish a different value for a specific stock or stock complex as merited by the best available scientific information. For tiers (2-4), a designation of the form “ $F_{X\%}$ ” refers to the F associated with an equilibrium level of spawning per recruit (SPR) equal to $X\%$ of the equilibrium level of spawning per recruit in the absence of any fishing. If reliable information sufficient to characterize the entire maturity schedule of a species is not available, the SSC may choose to view SPR calculations based on a knife-edge maturity assumption as reliable. For tier (3), the term $B_{40\%}$ refers to the long-term average biomass that would be expected under average recruitment and $F=F_{40\%}$.

Overfished or approaching an overfished condition is determined for all age-structured stock assessments by comparison of the stock level in relation to its MSY level according to the following two harvest scenarios (Note for Tier 3 stocks, the MSY level is defined as $B_{35\%}$):

Overfished (listed in each assessment as scenario 6):

In all future years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be 1) above its MSY level in 2008 or 2) above $\frac{1}{2}$ of its MSY level in 2008 and above its MSY level in 2018 under this scenario, then the stock is not overfished.)

Approaching an overfished condition (listed in each assessment as scenario 7):

In 2008 and 2009, F is set equal to $\max F_{ABC}$, and in all subsequent years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is expected to be above its MSY level in 2020 under this scenario, then the stock is not approaching an overfished condition.)

For stocks in Tiers 4-6, no determination can be made of overfished status or approaching an overfished condition as information is insufficient to estimate the MSY stock level.

Tier	
1)	Information available: <i>Reliable point estimates of B and B_{MSY} and reliable pdf of F_{MSY}.</i>
1a)	Stock status: $B/B_{MSY} > 1$ $F_{OFL} = \mu_A$, the arithmetic mean of the pdf $F_{ABC} \leq \mu_H$, the harmonic mean of the pdf
1b)	Stock status: $\alpha < B/B_{MSY} \leq 1$ $F_{OFL} = \mu_A \times (B/B_{MSY} - \alpha)/(1 - \alpha)$ $F_{ABC} \leq \mu_H \times (B/B_{MSY} - \alpha)/(1 - \alpha)$
1c)	Stock status: $B/B_{MSY} \leq \alpha$ $F_{OFL} = 0$ $F_{ABC} = 0$
2)	Information available: <i>Reliable point estimates of B, B_{MSY}, F_{MSY}, $F_{35\%}$, and $F_{40\%}$.</i>
2a)	Stock status: $B/B_{MSY} > 1$ $F_{OFL} = F_{MSY}$ $F_{ABC} \leq F_{MSY} \times (F_{40\%}/F_{35\%})$
2b)	Stock status: $\alpha < B/B_{MSY} \leq 1$ $F_{OFL} = F_{MSY} \times (B/B_{MSY} - \alpha)/(1 - \alpha)$ $F_{ABC} \leq F_{MSY} \times (F_{40\%}/F_{35\%}) \times (B/B_{MSY} - \alpha)/(1 - \alpha)$
2c)	Stock status: $B/B_{MSY} \leq \alpha$ $F_{OFL} = 0$ $F_{ABC} = 0$
3)	Information available: <i>Reliable point estimates of B, $B_{40\%}$, $F_{35\%}$, and $F_{40\%}$.</i>
3a)	Stock status: $B/B_{40\%} > 1$ $F_{OFL} = F_{35\%}$ $F_{ABC} \leq F_{40\%}$
3b)	Stock status: $\alpha < B/B_{40\%} \leq 1$ $F_{OFL} = F_{35\%} \times (B/B_{40\%} - \alpha)/(1 - \alpha)$ $F_{ABC} \leq F_{40\%} \times (B/B_{40\%} - \alpha)/(1 - \alpha)$
3c)	Stock status: $B/B_{40\%} \leq \alpha$ $F_{OFL} = 0$ $F_{ABC} = 0$
4)	Information available: <i>Reliable point estimates of B, $F_{35\%}$, and $F_{40\%}$.</i> $F_{OFL} = F_{35\%}$ $F_{ABC} \leq F_{40\%}$
5)	Information available: <i>Reliable point estimates of B and natural mortality rate M.</i> $F_{OFL} = M$ $F_{ABC} \leq 0.75 \times M$
6)	Information available: <i>Reliable catch history from 1978 through 1995.</i> $OFL =$ the average catch from 1978 through 1995, unless an alternative value is established by the SSC on the basis of the best available scientific information $ABC \leq 0.75 \times OFL$

Overview of "Stock Assessment" Section

The current status of individual groundfish stocks managed under the FMP is summarized in this section. Plan Team recommendations for 2008 and 2009 ABCs and OFLs are summarized in Tables 1, 5 and 6. The added year was included to assist NMFS management since the TAC setting process allows for a period of up to two years to review harvest specifications. The sum of the recommended ABCs for 2008 and 2009 are 2,440,000 t and 2,560,000 t, respectively. These are approximately 236,000 t and 118,000 t below the sum of the 2007 ABCs. However, these values still exceed the 2 million t cap employed by the Council as a conservation measure in setting TACs. Overall, the status of the stocks continues to appear relatively favorable, although many stocks are declining due to poor recruitment in recent years. Total biomass for 2008 (16.6 million t) is less than last year's estimate of 16.9 million t.

Overall groundfish exploitable biomass is high but declining, especially for pollock and Pacific cod. The bottom trawl survey biomass estimate for pollock in 2007 was 4.3 million t, only 87% of the long-term mean of the bottom-trawl survey. The 2007 echo-integration (EIT) survey biomass estimate was 1.88 million t, only 55% of the long-term mean for this survey. Both surveys indicate that the 2006 year class is strong and that the 2005 year class is now apparently below average. The biomass estimate from the 2007 bottom trawl survey for Pacific cod of 424,000 t is down about 18% from the 2006 estimate, and is the all-time low. Plan Team ABC recommendations are trending down for gadoids, but generally up for flatfishes.

The abundances of AI pollock, sablefish, all rockfishes, all flatfishes, and Atka mackerel are projected to be above target stock size. The abundances of EBS pollock and Pacific cod are projected to be below target stock size (Figure 2).

Summary and Use of Terms

Stock status is summarized and OFL and ABC recommendations are presented on a stock-by-stock basis in the remainder of this section, with the following conventions observed:

- "Fishing mortality rate" refers to the full-selection F (i.e., the rate that applies to fish of fully selected sizes or ages), except in the cases of the EBS pollock, yellowfin sole, and northern rock sole assessments. For these stocks, the fishing mortality rate consists of the ratio between catch (in biomass) and biomass at the start of the year. EBS pollock uses "fishable biomass" whereas the two flatfish stocks use age 6+ biomass for this calculation.
- "Projected age+ biomass" refers to the total biomass of all cohorts of ages greater than or equal to some minimum age, as projected for January 1 of the coming year. The minimum age varies from species to species. When possible, the minimum age corresponds to the age of recruitment listed in the respective stock assessment. Otherwise, the minimum age corresponds to the minimum age




Figure 2. Summary status of age-structured BSAI groundfish species relative to 2007 catch levels (vertical axis) and projected 2008 spawning biomass relative to B_{msy} levels. Note that the 2007 MSY level is taken as the 2007 OFL

included in the assessment model, or to some other early age traditionally used for a particular species. When a biomass estimate from the trawl survey is used as a proxy for projected age+ biomass, the minimum age is equated with the age of recruitment, even though the survey may not select that age fully and undoubtedly selects fish of younger ages to some extent.

- “Exploitation rate” refers to the ratio between catch (in numbers) and start-of-year stock size (also in numbers). Where information is lacking, the exploitation rate is sometimes multiplied by start-of-year biomass to compute ABC.
- Projected ABC, OFL, and biomass levels are reported to three significant digits, except when quoting a Council-approved value with more than three significant digits or when a stock-specific ABC is apportioned among areas on a percentage basis, in which case four significant digits may be used if necessary to avoid rounding error. Fishing mortality rates are reported to two significant digits.
- The reported ABCs and OFLs for the past year correspond to the values approved by the Council. Projected ABCs and OFLs listed for the next two years are the Plan Team’s recommendations.
- Reported catches are as of October 27, 2007

Two-Year OFL and ABC Projections

Amendment 48 to the BSAI Groundfish FMPs, implemented in 2005, made two significant changes with respect to the stock assessment process. First, annual assessments are no longer required for rockfishes since new data during years when no groundfish surveys are conducted are limited. For example, since 2007 was an off-year for the NMFS BSAI groundfish trawl survey, only summaries for these species were produced.

The second significant change is that the proposed and final specifications are to be specified for a period of up to two years. This requires providing ABC and OFL levels for 2008 and 2009 (Table 1). In the case of stocks managed under Tier 3, 2008 and 2009 ABC and OFL projections are typically based on the output for Scenarios 1 or 2 from the standard projection model using assumed (best estimates) of actual catch levels.

In the case of stocks managed under Tiers 4-6, 2009 projections are set equal to the Plan Team’s recommended values for 2008.

The 2009 ABC and OFL values recommended in next year’s SAFE report are likely to differ from this year’s projections for 2009, for the same reasons that the 2008 projections in this SAFE report differ from those in September.

Uncertainty

Statistical uncertainty is addressed in the individual assessments, and to some degree, by the tiers used to establish ABCs. Statistical uncertainty or natural variability in the stock led the Plan Team to recommend 2008 and 2009 ABC values lower than the maximum permissible level for walleye pollock and Greenland turbot. The Plan Team’s recommended 2008 and 2009 ABC for pollock is 15 and 7 percent below the maximum permissible level, respectively. The Plan Team’s recommended 2008 ABC for Greenland turbot is 82 percent below the maximum permissible level for 2008 and 2009.

Ecosystem Considerations / Research

Ecosystem considerations are addressed in the stock assessment chapters. Similar to past years, several assessments (walleye pollock, Pacific cod, yellowfin sole, arrowtooth flounder, northern rock sole, flathead sole) included discussions to estimate relationships between bottom trawl survey catchability and bottom temperature. In some assessments (rock sole, flathead sole, Alaska plaice), potential effects of a possible 1989 regime shift on the stock-recruitment relationship are investigated (the yellowfin sole assessment included a similar analysis for the 1978 regime shift). Assessments for BSAI arrowtooth

flounder, BS flathead sole, EBS walleye pollock, AI walleye pollock, BSAI cod, and AI Atka mackerel assessment incorporated results from ecosystem models. Ecosystem model results were also included in the squid and other species assessments for squids, skates, sculpins, and octopus. Linkages between BSAI arrowtooth and EBS pollock were reviewed again this year, and recent data indicate that arrowtooth are an increasingly important predator of pollock in the EBS (although not at the scale observed in the Gulf of Alaska (GOA)).

Last year, the BSAI Pacific cod ecosystem considerations reviewed the role of cod as a predator in the EBS and AI ecosystems. These results indicate that fishing mortality is the dominant source of explained cod mortality in both ecosystems, and that changing cod mortality in each ecosystem affects a different set of species with different magnitudes. This year, the Plan Team reiterated its suggestion that separate specifications for Pacific cod in the AI and EBS be considered as an ecosystem consideration in future assessments. During the discussion of natural mortality, it was discussed that “explained” sources of natural mortality (i.e., predation) only accounted for a natural mortality rate of 0.04 for cod, with the rest of the natural mortality being “unexplained” and perhaps due to disease, parasites, or migration. Thus predation mortality is uninformative in selecting a range of M for use in assessments.

A full ecosystem analyses for the EBS presented in the Ecosystem Assessment section of the Ecosystem Considerations chapter suggested that there may have been a decrease in forage species biomass in 1999-2007 relative to the 1980s-1990s. In general, the EBS was cold again this year, with a fuller ice extent and later ice retreat compared with recent years; it was remarked that this more “normal” condition was localized to the EBS, and that in general the Arctic continued its trend of warmer temperatures and lower ice coverage. The question of whether the changes in biota seen between 1998 and 2001 and the recent changes in ice cover represent a new or permanent “regime” was discussed in the context of whether pollock production was in a new regime, although no conclusion was reached or suggested. Although the Team was unable to identify any cases where these or other ecosystem considerations suggested an immediate need to adjust ABC, the Team encourages further development of ecosystem considerations. A full review of ecosystem status and trends is provided in the Ecosystem Considerations chapter.

Stock Status Summaries

1. Walleye Pollock

Status and catch specifications (t) of walleye pollock in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2007 and 2008 are those recommended by the Plan Team. Catch data are current through 10/27/07.

Area	Year	Age 3+ Biomass	OFL	ABC	TAC	Catch
Eastern Bering Sea	2006	8,050,000	2,090,000	1,930,000	1,485,000	1,486,413
	2007	6,360,000	1,640,000	1,394,000	1,394,000	1,350,530
	2008	4,360,000	1,440,000	1,000,000	n/a	n/a
	2009	n/a	1,320,000	1,000,000	n/a	n/a
Aleutian Islands	2006	130,000	39,100	29,400	19,000	1,735
	2007	229,000	54,500	44,500	19,000	2,488
	2008	197,000	34,000	28,200	n/a	n/a
	2009	n/a	26,100	22,700	n/a	n/a
Bogoslof	2006	253,000	50,600	5,500	10	0
	2007	240,000	48,000	5,220	10	0
	2008	292,000	58,400	7,970	n/a	n/a
	2009	n/a	58,400	7,970	n/a	n/a

Eastern Bering Sea

Changes from previous assessment

New data in this year's assessment include the following:

- Updated total catch for 2006 and a preliminary estimate of the 2007 catch.
- Biomass estimates from the 2007 bottom trawl survey and the 2007 echo-integration trawl (EIT) survey. The estimate from the bottom trawl survey was 4.3 million t, up 42% from the 2006 estimate. The estimate from the EIT survey was 1.88 million t, up 20% from last year's survey. Although both survey estimates are higher than last year's, both are substantially below the long-term means for their respective time series.
- Age composition data from the 2007 bottom trawl survey, updated age composition data from the 2006 EIT survey, and preliminary age composition data from the 2007 EIT survey (based on the age-length key from this year's bottom trawl survey). The 2007 survey age compositions give evidence of a large 2006 year class.
- Age and size composition data and weight-at-age data from the 2006 fishery. The new weight-at-age data resulted in a significant decrease in the 2006 mean weights at age relative to the values used in last year's assessment.
- Sample sizes specified in the model were re-evaluated using a bootstrap approach.

Changes in model structure include the following:

- Length composition data (not just age composition data) can now be used in the model.
- Relative abundance at age 1 in the EIT survey is now estimated separately (as an independent recruitment index) from the other age groups.
- The survey abundance index used for tuning the model was changed from age 1+ numbers to age 2+ numbers.
- The catch/biomass ratio used in applying the Tier 1 harvest control rules now uses "fishable biomass" (the sum of the product of selectivity-, weight-, and numbers-at-age) for the denominator instead of age 3+ biomass. This makes the computation less sensitive to fluctuations in incoming year class strength.

Spawning biomass and stock status trends

Consistent with the estimates produced in last year's assessment, abundance of EBS walleye pollock has declined steadily since 2003 due to poor recruitment from the 2001-2005 year classes. This string of five consecutive poor year classes is unprecedented in the known history of the stock. Spawning biomass is estimated to be 4% above B_{MSY} in 2007 but projected to be 28% below B_{MSY} in 2008. The age 3+ biomass for 2007 is estimated to be the lowest in the time series since 1980. Although preliminary indications are that the 2006 year class is well above average, spawning biomass is unlikely to exceed B_{MSY} until 2010.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that reliable estimates of B_{MSY} and the probability density function for F_{MSY} exist for this stock. Therefore, EBS walleye pollock qualify for management under Tier 1. The Plan Team concurs with the assessment authors' conclusion that the Tier 1 reference points continue to be reliably estimated.

The updated estimate of B_{MSY} from the present assessment is 1.88 million t, compared to 2.06 million t from last year's assessment. Projected spawning biomass for 2008 is 1.38 million t, placing EBS walleye pollock in sub-tier "b" of Tier 1. As in recent assessments, the maximum permissible ABC harvest rate was based on the ratio between MSY and the equilibrium biomass corresponding to MSY. The harmonic mean of this ratio from the present assessment is 0.341, significantly higher than last year's value of 0.243. The difference is due to a change in the biomass measure used in the denominator of the ratio,

from age 3+ biomass (in last year's assessment), to fishable biomass (in this year's assessment). The lead author noted that this method change results in the same average yield but with less inter-annual variability.

The harvest ratio of 0.341 is scaled according to the Tier 1b formula and then multiplied by the geometric mean of the projected fishable biomass for 2008 (4.77 million t) to obtain the maximum permissible ABC for 2008, which is 1.17 million t. This ABC is more than double the 2008 yield of 555,000 t that would correspond to a Tier 3b strategy based on a $B_{40\%}$ value of 2.63 million t and an $F_{40\%}$ value of 0.51.

A range of ABC values from 555,000 – 1,170,000 t was discussed by Plan Team, with arguments offered in support of candidate values spanning the full range. Arguments in support of setting the 2008 ABC at 1.17 million t included the following:

- The stock qualifies for management under Tier 1, so the maximum permissible Tier 1 ABC should have priority unless there is a compelling reason to set a lower ABC.
- The Tier 1 harvest control rules already have a built-in precautionary adjustment for stocks that fall below B_{MSY} .
- Uncertainty is already factored into the Tier 1 harvest control rules.
- A 2008 ABC of 1.17 million t would already constitute a very large (16%) reduction from the 2007 ABC of 1.394 million t and would result in greater short-term catch stability than a lower ABC.
- Biomass is expected to rebuild to B_{MSY} under the maximum permissible ABC about as fast as it would under more conservative strategies (Figure 1.43).

Arguments in support of an ABC lower than 1.17 million t included the following:

- A 2008 ABC of 1.17 million t would imply an all-time high spawning exploitation rate. Keeping the 2008 ABC at or below about 1 million t would not exceed the all-time high spawning exploitation rate.
- There are many examples of strong year classes being produced when biomass is near $B_{40\%}$, but only one strong year class has been produced at a biomass lower than the projected 2008 value, implying that it would be desirable to increase biomass sooner rather than later.
- The stock rebuilt successfully from a similarly low level in the late 1970s and early 1980s when catches were limited to 1 million t or less.
- The possibility of a retrospective bias calls for additional precaution.
- The five-year string of consecutive poor recruitments spawned between 2001 and 2005 is unprecedented, also calling for additional precaution.

After lengthy discussion, the Plan Team voted to support the authors' recommendation of a 1 million t ABC for 2008. The decision was not unanimous, with some Plan Team members voting for lower values (specifically, 976,000 t and 555,000 t). The Plan Team emphasized that its recommendation is intended to constitute a precautionary ABC, and that if next year's assessment does not confirm the current estimated strength of the 2006 year class, further reductions may be necessary. A 2008 ABC of 1 million t would correspond to a harvest ratio of 0.21 and an $F_{39\%}$ harvest rate (compared to Tier 1A ABC rate of $F_{32\%}$). The current projection for maximum permissible ABC in 2009 given a 2008 catch of 1 million t is 1.07 million t. However, the Team recommended 1 million t for 2009 ABC.

The OFL harvest ratio under Tier 1a is 0.422, the arithmetic mean of the ratio between MSY and the equilibrium fishable biomass corresponding to MSY. The product of this ratio, rescaled according to the Tier 1b formula, and the geometric mean of the projected fishable biomass for 2008 gives the OFL for 2008, which is 1.44 million t. A 2008 OFL of 1.44 million t would correspond to a harvest ratio of 0.30. The current projection for OFL in 2009 given a 2008 catch of 1 million t is 1.32 million t. The walleye pollock stock in the EBS is not overfished and is not approaching an overfished condition.

Aleutian Islands

Changes from previous assessment

Model 2B is similar to the model accepted by the SSC last year and is recommended by the authors again this year.

Spawning biomass and stock status trends

Age 2+ biomass is estimated to have increased from 1999 to 2004, after which it has been stable. Spawning biomass is estimated to have been increasing slowly since 1999. The 2000 year class is estimated to have been well above average (third largest in the time series), and preliminary indications are that the 2005 and 2006 year classes may be slightly above average. Spawning biomass for 2008 is projected to be 82,300 t.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that this stock qualifies for management under Tier 3. Given that spawning biomass has been increasing and is above the $B_{40\%}$ value of 51,500 t, the Plan Team concurs with the authors' recommendation to set 2008 ABC at the maximum permissible value (Tier 3a, with $F_{40\%}=0.20$) of 28,200 t. Assuming a 2008 catch equal to the ABC, the maximum permissible ABC for 2009 is projected to be 22,700 t.

Following the Tier 3a formula with $F_{35\%}=0.24$, OFL for 2008 is 34,000 t. The projected OFL for 2009 is 26,100 t. The walleye pollock stock in the Aleutian Islands is not overfished and is not approaching an overfished condition.

Bogoslof

Changes from previous assessment

No changes in analytic approach were made in this assessment.

Spawning biomass and stock status trends

Survey biomass estimates since 2000 have all been lower than estimates prior to 2000, ranging from a low of 198,000 t in 2003 to a high of 301,000 t in 2000. The 2007 estimate is the highest since the 2000 estimate.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that this stock qualifies for management under Tier 5. Traditionally, the ABC for this stock has been set using a formula similar to the Tier 3 formula, but substituting a reference biomass level of 2 million t for $B_{40\%}$. The Plan Team concurs with the authors' recommendation to continue this practice. Given $F_{40\%}=0.27$, this results in $F_{ABC}=0.022$ and a 2008 ABC of 7,970 t. The projected ABC for 2009 is the same.

Following the Tier 5 formula with $M=0.20$, OFL for 2008 is 58,400 t. The OFL for 2009 is the same. As a Tier 5 stock, it is not possible to determine whether Bogoslof pollock is overfished or is approaching an overfished condition.

2. Pacific Cod

Status and catch specifications (t) of Pacific cod in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through early Oct, 2007.						
Area	Year	Age 3+ Biomass	OFL	ABC	TAC	Catch
BSAI	2006	922,000	230,000	194,000	194,000	191,906
	2007	960,000	207,000	176,000	171,000	172,407
	2008	1,080,000	176,000	150,000	n/a	n/a
	2009	n/a	190,000	162,000	n/a	n/a

Changes from previous assessment

The assessment of the BSAI Pacific cod stock has been particularly challenging as more refinements are made to the modeling approaches and use of the data. The document described the evolution of analytical approaches. Following a series of modifications from 1993 through 1997, the base assessment model remained completely unchanged from 1997 through 2001. As data refinement and analytical techniques developed, a major change took place in 2005 when the model was migrated to the Stock Synthesis 2 (SS2) program. Difficulties encountered at the 2006 assessment resulted in a thorough review of the 2006 assessment model in April 2007 during a public work shop that brought together 44 participants. Many suggestions for changes and refinements of the analytical approaches were made.

The assessment authors took into consideration the results of the public work shop and preliminary SAFE Report in developing 4 versions of the analytical model for the 2007 assessment. The following points may be made to distinguish these 4 models:

Model 1 was developed to respond to requests of the SSC where, among all the other model parameters, the natural mortality (M) is to be based on external analyses of life history parameters which resulted in $M = 0.34$;

Model 2 is the same as Model 1 except that M is fixed at 0.37 (that was used in previous year's assessments);

Model 3 is the same as Model 1 except that M is estimated internally; and Model 4 which differs from Model 1 in several respects to respond to public comments on the use and fitting of the data.

While the assessment document elaborated on nine major categories of new input data, they are essentially necessary routine applications of updated data from the fisheries and surveys. The following new pieces of data stand out: 1) a new biomass for Pacific cod was estimated at 424,000 t from the 2007 EBS shelf bottom trawl survey (this biomass is 18% lower than the 2006 survey estimate and is the all-time low in the time series), and 2) the numbers of fish for 2007 was estimated by the survey at 713 million fish, up about 86% from the 2006 estimate. This dramatic increase in numbers of fish is due primarily to higher recruitment of the 2006 year class as age 1 fish in the survey. The addition of this new data point to the series of survey population numbers from 1979-2006 has a material impact on the projection of Pacific cod population numbers and biomass into the future.

The main features and challenges of the Pacific cod assessment, however, are still the fitting of the data to the four assessment models. Because all of the models seem to perform reasonably well in terms of fitting the data, the authors used the following three major criteria for selection of the best model to represent the dynamics of the stock. These criteria are:

- The model should use a reasonable value of M;
- The model should estimate mean trawl survey lengths for ages 1-3 that are close to the first 3 modes from the long-term average trawl survey size composition;

- The model should estimate a reasonable average for the product of trawl survey catchability and trawl survey selectivity for the 60-81 cm size range.

Based on these criteria, Model 1 was selected by the authors to best represent the population dynamics of the BSAI Pacific cod stock. The Plan Team agrees with this selection and added the following comments:

- All selectivity curves estimated by the 4 models appear reasonable, except for the IPHC longline survey selectivity curve. Increasing the number of parameters from 4 to 6 still did not overcome the stated problems with estimating selectivity.
- Model 4 ignores age data. It makes little sense to discard data that provides information on relative cohort strength, even if the fit is “not very good”. Relying on length data alone is not as informative as incorporating age data because there is not a one-to-one relationship between length and age, except for the youngest ages.
- Model 1, 2, 3 and 4 biomass (Fig. 2.3) and recruitment (Fig. 2.2) trends are similar. These models differ primarily in values of M (except that Model 4 has other structural differences). The implication of the model results is that the data are sufficient to estimate trends, but the biomass scalar is sensitive to values of M .
- The best estimate of M is 0.34. It is based on a derivation from standard functional relationships for the life history of Pacific cod reported by Jensen (1996). This value compares favorably with data and parameter estimates from other reports (e.g., Beverton and Holt (1963)).
- For model selection, Model 3 actually fits the data best, but the model was not selected because its M value (0.22) was too low. Model 2 ($M=0.37$) was not selected because it did not meet criterion 3. Model 4 was not selected because $M=0.46$ was judged to be too high.

Since Model 1 ($M=0.34$) was selected, the ABC and other parameters values for 2008 and 2009 are provided by the model results shown in Table 2.15 b and c. In accepting these values, the Plan Team made the following observations:

- The 2008 ABC of 150,000 t is a decline from 176,000 t in 2007. This ABC drop is consistent with the 18% decline in the NMFS survey biomass from 2006 to 2007, which currently is at its historic minimum.
- Five consecutive year classes of the EBS Pacific cod stock from 2001-2005 (that ranged from 204-399 million age zero fish) are noticeably below the 30-year average year class strength (658 million age zero fish from 1977-2006). However, the 2006 year class appears to be more than 2 ½ times higher than the average recruitment.
- The female spawning biomass for 2008 and 2009 are projected to be at about 29% of the unfished female spawning biomass; at least until the strong 2006 year class can contribute substantially to spawning from 2010.

Spawning biomass and stock status trends

A standard set of projections is required for each stock managed under Tiers 1, 2 or 3. The authors evaluated 7 harvest scenarios to make projections of the biomass and status trends to satisfy Amendment 56, NEPA, and the MSFCMA. The Plan Team has selected Scenario 1 as the most likely representation to determine how the spawning biomass and stock status would trend. Scenario 1 assumes that in all future years, F is set equal to $\max F_{abc}$ which is close to the Council’s decisions in the past to set TACs at or close to the maximum ABCs.

Under Scenario 1, the spawning biomass is projected to continue a slow decline from 2007 to 2009 before the strong 2006 year class would boost the female spawning biomass from 398,000 t during the 2008-09 period to 454,000 t in 2010 and 542,000 t in 2011 (Table 2.26). This projected increase in biomass and upward trend of stock status are predicated on the continued strength and contributions of the 2006 year class. This 2006 year class indicator is still an early indicator and it must be tracked carefully as the year

class follows through the population in time. The effect of such a strong year class following through is particularly important to the stock as it follows 5 consecutive weak year classes from 2001-2005. The average recruitment of age zero fish for the 30-year time series (1977-2006) was 658 million fish. The 4 years of consecutive weak year classes from 2001-05 ranged from 204-399 million fish. The 2006 year class at age zero was estimated to be 1.835 billion fish, the second highest on record in the 30 year time series that included the known strong 1977 year class that was estimated at 2.533 billion fish. The 2006 strong year class bodes well for the status of the stocks as it ages and moves into the fishery.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

According to criteria set by the SSC, this stock qualifies for management under Tier 3, where reliable estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist for the stock. The updated point estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ from the present assessment are 540,000 t, 0.31 and 0.37, respectively. Pacific cod specifically qualifies for management under sub-tier "b" of Tier 3 because the projected biomass for 2008 (398,000 t) is below $B_{40\%}$. Fishing at the adjusted Tier 3b rate of 0.22 is projected to result in a 2008 catch of 150,000 t, which is the maximum permissible ABC under Amendment 56.

The Plan Team recommends setting the 2008 ABC at the maximum permissible value of 150,000 t (which is 14.8% below the 2007 ABC of 176,000 t). This ABC is projected to increase to a maximum permissible ABC of 162,000 in 2009.

The OFL for 2008 under Tier 3b is 176,000 t ($F_{OFL}=0.26$) and the projected OFL for 2009 is 190,000 t.

The stock is not overfished nor approaching an overfishing condition.

Ecosystem Considerations summary

The Pacific cod chapter included a discussion paper on "results from ecosystem models on the role of Pacific cod in the eastern Bering Sea and Aleutian Islands ecosystems" as well as an extended discussion section on "ecosystem considerations". While these discussions provide useful clarifying information about the ecological dynamics of Pacific cod and its fisheries to the rest of the ecosystem, there was no special identification of ecosystem features that would require adjustments to the estimated ABCs and their attendant reference population parameters.

Area apportionment

At present, ABC of the BSAI Pacific cod is not allocated by area. However, the biomass distribution analysis made in 2006 using the Kalman filter approach estimated the biomass distribution at 84% EBS and 16% AI.

3. Sablefish

Status and catch specifications (t) of sablefish in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through 10/27/07.

Area	Year	Age 4+				
		Biomass	OFL	ABC	TAC	Catch
EBS	2006	34,000	3,680	3,060	3,060	1,055
	2007	34,000	3,520	2,980	2,980	1,088
	2008	41,000	3,380	2,860	n/a	n/a
	2009	n/a	2,910	2,610	n/a	n/a
AI	2006	32,000	3,740	3,100	3,100	1,130
	2007	32,000	3,320	2,810	2,810	1,078
	2008	34,000	2,890	2,440	n/a	n/a
	2009	n/a	2,510	2,230	n/a	n/a

Changes from previous assessment

Sablefish are assessed as a single stock in the BSAI and GOA using a split sex age structured model. The split sex model approach was fully implemented in 2006 and was deemed appropriate given differences in growth between males and females. The assessment model incorporates the following new data into the model: relative abundance and length data from the 2007 longline survey, relative abundance and length data from the 2006 longline fishery, length data from the 2006 trawl fishery, and age data from the 2006 longline survey and longline fishery. In addition, relative abundance and length data from the 2007 Gulf of Alaska trawl survey were included with the expectation of improving estimates of recruitment. New growth data were added (1996-2004) in the form of revised age-length transition matrices, and older growth data (1981-1993) were updated. Fishery CPUE data from observer data and logbooks were used in the catch rate analysis. The commercial CPUE observer data were screened to exclude sets where killer whale depredation and targeting of other species occurred. Logbook data were similarly screened to account for multiple gear configurations. The results showed a good agreement between observer and logbook fishery CPUE and the survey CPUE in most years. The only major changes to the model were the inclusion of informative priors on catchability for all abundance indices. The survey abundance index decreased 14% between 2006 and 2007, a change which follows the 13% increase between 2005 and 2006. The fishery abundance index was down 8% from 2005 to 2006.

Spawning biomass and stock status trends

The spawning biomass is projected to be similar from 2007 to 2008, but is expected to decline through 2012. The projected 2008 female spawning biomass is 37% of unfished biomass compared with about 29% of unfished biomass estimated during the 1998 to 2001 period. The 2000 year class now appears to be larger than the 1997 year class and is expected to comprise 18% of the spawning biomass in 2008.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

This stock qualifies for management under Tier 3. The updated point estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ from this assessment are 122,000 t (combined across the EBS, AI, and GOA), 0.093, and 0.111, respectively. Projected spawning biomass (combined areas) for 2008 is 112,000 t (91% of $B_{40\%}$), placing sablefish in sub-tier "b" of Tier 3. The maximum permissible value of F_{ABC} under Tier 3b is 0.084, which translates into a 2008 catch (combined areas) of 18,000 t and is the Plan Team's recommended combined 2008 ABC. The recommended 2008 ABC is approximately 10% lower than the 2007 ABC of 21,000 t. The OFL fishing mortality rate under Tier 3b is 0.101. This fishing mortality rate translates into a 2008 OFL (combined areas) of 21,300 t. The Team's recommendations for EBS and AI sablefish OFLs and ABCs are provided in the table above. Alaska sablefish are not overfished nor are they approaching an overfished condition.

Additional Plan Team recommendations

The combined ABC has been apportioned to regions using a weighted moving average method since 1993. Since 2000, both survey and fishery data have been used to apportion ABC. The current method is to compute a 5-year exponential weighting for each index which are then combined, with the survey data weighted twice as heavily as the fishery data. The original rationale for this was that the variance for the fishery data was twice that of the survey data. Recent improvements to the sample size of observer and logbook collections have reduced the variance on the fishery source and led to industry requests to weight the two data sets equally. The Plan Team heard no compelling evidence to switch from the present weighting scheme to the other and for this year has simply continued the recent method of double weighting the survey data, which is reflected in the recommended area apportionments. The Plan Team notes that the increase in fishery data has largely occurred due to voluntary submission of logbooks as well as the availability of funds to enable the IPHC to collect and process the fishery data. If equal weighting of the two data sets is envisioned, it becomes paramount that a more stable or permanent source of funding be found to ensure continued collection of logbooks. The Plan Team notes that the difference in apportionment between the two methods is relatively minor for 2007.

Area apportionment

A 5-year exponential weighting of longline survey and fishery relative abundance indices (the survey index is weighted double the fishery index) may be used to apportion the combined 2008 ABC among regions, resulting in the following values: 2,860 t for EBS and 2,440 t for AI. Relative to 2006, apportionments to the EBS and AI decreased.

Using the survey/fishery based apportionment scheme described above, 2008 OFL also may be apportioned among regions and results in the following values: 3,380 t for EBS and 2,890 t for AI. These values also represent a decrease from 2007 OFL levels for all regions.

4. Yellowfin Sole

Status and catch specifications (t) of yellowfin sole in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through 10/27/07.						
Area	Year	Age 2+ Bio.	OFL	ABC	TAC	Catch
BSAI	2006	1,680,000	144,000	121,000	95,700	99,068
	2007	2,000,000	240,000	225,000	136,000	119,332
	2008	2,200,000	265,000	248,000	n/a	n/a
	2009	n/a	296,000	276,000	n/a	n/a

Changes from previous assessment

Changes to the input data for this year's assessment are the inclusion of 2006 fishery and survey age compositions, the 2007 trawl survey biomass point estimate and standard error, estimates of the discarded and retained portions of the 2006 catch, catch through 8 September 2007 and an update of weight-at-age estimates using biological data through 2006

This year's EBS bottom trawl survey resulted in a biomass estimate of 2,150,000 t, about 1% greater than last year's survey biomass of 2,130,000 t.

Spawning biomass and stock status trends

The projected female spawning biomass estimate for 2008 is 550,000 t. Relative to the 585,000 t projected spawning biomass from the 2006 assessment; this continues the slight monotonic decline in model estimates of spawning biomass exhibited since 1994. However, above average recruitment from the 1995 and 1999 year-classes is expected to maintain the abundance of yellowfin sole at a level above B_{40} in the near future.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that reliable estimates of B_{MSY} and the probability density function for F_{MSY} exist for this stock. Accordingly, yellowfin sole qualify for management under Tier 1. The estimate of B_{MSY} from the present assessment is 303,000 t. As in last year's assessment, 1978-2002 spawner recruit data were used this year as the basis to determine the Tier 1 harvest recommendation. This provided an $F_{ABC} = F_{\text{harmonic mean } F_{msy}} = 0.19$. The $F_{OFL} = F_{MSY} = 0.2$. The product of the harmonic mean of F_{MSY} and the geometric mean of the 2008 biomass estimate produced the author- and PT-recommended ABC of 248,000 t and OFL of 265,000 t.

Model projections indicate that this stock is neither overfished nor approaching an overfished condition.

Ecosystem Considerations summary

The assessment contains an ecosystem component by representing catchability of the EBS shelf trawl survey as an exponential function of average annual bottom temperature during the EBS shelf trawl survey.

5. Greenland turbot

Status and catch specifications (t) of Greenland turbot in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through 10/27/07.

Year	Area	Age 1+ Bio.	OFL	Subarea	ABC	TAC	Catch
2006	BSAI	74,200	14,200		2,740	2,740	1,965
				EBS	1,890	1,890	1,440
				AI	850	850	525
2007	BSAI	119,000	15,600		2,440	2,440	1,946
				EBS	1,680	1,680	1,435
				AI	760	760	511
2008	BSAI	104,100	15,600		2,540	n/a	n/a
				EBS	1,750	n/a	n/a
				AI	787	n/a	n/a
2009	BSAI		16,000		2,540	n/a	n/a
				EBS	1,750	n/a	n/a
				AI	787	n/a	n/a

Changes from previous assessment

This year's Greenland turbot assessment model included updated 2006 and 2007 catch data, EBS shelf survey 2007 biomass and length composition estimates, and aggregated longline survey data index for the EBS and Aleutian Islands regions. A simplified Tier 5 approach was also provided for contrast to the Stock Synthesis 2 model results. As in last year's assessment, the slope-trawl survey was assumed to index 75% of the Greenland turbot stock inhabiting US waters.

Spawning biomass and stock status trends

The current estimate of 2008 female spawning biomass is 58,100 t. Compared to the 2007 spawning biomass of 56,900 t this represents a very slight increase, a departure from the decline prevalent since the mid 1970s.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that reliable estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist for this stock. Greenland turbot therefore qualify for management under Tier 3a. Updated point estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ from the present assessment are 38,200 t, 0.51, and 0.67, respectively. Projected spawning biomass for 2008 is 58,100 t. The maximum permissible value of F_{ABC} under this tier translates into a 2008 ABC of 12,200 t. Although there appear to be some favorable recruitments in recent years, fishing mortalities consistent with recent history are recommended for ABCs until another slope survey can be completed. Therefore the authors recommended setting the 2008 ABC at a value less than the maximum permissible. The Plan Team agrees with the authors' recommendation to use $F_{ABC} = 5\text{-year average catch}$, which results in a 2008 ABC of 2,500 t corresponding to a full selection fishing mortality rate of 0.09. The OFL fishing mortality rate is computed under Tier 3a, $F_{OFL} = F_{35\%} = 0.67$, and translates into a 2008 OFL of 15,600 t.

Additional Plan Team recommendations

The PT recommended conducting the EBS slope survey in 2008. This could provide information needed to determine whether abundance is sufficient to warrant recommending an ABC set to the maximum permissible level in the future.

6. Arrowtooth Flounder

Status and catch specifications (t) of arrowtooth flounder in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2007 and 2008 are those recommended by the Plan Team. Catch data are current through 10/27/07.						
Area	Year	Age 1+ Bio.	OFL	ABC	TAC	Catch
BSAI	2006	964,000	166,000	136,000	12,000	13,302
	2007	1,280,000	193,000	158,000	20,000	11,701
	2008	1,780,000	297,000	244,000	n/a	n/a
	2009	n/a	300,000	246,000	n/a	n/a

Changes from previous assessment

The present assessment is a straightforward update of last year's assessment. Input data were updated with the inclusion of fishery catch and discards through 8 September 2007, new data were also included for the 2007 shelf trawl survey size composition and biomass point-estimates and standard errors. Estimates of retained and discarded portions of the 2006 catch were added.

The stock assessment was expanded to include the 10 Aleutian Islands surveys and the survey size composition for arrowtooth flounder. Formerly 87% of the arrowtooth flounder stock was in shelf survey. With AI data added, only 73% of the stock is now from the Bering Sea shelf. The remaining fish come from the Bering Sea slope.

The ABC and OFL numbers appear to be greatly increased, but it is artifact of model because the Aleutian Islands data were included in this analysis.

Spawning biomass and stock status trends

The stock assessment model resulted in a biomass estimate of 1,780,000 t. This was an increase in the biomass estimate compared to the 2006 estimate of 1,280,000 t. The surveys in 2005 and 2006 had the highest biomasses of arrowtooth flounder recorded in the EBS to date. There is a long-term trend of increasing arrowtooth flounder biomass in the EBS. The trend in increasing biomass is expected to continue as recruitment has also been increasing for the last 10 years. Arrowtooth flounder in the AI leveled off in 1990s, but has been steadily increasing since then.

The authors noted that the ABC recommendation is for the combined harvest of arrowtooth flounder and Kamchatka flounder, which are difficult to distinguish and had similar biomass trends from the EBS trawl survey since 1991. Though Kamchatka flounder can now be identified and separated from arrowtooth flounder on AFSC surveys, it is not distinguished in the fishery. Therefore, the estimate is for one *Atheresthes* spp. stock.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The Plan Team concurred with the authors that there are structural problems with this model that must be resolved before arrowtooth flounder can be changed to Tier 1 status. Therefore, according to previous SSC determination that reliable estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist for this stock, arrowtooth flounder was assessed for management under Tier 3. The updated point estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ from the present assessment are 345,000 t, 0.24, and 0.30, respectively. Given that the projected 2008 spawning biomass of 994,000 t exceeds $B_{40\%}$, the Plan Team's ABC and OFL recommendations for 2008 were calculated under sub-tier "a" of Tier 3. The Plan Team recommends setting F_{ABC} at the $F_{40\%}$ (0.24) level, which is the maximum permissible level under Tier 3a. Projected harvesting at the $F_{40\%}$ level gives a 2007 ABC of 244,000 t. The OFL fishing mortality rate under Tier 3a is $F_{35\%}$ (0.30), which translates to a 2008 OFL of 297,000 t.

As there is little to no fishery for arrowtooth flounder the model is mostly driven by the survey data.

More female arrowtooth flounder are caught than males during the fisheries and the surveys. As in recent assessments, the base model that does not include the Aleutian Islands was evaluated using a range of male natural mortality rates between 0.27 and 0.34. It is believed males are not missed by the survey, but that they experience higher mortality than the females do. The best fit for males was $M=0.28$, but male selectivity is only 0.65.

To employ the base model without AI, the sex ratio must be changed. The authors recommend $M=0.33$, selectivity = 0.89. That leaves room for some male arrowtooth flounder to migrate down slope or be missed in survey. The age data from the GOA has few males >17 yrs, but lots of females that live to 25 yrs. Therefore, natural mortality of male arrowtooth flounder must be increased, as opposed to decreasing gear selectivity at older ages in the male population. If the fish were up in water column feeding and therefore not available to trawls, it would not be sex related. The authors tested a range of M , and determined $M = 0.33$ to be reasonable because 0.9 is a reasonable selectivity.

The catchability of some other flatfish species, notably yellow-fund sole, relates to water temperature. The authors examined the relationship of water temperature and catchability of arrowtooth flounder. However, at this time, the authors are uncertain as to how to interpret their findings. Therefore, this relationship was not included in the model.

Status determination

Arrowtooth flounder is an unexploited stock in the BSAI. Model projections indicate that this stock is neither overfished nor approaching an overfished condition.

Response to SSC comments

The AI surveys were added to the model.

Individual estimates were made for Kamchatka flounder, though the ABC and OFL values are reported for a combined arrowtooth-Kamchatka stock.

There are structural problems with this model that must be resolved before it is ready for Tier 1 status.

Ecosystem Considerations summary

Given the trend of increasing BSAI arrowtooth flounder stock size and in response to a request by the SSC, this chapter includes an enhanced ecosystem considerations section.

Arrowtooth flounder both eat pollock and are eaten by pollock. They prey on ages 0, 1, and 2, sizes not caught by trawl survey because the pollock are off bottom at those ages. Once pollock get to be >30 cm, outside range of being ATF food.

A 30-yr ecosystem simulation provided interesting results. Examining the effects of arrowtooth flounder on other species showed that a decrease of Arrowtooth flounder by 10% decreased Arrowtooth flounder adults and juveniles, slightly increased flathead sole and produced a negligible effect on pollock. However, it should be noted that those results were for a 10% increase. If Arrowtooth flounder increased by 50%, the result could be significantly different, as seen in the GOA simulation. Currently both Pacific cod and Arrowtooth flounder are negatively affecting pollock.

7. Northern Rock sole

Status and catch specifications (t) of rock sole in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2007 and 2008 are those recommended by the Plan Team. Catch data are current through 10/27/07.

Area	Year	Age 2+ Bio	OFL	ABC	TAC	Catch
BSAI	2006	1,490,000	150,000	126,000	41,500	36,452
	2007	1,670,000	200,000	198,000	55,000	37,013
	2008	1,880,000	304,000	301,000	n/a	n/a
	2009	n/a	379,000	375,000	n/a	n/a

Changes from previous assessment

The present assessment is a straightforward update of last year's assessment. Northern rock sole (*Lepidopsetta polyxystra*) is the dominant rock sole species in the Bering Sea. Only 2% of the commercial catch is estimated to be southern rock sole (*Lepidopsetta bilineata*). Therefore this assessment for 2008 is for northern rock sole only. No attempt was made to correct past catch or survey data by species.

Changes to input data in this analysis include addition of 2006 rock sole fishery age composition, 2006 northern rock sole survey age composition, and 2007 northern rock sole trawl survey biomass point estimate and standard error. Only Bering Sea survey data (no Aleutian Islands data, 3% of total rock sole) were used in calculations. Another change to the input data was inclusion of an estimate of retained and discarded portions of the 2006 rock sole catches. Analysis was conducted with fishery catch and discards through 8 September 2007.

Spawning biomass and stock status trends

The stock assessment model resulted in a biomass estimate of 1,880,000 t. This was an increase in the biomass estimate compared to the 2006 estimate of 1,670,000 t. The rock sole stock is expected to remain stable through 2009. However, good recruitment in 2001 through 2004 should increase the stock biomass at the beginning of the next decade.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that northern rock sole qualifies as a Tier 1 stock; therefore the 2008 assessment was calculated using Tier 1 methodology. In 2006, the SSC selected the 1978-2001 data set for the Tier 1 harvest recommendation. Increasing to a Tier 1 resulted in a large increase in ABC and OFC estimates over the 2006 assessment values.

The Plan Team agreed with the authors' recommendation (Model 1). Thus the Tier 1 2008 ABC harvest recommendation is 301,000 t ($F_{ABC} = 0.177$) and a 2008 OFL of 304,000 t ($F_{OFL} = F_{MSY} = 0.179$). Note that with this new Tier 1 assessment, there is only a 4,000 t difference between the ABC and OFL levels. This will require a more tightly managed fishery. The 2009 also used Tier 1 methodology. However, projections for 20 years into the future uses Tier 3.

M and q were estimated within the model yielded $M=0.15$ and $q=1.87$. This was very different from the previous estimates of $M=0.16$ and $q=1.52$. While experiments indicate herding, this estimate of q means that 40% of fish are herded into the net. The authors do not think this is a realistic estimate of herding. It is most likely biased because of the skewed age composition from low recruits in 1990s. Therefore, the q estimated in past assessments was used ($M=0.15$, $q=1.5$).

Status determination

This is a stable fishery that lightly exploits the stock because it is constrained by prohibited species catch limits and the BSAI optimum yield limit. Model projections indicate that this stock is neither overfished nor approaching an overfished condition. Usually the fishery only takes a small portion of the rock sole ABC, but there will be more room in cap this year because the pollock ABC is lower.

Response to SSC comments

The SSC requested continued management strategy evaluation (MSE) for Tier 1 management. The authors intend to continue this exploration for climate and productivity changes.

The authors will examine a split-sex model for the fall 2008 assessment.

8. Flathead Sole

Status and catch specifications (t) of flathead sole in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through November 10, 2007.

Area	Year	Age 3+ Bio.	OFL	ABC	TAC	Catch
BSAI	2006	636,000	71,800	59,800	19,500	17,947
	2007	875,000	95,300	79,200	30,000	19,477
	2008	820,000	86,000	71,700	n/a	n/a
	2009	814,000	83,700	69,700	n/a	n/a

Changes from previous assessment

The present assessment is a straightforward update of last year's assessment. The present assessment includes updated catch, survey biomass, length composition, and age composition data. Sex-specific length compositions from the 2006 and 2007 EBS trawl surveys were added and survey age compositions from previous years were recalculated. Normalization of age and length compositions was changed to being within-sex to summing over both sexes; this was found to aid model convergence.

Spawning biomass and stock status trends

According to the 2007 assessment, the age 3+ biomass decreased from 809,000 t in 2006 to 796,000 t in 2007, a 2% decrease. However, the trend between assessments represents a greater decrease, as the 2006 assessment estimate for 2007 was 875,000 t, so the decrease in 2007 estimated biomass estimated in 2006 vs. 2007 is 9%.

This year's survey biomass was 571,000t, a 12% decrease from 2006.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that reliable estimates of $B_{40\%}$ (140,000 t) $F_{40\%}$ (0.28) and $F_{35\%}$ (0.34) exist for this stock, thereby qualifying the stock for management under Tier 3. Given that the projected 2008 spawning biomass of 251,000 t exceeds $B_{40\%}$, the ABC and OFL recommendations for 2008 were calculated under sub-tier "a" of Tier 3. The Plan Team recommends setting F_{ABC} at the $F_{40\%}$ (0.28) level, which is the maximum permissible level under Tier 3a. Projected harvesting at the $F_{40\%}$ level gives a 2008 ABC of 71,700 t. The OFL was determined from the Tier 3a formula, where an $F_{35\%}$ value of 0.34 gives a 2008 OFL of 86,000 t. Model projections indicate that this stock is neither overfished nor approaching an overfished condition.

Additional Plan Team recommendations

In response to SSC and Plan Team comments, the author examined the distribution of Bering flounder with respect to the fishery. The northerly distribution of the species did not seem to overlap the spatial distribution of the fishery, although mismatch in seasonal timing of the survey versus the fishery means that this is not conclusive. The Plan Team recommended placing Bering flounder in the "other flatfish" category in future assessments.

Ecosystem Considerations

The assessment authors presented updated information on predators and prey of flathead sole, temperature-dependent habitat quality, and bycatch in the fishery. No specific concerns were noted by the author or the Plan Team.

9. Alaska plaice

Status and catch specifications (t) of Alaska plaice in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2007 and 2008 are those recommended by the Plan Team. Catch data are current through 10/27/07

Area	Year	Age 3 + Bio.	OFL	ABC	TAC	Catch
BSAI	2006	1,008,000	237,000	188,000	8,000	17,309
	2007	1,340,000	241,000	190,000	25,000	19,411
	2008	1,850,000	248,000	194,000	n/a	n/a
	2009	n/a	277,000	217,000	n/a	n/a

Changes from previous assessment

The present assessment is a straightforward update of last year's assessment. Input data were updated with 2007 catch data and inclusion of fishery catch through 8 September 2007. The 2007 trawl survey biomass estimate and standard error, and 2007 length composition of survey catch also were added to the model. The 2006 survey ages were read from otoliths and the 2006 survey age composition was added to the assessment. In response to a request from the SSC, length bins and the transition matrix were extended from 45 to 60 cm.

Spawning biomass and stock status trends

The female spawning biomass trend is similar to the overall biomass trend, It was slowly decreasing from 1985 to 2005 and has been increasing since then. The increase in total biomass is expected to continue. The shelf survey biomass has been fairly steady since the mid 1980s. Above-average recruitment year classes since 1998, with an exceptionally strong year class in 2003, will support the increasing trend in this stock.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

Reliable estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist for this stock, therefore qualifying it for management under Tier 3a of the BSAI Groundfish FMP. The updated point estimates are $B_{40\%} = 145,000$ t, $F_{ABC} = F_{40\%} = 0.59$, $F_{OFL} = F_{35\%} = 0.81$. These are high values for flatfishes, but these values are the consequence of Alaska plaice maturing before recruiting to the fishery. Given that the projected 2008 spawning biomass of 335,000 t exceeds $B_{40\%}$, the ABC and OFL recommendations for 2008 were calculated under sub-tier "a" of Tier 3. Projected harvesting at the $F_{40\%}$ level gives a 2008 ABC of 194,000 t. The OFL was determined from the Tier 3a formula, which gives a 2008 OFL of 248,000 t.

Status determination

Model projections indicate that this species is neither overfished nor approaching an overfished condition. There is not a targeted fishery for this species as there is no market. Alaska plaice is caught only as bycatch and is mostly discarded.

Response to SSC comments

The SSC asked the authors to extend the range of the length bins to mimic the dynamics of larger fish. The authors improved the modeling by extending the maximum length bin from 45 to 60 cm.

The authors will implement a split-sex model for the fall 2008 assessment.

10. Other flatfish complex

Status and catch specifications (t) of other flatfish in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2007 and 2008 are those recommended by the Plan Team. Catch data are current through 11/10/07.						
	Year	Total Bio.	OFL	ABC	TAC	Catch
BSAI	2006	121,000	24,200	18,100	3,500	3,157
	2007	149,000	28,500	21,400	10,000	5,838
	2008	149,000	28,800	21,600	n/a	n/a
	2009	n/a	28,800	21,600	n/a	n/a

Changes from previous assessment

With the removal of Alaska plaice from this category in 2002, the species currently collected in the “other flatfish” category in the Eastern Bering Sea survey are Arctic flounder, butter sole, curlfin sole, deepsea sole, Dover sole, English sole, longhead dab, Pacific sand dab, petrale sole, rex sole, roughscale sole, sand sole, slender sole, starry flounder, and Sakhalin sole. The species currently collected in the “other flatfish” category in the Aleutian Islands survey are Dover sole, rex sole, starry flounder, butter sole and English sole. Starry flounder, rex sole and butter sole comprise the majority of the species caught with a negligible amount of other species. Of those, starry flounder comprised 74 % in 2007, an increase in starry flounder in the shelf survey biomass from ~50 % in 2006. In 2007, 25 % was rex sole and butter sole.

The present assessment is a straightforward update of last year’s assessment. The assessment incorporates 2006 total catch and discard; catch through 8 September 2007 and 2007 EBS trawl survey information. As there was no AI survey in 2007, a linear regression was used to predict AI biomass for 2007. Together the 2006 EBS bottom trawl survey and AI prediction resulted in a biomass estimate of 149,000 t, the same value that resulted from the 2006 surveys.

Spawning biomass and stock status trends

Because this complex is managed under Tier 5, no models are available from which to predict future trends.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

With the removal of Alaska plaice from this category in 2002 the SSC reclassified “other flatfish” as a Tier 5 species complex with harvest recommendation calculated from estimates of biomass and natural mortality. Natural mortality values for rex and Dover sole in the GOA SAFE document and starry flounder on the west coast are used. For all other species, a natural mortality rate of 0.20 is assumed. Projected harvesting at the 0.75 *M* level ($F_{ABC} = 0.15$), gives a 2008 ABC of 21,600 t for the “other flatfish” species. The corresponding 2008 OFL (= 0.20) is 28,000 t.

Status determination

It is not possible to determine whether the “other flatfish” complex is overfished or approaching an overfished condition because it is Tier 5 and not managed under Tiers 1-3. Insufficient information about these species makes model analysis impossible.

This group of fisheries is usually closed prior to attainment of TAC because of the bycatch of Pacific halibut, a prohibited species. With the implementation of Amendment 80, higher TACs for other flatfishes are anticipated in the future.

Response to SSC comments

No progress was made on the assessment of bottom temperature effect on catchability of other flatfish species.

Estimated biomass of “other” flatfish species are available in Table 10.5

The catch numbers in the summary table were corrected. The numbers were lowered by subtracting flathead sole from the total catch reported by NMFS.

11. Pacific Ocean Perch (POP)

Status and catch specifications (t) of Pacific ocean perch. Biomass corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through 10/27/07.							
Year	Area	Age 3+ Bio	OFL	Subarea	ABC	TAC	Catch
2006	BSAI	385,000	17,600		14,800	12,600	12,856
				EBS	2,960	1,400	1,040
				Eastern AI	3,260	3,080	3,069
				Central AI	3,210	3,030	3,242
				Western AI	5,370	5,090	5,506
2007	BSAI	457,000	26,100		21,900	19,900	17,772
				EBS	4,170	2,160	811
				Eastern AI	4,980	4,970	5,116
				Central AI	5,060	5,050	4,423
				Western AI	7,730	7,720	7,421
2008	BSAI	453,000	25,700		21,700	n/a	n/a
				EBS	4,200	n/a	n/a
				Eastern AI	4,890	n/a	n/a
				Central AI	4,970	n/a	n/a
				Western AI	7,590	n/a	n/a
2009	BSAI	449,000	25,400		21,300	n/a	n/a
				EBS	4,140	n/a	n/a
				Eastern AI	4,820	n/a	n/a
				Central AI	4,900	n/a	n/a
				Western AI	7,490	n/a	n/a

Changes from previous assessment

Beginning in 2005, POP assessments are being conducted on a 2-year cycle. There is no new survey for 2007. Catch data were updated and the projection model was re-run using results from the 2006 assessment model as the starting point.

Spawning biomass and stock status trends

The estimated spawning biomass is projected to decline slightly from 155,000 t in 2007 to 153,000 t in 2008 and further to 150,000 t in 2009.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that reliable estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist for this stock, thereby qualifying Pacific ocean perch for management under Tier 3. The current estimates of $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ are 132,500 t, 0.059, and 0.070 respectively. There are reliable estimates of the 2007 spawning biomass (B), $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ and $B > B_{40\%}$ (153,000 t > 133,000 t). Therefore the POP reference fishing mortality is defined in Tier 3a. For this tier, F_{ABC} is constrained to be $\leq F_{40\%}$, and F_{OFL} is constrained to be equal to $F_{35\%}$. The ABC associated with the $F_{40\%}$ level of 0.059 is 21,700 t. Model projections indicate that this stock is neither overfished nor approaching an overfished condition. For 2008, the recommended ABC is 21,700 t, and the OFL is 25,700 t.

Area apportionment

The Team agrees with the author's recommendation that ABCs be set regionally based on the proportions in combined survey biomass as follows: BS = 4,200 t, Eastern Aleutians (Area 541) = 4,890 t, Central Aleutians (Area 542) = 4,970 t, Western Aleutians (Area 543) = 7,590 t. The OFL fishing mortality rate is computed under Tier 3a as 25,700 t, which is the author's and Plan Team's recommended OFL for the BSAI. The OFL for BSAI is not regionally apportioned.

12. Northern Rockfish

Status and catch specifications (t) of Northern rockfish. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch is reported through October 27, 2007.

Area	Year	Age 3+ Bio.	OFL	ABC	TAC	Catch
BSAI	2006	204,000	10,100	8,530	4,500	3,823
	2007	212,000	9,750	8,190	8,190	3,936
	2008	212,000	9,740	8,180	n/a	n/a
	2009	n/a	9,680	8,130	n/a	n/a

Changes from previous assessment

This was an "off-year" for a biennial assessment. This update was produced by updating the catch data and re-running the projection model using the results from the 2006 assessment model as a starting point.

Spawning biomass and stock status trends

Estimates of age 3+ biomass increased from 204,000 t in 2006 to 212,000 t in 2007 estimates (a 4% increase).

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has determined that this stock qualifies for management under Tier 3 due to the availability of reliable estimates for $B_{40\%}$ (52,000 t), $F_{40\%}$ (0.045), and $F_{35\%}$ (0.053). Because the female spawning biomass of 73,500 t is greater than $B_{40\%}$, sub-tier "a" would be applicable, with $F_{ABC} = F_{40\%}$ and $F_{OFL} = F_{35\%}$. Under Tier 3a, the maximum permissible ABC is 8,180 t, which is the authors' and Plan Team's recommendation for the 2008 ABC. Under Tier 3a, the 2008 OFL is 9,740 t for the Bering Sea/Aleutian Islands combined. The Plan Team continues to recommend setting a combined BSAI OFL and ABC. As the TAC has routinely been lower than the ABC, the TAC of the previous year was assumed as the 2008 catch, in order to make projections to 2009.

Model projections indicate that this stock is neither overfished nor approaching an overfished condition.

13. Shortraker/Rougheye Rockfish

Status and catch specifications (t) of shortraker/rougheye in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through October 27, 2007.							
Species	Area	Year	Survey Bio.	OFL	ABC	TAC	Catch
shortraker	BSAI	2006	20,500	774	580	580	208
		2007	18,900	564	424	424	318
		2008	18,900	564	424	n/a	n/a
		2009	n/a	564	424	n/a	n/a
rougheye	BSAI	2006	11,200	299	224	224	203
		2007	10,800	269	202	202	163
		2008	10,800	269	202	n/a	n/a
		2009	n/a	269	202	n/a	n/a

Changes from previous assessment

This was an “off-year” for a biennial assessment. This update was produced by updating the catch data and re-running the projection model using the results from the 2006 assessment model as a starting point.

Spawning biomass and stock status trends

The estimate of 2007 shortraker survey biomass is 18,900 t, down 8% from 2006. The estimate of 2007 rougheye survey biomass is 10,800 t, down 4% from 2006.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The SSC has previously determined that reliable estimates of biomass and natural mortality exist for shortraker and rougheye rockfish, qualifying the species for management under Tier 5. At the present time, the Plan Team recommends that the SSC retain Tier 5 management for these stocks. The Plan Team recommends setting F_{ABC} at the maximum permissible level under Tier 5, which is 75% of M . Accepted values for M for these stocks are 0.025 for rougheye rockfish and 0.030 for shortraker rockfish, resulting in F_{ABC} values of 0.019 and 0.023 for rougheye and shortraker, respectively.

The biomass estimates for 2007 are 18,900 t for shortraker rockfish and 10,800 t for rougheye rockfish, leading to BSAI OFLs of 564 t for shortraker and 269 t for rougheye, and ABCs of 424 t for shortraker and 202 t for rougheye. It is not possible to determine whether these species are overfished or whether they are approaching an overfished condition because they are managed under Tier 5.

14. Other Rockfish Complex

Status and catch specifications (t) of other rockfish (primarily thornyheads) in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2007 and 2008 are those recommended by the Plan Team. Catch data are current through 10/27/07. Values for 2009 reflect the removal of dark rockfish from the other rockfish complex per Amendment 77 to the BSAI FMP.

Area	Year	Survey Biomass	OFL	ABC	TAC	Catch
BSAI	2006	26,600	1,330			579
	2007	36,700	1,330	999	999	635
	2008	36,700	1,330	999	n/a	n/a
	2009	n/a	1,290	968	n/a	n/a
EBS	2006	15,400	n/a	414	414	157
	2007	18,100	n/a	414	414	205
	2008	18,100	n/a	414	414	n/a
	2009	n/a	n/a	414	414	n/a
AI	2006	11,200	n/a	585	585	422
	2007	18,600	n/a	585	585	430
	2008	18,600	n/a	585	585	n/a
	2009	n/a	n/a	554	554	n/a

Changes from previous assessment

This was an 'off-year' for a biennial assessment. Catches in 2006 have been revised and the 2007 catch through 10/27/07 has been included. The BSAI "other rockfish" assessment considers the 8 species that have been caught at least once during AFSC research surveys or appeared in more than 1% of observed fishery hauls between 1990 and 2001. Separate estimates of natural mortality (M) and biomass for shortspine thornyheads (SST; $M=0.03$), the most common species in the other rockfish complex, and the remaining species ($M=0.09$ based on dusky rockfish) in the complex were used. For 2009, dark rockfish will be removed from the other rockfish complex per Amendment 77.

Spawning biomass and stock status trends

Trends in spawning biomass are unknown. Stock biomass, as measured by trawl surveys of the EBS slope and in the Aleutian Islands, has increased since 1997.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The Plan Team agrees with the approach recommended by the author of setting F_{ABC} at the maximum allowable under Tier 5 ($F_{ABC} = 0.75 * M$). Multiplying these rates by 0.75 and the best estimates of SST and other "other rockfish" biomass yields 2008 ABCs of 414 t in the EBS and 585 t in the AI. Plan Team recommends that OFL be set for the entire BSAI area, which under Tier 5 is calculated by multiplying the best estimates of total biomass for the area by the separate Ms and adding the results, which yields an OFL of 1,330 t for 2008.

For 2009, dark rockfish will be removed from the other rockfish complex. The 2009 ABC in the EBS is 414 t, while in the AI is 554 t. The 2009 OFL in the BSAI = 1,290 t. The complex is not currently overfished, nor is it approaching an overfished condition.

15. Atka mackerel

Status and catch specifications (t) of Atka mackerel in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through 10/27/07.							
Area	Year	Sub-Area	Age 3+ Biomass	OFL	ABC	TAC	Catch
BSAI	2006		446,000	130,000	110,000	63,000	61,878
BSAI	2007		364,000	86,900	74,000	63,000	56,620
BSAI	2008		323,000	71,400	60,700	n/a	n/a
		541&EBS			19,500	n/a	n/a
		542			24,300	n/a	n/a
		543			16,900	n/a	n/a
BSAI	2009		n/a	50,600	47,500	n/a	n/a
		541&EBS			15,300	n/a	n/a
		542			19,000	n/a	n/a
		543			13,200	n/a	n/a

Changes from previous assessment

There were no changes in the assessment model methodology from that reviewed in 2006.

New input data were included and updated in the 2007 assessment.

Spawning biomass and stock status trends

The projected female spawning biomass for 2008 under an $F_{40\%}$ harvest strategy is estimated at 110,200 t which is 47% of unfished spawning biomass and above $B_{40\%}$ (94,100 t). The 2008 estimate of spawning biomass is down about 15% from last year's estimate for 2007. These results are consistent with the recent trend in survey biomass (18% decline in 2006 relative to 2004).

The projected age 3+ biomass at the beginning of 2008 is estimated at 323,400 t, down about 11% from last year's estimate for 2007. There is less of a decrease in projected biomass in this year's assessment relative to last year's projections. The addition of the 2006 fishery and survey age compositions changed the estimated magnitude of the 1999, 2003, and 2004 year-classes by 8, 46, and 68%, respectively; the magnitude of the 2001 and 2002 year-classes decreased 13, and 34%, respectively, relative to last year's assessment. However, year-classes spawned in 2002 through 2005 are all estimated to be below average in size. These follow the four above average 1998-2001 year-classes, which leads to short-term population declines.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

The projected female spawning biomass under an $F_{40\%}$ harvest strategy is estimated to be 47% of unfished spawning biomass in 2008 and above $B_{40\%}$, thereby placing BSAI Atka mackerel in Tier 3a. The projected 2008 yield (ABC) at $F_{40\%} = 0.331$ is 60,700 t, down about 18% from last year's estimate for 2007. The projected 2008 overfishing level at $F_{35\%}$ ($F = 0.40$) is 71,400 t, down about 18% from last year's estimate for 2007. The species is not currently overfished, nor is it approaching an overfished condition.

Atka mackerel female spawning biomass in 2009 (89,900 t) is projected to fall below $B_{40\%}$, putting the fishery in Tier 3b. The projected 2009 yield (ABC) at $F_{40\%} = 0.32$ is 47,500 t; the projected 2009 overfishing level at $F_{35\%}$ ($F = 0.36$) is 50,600 t. The population is projected to remain below $B_{40\%}$ from 2009-2012.

Ecosystem Considerations

Food habits data (from analysis of scats) from the Aleutian Islands indicate that Atka mackerel is the most common prey item of the endangered western Steller sea lion throughout the year. Analyses of historic fishery CPUE revealed that the fishery may create temporary localized depletions of Atka mackerel, and fishery harvest rates in localized areas may have been high enough to affect prey availability of Steller sea lions.

Bottom contact fisheries could have direct negative impacts on Atka mackerel by destroying egg nests and/or removing the males that are guarding nests. When trawl exclusion zones near Steller sea lion rookeries were implemented beginning in 1992, it was thought that these eliminated much of the overlap between bottom trawl fisheries and Atka mackerel nesting areas. However, nesting sites may be widespread across the continental shelf and found over a much broader depth range than just in nearshore areas. The use of bottom contact fishing gear, such as bottom trawls, pot gear, and longline gear, utilized in July to January could, therefore, still potentially affect Atka mackerel nesting areas, despite trawl closures in nearshore areas around Steller sea lion rookeries.

Area apportionment

Amendment 28 of the Bering Sea/Aleutian Islands Fishery Management Plan divided the Aleutian subarea into 3 districts at 177° E and 177° W longitude, providing the mechanism to apportion the Aleutian Atka mackerel TACs. The Council used a 4-survey (2000, 2002, 2004, and 2006) weighted average to apportion the 2007 ABC, and the authors recommend using the same method to apportion the 2008 and 2009 ABCs. The recommended ABC apportionment by subarea for both 2008 and 2009 is 32.2% for Area 541, 40.0% for 542, and 27.8% for Area 543.

16 – 20. Squid and Other Species Complex

Status and catch specifications (t) of squid and other species in recent years. Biomass for each year corresponds to the projection given in the SAFE report issued in the preceding year. The OFL and ABC for 2008 and 2009 are those recommended by the Plan Team. Catch data are current through 10/27/07.

Squid	Area	Year	Biomass	OFL	ABC	TAC	Catch
	BSAI	2006	n/a	2,620	1,970	1,275	1,416
		2007	n/a	2,620	1,970	1,970	1,193
		2008	n/a	2,620	1,970	n/a	n/a
		2009	n/a	2,620	1,970	n/a	n/a
Other species	Area	Year	Biomass	OFL	ABC	TAC	Catch
BSAI Council	Approved	2006	724,000	93,800	70,400	29,000	26,562
		2007	734,000	91,700	68,800	37,355	26,467
Plan Team	Recommend.:	2008					
	Sharks		18,100	617	463	n/a	n/a
	Skates		574,801	37,200	31,300	n/a	n/a
	Sculpins		228,995	53,100	39,800	n/a	n/a
	Octopus		n/a	324	243	n/a	n/a
	Total		n/a	91,300	71,800	n/a	n/a
		2009					
	Sharks		18,100	617	463	n/a	n/a
	Skates		567,134	36,800	30,900	n/a	n/a
	Sculpins		228,995	53,100	39,800	n/a	n/a
	Octopus		n/a	324	243	n/a	n/a
	Total		n/a	91,300	71,800	n/a	n/a

Changes from previous assessment

There were no changes in assessment methods for the squid and other species complex, except that the Stock Synthesis 2 assessment framework was used for Alaska skate within the skate component. Catch and survey data were updated in these assessments.

Spawning biomass and stock status trends

No spawning biomass trend is available.

Tier determination/Plan Team discussion and resulting ABCs and OFLs

Squid - Using Tier 6 criteria, the recommended ABC for BSAI squid for 2008-2009 is calculated as 0.75 times the average catch from 1978-1995, or 1,970 t; the recommended overfishing level for squid in 2008-2009 is calculated as the average catch from 1978-1995, or 2,620 t. The groundfish surveys do not provide a reliable biomass estimate. The authors reviewed some life history information and indicated that the predominant species in this group may experience multiple cohorts within a single year within the context of a discussion of whether this species group may more appropriately reside in the forage fish category.

Sharks – The Plan team recommended Tier 6 ABC (463 t) and OFL (617 t) for 2008 and 2009. The 2007 SAFE has been updated with life history information and length frequency information. In a departure from the November 2006 meeting the Plan Team agreed with the SSC that the survey information was not reliable enough to promote the Tier 5 assessment.

Sculpin – In contrast to the Tier 5 assessment in 2006, where a single M (0.19) and two biomass estimates (one for the EBS and one for the AI) were used, the 2007 assessment applies distinct M estimates from the literature and species or species group biomass estimates for the five most predominate sculpin species/species groups. The assessment authors noted that species composition differs among the BS shelf, BS slope, and the AI. The survey biomass is stable. The ratio of fishing mortality to total mortality is low. The Plan Team recommended the alternative approach, and the ABC and OFL for 2008 – 2009 to be 39,800 t and 53,100 t, respectively.

Octopus – The author thinks the octopus biomass estimate is not reliable and therefore does not support a Tier 5 assessment. The author also noted the Tier 6 estimate based on average catch likely underestimates the population and would unnecessarily constrain primarily the pot fishery for Pacific cod if octopus were managed as a single species group or if it were broken out into two groups external from the ‘other species’ category. The author recommended a revised Tier 6 based on recent maximum catch. On going research includes species identification discard mortality estimates. The Plan Team recommended Tier 6 average for ABC 243 t and OFL 324 t calculation.

Skate – The 2007 assessment differs from the 2006 estimate. The Alaska skate component is estimated within Tier 3 rather than Tier 5. The ‘other skate’ component of the estimate continues to be estimated within Tier 5. Plan Team agreed with the author’s recommendation to assess Alaska skate within Tier 3 and other skate stay within Tier 5. The authors presented several Tier 3 Alaska skate models based on historic biomass and catch estimates. Their recommended Tier 3 ‘base’ model utilized data that began in 1992. The team looks forward to the author’s review of a discrepancy between increasing survey biomass estimate in recent years relative to the model projections of declining biomass.

Appendix A: Pacific Halibut Discard Mortality Rates

Halibut discard mortality rates (DMRs) are set by the Council on a 3-year cycle for non-CDQ fisheries based on an average of the past 10 years and annually for CDQ fisheries based on available data. Rates for non-CDQ fisheries for 2008 are included in rates previously adopted by the Council for use in 2007-2009. International Pacific Halibut Commission staff recommendations for DMRs for the BSAI CDQ fisheries for 2008 were reviewed by the BSAI Plan Team during its September meeting. The Team endorsed the IPHC recommendations for revised halibut DMRs for the CDQ fisheries. The rates were adopted by the Council in October 2007 and are shown in the summary table below. It is anticipated that the rates for CDQ fisheries will likely be set on a 3-year cycle when the next 3-year cycle commences for the non-CDQ fisheries in 2009 for 2010-2012.

CDQ Fisheries	
Gear/Target	Recommended DMR
<i>Trawl</i>	
Atka mackerel	85
Bottom pollock	86
Rockfish	82
Flathead sole	87
Pelagic pollock	90
Rock sole	86
Yellowfin sole	86
<i>Pot</i>	
Sablefish	34
<i>Longline</i>	
Pacific cod	10
Turbot	4

Table 1. Bering Sea Aleutian Islands Groundfish Plan Team OFL, ABC, and TAC Recommendations for the 2008-2009 Fisheries.

Species	Area	2007				2008			2009		
		OFL	ABC	TAC	Catch	OFL	ABC	TAC	OFL	ABC	TAC
Pollock	EBS	1,640,000	1,394,000	1,394,000	1,350,000	1,440,000	1,000,000		1,320,000	1,000,000	
	Aleutian Islands	54,500	44,500	19,000	2,488	34,000	28,200		26,100	22,700	
	Bogoslof District	48,000	5,220	10	0	58,400	7,970		58,400	7,970	
Pacific cod	BSAI	207,000	176,000	171,000	172,000	176,000	150,000		190,000	162,000	
Sablefish	BS	3,520	2,980	2,980	1,090	3,380	2,860		2,910	2,610	
	AI	3,320	2,810	2,810	1,080	2,890	2,440		2,510	2,230	
Yellowfin sole	BSAI	240,000	225,000	136,000	119,332	265,000	248,000		296,000	276,000	
Greenland turbot	Total	15,600	2,440	2,440	1,946	15,600	2,540		16,000	2,540	
	BS		1,680	1,680	1,435		1,750			1,750	
	AI		760	760	511		787			787	
Arrowtooth flounder	BSAI	193,000	158,000	20,000	11,700	297,000	244,000		300,000	246,000	
Northern rock sole	BSAI	200,000	198,000	55,000	37,013	304,000	301,000		379,000	375,000	
Flathead sole	BSAI	95,300	79,200	30,000	19,500	86,000	71,700		83,700	69,700	
Alaska plaice	BSAI	241,000	190,000	25,000	19,411	248,000	194,000		277,000	217,000	
Other flatfish	BSAI	28,500	21,400	10,000	25,176	28,800	21,600		28,800	21,600	
Pacific Ocean perch	BSAI	26,100	21,900	19,900	17,800	25,700	21,700		25,400	21,300	
	BS		4,160	2,160			4,200			4,130	
	AI total		15,390	17,740			17,590			17,200	
	WAI		5,370	7,720			7,590			7,490	
	CAI		5,050	5,050			4,970			4,900	
	EAI		4,970	4,970			4,890			4,820	
Northern rockfish	BSAI	9,750	8,190	8,190	3,940	9,740	8,180		9,680	8,130	
Shorthead	BSAI	564	424	424	318	564	424		564	424	
Rougheye	BSAI	269	202	202	163	269	202		269	202	
Other rockfish	BSAI	1,330	999	999	635	1,330	999		1,290	968	
	BS		414	414			414			414	
	AI		585	585			585			554	
Atka mackerel	Total	86,900	74,000	63,000	56,620	71,400	60,700		50,600	47,500	
	WAI		20,600	9,600			16,900			13,200	
	CAI		29,600	29,600			24,300			19,000	
	EAI/BS		23,800	23,800			19,500			15,300	
Squid	BSAI	2,620	1,970	1,970	1,190	2,620	1,970		2,620	1,970	
Other species	BSAI	91,700	68,800	37,400	26,500	91,200	71,800		91,200	71,800	
	Sharks					617	463		617	463	
	Skates					37,200	31,300		36,800	30,900	
	Sculpins					53,100	39,800		53,100	39,800	
	Octopus					324	243		324	243	
Total	BSAI	3,188,973	2,676,035	2,000,000	1,867,902	3,161,893	2,440,285		3,162,043	2,557,644	

Table 2. Groundfish catches (metric tons) in the eastern Bering Sea, 1954-2007.

Year	Pacific Pollock	Sable Cod	Yellowfin Fish	Greenland Sole	Arrowtooth Turbot	Rock Flounder/a	Other Sole/c	Alaska Flatfish	Pacific Plaice	Northern Perch	Pacific Ocean Rockfish	Northern Rockfish	Shortraker Rockfish	Rougheyc Rockfish	Other Rockfish	Atka Mackerel	Other Squid	Total (All Species)		
1954				12,562														12,562		
1955				14,690														14,690		
1956				24,697														24,697		
1957				24,145														24,145		
1958		6,924	171	6	44,153												147	51,401		
1959	32,793	2,864	289	185,321													380	221,647		
1960			1,861	456,103	36,843						6,100							500,907		
1961			15,627	553,742	57,348						47,000							673,717		
1962			25,989	420,703	58,226						19,900							524,818		
1963			13,706	85,810	31,565			35,643			24,500							191,224		
1964	174,792	13,408	3,545	111,177	33,729						30,604						736	393,891		
1965	230,551	14,719	4,838	53,810	9,747			11,686			16,800						2,218	344,369		
1966	261,678	18,200	9,505	102,353	13,042			24,864			20,200						2,239	452,081		
1967	550,362	32,064	11,698	162,228	23,869						19,600						4,378	836,308		
1968	702,181	57,902	4,374	84,189	35,232						31,500						22,058	967,083		
1969	862,789	50,351	16,009	167,134	36,029						14,500						10,459	1,192,020		
1970	1,256,565	70,094	11,737	133,079	19,691	12,598		64,690			9,900						15,295	1,593,649		
1971	1,743,763	43,054	15,106	160,399	40,464	18,792		92,452			9,800						13,496	2,137,326		
1972	1,874,534	42,905	12,758	47,856	64,510	13,123		76,813			5,700						10,893	2,149,092		
1973	1,758,919	53,386	5,957	78,240	55,280	9,217		43,919			3,700						55,826	2,064,444		
1974	1,588,390	62,462	4,258	42,235	69,654	21,473		37,357			14,000						60,263	1,900,092		
1975	1,356,736	51,551	2,766	64,690	64,819	20,832		20,393			8,600						54,845	1,645,232		
1976	1,177,822	50,481	2,923	56,221	60,523	17,806		21,746			14,900						26,143	1,428,565		
1977	978,370	33,335	2,718	58,373	27,708	9,454		14,393			2,654					311	4,926	1,168,144		
1978	979,431	42,543	1,192	138,433	37,423	8,358		21,040			2,221					2,614	831	6,886	1,302,509	
1979	913,881	33,761	1,376	99,017	34,998	7,921		19,724			1,723					2,108	1,985	4,286	1,159,547	
1980	958,279	45,861	2,206	87,391	48,856	13,761		20,406			1,097					459	4,955	4,040	34,633	1,221,944
1981	973,505	51,996	2,604	97,301	52,921	13,473		23,428			1,222					356	3,027	4,182	35,651	1,259,666
1982	955,964	55,040	3,184	95,712	45,805	9,103		23,809			224					276	328	3,838	18,200	1,211,483
1983	982,363	83,212	2,695	108,385	43,443	10,216		30,454			221					220	141	3,470	15,465	1,280,285
1984	1,098,783	110,944	2,329	159,526	21,317	7,980		44,286			1,569					176	57	2,824	8,508	1,458,299
1985	1,179,759	132,736	2,348	227,107	14,698	7,288		71,179			784					92	4	1,611	11,503	1,649,109
1986	1,188,449	130,555	3,518	208,597	7,710	6,761		76,328			560					102	12	848	10,471	1,633,911
1987	1,237,597	144,539	4,178	181,429	6,533	4,380		50,372			930					474	12	108	8,569	1,639,121
1988	1,228,000	192,726	3,193	223,156	6,064	5,477		137,418			1,047					341	428	414	12,206	1,810,470
1989	1,230,000	164,800	1,252	153,165	4,061	3,024		63,452			2,017					192	3,126	300	4,993	1,630,382
1990	1,353,000	162,927	2,329	80,584	7,267	2,773		22,568			5,639					384	480	460	5,698	1,644,109
1991	1,268,360	165,444	1,128	94,755	3,704	12,748	46,681	30,401			4,744					396	2,265	544	16,285	1,647,455
1992	1,384,376	163,240	558	146,942	1,875	11,080	51,720	34,757			3,309					675	2,610	819	29,993	1,831,954
1993	1,301,574	133,156	669	105,809	6,330	7,950	63,942	28,812			3,763					190	201	597	21,413	1,674,406
1994	1,362,694	174,151	699	144,544	7,211	13,043	60,276	29,720			1,907					261	190	502	23,430	1,818,628
1995	1,264,578	228,496	929	124,746	5,855	8,282	54,672	34,861			1,210					629	340	364	20,928	1,745,890
1996	1,189,296	209,201	629	129,509	4,699	13,280	46,775	35,390			2,635					364	780	1,080	19,717	1,653,355
1997	1,115,268	209,475	547	166,681	6,589	8,580	67,249	42,374			1,060					161	171	1,438	20,997	1,640,590
1998	1,101,428	160,681	586	101,310	8,303	14,985	33,221	39,940			1,134					203	901	891	23,156	1,486,739
1999	889,589	134,647	646	67,307	5,205	9,827	39,934	33,042			609					135	2,008	393	17,045	1,200,387
2000	1,132,736	151,372	742	84,057	5,888	12,071	49,186	36,813			704					239	239	375	23,098	1,497,520
2001	1,387,452	142,452	863	63,563	4,252	12,836	28,949	27,693			1,148					296	264	1,761	23,148	1,694,677
2002	1,481,815	166,552	1,143	74,956	3,150	10,821	40,700	30,229			858					401	572	1,334	26,639	1,839,170
2003	1,492,039	180,592	1,039	81,050	2,565	13,667	36,375	17,507	10,118		1,391					336	6,362	1,246	26,986	1,871,273
2004	1,480,543	183,283	1,038	75,501	1,805	17,327	47,862	22,109	7,888			731	116	119	24	318	7,157	1,000	27,493	1,874,315
2005	1,483,274	182,797	1,075	94,382	2,120	13,404	36,814	20,640	11,194			879	112	108	12	178	3,538	1,168	28,041	1,879,735
2006	1,486,413	167,725	1,055	99,064	1,440	11,852	35,874	21,029	17,309			1,039	246	46	7	157	3,170	1,401	24,627	1,872,455
2007/d	1,350,530	138,456	1,088	119,329	1,435	10,902	36,261	25,117	19,411			811	69	109	9	205	2,961	1,182	24,516	1,732,391

a/ Arrowtooth flounder included in Greenland turbot catch statistics, 1960-69. c/ Rock sole prior to 1991 is included in other flatfish catch statistics.

b/ Includes POP shortraker, rougheyc, northern and sharpchin.

d/ Data through October 27, 2007.

Note: Numbers don't include fish taken for research.

Table 3. Groundfish catches (metric tons) in the Aleutian Islands, 1962-2007 (no catches were recorded from this region prior to 1962).

Year	Pacific Pollock	Sable Cod	Yellowfin Fish	Greenland Sole	Greenland Turbot	Arrowtooth Flounder/a	Rock Sole/c	Other Flatfish	Alaska Plaice	Pacific Ocean Perch	Pacific Ocean Perch	Northern Rockfish	Shortraker Rockfish	Roughye Rockfish	Other Rockfish	Atkn Mackerel	Squid	Other Species	Total (All Species)
1962										200									200
1963					7					20,800									21,471
1964		241	1,541		504					90,300								66	92,652
1965		451	1,249		300					109,100								768	111,868
1966		154	1,341		63					85,900								131	87,589
1967		293	1,652		394					55,900								8,542	66,781
1968		289	1,673		213					44,900								8,948	56,023
1969		220	1,673		228					38,800								3,088	44,009
1970		283	1,248		285	274				66,900						949		10,671	80,610
1971		2,078	2,936		1,750	581				21,800								2,973	32,118
1972		435	3,531		12,874	1,323				33,200						5,907		22,447	79,717
1973		977	2,902		8,666	3,705				11,800						1,712		4,244	34,006
1974		1,379	2,477		8,788	3,195				22,400						1,377		9,724	49,340
1975		2,838	1,747		2,970	784				16,600						13,326		8,288	46,553
1976		4,190	1,659		2,067	1,370				14,000						13,126		7,053	43,465
1977	7,625	3,262	1,897		2,453	2,035				8,080					3,043	20,975	1,808	16,170	67,348
1978	6,282	3,295	821		4,766	1,782				5,286					921	23,418	2,085	12,436	61,092
1979	9,504	5,593	782		6,411	6,436				5,487					4,517	21,279	2,252	12,934	75,195
1980	58,156	5,788	274		3,697	4,603				4,700					420	15,533	2,332	13,028	108,531
1981	55,516	10,462	533		4,400	3,640				3,622					328	16,661	1,763	7,274	104,199
1982	57,978	1,526	955		6,317	2,415				1,014					2,114	19,546	1,201	5,167	98,233
1983	59,026	9,955	673		4,115	3,753				280					1,045	11,585	510	3,675	94,617
1984	81,834	22,216	999		1,803	1,472				631					56	35,998	343	1,670	147,022
1985	58,730	12,690	1,448		33	87				308					99	37,856	9	2,050	113,310
1986	46,641	10,332	3,028		2,154	142				286					169	31,978	20	1,509	96,259
1987	28,720	13,207	3,834		3,066	159				1,004					147	30,049	23	1,155	81,364
1988	43,000	5,165	3,415		1,044	406				1,979					278	21,656	3	437	77,383
1989	156,000	4,118	3,248		4,761	198				2,706					481	14,868	6	108	186,494
1990	73,000	8,081	2,116		2,353	1,459				14,650					864	21,725	11	627	124,886
1991	78,104	6,714	2,071	1,380	3,174	938		88		2,545					549	22,258	30	91	117,942
1992	54,036	42,889	1,546	4	895	900	236	68		10,277					3,689	46,831	61	3,081	164,513
1993	57,184	34,234	2,078	0	2,138	1,348	318	59		13,375					495	65,805	85	2,540	179,659
1994	58,708	22,421	1,771	0	3,168	1,334	308	55		16,959					301	69,401	86	1,102	175,614
1995	64,925	16,534	1,119	6	2,338	1,001	356	47		14,734					220	81,214	95	1,273	183,862
1996	28,933	31,389	720	654	1,677	1,330	371	61		20,443					278	103,087	87	1,720	190,750
1997	26,872	25,166	779	234	1,077	1,071	271	39		15,687					307	65,668	323	1,555	139,049
1998	23,821	34,964	595	5	821	694	446	54		13,729					385	56,195	25	2,448	134,182
1999	965	27,714	565	13	422	746	577	53		17,619					630	51,636	9	1,633	102,582
2000	1,244	39,684	1,048	13	1,086	1,157	480	113		14,893					601	46,990	8	3,010	110,327
2001	824	34,207	1,074	15	1,060	1,220	526	97		15,587					610	61,296	5	4,029	120,550
2002	1,177	30,801	1,118	29	485	1,032	1,165	150		14,996					551	44,722	10	1,980	98,216
2003	1,653	32,193	1,009	0	965	913	964	76		18,765					401	52,988	36	1,326	111,289
2004	1,158	28,869	955	9	414	806	818	71	0		11,165	4,567	121	184	337	53,405	14	1,851	104,744
2005	1,621	22,627	1,475	2	439	829	548	59	0		9,548	3,852	61	78	286	58,474	17	1,401	101,315
2006	1,735	24,181	1,130	4	525	1,450	578	75	0		11,817	3,577	162	196	422	58,708	15	1,935	106,510
2007d	2,488	33,951	1,078	3	511	799	752	59	0		16,961	3,867	209	154	430	53,659	11	1,951	116,883

a/ Arrowtooth flounder included in Greenland turbot catch statistics, 1964-69.

b/ Includes POP shortraker, roughycy, northern and sharpchin rockfish until 2004.

c/ Rocksole prior to 1991 is included in other flatfish catch statistics.

d/ Data through October 27, 2007.

Note: Numbers don't include fish taken for research.

Table 4. Groundfish catches (metric tons) in the Bering Sea and Aleutian Islands, 1954-2007.

Year	Pacific			Sable	Yellowfin	Greenland	Arrowtooth	Rock	Other	Alaska	Pacific Ocean		Pacific	Northern	Shortraker	Rougheye	Other	Alaska	Other	Total	
	Pollock	Cod	Fish	Sole	Turbot	Flounder/a	Sole/c	Flatfish	Plaice	Perch	Complex/b	Ocean	Perch	Rockfish	Rockfish	Rockfish	Rockfish	Mackerel	Squid	Species	(All Species)
1954	0	0	0	12,562	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12,562
1955	0	0	0	14,690	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	14,690
1956	0	0	0	24,697	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	24,697
1957	0	0	0	24,145	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	24,145
1958	6,924	171	6	44,153	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	147	51,401
1959	32,793	2,864	289	185,321	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	380	221,647
1960	0	0	1,861	456,103	36,843	0	0	0	0	0	6,100	0	0	0	0	0	0	0	0	0	500,907
1961	0	0	15,627	553,742	57,348	0	0	0	0	0	47,000	0	0	0	0	0	0	0	0	0	673,717
1962	0	0	25,989	420,703	58,226	0	0	0	0	0	20,100	0	0	0	0	0	0	0	0	0	525,018
1963	0	0	14,370	85,810	31,572	0	0	35,643	0	0	45,300	0	0	0	0	0	0	0	0	0	212,695
1964	174,792	13,649	5,086	111,177	34,233	0	0	30,604	0	0	116,200	0	0	0	0	0	0	0	0	802	486,543
1965	230,551	15,170	6,087	53,810	10,047	0	0	11,686	0	0	125,900	0	0	0	0	0	0	0	0	2,986	456,237
1966	261,678	18,354	10,846	102,353	13,105	0	0	24,864	0	0	106,100	0	0	0	0	0	0	0	0	2,370	539,670
1967	550,362	32,357	13,350	162,228	24,263	0	0	32,109	0	0	75,500	0	0	0	0	0	0	0	0	12,920	903,089
1968	702,181	58,191	6,047	84,189	35,445	0	0	29,647	0	0	76,400	0	0	0	0	0	0	0	0	31,006	1,023,106
1969	862,789	50,571	17,682	167,134	36,257	0	0	34,749	0	0	53,300	0	0	0	0	0	0	0	0	13,547	1,236,029
1970	1,256,565	70,377	12,985	133,079	19,976	12,872	0	64,690	0	0	76,800	0	0	0	0	0	0	949	0	25,966	1,674,259
1971	1,743,763	45,132	18,042	160,399	42,214	19,373	0	92,452	0	0	31,600	0	0	0	0	0	0	0	0	16,469	2,169,444
1972	1,874,534	43,340	16,289	47,856	77,384	14,446	0	76,813	0	0	38,900	0	0	0	0	0	0	5,907	0	33,340	2,228,809
1973	1,758,919	54,363	8,859	78,240	63,946	12,922	0	43,919	0	0	15,500	0	0	0	0	0	0	1,712	0	60,070	2,098,450
1974	1,588,390	63,841	6,735	42,235	78,442	24,668	0	37,357	0	0	36,400	0	0	0	0	0	0	1,377	0	69,987	1,949,432
1975	1,356,736	54,389	4,513	64,690	67,789	21,616	0	20,393	0	0	25,200	0	0	0	0	0	0	13,326	0	63,133	1,691,785
1976	1,177,822	54,671	4,582	56,221	62,590	19,176	0	21,746	0	0	28,900	0	0	0	0	0	0	13,126	0	33,196	1,472,030
1977	985,995	36,597	4,615	58,373	30,161	11,489	0	14,393	0	0	10,734	0	0	0	0	0	3,354	20,975	6,734	52,072	1,235,492
1978	985,713	45,838	2,013	138,433	42,189	10,140	0	21,040	0	0	7,507	0	0	0	0	0	3,535	24,249	8,971	73,973	1,363,601
1979	923,385	39,354	2,158	99,017	41,409	14,357	0	19,724	0	0	7,210	0	0	0	0	0	6,625	23,264	6,538	51,701	1,234,742
1980	1,016,435	51,649	2,480	87,391	52,553	18,364	0	20,406	0	0	5,797	0	0	0	0	0	879	20,488	6,372	47,661	1,330,475
1981	1,029,021	62,458	3,137	97,301	57,321	17,113	0	23,428	0	0	4,844	0	0	0	0	0	684	19,688	5,945	42,925	1,363,865
1982	1,013,942	56,566	4,139	95,712	52,122	11,518	0	23,809	0	0	1,238	0	0	0	0	0	2,390	19,874	5,039	23,367	1,309,716
1983	1,041,389	93,167	3,368	108,385	47,558	13,969	0	30,454	0	0	501	0	0	0	0	0	1,265	11,726	3,980	19,140	1,374,902
1984	1,180,617	133,160	3,328	159,526	23,120	9,452	0	44,286	0	0	2,200	0	0	0	0	0	232	36,055	3,167	10,178	1,605,321
1985	1,238,489	145,426	3,796	227,107	14,731	7,375	0	71,179	0	0	1,092	0	0	0	0	0	191	37,860	1,620	13,553	1,762,419
1986	1,235,090	140,887	6,546	208,597	9,864	6,903	0	76,328	0	0	846	0	0	0	0	0	271	31,990	868	11,980	1,730,170
1987	1,266,317	157,746	8,012	181,429	9,599	4,539	0	50,372	0	0	1,934	0	0	0	0	0	621	30,061	131	9,724	1,720,485
1988	1,271,000	197,891	6,608	223,156	7,108	5,883	0	137,418	0	0	3,026	0	0	0	0	0	619	22,084	417	12,643	1,887,853
1989	1,386,000	168,918	4,500	153,165	8,822	3,222	0	63,452	0	0	4,723	0	0	0	0	0	673	17,994	306	5,101	1,816,876
1990	1,426,000	171,008	4,445	80,584	9,620	4,232	0	22,568	0	0	20,289	0	0	0	0	0	1,248	22,205	471	6,325	1,768,995
1991	1,346,464	172,158	3,199	96,135	6,878	13,686	46,681	30,489	0	0	7,289	0	0	0	0	0	945	24,523	574	16,376	1,765,397
1992	1,438,412	206,129	2,104	146,946	2,770	11,980	51,956	34,825	0	0	13,586	0	0	0	0	0	4,364	49,441	880	33,074	1,996,467
1993	1,358,758	167,390	2,747	105,809	8,468	9,298	64,260	28,871	0	0	17,138	0	0	0	0	0	685	66,006	682	23,953	1,854,065
1994	1,421,402	196,572	2,470	144,544	10,379	14,377	60,584	29,775	0	0	18,866	0	0	0	0	0	562	69,591	588	24,532	1,994,242
1995	1,329,503	245,030	2,048	124,752	8,193	9,283	55,028	34,908	0	0	15,944	0	0	0	0	0	849	81,554	459	22,201	1,929,752
1996	1,218,229	240,590	1,349	130,163	6,376	14,610	47,146	35,451	0	0	23,078	0	0	0	0	0	642	103,867	1,167	21,437	1,844,105
1997	1,142,140	234,641	1,326	166,915	7,666	9,651	67,520	42,413	0	0	16,747	0	0	0	0	0	468	65,839	1,761	22,552	1,779,639
1998	1,125,249	195,645	1,181	101,315	9,124	15,679	33,667	39,994	0	0	14,863	0	0	0	0	0	588	57,096	916	25,604	1,620,921
1999	890,554	162,361	1,211	67,320	5,627	10,573	40,511	33,095	0	0	18,228	0	0	0	0	0	765	53,644	402	18,678	1,302,969
2000	1,133,980	191,056	1,790	84,070	6,974	13,228	49,666	36,926	0	0	15,597	0	0	0	0	0	840	47,229	383	26,108	1,607,847
2001	1,388,276	176,659	1,937	63,578	5,312	14,056	29,475	27,790	0	0	16,735	0	0	0	0	0	906	61,560	1,766	27,177	1,815,227
2002	1,482,992	197,353	2,261	74,985	3,635	11,853	41,865	30,379	0	0	15,854	0	0	0	0	0	952	45,294	1,344	28,619	1,937,386
2003	1,493,692	212,785	2,048	81,050	3,530	14,580	37,339	17,583	10,118	0	20,156	0	0	0	0	0	737	59,550	1,282	28,312	1,982,562
2004	1,481,701	212,152	1,993	75,510	2,219	18,133	48,680	22,180	7,888	0	11,896	4,684	240	208	655	60,562	1,014	29,344	0	1,979,058	
2005	1,484,895	205,424	2,550	94,384	2,559	14,233	37,361	20,699	11,194	0	10,426	3,964	169	90	464	62,012	1,185	29,442	0	1,981,050	
2006	1,488,148	191,906	2,185	99,068	1,965	13,302	36,452	21,104	17,309	0	12,856	3,823	208	203	579	61,878	1,416	26,562	0	1,978,964	
2007/d	1,353,018	172,407	2,166	119,332	1,946	11,701	37,013	25,176	19,411	0	17,772	3,936	318	163	635	56,620	1,193	26,467	0	1,849,274	

a/ Arrowtooth flounder included in Greenland turbot catch statistics, 1964-69.
 b/ Includes POP shortraker, rougheye, northern and sharpchin rockfish until 2004.
 for research.

c/ Rocksole prior to 1991 is included in other flatfish catch statistics.
 d/ Data through October 27, 2007.

Note: Numbers don't include fish taken

Table 5. Summary of stock abundance (biomass), overfishing level (OFL), acceptable biological catch (ABC), the fishing mortality rate corresponding to ABC (FABC), and the fishing mortality rate corresponding to OFL (FOFL) for the eastern Bering Sea (EBS), Aleutian Islands (AI), and Bogoslof district as projected for 2008 and 2009. "Biomass" corresponds to projected January abundance for the age+ range reported in the summary. Stock-specific biomass, OFL, and ABC are in metric tons, reported to three significant digits (four digits are used when a stock-specific ABC is apportioned among areas on a percentage basis). Fishing mortality rates are reported to two.

		Biomass	2008				2009				
			OFL	ABC	FOFL	F _{ABC}	OFL	ABC	FOFL	FABC	
Pollock	EBS*	4,360,000	1,440,000	1,000,000	0.30	0.21	1,320,000	1,000,000	0.35	0.27	
	Aleutian Islands	197,000	34,000	28,200	0.24	0.2	26,100	22,700	0.24	0.2	
	Bogoslof District	292,000	58,400	7,970	0.20	0.027	58,400	7,970	0.20	0.027	
Pacific cod	BSAI	1,080,000	176,000	150,000	0.26	0.22	190,000	162,000	0.26	0.22	
Sablefish	BS	41,000	3,380	2,860	0.101	0.084	2,910	2,610	0.101	0.084	
	AI	34,000	2,890	2,440	0.101	0.084	2,510	2,230	0.101	0.084	
Yellowfin sole	BSAI	2,200,000	265,000	248,000	0.2	0.19	296,000	276,000	0.2	0.19	
Greenland turbot	Total	104,000	15,600	2,540	0.67	0.51	16,000	2,540	0.67	0.51	
	BS			1,750			0	1,750			
	AI			787			0	787			
Arrowtooth flounder	BSAI	1,780,000	297,000	244,000	0.3	0.24	300,000	246,000	0.3	0.24	
Northern rock sole	BSAI	1,880,000	304,000	301,000	0.22	0.19	379,000	375,000	0.22	0.19	
Flathead sole	BSAI	820,000	86,000	71,700	0.34	0.28	83,700	69,700	0.34	0.28	
Alaska plaice	BSAI	1,850,000	248,000	194,000	0.81	0.59	277,000	217,000	0.81	0.59	
Other flatfish	BSAI	150,000	28,800	21,600	.17/.09/.15	.13/.06/.2	28,800	21,600	.17/.09/.15	.13/.06/.2	
Pacific Ocean perch	BSAI	453,000	25,700	21,700	0.07	0.059	25,400	21,300	0.07	0.059	
	BS			4,200				4,140			
	AI total			17,500				17,200			
	WAI			7,600				7,490			
	CAI			4,990				4,900			
	EAI			4,900				4,820			
Northern rockfish	BSAI	212,000	9,740	8,180	0.053	0.045	9,680	8,130	0.053	0.045	
Shortraker	BSAI	18,900	564	424	0.030	0.023	564	424	0.03	0.023	
Rougheye	BSAI	10,800	269	202	0.025	0.019	269	202	0.025	0.019	
Other rockfish	BSAI	36,700	1,330	999	.03/.09	.023/.068	1,290	968	.03/.09	.023/.068	
	BS			414			0	414			
	AI			585			0	554			
Atka mackerel	Total	323,000	71,400	60,700	0.4	0.33	50,600	47,500	0.36	0.32	
	WAI			16,900				13,200			
	CAI			24,300				19,000			
	EAI/BS			19,500				15,300			
Squid	BSAI	n/a	2,620	1,970	n/a	n/a	2,620	1,970	n/a	n/a	
Other species	BSAI	725,600	91,200	71,800			91,200	71,800			
	Sharks	BSAI	17,600	617	463		617	463			
	Skates	BSAI	491,000	37,200	31,300	0.10	0.075	36,800	30,900	0.1	0.075
	Sculpins	BSAI	217,000	53,100	39,800	0.19	0.14	53,100	39,800	0.19	0.14
	Octopus	BSAI		324	243			324	243		
Total	BSAI	16,565,000	3,161,893	2,440,285			3,162,043	2,557,644			

*spawning biomass

Table 6. Summary of groundfish tier designations under Amendment 56, maximum permissible ABC fishing mortality rate (max FABC), the Plan Team's recommended tier designation, ABC fishing mortality rate (FABC), the maximum permissible value of ABC (max ABC), the Plan Team's recommended ABC, and the percentage reduction (% Red.) between max ABC and the Plan Team's recommended ABC for 2008-2009. Stock-specific max ABC and ABC are in metric tons, reported to three significant digits (four significant digits are used when a stock-specific ABC is apportioned among areas on a percentage basis). Fishing mortality rates are reported to two significant digits.

Species or Complex	Area	Tier	2008					2009					
			max FABC	FABC	max ABC	ABC	% Red.	max FABC	FABC	max ABC	ABC	% Red.	
Pollock	EBS	1b	0.22	0.22	1,170,000	1,000,000	15%	1b	0.22	0.22	1,070,000	1,000,000	7%
	Aleutian Islands	3a	0.2	0.2	28,200	28,200		3a	0.2	0.2	22,700	22,700	
	Bogoslof District	5	0.017	0.017	43,800	7,970	82%	5	0.017	0.017	43,800	7,970	82%
Pacific cod	BSAI	3b	0.22	0.22	150,000	150,000		3b	0.22	0.22	162,000	162,000	
Sablefish	BS	3b	0.084	0.084	2,860	2,860		3b	0.084	0.084	2,610	2,610	
	AI	3b	0.084	0.084	2,440	2,440		3b	0.084	0.084	2,230	2,230	
Yellowfin sole	BSAI	1	0.19	0.19	248,000	248,000		1	0.19	0.19	276,000	276,000	
Greenland turbot	Total	3a	0.51	0.51	2,540	2,540		3a	0.51	0.51	2,540	2,540	
	BS												
	AI												
Arrowtooth flounder	BSAI	3a	0.24	0.24	244,000	244,000		3a	0.24	0.24	246,000	246,000	
Northern rock sole	BSAI	1	0.19	0.19	301,000	301,000		1	0.19	0.19	375,000	375,000	
Flathead sole	BSAI	3a	0.28	0.28	71,700	71,700		3a	0.28	0.28	69,700	69,700	
Alaska plaice	BSAI	3a	0.59	0.59	194,000	194,000		3a	0.59	0.59	217,000	217,000	
Other flatfish	BSAI	5	.13/.06/.2	.13/.06/.2	21,600	21,600		5	.13/.06/.2	.13/.06/.2	21,600	21,600	
Pacific Ocean perch	BSAI	3a	0.059	0.059	21,700	21,700		3a	0.059	0.059	21,300	21,300	
	BS				4,200	4,200		0	0	0	4,130	4,130	
	AI total				17,500	17,500		0	0	0	17,200	17,200	
	WAI				7,600	7,590		0	0	0	7,490	7,490	
	CAI				4,990	4,970		0	0	0	4,900	4,900	
	EAI				4,900	4,890		0	0	0	4,820	4,820	
	Northern rockfish	BSAI	3a	0.045	0.045	8,180	8,180		3a	0.045	0.045	8,130	8,130
Shortraker	BSAI	5	0.023	0.023	424	424		5	0.023	0.023	424	424	
Roughycye	BSAI	5	0.019	0.019	202	202		5	0.019	0.019	202	202	
Other rockfish	BSAI	5	.023/.068	.023/.068	999	999		5	.023/.068	.023/.068	968	968	
	BS				414	414		0			414	414	
	AI				585	585		0			554	554	
Atka mackerel	Total	3a	0.33	0.33	60,700	60,700		3b	0.33	0.33	47,500	47,500	
	WAI		0		16,900	16,900		0	0	0	13,200	13,200	
	CAI		0		24,300	24,300		0	0	0	19,000	19,000	
	EAI/BS		0		19,500	19,500		0	0	0	15,300	15,300	
Squid	BSAI	6	n/a	n/a	1,970	1,970		6	n/a	n/a	1,970	1,970	
Other Species		5	n/a	n/a	71,800	71,800		5	n/a	n/a	71,800	71,800	
	Sharks	BSAI	5		463	463		5			463	463	
Skates	BSAI	3	0.075	0.075	31,300	31,300		3	0.075	0.075	30,900	30,900	
Sculpins	BSAI	5	0.14	0.14	39,800	39,800		5	0.14	0.14	39,800	39,800	
Octopus	BSAI	6			243	243		6			243	243	
Total	BSAI				2,646,115	2,440,285					2,663,474	2,557,644	

Yellowfin Sole Product Quantities and Values 2003-2006

Year	Quantity (Tons)	Total Value (\$)	Unit Value (\$/Ton)
2003	39,443	38,138,876	967
2004	38,730	38,747,700	1,000
2005	52,221	69,132,577	1,324
2006	50,938	72,685,321	1,427
Increase 03-06	29%	91%	49%

Source: Table 2-6, NPFMC Agenda Item C-5(a), December, 2007.

**Summary Review Report for CIE Review
of Alaska flatfish stock assessments, June 11-15, 2007**

(excerpts from page 13)

The tier system was designed to allow greater levels of exploitation as data on the stocks is improved and all three reviewers noted that the tier system encourages better data collection and reporting. As noted in this report by several reviewers, "This system has multiple and cumulative layers of conservation for a high levels (sic) of biomass." Strengths of this management system include inclusion of precautionary measures, encouragement of the industry to improve data quality, and excellence of the stock assessment for several of the species. The reviewers believed that the tier system applies "precaution in dealing with structural uncertainty, such as on the reliability of the stock recruitment relationship." They believed that the principles behind the tier system were excellent and that the harvest rules were appropriate.

The reviewers specifically commented that "Tier 1 is supported by sound research with reliable point estimate of B and Bmsy and reliable pdf of Fmsy." Based on the quality of data the all agreed that the stock assessment for BSAI yellowfin sole and rock sole showed as "good a fit as possible to the data available" and that there were reliable estimates of B, BMSY, and FMSY. Thus there was consensus that this (sic) species should be managed under Tier 1.

Mike Hyde
American Seaf
Item D-1
Dec 2007

NPFMC


Transcript of Amendment 80 Discussions June 2005

[Discussion of basis for stair step allocations of yellowfin sole in Amendment 80 motion]

Arne Fuglvog:

Thank you, Madam Chair. Just one last comment. The idea of the threshold fishery is that if the pollock TAC comes down and those vessels who in past history had more participation in flatfish would then have opportunities. I think that we need to remember right now while the pollock TAC is very high there is less need to participate in the yellowfin sole fishery and much less dependency. While when the threshold fishery kicks in these vessels are going to have a much greater participation in yellowfin and they're going to make up for lost revenues from pollock; they are going to benefit from the threshold fishery.

Yellowfin Allocation at Various TAC Levels



St. George Kayumixtax Eco Office
St. George Traditional Council
PO Box 940
St. George Island, AK 99591

December 4, 2007

Mr. Eric Olsen, Chair
North Pacific Fisheries Management Council
605 W. 4th Avenue, Suite 306
Anchorage, AK 99501

Re: Agenda Item D-4, Groundfish Management

Dear Mr. Olsen and Council Members:

The Kayumixtax Ecosystem Conservation Office of St. George Island urges the Council to take a more precautionary, ecosystem-based approach to groundfish management. Pollock stocks are at an all time low and our community is worried about the overall health of the Bering Sea.

The Aleut community of St. George is dependant upon the sea for our source of food. Subsistence foods have been part of our culture for thousands of years from Steller Sea lions, Northern fur seals and sea birds, fish and many other species in the Bering Sea that are now being depleted at an unprecedented rate with no known solution in sight.

The council in a recent report states there is not enough scientific evidence to preclude large trawl ships from dragging the sensitive sea floor and catching millions of tons of by-catch annually this statement and fact does not give justification for blindly doing so. Until science can justify that dragging gear along the sea floor will cause no harm and that by-catch rates do not harm marine mammal populations and all our subsistence foods we urge you to reconsider your approach to management of groundfish stocks.

Respectfully,


Andrew Malavansky