

## BRISTOL BAY RED KING CRAB STOCK ASSESSMENT IN SPRING 2015

J. Zheng and M.S.M. Siddeek  
Alaska Department of Fish and Game  
Division of Commercial Fisheries  
P.O. Box 115526  
Juneau, AK 99811-5526, USA  
Phone: (907) 465-6102  
Fax: (907) 465-2604  
Email: jie.zheng@alaska.gov

### Executive Summary

1. Stock: red king crab (RKC), *Paralithodes camtschaticus*, in Bristol Bay, Alaska.
2. Catches: The domestic RKC fishery began to expand in the late 1960s and peaked in 1980 with a catch of 129.95 million lbs (58,943 t). The catch declined dramatically in the early 1980s and remained at low levels during the last three decades. Catches during recent years until 2010/11 were among the high catches in last 15 years. The retained catch in 2013/14 was about 7 million lbs (3,154 t) less than it was in 2009/10. The magnitude of bycatch from groundfish trawl fisheries has been stable and small relative to stock abundance during the last 10 years.
3. Stock biomass: Estimated mature biomass increased dramatically in the mid 1970s and decreased precipitously in the early 1980s. Estimated mature crab abundance has increased during the last 25 years with mature females being 3.4 times more abundant in 2009 than in 1985 and mature males being 2.3 times more abundant in 2009 than in 1985. Estimated mature abundance has steadily declined since 2009.
4. Recruitment: Estimated recruitment was high during 1970s and early 1980s and has generally been low since 1985 (1979 year class). During 1984-2014, only in 1984, 1995, 2002 and 2005 was estimated recruitment above the historical average for 1969-2014. Estimated recruitment was extremely low during the last 8 years.
5. Management performance:

Status and catch specifications (1000 t) (scenario 1):

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch	OFL	ABC
2010/11	13.63 <sup>A</sup>	32.64 <sup>A</sup>	6.73	6.76	7.71	10.66	N/A
2011/12	13.77 <sup>B</sup>	30.88 <sup>B</sup>	3.55	3.61	4.09	8.80	7.92
2012/13	13.19 <sup>C</sup>	29.05 <sup>C</sup>	3.56	3.62	3.90	7.96	7.17
2013/14	12.85 <sup>D</sup>	27.12 <sup>D</sup>	3.90	3.99	4.56	7.07	6.36
2014/15		24.69 <sup>D</sup>				6.82	6.14

The stock was above MSST in 2013/14 and is hence not overfished. Overfishing did not occur.

Status and catch specifications (million lbs):

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch	OFL	ABC
2010/11	30.0 <sup>A</sup>	72.0 <sup>A</sup>	14.84	14.91	17.00	23.52	N/A
2011/12	30.4 <sup>B</sup>	68.1 <sup>B</sup>	7.83	7.95	9.01	19.39	17.46
2012/13	29.1 <sup>C</sup>	64.0 <sup>C</sup>	7.85	7.98	8.59	17.55	15.80
2013/14	28.3 <sup>D</sup>	59.9 <sup>D</sup>	8.60	8.80	10.05	15.58	14.02
2014/15		54.4 <sup>D</sup>				15.04	13.53

Notes:

A – Calculated from the assessment reviewed by the Crab Plan Team in September 2011

B – Calculated from the assessment reviewed by the Crab Plan Team in September 2012

C – Calculated from the assessment reviewed by the Crab Plan Team in September 2013

D – Calculated from the assessment reviewed by the Crab Plan Team in September 2014

6. Basis for the OFL: All table values are in 1000 t (Scenario 1).

Year	Tier	B <sub>MSY</sub>	Current MMB	B/B <sub>MSY</sub> (MMB)	F <sub>OFL</sub>	Years to define B <sub>MSY</sub>	Natural Mortality
2010/11	3a	28.4	37.7	1.33	0.32	1995-2010	0.18
2011/12	3a	27.3	29.8	1.09	0.32	1984-2011	0.18
2012/13	3b	27.5	26.3	0.96	0.31	1984-2012	0.18
2013/14	3b	26.4	25.0	0.95	0.27	1984-2013	0.18
2014/15	3b	25.7	24.7	0.96	0.28	1984-2014	0.18

Basis for the OFL: All table values are in million lbs.

Year	Tier	$B_{MSY}$	Current MMB	$B/B_{MSY}$ (MMB)	$F_{OFL}$	Years to define $B_{MSY}$	Natural Mortality
2010/11	3a	62.7	83.1	1.33	0.32	1995-2010	0.18
2011/12	3a	60.1	65.6	1.09	0.32	1984-2011	0.18
2012/13	3b	60.7	58.0	0.96	0.31	1984-2012	0.18
2013/14	3b	58.2	55.0	0.95	0.27	1984-2013	0.18
2014/15	3b	56.7	54.4	0.96	0.28	1984-2014	0.18

## ***A. Summary of Major Changes***

**1. Change to management of the fishery:** None.

**2. Changes to the input data:**

- a. An alternative scenario is run with the new time series of NMFS trawl survey area-swept estimates provided by NMFS in February 2015.
- b. An alternative scenario is run with new female growth data from Kodiak red king crab.

**3. Changes to the assessment methodology:**

Three model scenarios are evaluated in this report (See Section E.3.a for details):

Scenarios 1 and 1n: Scenario 1 is renamed from scenario 4nb in the SAFE report in September 2014 for simplicity. Scenario 1n differs with scenario 1 by using the new time series of NMFS trawl survey area-swept estimates.

Scenario 2: Scenario 2 is the same as scenario 1 except that growth of immature and mature female are modeled differently and immature female growth data from Kodiak red king crab are used to estimate initial parameter values of immature female growth increments per molt function. Initial parameter values for three growth increments-per-molt functions are estimated using the growth increments per molt data: immature females, mature females, and males. Parameters for growth increments per molt are estimated inside the model with these initial estimates as a prior. A random walk approach is used to model the annual changes of sizes at the 50% maturity for females. A two-parameter logistic function is used to separate the immature and mature female length compositions for the initial year (1975) to reduce the number of parameter estimates.

We also try the approach of scenario 2 with the new time series of NMFS trawl survey data, but were unable to achieve model convergence. The model also failed to converge with the fixed initial parameter values for growth increments per molt with the current time series of trawl survey area-swept estimates for scenario 2. We will need further work on scenario 2 and some future Bristol Bay juvenile female growth data will help as well.

**4. Changes to assessment results:**

The new time series of area-swept abundance estimates provided by NMFS in February 2015 are similar to those provided in August 2014 except for a few years. The additional tow data are not used in the hot spot stations and area-swept estimates in some early years are also affected in the new time series, especially in 1979.

Model estimated relative survey biomasses are very similar between scenarios 1 and 1n except during mid- and late 1970s; the much lower area-swept biomass estimate in 1979 with the new time series causes the lower relative survey biomass estimates for scenario 1n. The absolute population biomass estimates are slightly higher for scenario 1n than for scenario 1 due to a slightly lower estimate of trawl survey catchability for the new time series. Model estimated relative survey biomasses are very similar between scenarios 1 and 2 except the most recent four years. Recent model relative biomass estimates for scenario 2 are lower than scenario 1. Modeling immature and mature female growths separately (scenario 2) improves the residual patterns for female length compositions. However, it is very difficult to get the model converge with scenario 2. For the new time series of survey area-swept estimates, we cannot get it converge when modeling immature and mature female growths separately with the same approach of scenario 2.

## ***B. Responses to SSC and CPT Comments***

### **1. Responses to the most recent two sets of SSC and CPT comments on assessments in general:**

None.

### **2. Responses to the most recent two sets of SSC and CPT comments specific to this assessment:**

#### **Response to CPT Comments (from September 2014)**

*“The CPT recommended that the assessment authors consider the affects of the final size bin used in the retained size composition data on model fitting (including the effects of the assumption of fixed sample size in the final bin) and consider the possibility of subdividing the final size bin into more than one bin.”*

This question comes from the difficulty of GMACS to fit the length composition data. The primary reason for the difficult fit is due to problems of estimating the growth transition matrix and survey selectivities. However, the final plus-length group does have impacts on the results. It is a trade-off between the numbers of empty length groups and relatively low impacts of large plus groups. We may consider examining this in the future.

#### **Response to CPT Comments (from May 2014)**

*“1. Drop Scenarios 4 and 4b because these use the old data.”*

Done.

*“2. Move forward with Scenarios 4na, 4nb for September 2014.”*

Done.

*“3. Although it appears to result in improved model fits, drop Scenario 4nb7 from consideration until a mechanism for the estimated higher  $M$  can be established; this scenario can be presented for reconsideration once a plausible mechanism has been identified.”*

SSC asked to continue 4nb7, which has been changed to 4n7. So scenario 4n7 is still in the SAFE report for September 2014. We drop it in the May 2015 SAFE report.

*“4. Add the number of estimated parameters to tables that compare values for likelihood components from different Scenarios so that the degree of improved fit can be more easily evaluated. Also, express the values of log-likelihood components between the base and alternative models as differences (e.g., base less alternative), rather than reporting the actual values because it is the differences in log-likelihood values that are informative.”*

Done.

#### **Response to SSC Comments specific to this assessment (from October 2014)**

*“The SSC recommends that if Model 4n7 is brought forward in 2015 as an alternative model, that reference points for Model 4n7 be recalculated with the higher  $M = 0.27$  estimated for 2006 – 2010. The SSC looks forward to the additional work planned by the author: implementing a random walk for natural mortality, investigation of recruitment dynamics, and investigation of survey weighting.”*

A scenario with random walk may be added in September-2015 assessments in the future. If this is the case, we will compute the reference point as suggested by the SSC.

We focus our effort to examine the female maturity as suggested by the SSC in 2014. Currently, we use a step curve to model changes in female size-at-maturity over time (see Figure A3). It would be better to fit the data with a continuous curve over time or model immature and mature female growths separately. However, the reason for modeling the change is to improve estimation of growth increments per molt. There are very little growth increment data for immature females in the eastern Bering Sea. Limited availability of growth increment data is the main reason for using a simple step curve.

In this report (May 2015), we use the immature female growth increments per molt data from Kodiak red king crab to estimate the parameters of the linear function of growth increments per molt. It is difficult because Kodiak red king crab grow faster with each molt than Bristol Bay red king crab. We assume that the slope parameter is the same for these two stocks and use the

limited Bristol Bay data to estimate the intercept parameter for immature females. With two separate growth increments per molt functions for immature and mature females, we are able to model the growth of females by maturity status inside the model.

Thank SSC for the good suggestions on spatial statistical analysis similar to that conducted by Kotwicki and Lauth (2012) and incorporating bottom temperature as a covariate on survey Q using the method in Wilderbuer et al. (2013). We will conduct these studies in the future. We will look at the survey Q with temperature data at the same time when we analyze the BSFRF's side-by-side tow survey data once the BSFRF study is completed.

### **Response to SSC Comments specific to this assessment (from June 2014)**

*“The SSC concurs with the PT recommendations, except that it would like Model 4nb7 or similar models to be investigated further for September 2014, if time permits. Similar models include the random walk model investigated in June 2013 or a model that uses environmental (e.g., SST) or biological (e.g., Pacific cod abundance) covariates. These models may provide insights into processes influencing natural mortality rates. The SSC agrees with the CPT that new procedures would be needed to accommodate estimation of biological reference points under assumptions of time varying M. A critical issue is to consider what “equilibrium” means under time varying M (especially when M is increasing in the most recent time period).”*

Scenario 4nb7, renamed as 4n7, is included in the September 2014 assessment.

*“The SSC found that the nomenclature for models was confusing and recommends that a more straightforward system be used. Also, the SSC encourages authors to continue to investigate whether recruitment is related to environmental or biological variables.”*

In the May 2015 SAFE report, we rename Scenario 4nb as scenario 1 for simplicity. In the September 2014 report, we still used the names similar to those in May 2014 for continuity. Scenario 4nb7 was shortened as 4n7.

Recruitment dynamics is the top priority for our research. We will continue to investigate factors that impact recruitment strength.

## ***C. Introduction***

### **1. Species**

Red king crab (RKC), *Paralithodes camtschaticus*, in Bristol Bay, Alaska.

## **2. General distribution**

Red king crab inhabit intertidal waters to depths >200 m of the North Pacific Ocean from British Columbia, Canada, to the Bering Sea, and south to Hokkaido, Japan. RKC are found in several areas of the Aleutian Islands and eastern Bering Sea.

## **3. Stock Structure**

The State of Alaska divides the Aleutian Islands and eastern Bering Sea into three management registration areas to manage RKC fisheries: Aleutian Islands, Bristol Bay, and Bering Sea (Alaska Department of Fish and Game, ADF&G, 2012). The Bristol Bay area includes all waters north of the latitude of Cape Sarichef (54°36' N lat.), east of 168°00' W long., and south of the latitude of Cape Newenham (58°39' N lat.) and the fishery for red king crab in this area is managed separately from fisheries for red king crab outside of this area; i.e., the red king crab in the Bristol Bay area are assumed to be a separate stock from red king crab outside of this area. This report summarizes the stock assessment results for the Bristol Bay RKC stock.

## **4. Life History**

Red king crab have a complex life history. Fecundity is a function of female size, ranging from several tens of thousands to a few hundreds of thousands (Haynes 1968, Swiney et al. 2012). The eggs are extruded by females and fertilized in the spring and are held by females for about 11 months (Powell and Nickerson 1965). Fertilized eggs are hatched in spring, most during the April to June period (Weber 1967). Primiparous females are bred a few weeks earlier in the season than multiparous females.

Larval duration and juvenile crab growth depend on temperature (Stevens 1990; Stevens and Swiney 2007). Male and female RKC mature at 5–12 years old, depending on stock and temperature (Loher et al. 2001, Stevens 1990) and may live >20 years (Matsuura and Takeshita 1990). Males and females attain a maximum size of 227 and 195 mm carapace length (CL), respectively (Powell and Nickerson 1965). Female maturity is evaluated by the size at which females are observed to carry egg clutches. Male maturity can be defined by multiple criteria including spermatophore production and size, chelae vs. carapace allometry, and participation in mating *in situ* (reviewed by Webb 2014). For management purposes, females >89 mm CL and males >119 mm CL are assumed to be mature for Bristol Bay RKC. Juvenile RKC molt multiple times per year until age 3 or 4; thereafter, molting continues annually in females for life and in males until maturity. Male molting frequency declines after attaining functional maturity.

## **5. Fishery**

The RKC stock in Bristol Bay, Alaska, supports one of the most valuable fisheries in the United States. A review of the history of the Bristol Bay red king crab fishery is provided in Fitch et al. (2012) and Otto (1989). The Japanese fleet started the fishery in the early 1930s, stopped fishing from 1940 to 1952, and resumed the fishery from 1953 until 1974. The Russian fleet fished for RKC from 1959 through 1971. The Japanese fleet employed primarily tanglenets with a very small proportion of catch from trawls and pots. The Russian fleet used only tanglenets. United States trawlers started to fish for Bristol Bay RKC in 1947, but their effort and catch declined in the 1950s. The domestic RKC fishery began to expand in the late 1960s and peaked in 1980 with a catch of 129.95 million lbs (58,943 t), worth an estimated \$115.3 million ex-vessel value. The catch declined dramatically in the early 1980s and has continued at low levels during the last two decades (Table

1). After the stock collapse in the early 1980s, the Bristol Bay RKC fishery took place during a short period in the fall (usually lasting about a week) with the catch quota based on the stock assessment conducted the previous summer (Zheng and Kruse 2002). Beginning with the 2005/2006 season, new regulations associated with fishery rationalization resulted in an increase in the duration of the fishing season (October 15 to January 15). With the implementation of crab rationalization, historical guideline harvest levels (GHL) were changed to a total allowable catch (TAC). Before rationalization, the implementation errors were quite high for some years and total actual catch from 1980 to 2007 was about 6% less than the sum of GHL/TAC over that period.

## **6. Fisheries Management**

King and Tanner crab stocks in the Bering Sea and Aleutian Islands are managed by the State of Alaska through a federal king and Tanner crab fishery management plan (FMP). Under the FMP, management measures are divided into three categories: (1) fixed in the FMP, (2) frameworked in the FMP, and (3) discretion of the State of Alaska. The State of Alaska is responsible for determining and establishing the GHL/TAC under the framework in the FMP.

Harvest strategies for the Bristol Bay RKC fishery have changed over time. Two major management objectives for the fishery are to maintain a healthy stock that ensures reproductive viability and to provide for sustained levels of harvest over the long term (ADF&G 2012). In attempting to meet these objectives, the GHL/TAC is coupled with size-sex-season restrictions. Only males  $\geq 6.5$ -in carapace width (equivalent to 135-mm carapace length, CL) may be harvested and no fishing is allowed during molting and mating periods (ADF&G 2012). Specification of TAC is based on a harvest rate strategy. Before 1990, harvest rates on legal males were based on population size, abundance of prerecruits to the fishery, and postrecruit abundance, and rates varied from less than 20% to 60% (Schmidt and Pengilly 1990). In 1990, the harvest strategy was modified, and a 20% mature male harvest rate was applied to the abundance of mature-sized ( $\geq 120$ -mm CL) males with a maximum 60% harvest rate cap of legal ( $\geq 135$ -mm CL) males (Pengilly and Schmidt 1995). In addition, a minimum threshold of 8.4 million mature-sized females ( $\geq 90$ -mm CL) was added to existing management measures to avoid recruitment overfishing (Pengilly and Schmidt 1995). Based on a new assessment model and research findings (Zheng et al. 1995a, 1995b, 1997a, 1997b), the Alaska Board of Fisheries adopted a new harvest strategy in 1996. That strategy had two mature male harvest rates: 10% when effective spawning biomass (ESB) is between 14.5 and 55.0 million lbs and 15% when ESB is at or above 55.0 million lbs (Zheng et al. 1996). The maximum harvest rate cap of legal males was changed from 60% to 50%. An additional threshold of 14.5 million lbs of ESB was also added. In 1997, a minimum threshold of 4.0 million lbs was established as the minimum GHL for opening the fishery and maintaining fishery manageability when the stock abundance is low. The Board modified the current harvest strategy by adding a mature harvest rate of 12.5% when the ESB is between 34.75 and 55.0 million lbs in 2003 and eliminated the minimum GHL threshold in 2012. The current harvest strategy is illustrated in Figure 1.

## ***D. Data***

### **1. Summary of New Information**

A new time series of NMFS trawl survey results provided by NMFS in 2015. Different scenarios are used to compare the current and new time series of the NMFS trawl survey results.



Some new female growth data from Kodiak red king crab are used to estimate growth increments per molt for females. Scenario 2 is using these new data.

## **2. Catch Data**

Data on landings of Bristol Bay RKC by length and year and catch per unit effort were obtained from annual reports of the International North Pacific Fisheries Commission from 1960 to 1973 (Hoopes et al. 1972; Jackson 1974; Phinney 1975) and from the ADF&G from 1974 to 2012. Bycatch data are available starting from 1990 and were obtained from the ADF&G observer database and reports (Gaeuman 2013). Sample sizes for catch by length and shell condition are summarized in Table 2. Relatively large samples were taken from the retained catch each year. Sample sizes for trawl bycatch were the annual sums of length frequency samples in the National Marine Fisheries Service (NMFS) database.

### ***(i). Catch Biomass***

Retained catch and estimated bycatch biomasses are summarized in Table 1 and illustrated in Figure 2. Retained catch and estimated bycatch from the directed fishery include the general, open-access fishery (prior to rationalization) or the individual fishery quota (IFQ) fishery (after rationalization) as well as the Community Development Quota (CDQ) fishery and the ADF&G cost-recovery harvest. Starting in 1973, the fishery generally occurred during the late summer and fall. Before 1973, a small portion of retained catch in some years was caught from April to June. Because most crab bycatch from the groundfish trawl fisheries occurred during the spring, the years in Table 1 are one year less than those from the NMFS trawl bycatch database to approximate the annual bycatch for reporting years defined as June 1 to May 31; e.g., year 2002 in Table 1 corresponds to what is reported for year 2003 in the NMFS database. Catch biomass is shown in Figure 2. Bycatch data for the cost-recovery fishery before 2006 were not available. In this report, pot fisheries include both the directed fishery and RKC bycatch in the Tanner crab pot fishery and trawl fisheries are groundfish trawl fisheries.

### ***(ii). Catch Size Composition***

Retained catch by length and shell condition and bycatch by length, shell condition, and sex were obtained for stock assessments. From 1960 to 1966, only retained catch length compositions from the Japanese fishery were available. Retained catches from the Russian and U.S. fisheries were assumed to have the same length compositions as the Japanese fishery during this period. From 1967 to 1969, the length compositions from the Russian fishery were assumed to be the same as those from the Japanese and U.S. fisheries. After 1969, foreign catch declined sharply and only length compositions from the U.S. fishery were used to distribute catch by length.

### ***(iii). Catch per Unit Effort***

Catch per unit effort (CPUE) is defined as the number of retained crab per tan (a unit fishing effort for tanglenets) for the Japanese and Russian tanglenet fisheries and the number of retained crab per potlift for the U.S. fishery (Table 3). Soak time, while an important factor influencing CPUE, is difficult to standardize. Furthermore, complete historical soak time data from the U.S. fishery are not available. Based on the approach of Balsiger (1974), all fishing effort from Japan, Russia, and U.S. were standardized to the Japanese tanglenet from 1960 to 1971, and the CPUE was standardized as crab per tan. Except for the peak-to-crash years of late 1970s and early 1980s the

correspondence between U.S. fishery CPUE and area-swept survey abundance is poor (Figure 3). Due to the difficulty in estimating commercial fishing catchability and crab availability to the NMFS annual trawl survey data, commercial CPUE data were not used in the model.

### 3. NMFS Survey Data

The NMFS has performed annual trawl surveys of the eastern Bering Sea since 1968. Two vessels, each towing an eastern otter trawl with an 83 ft headrope and a 112 ft footrope, conduct this multispecies, crab-groundfish survey during the summer. Stations are sampled in the center of a systematic 20 X 20 nm grid overlaid in an area of  $\approx 140,000 \text{ nm}^2$ . Since 1972 the trawl survey has covered the full stock distribution except in nearshore waters. The survey in Bristol Bay occurs primarily during late May and June. Tow-by-tow trawl survey data for Bristol Bay RKC during 1975-2014 were provided by NMFS.

Abundance estimates by sex, carapace length, and shell condition were derived from survey data using an area-swept approach (Figures 4 and 5). Spatial distributions of crab from the standard trawl surveys during recent years are shown in Appendix B. Until the late 1980s, NMFS used a post-stratification approach, but subsequently treated Bristol Bay as a single stratum; the estimates shown in Figures 4 and 5 were made without post-stratification. If multiple tows were made for a single station in a given year, the average of the abundances from all tows within that station was used as the estimate of abundance for that station. If more than one tow was conducted in a station because of high RKC abundance (i.e., the station is a “hot spot”), NMFS regards the station as a separate stratum. A “hot spot” was not surveyed with multiple tows during the early years. Two such “hot spots” affected the survey abundance estimates greatly: station H13 in 1984 (mostly juvenile crab 75-90 mm CL) and station F06 in 1991 (mostly newshell legal males). The tow at station F06 was discarded in the older NMFS abundance estimates (Stevens et al. 1991). In this study, all tow data were used. NMFS re-estimated the historic area-swept by tow using variable versus fixed net width and re-estimated area-swept abundance in 2008, using all tow data and standardized the survey time series estimates in 2014. We used the new area-swept estimates provided by NMFS in 2014.

In addition to standard surveys, NMFS also conducted some surveys after the standard surveys to assess mature female abundance. In addition to the standard survey conducted in early June (late May to early June in 1999 and 2000), a portion of the distribution of Bristol Bay RKC was re-surveyed in 1999, 2000, and 2006-2012. Resurveys performed in late July, about six weeks after the standard survey, included 31 stations (1999), 23 stations (2000), 31 stations (2006, 1 bad tow and 30 valid tows), 32 stations (2007-2009), 23 stations (2010) and 20 stations (2011 and 2012) with high female density. The resurveys were necessary because a high proportion of mature females had not yet molted or mated when sampled by the standard survey. Differences in area-swept estimates of abundance between the standard surveys and resurveys of these same stations are attributed to survey measurement errors or to seasonal changes in distribution between survey and resurvey. More large females were observed in the resurveys than during the standard surveys in 1999 and 2000 because most mature females had not molted prior to the standard surveys. As in 2006, area-swept estimates of males  $>89$  mm CL, mature males, and legal males within the 32 resurvey stations in 2007 were not significantly different between the standard survey and resurvey ( $P=0.74$ ,  $0.74$  and  $0.95$ ) based on paired  $t$ -tests of sample means. However, similar to 2006, area-swept estimates of mature females within the 32 resurvey stations in 2007 were significantly different between the standard survey and resurvey ( $P=0.03$ ) based on the  $t$ -test. Resurvey stations were close

to shore during 2010-2012 and mature and legal male abundance estimates were lower for the re-tow than the standard survey. Following the CPT recommendation, we used the standard survey data for male abundance estimates and only the resurvey data, plus the standard survey data outside the resurveyed stations, to assess female abundance during these resurvey years.

#### **4. Bering Sea Fisheries Research Foundation Survey Data**

The BSFRF conducted trawl surveys for Bristol Bay red king crab in 2007 and 2008 with a small-mesh trawl net and 5-minute tows. The surveys occurred at similar times with the NMFS standard surveys and covered about 97% of the Bristol Bay area. Few Bristol Bay red king crab were outside of the BSFRF survey area. Because of small mesh size, the BSFRF surveys were expected to catch nearly all red king crab within the swept area. Crab abundances of different size groups were estimated by the kriging method. Mature male abundances were estimated to be 22.331 in 2007 and 19.747 million in 2008 with associated CVs of 0.0634 and 0.0765.

### ***E. Analytic Approach***

#### **1. History of Modeling Approaches**

To reduce annual measurement errors associated with abundance estimates derived from the area-swept method, ADF&G developed a length-based analysis (LBA) in 1994 that incorporates multiple years of data and multiple data sources in the estimation procedure (Zheng et al. 1995a). Annual abundance estimates of the Bristol Bay RKC stock from the LBA have been used to manage the directed crab fishery and to set crab bycatch limits in the groundfish fisheries since 1995 (Figure 1). An alternative LBA (research model) was developed in 2004 to include small size groups for federal overfishing limits. The crab abundance declined sharply during the early 1980s. The LBA estimated natural mortality for different periods of years, whereas the research model estimated additional mortality beyond a basic constant natural mortality during 1976-1993. In this report, we present only the research model that was fit to the data from 1975 to 2014.

#### **2. Model Description**

The original LBA model was described in detail by Zheng et al. (1995a, 1995b) and Zheng and Kruse (2002). The model combines multiple sources of survey, catch, and bycatch data using a maximum likelihood approach to estimate abundance, recruitment, selectivities, catches, and bycatch of the commercial pot fisheries and groundfish trawl fisheries. A full model description is provided in Appendix A.

a-f. See appendix A.

g. Critical assumptions of the model:

- i. The base natural mortality is constant over shell condition and length and was estimated assuming a maximum age of 25 and applying the 1% rule (Zheng 2005).
- ii. Survey and fisheries selectivities are a function of length and were constant over shell condition. Selectivities are a function of sex except for trawl bycatch selectivities, which are the same for both sexes. Two different survey selectivities

were estimated: (1) 1975-1981 and (2) 1982-2014 based on modifications to the trawl gear used in the assessment survey.

- iii. Growth is a function of length and is assumed to not change over time for males. For females with scenarios 1 and 1n, growth-per-molt increments as a function of length were estimated for three periods (1975-1982, 1983-1993, and 1994-2014) based on sizes at maturity. For females with scenario 2, sizes at 50% maturity change annually with a random walk approach. Once mature, female red king crab grow with a much smaller growth increment per molt.
- iv. Molting probabilities are an inverse logistic function of length for males. Females molt annually.
- v. Annual fishing seasons for the directed fishery are short.
- vi. The prior of survey catchability ( $Q$ ) was estimated to be 0.896, based on a trawl experiment by Weinberg et al. (2004) with a standard deviation of 0.025.  $Q$  is assumed to be constant over time and is estimated in the model.
- vii. Males mature at sizes  $\geq 120$  mm CL. For convenience, female abundance was summarized at sizes  $\geq 90$  mm CL as an index of mature females.
- viii. Measurement errors were assumed to be normally distributed for length compositions and were log-normally distributed for biomasses.
- h. Changes to the above since previous assessment: see Section A.3. Changes to the assessment methodology.
- i. Outline of methods used to validate the code used to implement the model and whether the code is available: The code is available.

### 3. Model Selection and Evaluation

- a. Alternative model configurations:

Several scenarios were compared for this report:

Scenario 1 (renamed from previous scenario 4nb): base scenario. Scenario 1 includes:

- (1) Basic  $M = 0.18$ , with an additional mortality level during 1980-1984 for males and two additional mortality levels (one for 1980-1984 and the other for 1976-1979 and 1985-1993) for females. For scenario 2, the additional mortality level for 1976-1979 and 1985-1993 is 0 based on the model estimate, and thus is fixed to 0.
- (2) Including BSFRF survey data in 2007 and 2008.
- (3) Survey catchability is estimated in the model and is assumed to be constant over time.
- (4) Two levels of molting probabilities for males: one before 1980 and one after 1979, based on survey shell condition data. Each level has two parameters.
- (5) Estimating effective sample size from observed sample sizes. Effective sample sizes are estimated as  $\min(0.5 \cdot \text{observed-size}, N)$  for trawl surveys and  $\min(0.1 \cdot \text{observed-size}, N)$  for catch and bycatch, where  $N$  is the maximum sample size (200

for trawl surveys, 100 for males from the pot fishery and 50 for females from pot fishery and both males and females from the trawl fisheries. The effective sample sizes are plotted against the implied effective sample sizes in Figures 6 and 7, where the implied effective sample sizes are estimated as follows:

$$n_y = \sum_l \hat{P}_{y,l}(1-\hat{P}_{y,l}) / \sum_l (P_{y,l} - \hat{P}_{y,l})^2$$

where  $\hat{P}_{y,l}$  and  $P_{y,l}$  are estimated and observed size compositions in year  $y$  and length group  $l$ , respectively.

(6) Standard survey data for males and retow data for females.

(7) Estimating initial year length compositions.

Scenario 1n: Scenario 1n differs with scenario 1 by using the new time series of NMFS trawl survey results provided in February 2015.

Scenario 2: Scenario 2 is the same as scenario 1 except that growth of immature and mature female are modeled differently and immature female growth data from Kodiak red king crab are used to estimate initial parameter values of immature female growth increments per molt function. Initial parameter values for three growth increments-per-molt functions are estimated using the growth increments per molt data: immature females, mature females, and males. Parameters for growth increments per molt are estimated inside the model with these initial estimates as a prior. A random walk approach is used to model the annual changes of sizes at the 50% maturity for females. A two-parameter logistic function is used to separate the immature and mature female length compositions for the initial year (1975) to reduce the number of parameter estimates. We also try the approach of scenario 2 with the new time series of NMFS trawl survey data, but we are not able to get it converge.

Only the full results for scenarios 1 and 1n are presented in this report, although most results for scenario 2 are also shown. Each figure or table is indicated with a scenario.

- b. Progression of results: See the new results at the beginning of the report.
- c. Evidence of search for balance between realistic and simpler models: NA.
- d. Convergence status/criteria: ADMB default convergence criteria.
- e. Sample sizes for length composition data. Estimated sample sizes and effective sample sizes are summarized in tables.
- f. Credible parameter estimates: All estimated parameters seem to be credible.
- g. Model selection criteria. The likelihood values were used to select among alternatives that could be legitimately compared by that criterion.
- h. Residual analysis. Residual plots are illustrated in figures.
- i. Model evaluation is provided under Results, below.

## 4. Results

### a. Effective sample sizes and weighting factors.

#### i. The effective sample sizes are:

- (1) Trawl surveys: 200 for males and females except for females: 184 in 1986, 180 in 1992 and 133 in 1994.
- (2) Retained catch: 100.
- (3) Pot male discard: 100 except 87 in 1990 and 23 in 1996.
- (4) Pot female discard: 50 except 38 in 1991, 1 in 1996, 4 in 1999, and 30 in 2002.
- (5) Trawl bycatch: 50 for males and females except for males 44 in 1988, 21 in 1991 and 1992, 33 in 1994, 10 in 1995, and for females 28 in 1986 and 1988, 19 in 1989, 40 in 1991, 11 in 1992, 25 in 1994, 5 in 1995, 48 in 1997.
- (6) Tanner fishery bycatch: 50 for males and females except for males 28 in 1992, 23 in 1993 and 22 in 2013, and for females 27 in 1993.
- (7) BSFRF survey: 200 for the BSFRF survey males and females.

For scenario 1, effective sample sizes are illustrated in Figures 6 and 7.

#### ii. Weights are assumed to be 500 for retained catch biomass, and 100 for all bycatch biomasses, 2 for recruitment variation, and 10 for recruitment sex ratio.

iii. Initial trawl survey catchability ( $Q$ ) is estimated to be 0.896 with a standard deviation of 0.025 (CV about 0.03) based on the double-bag experiment results. These values are used as a prior for estimating  $Q$  in the model. For scenario 2, initial parameter estimates (standard deviations) of  $a$  and  $b$  for the growth increments-per-molt functions are: immature females: 19.1291 (3.7880) and -0.0543 (0.0454), mature females: 13.9552 (0.9957) and -0.0787 (0.0077), and males: 17.2575 (0.6364) and -0.0111 (0.00476).

### b. Tables of estimates.

- i. Parameter estimates for scenarios 1 and 1n are summarized in Tables 4 and 5.
- ii. Abundance and biomass time series are provided in Table 6 for scenarios 1 and 1n.
- iii. Recruitment time series for scenarios 1 and 1n are provided in Table 6.
- iv. Time series of catch biomass is provided in Table 1.

Negative log-likelihood values and parameter estimates are summarized in Tables 4 and 5, respectively. Length-specific fishing mortality is equal to selectivity-at-length times the full fishing mortality. Estimated full pot fishing mortalities for females and full fishing mortalities for trawl bycatch were very low due to low bycatch as well as handling mortality rates less than 1.0. Estimated recruits varied greatly from year to year (Table 6). Estimated low selectivities for male pot bycatch, relative to the retained catch, reflected the 20% handling mortality rate (Figure 8). Both selectivities were applied to the same level of full fishing mortality. Estimated selectivities for female pot bycatch were close to 1.0 for all mature females, and the estimated full fishing mortalities for female pot bycatch were lower than for male retained catch and bycatch (Table 5).

c. Graphs of estimates.

- i. Selectivities and molting probabilities by length are provided in Figures 8 and 9 for scenarios 1 and 1n.

One of the most important results is estimated trawl survey selectivity/catchability (Figure 8). Survey selectivity affects not only the fitting of the data but also the absolute abundance estimates. Estimated survey selectivities in Figure 8 are generally smaller than the capture probabilities in Figure A1 because survey selectivities include capture probabilities and crab availability. NMFS survey catchability was estimated to be 0.896 from the trawl experiment, which is higher than that estimated from the BSFRF surveys (0.854). The reliability of estimated survey selectivities will greatly affect the application of the model to fisheries management. Under- or overestimates of survey selectivities will cause a systematic upward or downward bias of abundance estimates. Information about crab availability to the survey area at survey times will help estimate the survey selectivities.

For scenarios 1 and 1n, estimated molting probabilities during 1975-2014 (Figure 9) were generally lower than those estimated from the 1954-1961 and 1966-1969 tagging data (Balsiger 1974). Lower molting probabilities mean more oldshell crab, possibly due to changes in molting probabilities over time or shell aging errors. Overestimates or underestimates of oldshell crab will result in lower or higher estimates of male molting probabilities.

- ii. Estimated total survey biomass and mature male and female abundances are plotted in Figure 10.

Model estimated relative survey biomasses are very similar between scenarios 1 and 1n except during mid- and late 1970s; the much lower area-swept biomass estimate in 1979 with the new time series causes the lower relative survey biomass estimates for scenario 1n. The absolute population biomass estimates are slightly higher for scenario 1n than for scenario 1 due to a slightly lower estimate of trawl survey catchability for the new time series. Model estimated relative survey biomasses are very similar between scenarios 1 and 2 except the most recent four years. Recent model relative biomass estimates for scenario 2 are lower than scenario 1.

Although the model did not fit the mature crab abundance directly, trends in the mature abundance estimates agree well with observed survey values except in 2014 (Figure 10b). Estimated mature crab abundance increased dramatically in the mid 1970s then decreased precipitously in the early 1980s. Estimated mature crab abundance has increased during the last 27 years with mature females being 3.4 times more abundant in 2009 than in 1985 and mature males being 2.3 times more abundant in 2009 than in 1985 for scenario 1 (Figure 10b). Model estimates of mature abundances have declined since the late 2000s.

The fit to BSFRF survey data and estimated survey selectivities are illustrated in Figures 10c-e.

- iii. Estimated recruitment time series are plotted in Figure 11 for scenarios 1, 1n and 2.
- iv. Estimated fishing mortality rates are plotted against mature male biomass in Figure 12 for scenarios 1, 1n and 2.

The average of estimated male recruits from 1984 to 2014 (Figure 11) and mature male biomass per recruit were used to estimate  $B_{35\%}$ . Alternative periods of 1976-present and 1976-1983 were compared in our report. The full fishing mortalities for the directed pot fishery at the time of fishing were plotted against mature male biomass on Feb. 15 (Figure 12). Estimated fishing mortalities in most years before the current harvest strategy was adopted in 1996 were above  $F_{35\%}$  (Figure 12). Under the current harvest strategy, estimated fishing mortalities were at or above the  $F_{35\%}$  limits in 1998, 2003, 2005-2010 for scenario 1 but below the  $F_{35\%}$  limits in the other post-1995 years. The estimated higher survey catchabilities with scenarios 1 and 2 result in relatively higher fishing mortalities than those with scenario 1n.

For scenario 1, estimated full pot fishing mortalities ranged from 0.00 to 1.58 during 1975-2013, with estimated values over 0.40 during 1975-1981, 1986-1987, 1993, and 2007-2008 (Table 5, Figure 12). For scenario 1n, estimated full pot fishing mortalities ranged from 0.00 to 1.52 during 1975-2013, with estimated values over 0.40 during 1975-1981, 1986-1987, and 2008 (Table 5, Figure 12). For scenario 2, estimated full pot fishing mortalities ranged from 0.00 to 1.77 during 1975-2013, with estimated values over 0.40 during 1975-1981, 1984-1987, 1993, 1998, 2005, and 2007-2008 (Figure 12). Estimated fishing mortalities for pot female bycatch and trawl bycatch were generally less than 0.06.

- v. Estimated mature male biomass and recruitment are plotted to illustrate their relationships with scenario 1 (Figure 13a). Annual stock productivities are illustrated in Figure 13b.

Stock productivity (recruitment/mature male biomass) was generally lower during the last 20 years (Figure 13c).

Egg clutch data collected during summer surveys may provide information about mature female reproductive conditions. Although egg clutch data are subject to rating errors as well as sampling errors, data trends over time may be useful. Proportions of empty clutches for newshell mature females >89 mm CL were high in some years before 1990, but have been low since 1990 (Figure 14). The highest proportion of empty clutches (0.2) was in 1986, and primarily involved soft shell females (shell condition 1). Clutch fullness fluctuated annually around average levels during two periods: before 1991 and after 1990 (Figure 14). The average clutch fullness was close for these two periods (Figure 14).

- d. Graphic evaluation of the fit to the data.
  - i. Observed vs. estimated catches are plotted in Figure 15.
  - ii. Model fits to total survey biomass are shown in Figure 10 with a standardized residual plot in Figure 16.



iii. Model fits to catch and survey proportions by length are illustrated in Figures 17-24 and residual bubble plots are shown in Figures 25-27.

The model (scenarios 1, 1n and 2) fit the fishery biomass data well and the survey biomass reasonably well (Figures 10 and 15). Because the model estimates annual fishing mortality for directed pot male catch, undirected pot male bycatch, pot female bycatch, and trawl bycatch, the deviations of observed and predicted (estimated) fishery biomass are mainly due to size composition differences.

The model also fit the length composition data well (Figures 17-24). It is surprising that the model fit the length proportions of the pot male bycatch well with two simple linear selectivity functions (Figure 21). We explored a logistic selectivity function, but due to the long left tail of the pot male bycatch selectivity, the logistic selectivity function did not fit the data well.

Modal progressions are tracked well in the trawl survey data, particularly beginning in the mid-1990s (Figures 17 and 19). Cohorts first seen in the trawl survey data in 1975, 1986, 1990, 1995, 1999, 2002 and 2005 can be tracked over time. Some cohorts can be tracked over time in the pot bycatch as well (Figure 21), but the bycatch data did not track the cohorts as well as the survey data. Groundfish trawl bycatch data provide little information to track modal progression (Figures 23 and 24).

Standardized residuals of total survey biomass and proportions of length are plotted to examine their patterns. Residuals were calculated as observed minus predicted and standardized by the estimated standard deviation. Standardized residuals of total survey biomass did not show any consistent patterns (Figure 16). Standardized residuals of proportions of survey males appear to be random over length and year (Figures 25 and 26). There is an interesting pattern for residuals of proportions of survey females. Residuals were generally negative for large-sized mature females during 1975-1987 for scenarios 1 and 1n (Figure 27). Changes in growth over time or increased mortality may cause this pattern. The inadequacy of the model can be corrected by adding parameters to address these factors. Molding immature and mature female growths separately (scenario 2) improves the residual patterns for female length compositions. However, it is very difficult to get the model converge with scenario 2.

e. Retrospective and historic analyses.

Two kinds of retrospective analyses were conducted for this report: (1) the 2014 model (scenario 1) hindcast results and (2) historical results. The 2014 model results are based on sequentially excluding one-year of data to evaluate the current model performance with fewer data. The historical results are the trajectories of biomass and abundance from previous assessments that capture both new data and changes in methodology over time. Treating the 2014 estimates as the baseline values, we can also evaluate how well the model had done in the past.

i. Retrospective analysis (retrospective bias in base model or models).

The performance of the 2014 model includes sequentially excluding one-year of data. The model with scenario 4nb performed reasonably well during 2008-2013

with a lower terminal year estimates in 2012 and 2013 and higher estimates during 2008-2010 (Figure 28).

ii. Historic analysis (plot of actual estimates from current and previous assessments).

The model first fit the data from 1985 to 2004 in the terminal year of 2004. Thus, 10 historical assessment results are available for comparison with the 2014 assessment model results (Figure 29). The main differences of the 2004 model were weighting factors and effective sample sizes for the likelihood functions. In 2004, the weighting factors were 1000 for survey biomass, 2000 for retained catch biomass and 200 for bycatch biomasses. The effective sample sizes were set to be 200 for all proportion data but weighting factors of 5, 2, and 1 were also applied to retained catch proportions, survey proportions and bycatch proportions. Estimates of time series of abundance in 2004 were generally higher than those estimated after 2004 (Figure 29).

In 2005, to improve the fit for retained catch data, the weight for retained catch biomass was increased to 3000 and the weight for retained catch proportions was increased to 6. All other weights were not changed. In 2006, all weights were re-configured. No weights were used for proportion data, and instead, effective sample sizes were set to 500 for retained catch, 200 for survey data, and 100 for bycatch data. Weights for biomasses were changed to 800 for retained catch, 300 for survey and 50 for bycatch. The weights in 2007 were the same as 2006. Generally, estimates of time series of abundance in 2005 were slightly lower than in 2006 and 2007, and there were few differences between estimates in 2006 and 2007 (Figure 29).

In 2008, estimated coefficients of variation for survey biomass were used to compute likelihood values as suggested by the CPT in 2007. Thus, weights were re-configured to: 500 for retained catch biomass, 50 for survey biomass, and 20 for bycatch biomasses. Effective sample size was lowered to 400 for the retained catch data. These changes were necessary for the estimation to converge and for a relatively good balanced fit to both biomasses and proportion data. Also, sizes at 50% selectivities for all fisheries data were allowed to change annually, subject to a random walk pattern, for all assessments before 2008. The 2008 model does not allow annual changes in any fishery selectivities. Except for higher estimates of abundance during the late 1980s and early 1990s, estimates of time series of abundance in 2008 were generally close to those in 2006 and 2007 (Figure 29).

During 2009-2013, the model was extended to the data through 1968. No weight factors were used for the NMFS survey biomass during 2009-2013 assessments. Since 2013, the model has fitted the data only back to 1975 for consistence of trawl survey data. Two levels of molting probabilities over time were used, shell conditions for males were combined, and length composition data of the BSFRF survey were used as well. In 2014, the trawl survey time series were re-estimated and a trawl survey catchability was estimated for some scenarios.

Overall, both historical results (historic analysis) and the 2014 model results (retrospective analysis) performed reasonably well. No great overestimates or

underestimates occurred as was observed in assessments for Pacific halibut (*Hippoglossus stenolepis*) (Parma 1993) and some eastern Bering Sea groundfish stocks (Zheng and Kruse 2002, Ianelli et al. 2003). Since the most recent model was not used to set TAC or overfishing limits until 2009, historical implications for management from the stock assessment errors cannot be evaluated at the current time. However, management implications of the ADF&G stock assessment model were evaluated by Zheng and Kruse (2002).

f. Uncertainty and sensitivity analyses

- i. Estimated standard deviations of parameters are summarized in Table 5 for scenarios 1 and 1n. Estimated standard deviations of mature male biomass are listed in Table 6.
- ii. Probabilities for trawl survey catchability  $Q$  are illustrated in Figure 30 for scenario 1 using the mcmc approach; estimated  $Q$ s are generally less than 1.0. Probabilities for mature male biomass and OFL in 2014 are illustrated in Figure 31 for scenario 1 using the mcmc approach. The confidence intervals are quite narrow.
- iii. Sensitivity analysis for handling mortality rate was reported in the SAFE report in May 2010. The baseline handling mortality rate for the directed pot fishery was set at 0.2. A 50% reduction and 100% increase resulted in 0.1 and 0.4 as alternatives. Overall, a higher handling mortality rate resulted in slightly higher estimates of mature abundance, and a lower rate resulted in a minor reduction of estimated mature abundance. Differences of estimated legal abundance and mature male biomass were small among these handling mortality rates.
- iv. Sensitivity of weights. Sensitivity of weights was examined in the SAFE report in May 2010. Weights to biomasses (trawl survey biomass, retained catch biomass, and bycatch biomasses) were reduced to 50% or increased to 200% to examine their sensitivity to abundance estimates. Weights to the penalty terms (recruitment variation and sex ratio) were also reduced or increased. Overall, estimated biomasses were very close under different weights except during the mid-1970s. The variation of estimated biomasses in the mid-1970s was mainly caused by the changes in estimates of additional mortalities in the early 1980s.

g. Comparison of alternative model scenarios

These comparisons, based on the data through 2010, were reported in the SAFE report in May 2011. Estimating length proportions in the initial year (scenario 1a) results in a better fit of survey length compositions at an expense of 36 more parameters than scenario 1. Abundance and biomass estimates with scenario 1a are similar between scenarios. Using only standard survey data (scenario 1b) results in a poorer fit of survey length compositions and biomass than scenarios using both standard and re-tow data (scenarios 1, 1a, and 1c) and has the lowest likelihood value. Although the likelihood value is higher for using both standard survey and re-tow data for males (scenario 1) than using only standard survey for males (scenario 1c), estimated abundances and biomasses are almost identical. The higher likelihood value for scenario 1 over scenario 1c is due to trawl bycatch length compositions.

In this report (May 2015), three scenarios are compared. Model estimated relative survey biomasses are very similar between scenarios 1 and 1n and differ with those of 2. Scenario 2 provides a better fit of trawl survey data during 2011-2013 and results in a much lower OFL than scenarios 1 and 1n.

## ***F. Calculation of the OFL and ABC***

1. Bristol Bay RKC is currently placed in Tier 3b (NPFMC 2007).
2. For Tier 3 stocks, estimated biological reference points include  $B_{35\%}$  and  $F_{35\%}$ . Estimated model parameters were used to conduct mature male biomass-per-recruit analysis.
3. Specification of the OFL:

The Tier 3 can be expressed by the following control rule:

$$\begin{aligned}
 \text{a) } \frac{B}{B^*} > 1 & \quad F_{OFL} = F^* \\
 \text{b) } \beta < \frac{B}{B^*} \leq 1 & \quad F_{OFL} = F^* \left( \frac{B/B^* - \alpha}{1 - \alpha} \right) \quad (1) \\
 \text{c) } \frac{B}{B^*} \leq \beta & \quad \text{directed fishery } F = 0 \text{ and } F_{OFL} \leq F^*
 \end{aligned}$$

Where

$B$  = a measure of the productive capacity of the stock such as spawning biomass or fertilized egg production. A proxy of  $B$ , MMB estimated at the time of primiparous female mating (February 15) is used as a default in the development of the control rule.

$F^* = F_{35\%}$ , a proxy of  $F_{MSY}$ , which is a full selection instantaneous  $F$  that will produce MSY at the MSY producing biomass,

$B^* = B_{35\%}$ , a proxy of  $B_{MSY}$ , which is the value of biomass at the MSY producing level,

$\beta$  = a parameter with restriction that  $0 \leq \beta < 1$ . A default value of 0.25 is used.

$\alpha$  = a parameter with restriction that  $0 \leq \alpha \leq \beta$ . A default value of 0.1 is used.

Because trawl bycatch fishing mortality was not related to pot fishing mortality, average trawl bycatch fishing mortality during 2004 to 2013 was used for the per recruit analysis as well as for projections in the next section. Pot female bycatch fishing mortality was set equal to pot male fishing mortality times 0.02, an intermediate level during 1990-2013. Some discards of legal males occurred since the IFQ fishery started in 2005, but the discard rates were much lower during 2007-2013 than in 2005 after the fishing industry minimized discards of legal males. Thus, the average of retained selectivities and discard male selectivities during 2012-2013 were used to represent current trends for per recruit analysis

and projections. Average molting probabilities during 2004-2013 were used for per recruit analysis and projections.

Average recruitments during three periods were used to estimate  $B_{35\%}$ : 1976-1983, 1976-2013, and 1984-2013 (Figure 11). Estimated  $B_{35\%}$  is compared with historical mature male biomass in Figure 13a. We recommend using the average recruitment during 1984-present, corresponding to the 1976/77 regime shift. Note that recruitment period 1984-present has been used since 2011 to set the overfishing limits. Several factors support our recommendation. First, estimated recruitment was lower after 1983 than before 1984, which corresponded to brood years 1978 and later, after the 1976/77 regime shift. Second, high recruitments during the late 1960s and 1970s generally occurred when the spawning stock was primarily located in the southern Bristol Bay, whereas the current spawning stock is mainly in the middle of Bristol Bay. The current flows favor larvae hatched in the southern Bristol Bay (see the section on Ecosystem Considerations for SAFE reports in 2008 and 2009). Finally, stock productivity (recruitment/mature male biomass) was higher before the 1976/1977 regime shift.

If we believe that the productivity differences and differences of other population characteristics before 1978 were caused by fishing, not by the regime shift, then we should use the recruitment from 1976-1983 (corresponding to brood years before 1978) as the baseline to estimate  $B_{35\%}$ . If we believe that the regime shift during 1976/77 caused the productivity differences, then we should select the recruitments from period 1984-2014 as the baseline.

The control rule is used for stock status determination. If total catch exceeds OFL estimated at  $B$ , then “overfishing” occurs. If  $B$  equals or declines below  $0.5 B_{MSY}$  (i.e., MSST), the stock is “overfished.” If  $B$  equals or declines below  $\beta^*B_{MSY}$  or  $\beta^*$ a proxy  $B_{MSY}$ , then the stock productivity is severely depleted and the fishery is closed.

The estimated probability distribution of MMB in 2014 is illustrated in Figure 30. The normal approximation is used to estimate the 49<sup>th</sup> percentile for the OFL in 2014 (Figure 31). Based the SSC suggestion in 2011,  $ABC = 0.9*OFL$  is used to estimate ABC.

Status and catch specifications (1000 t) (scenario 1):

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch	OFL	ABC
2010/11	13.63 <sup>A</sup>	32.64 <sup>A</sup>	6.73	6.76	7.71	10.66	N/A
2011/12	13.77 <sup>B</sup>	30.88 <sup>B</sup>	3.55	3.61	4.09	8.80	7.92
2012/13	13.19 <sup>C</sup>	29.05 <sup>C</sup>	3.56	3.62	3.90	7.96	7.17
2013/14	12.85 <sup>D</sup>	27.12 <sup>D</sup>	3.90	3.99	4.56	7.07	6.36
2014/15		24.69 <sup>D</sup>				6.82	6.14

The stock was above MSST in 2013/14 and is hence not overfished. Overfishing did not occur.

Status and catch specifications (million lbs):

Year	MSST	Biomass (MMB)	TAC	Retained Catch	Total Catch	OFL	ABC
2010/11	30.0 <sup>A</sup>	72.0 <sup>A</sup>	14.84	14.91	17.00	23.52	N/A
2011/12	30.4 <sup>B</sup>	68.1 <sup>B</sup>	7.83	7.95	9.01	19.39	17.46
2012/13	29.1 <sup>C</sup>	64.0 <sup>C</sup>	7.85	7.98	8.59	17.55	15.80
2013/14	28.3 <sup>D</sup>	59.9 <sup>D</sup>	8.60	8.80	10.05	15.58	14.02
2014/15		54.4 <sup>D</sup>				15.04	13.53

Notes:

- A – Calculated from the assessment reviewed by the Crab Plan Team in September 2011
- B – Calculated from the assessment reviewed by the Crab Plan Team in September 2012
- C – Calculated from the assessment reviewed by the Crab Plan Team in September 2013
- D – Calculated from the assessment reviewed by the Crab Plan Team in September 2014

4. Based on the  $B_{35\%}$  estimated from the average male recruitment during 1984-2014, the biological reference points and OFL were estimated as follows:

	Scenario 1		Scenario 1n		Scenario 2	
	1000t	Million lbs	1000t	Million lbs	1000t	Million lbs
$B_{35\%}$	25.703	56.665	26.212	57.788	24.590	54.212
$F_{35\%}$	0.29		0.29		0.29	
MMB <sub>2014</sub>	24.687	54.443	25.622	56.486	21.249	46.846
OFL <sub>2014</sub>	6.820	15.036	7.296	16.085	5.213	11.492
ABC <sub>2014</sub>	6.138	13.532	6.567	14.477	4.691	10.343

5. Based on the 10% buffer rule used last year,  $ABC = 0.9 * OFL$ . If  $P^* = 49\%$  is used, the ABC would be higher.

### G. Rebuilding Analyses

NA.

### H. Data Gaps and Research Priorities

1. The following data gaps exist for this stock:
  - a. Information about changes in natural mortality in the early 1980s;
  - b. Un-observed trawl bycatch in the early 1980s;
  - c. Natural mortality;
  - d. Crab availability to the trawl surveys;
  - e. Juvenile crab abundance;
  - f. Female growth per molt as a function of size and maturity;
  - g. Changes in male molting probability over time.

## 2. Research priorities:

- a. Estimating natural mortality;
- b. Estimating crab availability to the trawl surveys;
- c. Surveying juvenile crab abundance in nearshore;
- d. Studying environmental factors that affect the survival rates from larvae to recruitment.

## ***I. Projections and Future Outlook***

### **1. Projections**

Future population projections primarily depend on future recruitment, but crab recruitment is difficult to predict. Therefore, annual recruitment for the projections was a random selection from estimated recruitments during 1984-2014. Besides recruitment, the other major uncertainty for the projections is estimated abundance in 2014. The 2014 abundance was randomly selected from the estimated normal distribution of the assessment model output for each replicate. Three scenarios of fishing mortality for the directed pot fishery were used in the projections:

- (1) No directed fishery. This was used as a base projection.
- (2)  $F_{40\%}$ . This fishing mortality creates a buffer between the limits and target levels.
- (3)  $F_{35\%}$ . This is the maximum fishing mortality allowed under the current overfishing definitions.

Each scenario was replicated 1000 times and projections made over 10 years beginning in 2014 (Table 7).

As expected, projected mature male biomasses are much higher without the directed fishing mortality than under the other scenarios. At the end of 10 years, projected mature male biomass is above  $B_{35\%}$  for all scenarios (Table 7; Figure 32). Projected retained catch for the  $F_{35\%}$  scenario is higher than those for the  $F_{40\%}$  scenario (Table 7, Figure 33). Due to the poor recruitment during recent years, the projected biomass and retained catch are expected to decline during the next few years.

### **2. Near Future Outlook**

The near future outlook for the Bristol Bay RKC stock is a declining trend. The three recent above-average year classes (hatching years 1990, 1994, and 1997) had entered the legal population by 2006 (Figure 34). Most individuals from the 1997 year class will continue to gain weight to offset loss of the legal biomass to fishing and natural mortalities. The above-average year class (hatching year 2000) with lengths centered around 87.5 mm CL for both males and females in 2006 and with lengths centered around 112.5-117.5 mm CL for males and around 107.5 mm CL for females in 2008 has largely entered the mature male population in 2009 and the legal population by this year (Figure 34). No strong cohorts have been observed in the survey data after this cohort through 2010 (Figure 34). There was a huge tow of juvenile crab of size 45-55 mm in 2011, but these juveniles were not observed during 2012-2014 surveys. This single tow is unlikely to be an indicator for a

strong cohort. The high survey abundance of large males and mature females in 2014 cannot be explained by the survey data during the previous years (Figure 34). Due to lack of recruitment, mature and legal crab should continue to decline next year. Current crab abundance is still low relative to the late 1970s, and without favorable environmental conditions, recovery to the high levels of the late 1970s is unlikely.

### ***J. Acknowledgements***

We thank the Crab Plan Team and Joel Webb for reviewing the earlier draft of this manuscript.

### ***K. Literature Cited***

- Alaska Department of Fish and Game (ADF&G). 2012. Commercial king and Tanner crab fishing regulations, 2012-2013. Alaska Department of Fish and Game, Division of Commercial Fisheries, Juneau. 170 pp.
- Balsiger, J.W. 1974. A computer simulation model for the eastern Bering Sea king crab. Ph.D. dissertation, Univ. Washington, Seattle, WA. 198 pp.
- Fitch, H., M. Deiman, J. Shaishnikoff, and K. Herring. 2012. Annual management report for the commercial shellfish fisheries of the Bering Sea, 2010/11. *In* Fitch, H. M. Schwenzfeier, B. Baechler, T. Hartill, M. Salmon, M. Deiman, E. Evans, E. Henry, L. Wald, J. Shaishnikoff, K. Herring, and J. Wilson. 2012. Annual management report for the commercial and subsistence fisheries of the Aleutian Islands, Bering Sea and the Westward Region's shellfish observer program, 2010/11. Alaska Department of Fish and Game, Fishery Management report No. 12-22, Anchorage.
- Gaeuman, W.G. 2013. Summary of the 2012/13 mandatory crab observer program database for the Bering Sea/Aleutian Islands commercial crab fisheries. Alaska Department of Fish and Game, Fishery Data Series No. 13-54, Anchorage.
- Gray, G.W. 1963. Growth of mature female king crab *Paralithodes camtschaticus* (Tilesius). Alaska Dept. Fish and Game, Inf. Leaflet. 26. 4 pp.
- Griffin, K. L., M. F. Eaton, and R. S. Otto. 1983. An observer program to gather in-season and post-season on-the-grounds red king crab catch data in the southeastern Bering Sea. Contract 82-2, North Pacific Fishery Management Council, 605 West 4<sup>th</sup> Avenue, Suite 306, Anchorage, Alaska 99501. 39 pp.
- Haynes, E.B. 1968. Relation of fecundity and egg length to carapace length in the king crab, *Paralithodes camtschaticus*. Proc. Nat. Shellfish Assoc. 58: 60-62.
- Hoopes, D.T., J.F. Karinen, and M. J. Pelto. 1972. King and Tanner crab research. Int. North Pac. Fish. Comm. Annu. Rep. 1970:110-120.
- Ianelli, J.N., S. Barbeaux, G. Walters, and N. Williamson. 2003. Eastern Bering Sea walleye Pollock stock assessment. Pages 39-126 *In* Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pacific Fishery Management Council, Anchorage.



- Jackson, P.B. 1974. King and Tanner crab fishery of the United States in the Eastern Bering Sea, 1972. *Int. North Pac. Fish. Comm. Annu. Rep.* 1972:90-102.
- Loher, T., D.A. Armstrong, and B.G. Stevens. 2001. Growth of juvenile red king crab (*Paralithodes camtschaticus*) in Bristol Bay (Alaska) elucidated from field sampling and analysis of trawl-survey data. *Fish. Bull.* 99:572-587.
- Matsuura, S., and K. Takeshita. 1990. Longevity of red king crab, *Paralithodes camtschaticus*, revealed by long-term rearing study. In *Proceedings of the International Symposium on King and Tanner Crabs*, pp. 181–188. University Alaska Fairbanks, Alaska Sea Grant College Program Report 90-04, Fairbanks. 633 pp.
- McCaughran, D.A., and G.C. Powell. 1977. Growth model for Alaskan king crab (*Paralithodes camtschaticus*). *J. Fish. Res. Board Can.* 34:989-995.
- North Pacific Fishery Management Council (NPFMC). 2007. Environmental assessment for proposed amendment 24 to the fishery management plan for Bering Sea and Aleutian Islands king and Tanner crabs to revise overfishing definitions. A review draft.
- Otto, R.S. 1989. An overview of eastern Bering Sea king and Tanner crab fisheries. Pages 9–26 *In Proceedings of the International Symposium on King and Tanner Crabs*, Alaska Sea Grant College Program Report No. 90-04.
- Parma, A.M. 1993. Retrospective catch-at-age analysis of Pacific halibut: implications on assessment of harvesting policies. Pages 247-266 *In* G. Kruse, D.M. Eggers, R.J. Marasco, C. Pautzke, and T.J. Quinn II (eds.). *Proceedings of the international symposium on management strategies for exploited fish populations*. University of Alaska Fairbanks, Alaska Sea Grant Rep. 90-04.
- Paul, J.M., and A.J. Paul. 1990. Breeding success of sublegal size male red king crab *Paralithodes camtschaticus* (Tilesius, 1815) (Decapoda, Lithodidae). *J. Shellfish Res.* 9:29-32.
- Paul, J.M., A.J. Paul, R.S. Otto, and R.A. MacIntosh. 1991. Spermatophore presence in relation to carapace length for eastern Bering Sea blue king crab (*Paralithodes platypus*, Brandt, 1850) and red king crab (*P. camtschaticus*, Tilesius, 1815). *Journal of Shellfish research*, Vol. 10, No. 1, 157-163.
- Pengilly, D., S.F. Blau, and J.E. Blackburn. 2002. Size at maturity of Kodiak area female red king crab. Pages 213-224 *In* A.J. Paul, E.G. Dawe, R. Elner, G.S. Jamieson, G.H. Kruse, R.S. Otto, B. Sainte-Marie, T.C. Shirley, and D. Woodby (eds.). *Crabs in Cold Water Regions: Biology, Management, and Economics*. University of Alaska Sea Grant, AK-SG-02-01, Fairbanks.
- Pengilly, D., and D. Schmidt. 1995. Harvest strategy for Kodiak and Bristol Bay red king crab and St. Matthew Island and Pribilof Islands blue king crab. Alaska Dep. Fish and Game, Comm. Fish. Manage. and Dev. Div., Special Publication 7. Juneau, AK. 10 pp.
- Phinney, D.E. 1975. United States fishery for king and Tanner crabs in the eastern Bering Sea, 1973. *Int. North Pac. Fish. Comm. Annu. Rep.* 1973: 98-109.
- Powell, G.C. 1967. Growth of king crabs in the vicinity of Kodiak, Alaska. Alaska Dept. Fish and Game, Inf. Leaflet. 92. 106 pp.

- Powell, G. C., and R.B. Nickerson. 1965. Aggregations among juvenile king crab (*Paralithodes camtschaticus*, Tilesius) Kodiak, Alaska. *Animal Behavior* 13: 374–380.
- Schmidt, D., and D. Pengilly. 1990. Alternative red king crab fishery management practices: modeling the effects of varying size-sex restrictions and harvest rates, p.551-566. *In Proc. Int. Symp. King & Tanner Crabs*, Alaska Sea Grant Rep. 90-04.
- Sparks, A.K., and J.F. Morado. 1985. A preliminary report on diseases of Alaska king crabs, p.333-340. *In Proc. Int. Symp. King & Tanner Crabs*, Alaska Sea Grant Rep. 85-12.
- Stevens, B.G. 1990. Temperature-dependent growth of juvenile red king crab (*Paralithodes camtschaticus*), and its effects on size-at-age and subsequent recruitment in the eastern Bering Sea. *Can. J. Fish. Aquat. Sci.* 47: 1307-1317.
- Stevens, B.G., R.A. MacIntosh, and J.A. Haaga. 1991. Report to industry on the 1991 eastern Bering Sea crab survey. Alaska Fisheries Science Center, Processed Rep. 91-17. 51 pp. NMFS, NOAA, 7600 Sand Point Way NE, Seattle, WA 99115.
- Stevens, B.G., and K. Swiney. 2007. Hatch timing, incubation period, and reproductive cycle for primiparous and multiparous red king crab, *Paralithodes camtschaticus*. *J. Crust. Bio.* 27(1): 37-48.
- Swiney, K. M., W.C. Long, G.L. Eckert, and G.H. Kruse. 2012. Red king crab, *Paralithodes camtschaticus*, size-fecundity relationship, and interannual and seasonal variability in fecundity. *Journal of Shellfish Research*, 31:4, 925-933.
- Webb, J. 2014. Reproductive ecology of commercially important Lithodid crabs. Pages 285-314 *In B.G. Stevens (ed.): King Crabs of the World: Biology and Fisheries Management.* CRC Press, Taylor & Francis Group, New York.
- Weber, D.D. 1967. Growth of the immature king crab *Paralithodes camtschaticus* (Tilesius). *Int. North Pac. Fish. Comm. Bull.* 21:21-53.
- Weber, D.D., and T. Miyahara. 1962. Growth of the adult male king crab, *Paralithodes camtschaticus* (Tilesius). *Fish. Bull. U.S.* 62:53-75.
- Weinberg, K.L., R.S. Otto, and D.A. Somerton. 2004. Capture probability of a survey trawl for red king crab (*Paralithodes camtschaticus*). *Fish. Bull.* 102:740-749.
- Zheng, J. 2005. A review of natural mortality estimation for crab stocks: data-limited for every stock? Pages 595-612 *in G.H. Kruse, V.F. Gallucci, D.E. Hay, R.I. Perry, R.M. Peterman, T.C. Shirley, P.D. Spencer, B. Wilson, and D. Woodby (eds.). Fisheries Assessment and Management in Data-limited Situation.* Alaska Sea Grant College Program, AK-SG-05-02, Fairbanks.
- Zheng, J., and G.H. Kruse. 2002. Retrospective length-based analysis of Bristol Bay red king crabs: model evaluation and management implications. Pages 475-494 *In A.J. Paul, E.G. Dawe, R. Elnor, G.S. Jamieson, G.H. Kruse, R.S. Otto, B. Sainte-Marie, T.C. Shirley, and D. Woodby (eds.). Crabs in Cold Water Regions: Biology, Management, and Economics.* University of Alaska Sea Grant, AK-SG-02-01, Fairbanks.

- Zheng, J., M.C. Murphy, and G.H. Kruse. 1995a. A length-based population model and stock-recruitment relationships for red king crab, *Paralithodes camtschaticus*, in Bristol Bay, Alaska. *Can. J. Fish. Aquat. Sci.* 52:1229-1246.
- Zheng, J., M.C. Murphy, and G.H. Kruse. 1995b. Updated length-based population model and stock-recruitment relationships for red king crab, *Paralithodes camtschaticus*, in Bristol Bay, Alaska. *Alaska Fish. Res. Bull.* 2:114-124.
- Zheng, J., M.C. Murphy, and G.H. Kruse. 1996. Overview of population estimation methods and recommended harvest strategy for red king crabs in Bristol Bay. Alaska Department of Fish and Game, Reg. Inf. Rep. 5J96-04, Juneau, Alaska. 37 pp.
- Zheng, J., M.C. Murphy, and G.H. Kruse. 1997a. Analysis of the harvest strategies for red king crab, *Paralithodes camtschaticus*, in Bristol Bay, Alaska. *Can. J. Fish. Aquat. Sci.* 54:1121-1134.
- Zheng, J., M.C. Murphy, and G.H. Kruse. 1997b. Alternative rebuilding strategies for the red king crab *Paralithodes camtschaticus* fishery in Bristol Bay, Alaska. *J. Shellfish Res.* 16:205-217.

Table 1. Bristol Bay red king crab annual catch and bycatch mortality biomass (t) from June 1 to May 31. A handling mortality rate of 20% for the directed pot, 25% for the Tanner fishery, and 80% for trawl was assumed to estimate bycatch mortality biomass.

Year	Retained Catch			Pot Bycatch		Trawl Bycatch	Tanner Fishery Bycatch	Total Catch
	U.S.	Cost-Recovery	Foreign	Total	Males			
1953	1331.3		4705.6	6036.9				6036.9
1954	1149.9		3720.4	4870.2				4870.2
1955	1029.2		3712.7	4741.9				4741.9
1956	973.4		3572.9	4546.4				4546.4
1957	339.7		3718.1	4057.8				4057.8
1958	3.2		3541.6	3544.8				3544.8
1959	0.0		6062.3	6062.3				6062.3
1960	272.2		12200.7	12472.9				12472.9
1961	193.7		20226.6	20420.3				20420.3
1962	30.8		24618.7	24649.6				24649.6
1963	296.2		24930.8	25227.0				25227.0
1964	373.3		26385.5	26758.8				26758.8
1965	648.2		18730.6	19378.8				19378.8
1966	452.2		19212.4	19664.6				19664.6
1967	1407.0		15257.0	16664.1				16664.1
1968	3939.9		12459.7	16399.6				16399.6
1969	4718.7		6524.0	11242.7				11242.7
1970	3882.3		5889.4	9771.7				9771.7
1971	5872.2		2782.3	8654.5				8654.5
1972	9863.4		2141.0	12004.3				12004.3
1973	12207.8		103.4	12311.2				12311.2
1974	19171.7		215.9	19387.6				19387.6
1975	23281.2		0	23281.2				23281.2
1976	28993.6		0	28993.6			682.8	29676.4
1977	31736.9		0	31736.9			1249.9	32986.8
1978	39743.0		0	39743.0			1320.6	41063.6
1979	48910.0		0	48910.0			1331.9	50241.9
1980	58943.6		0	58943.6			1036.5	59980.1
1981	15236.8		0	15236.8			219.4	15456.2
1982	1361.3		0	1361.3			574.9	1936.2
1983	0.0		0	0.0			420.4	420.4
1984	1897.1		0	1897.1			1094.0	2991.1
1985	1893.8		0	1893.8			390.1	2283.8
1986	5168.2		0	5168.2			200.6	5368.8
1987	5574.2		0	5574.2			186.4	5760.7
1988	3351.1		0	3351.1			597.8	3948.9
1989	4656.0		0	4656.0			174.1	4830.1
1990	9236.2	36.6	0	9272.8	526.9	651.5	247.6	10698.7
1991	7791.8	93.4	0	7885.1	407.8	75.0	316.0	10085.7
1992	3648.2	33.6	0	3681.8	552.0	418.5	335.4	5232.2
1993	6635.4	24.1	0	6659.6	763.2	637.1	426.6	8541.0
1994	0.0	42.3	0	42.3	3.8	1.9	88.9	147.8
1995	0.0	36.4	0	36.4	3.3	1.6	194.2	235.5
1996	3812.7	49.0	0	3861.7	164.6	1.0	106.5	4133.9
1997	3971.9	70.2	0	4042.1	244.7	19.6	73.4	4379.8
1998	6693.8	85.4	0	6779.2	959.7	864.9	159.8	8763.7
1999	5293.5	84.3	0	5377.9	314.2	8.8	201.6	5902.4
2000	3698.8	39.1	0	3737.9	360.8	40.5	100.4	4239.5
2001	3811.5	54.6	0	3866.2	417.9	173.5	164.6	4622.1
2002	4340.9	43.6	0	4384.5	442.7	7.3	155.1	4989.6

2003	7120.0	15.3	0	7135.3	918.9	430.4	172.3	0.0	8656.9
2004	6915.2	91.4	0	7006.7	345.5	187.0	119.6	0.0	7658.8
2005	8305.0	94.7	0	8399.7	1359.5	498.3	155.2	0.0	10412.8
2006	7005.3	137.9	0	7143.2	563.8	37.0	116.7	3.8	7864.4
2007	9237.9	66.1	0	9303.9	1001.3	186.1	138.5	1.8	10631.6
2008	9216.1	0.0	0	9216.1	1165.5	148.4	159.5	4.0	10693.5
2009	7226.9	45.5	0	7272.5	888.1	85.2	103.7	1.6	8351.2
2010	6728.5	33.0	0	6761.5	797.5	122.6	89.0	0.0	7770.7
2011	3553.3	53.8	0	3607.1	395.0	24.0	69.2	0.0	4095.3
2012	3560.6	61.1	0	3621.7	205.2	12.3	62.2	0.0	3901.4
2013	3901.1	89.9	0	3991.0	310.6	99.8	126.8	28.5	4556.6

---

. Annual sample sizes (>64 mm CL) for catch by length and shell condition for retained catch and bycatch of Bristol Bay red king crab.

Year	Trawl Survey		Retained Catch	Pot Bycatch		Trawl Bycatch		Tanner Fishery Bycatch	
	Males	Females		Males	Females	Males	Females	Males	Females
1968	3,684	2,165	18,044						
1969	6,144	4,992	22,812						
1970	1,546	1,216	3,394						
1971			10,340						
1972	1,106	767	15,046						
1973	1,783	1,888	11,848						
1974	2,505	1,800	27,067						
1975	2,943	2,139	29,570						
1976	4,724	2,956	26,450			2,327	676		
1977	3,636	4,178	32,596			14,014	689		
1978	4,132	3,948	27,529			8,983	1,456		
1979	5,807	4,663	27,900			7,228	2,821		
1980	2,412	1,387	34,747			47,463	39,689		
1981	3,478	4,097	18,029			42,172	49,634		
1982	2,063	2,051	11,466			84,240	47,229		
1983	1,524	944	0			204,464	104,910		
1984	2,679	1,942	4,404			357,981	147,134		
1985	792	415	4,582			169,767	30,693		
1986	1,962	367	5,773			1,199	284		
1987	1,168	1,018	4,230			723	927		
1988	1,834	546	9,833			437	275		
1989	1,257	550	32,858			3,147	194		
1990	858	603	7,218	873	699	761	1,570		
1991	1,378	491	36,820	1,801	375	208	396	885	2,198
1992	513	360	23,552	3,248	2,389	214	107	280	685
1993	1,009	534	32,777	5,803	5,942			232	265
1994	443	266	0	0	0	330	247		
1995	2,154	1,718	0	0	0	103	35		
1996	835	816	8,896	230	11	1,025	968		
1997	1,282	707	15,747	4,102	906	1,202	483		
1998	1,097	1,150	16,131	11,079	9,130	1,627	915		
1999	764	540	17,666	1,048	36	2,154	858		
2000	731	1,225	14,091	8,970	1,486	994	671		
2001	611	743	12,854	9,102	4,567	4,393	2,521		
2002	1,032	896	15,932	9,943	302	3,372	1,464		
2003	1,669	1,311	16,212	17,998	10,327	1,568	1,057		
2004	2,871	1,599	20,038	8,258	4,112	1,689	1,506		
2005	1,283	1,682	21,938	55,019	26,775	1,815	1,872		
2006	1,171	2,672	18,027	32,252	3,980	1,481	1,983		
2007	1,219	2,499	22,387	59,769	12,661	1,011	1,097		
2008	1,221	3,352	14,567	49,315	8,488	1,867	1,039		
2009	830	1,857	16,708	52,359	6,041	1,482	870		
2010	705	1,633	20,137	36,654	6,868	734	876		
2011	525	994	10,706	20,629	1,920	600	1,094		
2012	580	707	8,956	7,206	561	1,577	1,770		
2013	633	560	10,197	13,828	6,048	4,681	4,174	218	596
2014	1,106	1,255							

Table 3. Annual retained catch (million crab) and catch per unit effort of the Bristol Bay red king crab fishery.

Year	Japanese Tanglenet		Russian Tanglenet		U.S. Pot/Trawl		Standardized Crab/tan
	Catch	Crab/tan	Catch	Crab/tan	Catch	Crab/Potlift	
1960	1.949	15.2	1.995	10.4	0.088		15.8
1961	3.031	11.8	3.441	8.9	0.062		12.9
1962	4.951	11.3	3.019	7.2	0.010		11.3
1963	5.476	8.5	3.019	5.6	0.101		8.6
1964	5.895	9.2	2.800	4.6	0.123		8.5
1965	4.216	9.3	2.226	3.6	0.223		7.7
1966	4.206	9.4	2.560	4.1	0.140	52	8.1
1967	3.764	8.3	1.592	2.4	0.397	37	6.3
1968	3.853	7.5	0.549	2.3	1.278	27	7.8
1969	2.073	7.2	0.369	1.5	1.749	18	5.6
1970	2.080	7.3	0.320	1.4	1.683	17	5.6
1971	0.886	6.7	0.265	1.3	2.405	20	5.8
1972	0.874	6.7			3.994	19	
1973	0.228				4.826	25	
1974	0.476				7.710	36	
1975					8.745	43	
1976					10.603	33	
1977					11.733	26	
1978					14.746	36	
1979					16.809	53	
1980					20.845	37	
1981					5.308	10	
1982					0.541	4	
1983					0.000		
1984					0.794	7	
1985					0.796	9	
1986					2.100	12	
1987					2.122	10	
1988					1.236	8	
1989					1.685	8	
1990					3.130	12	
1991					2.661	12	
1992					1.208	6	
1993					2.270	9	
1994					0.015		
1995					0.014		
1996					1.264	16	
1997					1.338	15	
1998					2.238	15	
1999					1.923	12	
2000					1.272	12	
2001					1.287	19	
2002					1.484	20	
2003					2.510	18	
2004					2.272	23	
2005					2.763	30	
2006					2.477	31	
2007					3.154	28	
2008					3.064	22	
2009					2.553	21	
2010					2.410	18	
2011					1.298	28	
2012					1.176	30	
2013					1.272	27	

Table 4(1). Summary of statistics for the model (Scenario 1).

**Parameter counts**

Fixed growth parameters	9
Fixed recruitment parameters	2
Fixed length-weight relationship parameters	6
Fixed mortality parameters	4
Fixed survey catchability parameter	1
Fixed high grading parameters	9
Total number of fixed parameters	31
Free survey catchability parameter	1
Free growth parameters	6
Initial abundance (1975)	1
Recruitment-distribution parameters	2
Mean recruitment parameters	1
Male recruitment deviations	40
Female recruitment deviations	40
Natural and fishing mortality parameters	4
Pot male fishing mortality deviations	41
Bycatch mortality from the Tanner crab fishery	8
Pot female bycatch fishing mortality deviations	26
Trawl bycatch fishing mortality deviations	40
Initial (1975) length compositions	35
Free selectivity parameters	22
Total number of free parameters	267
Total number of fixed and free parameters	298

**Negative log likelihood components (see table 4)**

Length compositions---retained catch
Length compositions---pot male discard
Length compositions---pot female discard
Length compositions---survey
Length compositions---trawl discard
Length compositions---Tanner crab discards
Pot discard male biomass
Retained catch biomass
Pot discard female biomass
Trawl discard
Survey biomass
Recruitment variation
Others
Total



Table 4(1n). Summary of statistics for the model (Scenario 1n).

**Parameter counts**

Fixed growth parameters	9
Fixed recruitment parameters	2
Fixed length-weight relationship parameters	6
Fixed mortality parameters	4
Fixed survey catchability parameter	1
Fixed high grading parameters	9
Total number of fixed parameters	31
Free survey catchability parameter	1
Free growth parameters	6
Initial abundance (1975)	1
Recruitment-distribution parameters	2
Mean recruitment parameters	1
Male recruitment deviations	40
Female recruitment deviations	40
Natural and fishing mortality parameters	4
Pot male fishing mortality deviations	41
Bycatch mortality from the Tanner crab fishery	8
Pot female bycatch fishing mortality deviations	26
Trawl bycatch fishing mortality deviations	40
Initial (1975) length compositions	35
Free selectivity parameters	22
Total number of free parameters	267
Total number of fixed and free parameters	298

**Negative log likelihood components (see table 4)**

Length compositions---retained catch
Length compositions---pot male discard
Length compositions---pot female discard
Length compositions---survey
Length compositions---trawl discard
Length compositions---Tanner crab discards
Pot discard male biomass
Retained catch biomass
Pot discard female biomass
Trawl discard
Survey biomass
Recruitment variation
Others
Total

Table 4. Negative log likelihood components for scenarios 1 and 1n and differences in negative log-likelihood components among model scenarios.

	Scenario				
	1	1n	1n - 1	2 - 1	2 - 1n
Negative log likelihood					
R-variation	78.01	80.86	2.84	3.29	0.44
Length-like-retained	-949.47	-951.44	-1.97	-4.21	-2.25
Length-like-discmale	-953.27	-953.31	-0.04	-2.53	-2.49
Length-like-discfemale	-2251.11	-2254.15	-3.04	-26.43	-23.39
Length-like-survey	-44873.70	-44971.70	-98.00	-297.40	-199.40
Length-like-disctrawl	-1966.13	-1967.25	-1.12	15.82	16.94
Length-like-discTanner	-330.79	-330.65	0.14	-4.62	-4.76
Length-like-bsfrfsurvey	-237.30	-237.99	-0.69	3.49	4.17
Catchbio_retained	46.63	46.73	0.09	3.38	3.29
Catchbio_discmale	210.27	216.59	6.32	4.66	-1.65
Catchbio-discfemale	0.14	0.13	0.00	0.00	0.01
Catchbio-disctrawl	0.86	0.91	0.05	-0.02	-0.07
Biomass-trawl survey	85.37	92.91	7.54	2.60	-4.94
Biomass-bsfrfsurvey	-5.42	-4.42	1.00	2.00	1.00
Others	23.62	21.19	-2.43	11.37	13.80
Total	-51122.30	-51211.60	-89.30	-288.60	-199.30
Free parameters	266	266	0	49	49

Table 5(1). Summary of model parameter estimates (scenario 1) for Bristol Bay red king crab. Estimated values and standard deviations. All values are on a log scale. Male recruit is  $\exp(\text{mean}+\text{males})$ , and female recruit is  $\exp(\text{mean}+\text{males}+\text{females})$ .

Year	Recruits				F for Directed Pot Fishery				F for Trawl	
	Females	SD	Males	SD	Males	SD	Females	SD	Estimate	SD
Mean	15.910	0.024	15.910	0.024	-1.970	0.042	0.011	0.001	-5.205	0.064
Limits↑	13,18		13,18		-4.0,0.0		.001,0.1		-8.5,-1.0	
Limits↓	-15,15		-15,15		-15,2.43		-6.0,3.5		-10,10	
1975					1.095	0.102				
1976	-0.387	0.302	0.769	0.133	1.111	0.072			0.154	0.107
1977	0.725	0.133	0.688	0.096	1.137	0.063			0.683	0.105
1978	0.598	0.112	0.907	0.078	1.369	0.057			0.677	0.104
1979	0.311	0.111	1.062	0.075	1.626	0.053			0.704	0.104
1980	0.315	0.106	1.265	0.074	2.405	0.050			0.734	0.104
1981	0.461	0.117	0.624	0.094	2.425	0.007			0.315	0.104
1982	-0.099	0.049	2.244	0.044	0.551	0.047			2.053	0.106
1983	0.028	0.073	1.376	0.050	-10.21	0.709			1.934	0.105
1984	0.414	0.062	1.254	0.045	0.951	0.057			2.908	0.104
1985	0.186	0.157	-0.561	0.103	1.028	0.064			1.834	0.105
1986	0.473	0.058	0.649	0.045	1.479	0.059			0.755	0.104
1987	-0.092	0.136	-0.253	0.072	1.085	0.055			0.444	0.104
1988	0.371	0.166	-1.009	0.108	0.182	0.049			1.425	0.102
1989	0.049	0.148	-0.738	0.083	0.311	0.047			0.021	0.102
1990	-0.071	0.068	0.333	0.045	0.927	0.043	2.101	0.102	0.313	0.102
1991	-0.122	0.095	-0.123	0.055	0.912	0.045	-0.046	0.102	0.653	0.103
1992	-0.427	0.357	-1.790	0.159	0.401	0.046	2.243	0.102	0.834	0.103
1993	-0.278	0.099	-0.347	0.055	1.055	0.048	2.118	0.103	1.097	0.103
1994	-0.134	0.387	-2.124	0.187	-4.085	0.048	1.484	0.130	-0.360	0.104
1995	0.027	0.039	1.197	0.035	-4.429	0.045	1.611	0.135	0.264	0.103
1996	-0.681	0.235	-0.559	0.104	0.119	0.043	-3.612	0.152	-0.450	0.103
1997	-0.759	0.361	-1.347	0.150	0.232	0.043	-0.959	0.104	-0.828	0.103
1998	-0.244	0.119	-0.222	0.067	0.935	0.044	2.109	0.101	-0.097	0.102
1999	0.068	0.058	0.648	0.041	0.491	0.043	-2.023	0.106	0.130	0.102
2000	-0.118	0.139	-0.303	0.079	0.116	0.043	-0.233	0.102	-0.629	0.102
2001	0.788	0.168	-0.935	0.132	0.135	0.042	1.128	0.101	-0.185	0.102
2002	0.252	0.056	1.008	0.042	0.238	0.042	-2.201	0.107	-0.284	0.101
2003	-0.023	0.208	-0.501	0.124	0.751	0.042	1.202	0.101	-0.226	0.101
2004	-0.043	0.140	0.056	0.083	0.610	0.042	0.413	0.101	-0.573	0.102
2005	0.345	0.061	0.952	0.047	1.037	0.043	0.928	0.101	-0.341	0.101
2006	-0.582	0.160	0.271	0.069	0.762	0.043	-1.498	0.103	-0.624	0.102
2007	-0.366	0.148	-0.107	0.077	1.094	0.044	-0.280	0.101	-0.506	0.102
2008	0.124	0.161	-0.708	0.106	1.191	0.047	-0.595	0.102	-0.367	0.103
2009	0.206	0.142	-0.663	0.096	0.903	0.050	-0.828	0.103	-0.806	0.104
2010	-0.040	0.106	-0.116	0.068	0.767	0.053	-0.291	0.103	-0.988	0.105
2011	0.025	0.110	-0.117	0.073	0.090	0.055	-1.213	0.105	-1.230	0.106
2012	-0.112	0.140	-0.308	0.085	-0.015	0.057	-1.749	0.107	-1.349	0.107
2013	-0.548	0.206	-0.516	0.105	0.165	0.061	0.193	0.105	-0.631	0.107
2014	-0.641	0.458	-1.960	0.239						

Table 5(1) (continued). Summary of model parameter estimates for Bristol Bay red king crab (scenario 1). Estimated values and standard deviations. For initial year length composition deviations, the first 20 length groups are for males and the last 16 length groups are for females.

Parameter	Value	SD	Limits	Initial Length Composition 1975			
				Length	Value	SD	Limits
Mm80-84	0.466	0.016	0.184, 1.0	68	1.235	0.095	-5, 5
Mf80-84	0.816	0.020	0.276, 1.5	73	1.266	0.087	-5, 5
Mf76-79,85-93	0.082	0.006	0.0, 0.108	78	0.485	0.111	-5, 5
log_betal, females	0.177	0.055	-0.67, 1.32	83	0.461	0.097	-5, 5
log_betal, males	0.523	0.084	-0.67, 1.32	88	0.421	0.090	-5, 5
log_betar, females	-0.724	0.062	-1.14, 0.5	93	0.115	0.102	-5, 5
log_betar, males	-0.652	0.047	-1.14, 0.5	98	0.141	0.099	-5, 5
Bsfrf_CV	0.941	0.021	0.00, 0.40	103	-0.089	0.114	-5, 5
moltp_slope, 75-78	0.135	0.025	0.01, 0.207	108	-0.032	0.113	-5, 5
moltp_slope, 79-14	0.100	0.004	0.01, 0.207	113	0.083	0.112	-5, 5
log_moltp_L50, 75-78	4.969	0.014	4.47, 5.62	118	-0.066	0.129	-5, 5
log_moltp_L50, 79-14	4.948	0.004	4.47, 5.62	123	-0.081	0.138	-5, 5
log_N75	20.028	0.033	15.0, 21.0	128	-0.065	0.147	-5, 5
log_avg_L50_ret	4.921	0.002	4.78, 5.05	133	-0.120	0.161	-5, 5
ret_fish_slope	0.529	0.032	0.05, 0.70	138	-0.210	0.146	-5, 5
pot disc.males, $\varphi$	-0.328	0.014	-0.40, 0.00	143	-0.310	0.147	-5, 5
pot disc.males, $\kappa$	0.004	0.000	0.0, 0.005	148	-0.465	0.157	-5, 5
pot disc.males, $\gamma$	-0.015	0.001	-0.025, 0.0	153	-0.824	0.192	-5, 5
pot disc.fema., slope	0.240	0.068	0.05, 0.69	158	-1.319	0.258	-5, 5
log_pot disc.fema., L50	4.424	0.019	4.24, 4.61	163	-1.354	0.273	-5, 5
trawl disc slope	0.061	0.003	0.01, 0.20	68	1.661	0.096	-5, 5
log_trawl disc L50	4.974	0.032	4.40, 5.20	73	1.592	0.095	-5, 5
log_srv_L50, m, bsfrf	4.393	0.042	3.59, 5.49	78	1.408	0.094	-5, 5
srv_slope, f, bsfrf	0.015	0.007	0.01, 0.435	83	1.161	0.097	-5, 5
log_srv_L50, f, bsfrf	5.083	0.460	4.09, 5.54	88	1.153	0.088	-5, 5
log_srv_L50, m, 75-81	4.324	0.010	4.09, 5.54	93	0.764	0.101	-5, 5
srv_slope, f, 75-81	0.066	0.004	0.01, 0.33	98	0.480	0.115	-5, 5
log_srv_L50, f, 75-81	4.443	0.018	4.09, 4.70	103	0.398	0.117	-5, 5
log_srv_L50, m, 82-14	4.478	0.008	4.09, 5.10	108	0.203	0.130	-5, 5
srv_slope, f, 82-14	0.062	0.002	0.01, 0.30	113	0.026	0.145	-5, 5
log_srv_L50, f, 82-14	4.517	0.011	4.09, 4.90	118	-0.512	0.215	-5, 5
TC_slope, females	0.365	0.139	0.02, 0.40	123	-0.698	0.261	-5, 5
log_TC_L50, females	4.543	0.015	4.24, 4.90	128	-1.119	0.387	-5, 5
TC_slope, males	0.253	0.111	0.05, 0.90	133	-1.922	0.795	-5, 5
log_TC_L50, males	4.586	0.022	4.25, 5.14	138	-2.354	1.271	-5, 5
log_TC_F, males, 91	-4.116	0.086	-10.0, 1.00	143	NA	NA	
log_TC_F, males, 92	-6.083	0.088	-10.0, 1.00				
log_TC_F, males, 93	-6.807	0.090	-10.0, 1.00	Q	0.941	0.021	0.6, 1.2
log_TC_F, males, 13	-8.202	0.098	-10.0, 1.00				
log_TC_F, females, 91	-2.848	0.086	-10.0, 1.00				
log_TC_F, females, 92	-4.508	0.086	-10.0, 1.00				
log_TC_F, females, 93	-6.407	0.088	-10.0, 1.00				
log_TC_F, females, 13	-7.693	0.084	-10.0, 1.00				

Table 5(1n). Summary of model parameter estimates (scenario 1n) for Bristol Bay red king crab. Estimated values and standard deviations (SD). All values are on a log scale. Male recruit is  $\exp(\text{mean}+\text{males})$ , and female recruit is  $\exp(\text{mean}+\text{males}+\text{females})$ .

Year	Recruits				F for Directed Pot Fishery				F for Trawl	
	Females	SD	Males	SD	Males	SD	Females	SD	Estimate	SD
Mean	15.909	0.026	15.909	0.026	-2.008	0.043	0.012	0.001	-5.265	0.064
Limits↑	13,18		13,18		-4.0,0.0		.001,0.1		-8.5,-1.0	
Limits↓	-15,15		-15,15		-15,2.43		-6.0,3.5		-10,10	
1975					1.135	0.100				
1976	0.134	0.247	0.748	0.142	1.138	0.071			0.172	0.107
1977	0.550	0.158	0.654	0.104	1.136	0.061			0.696	0.105
1978	0.479	0.134	0.871	0.086	1.344	0.056			0.690	0.104
1979	0.745	0.102	1.143	0.077	1.616	0.053			0.724	0.104
1980	0.278	0.115	1.335	0.078	2.413	0.050			0.765	0.105
1981	0.164	0.145	0.512	0.106	2.425	0.007			0.335	0.104
1982	0.004	0.050	2.150	0.050	0.565	0.047			2.055	0.106
1983	-0.045	0.071	1.431	0.051	-10.184	0.705			1.937	0.105
1984	0.416	0.060	1.399	0.053	0.951	0.057			2.906	0.103
1985	0.163	0.182	-0.672	0.123	1.054	0.064			1.853	0.105
1986	0.511	0.059	0.664	0.048	1.577	0.063			0.781	0.105
1987	-0.066	0.136	-0.217	0.074	1.182	0.059			0.464	0.104
1988	0.286	0.167	-0.909	0.108	0.232	0.051			1.436	0.102
1989	0.095	0.152	-0.759	0.090	0.332	0.047			0.027	0.102
1990	-0.090	0.068	0.378	0.046	0.937	0.043	2.038	0.102	0.317	0.102
1991	-0.120	0.095	-0.086	0.057	0.912	0.045	-0.095	0.103	0.652	0.103
1992	-0.346	0.348	-1.826	0.171	0.392	0.046	2.205	0.102	0.825	0.103
1993	-0.307	0.100	-0.311	0.056	1.038	0.049	2.094	0.103	1.083	0.103
1994	-0.099	0.382	-2.189	0.199	-4.104	0.048	1.465	0.130	-0.384	0.104
1995	-0.021	0.039	1.250	0.036	-4.440	0.045	1.583	0.135	0.249	0.103
1996	-0.653	0.235	-0.579	0.114	0.108	0.043	-3.638	0.153	-0.458	0.103
1997	-0.718	0.364	-1.451	0.171	0.217	0.043	-0.973	0.104	-0.839	0.103
1998	-0.311	0.122	-0.187	0.069	0.911	0.044	2.112	0.101	-0.114	0.102
1999	0.036	0.061	0.646	0.044	0.466	0.043	-2.019	0.106	0.107	0.102
2000	-0.105	0.141	-0.318	0.082	0.096	0.042	-0.229	0.102	-0.647	0.102
2001	0.712	0.182	-0.972	0.142	0.118	0.042	1.138	0.101	-0.198	0.102
2002	0.186	0.055	1.085	0.042	0.222	0.042	-2.191	0.107	-0.293	0.101
2003	0.082	0.227	-0.688	0.148	0.747	0.042	1.204	0.102	-0.229	0.101
2004	-0.176	0.148	0.070	0.083	0.608	0.042	0.411	0.101	-0.574	0.102
2005	0.314	0.061	0.987	0.048	1.030	0.043	0.938	0.101	-0.342	0.101
2006	-0.679	0.159	0.367	0.068	0.750	0.043	-1.484	0.103	-0.628	0.102
2007	-0.330	0.156	-0.188	0.086	1.076	0.044	-0.258	0.101	-0.509	0.102
2008	0.055	0.162	-0.660	0.106	1.171	0.048	-0.569	0.102	-0.373	0.103
2009	0.186	0.147	-0.673	0.101	0.877	0.051	-0.796	0.103	-0.815	0.104
2010	-0.089	0.108	-0.094	0.070	0.739	0.054	-0.255	0.103	-0.997	0.105
2011	-0.018	0.112	-0.105	0.075	0.066	0.056	-1.182	0.105	-1.237	0.107
2012	-0.174	0.144	-0.293	0.087	-0.035	0.059	-1.720	0.107	-1.351	0.108
2013	-0.598	0.210	-0.529	0.108	0.147	0.063	0.222	0.105	-0.630	0.108
2014	-0.453	0.430	-1.986	0.243						

Table 5(1n) (continued). Summary of model parameter estimates for Bristol Bay red king crab (scenario 1n). Estimated values and standard deviations. For initial year length composition deviations, the first 20 length groups are for males and the last 16 length groups are for females.

Parameter	Value	SD	Limits	Initial Length Composition 1975			
				Length	Value	SD	Limits
Mm80-84	0.468	0.017	0.184, 1.00	68	1.159	0.103	-5, 5
Mf80-84	0.811	0.021	0.276, 1.50	73	1.182	0.089	-5, 5
Mf76-79,85-93	0.085	0.006	0.0, 0.108	78	0.516	0.108	-5, 5
log_betal, females	0.208	0.055	-0.67, 1.32	83	0.591	0.090	-5, 5
log_betal, males	0.647	0.082	-0.67, 1.32	88	0.401	0.089	-5, 5
log_betar, females	-0.632	0.062	-1.14, 0.50	93	0.209	0.094	-5, 5
log_betar, males	-0.606	0.051	-1.14, 0.50	98	0.215	0.093	-5, 5
Bsfrf_CV	0.067	0.070	0.00, 0.40	103	0.001	0.105	-5, 5
moltp_slope, 75-79	0.134	0.022	0.01, 0.168	108	0.078	0.104	-5, 5
moltp_slope, 80-14	0.103	0.004	0.01, 0.168	113	0.208	0.101	-5, 5
log_moltp_L50, 75-79	4.968	0.013	4.47, 5.52	118	0.008	0.119	-5, 5
log_moltp_L50, 80-14	4.948	0.004	4.47, 5.52	123	0.050	0.124	-5, 5
log_N75	20.008	0.034	15.0, 21.00	128	-0.034	0.140	-5, 5
log_avg_L50_ret	4.921	0.002	4.78, 5.05	133	-0.046	0.149	-5, 5
ret_fish_slope	0.531	0.032	0.05, 0.70	138	-0.144	0.139	-5, 5
pot disc.males, $\phi$	-0.336	0.015	-0.40, 0.00	143	-0.263	0.143	-5, 5
pot disc.males, $\kappa$	0.004	0.000	0.0, 0.005	148	-0.450	0.154	-5, 5
pot disc.males, $\gamma$	-0.016	0.001	-0.025, 0.0	153	-0.791	0.189	-5, 5
pot disc.fema., slope	0.247	0.080	0.05, 0.69	158	-1.322	0.262	-5, 5
log_pot disc.fema., L50	4.420	0.023	4.24, 4.61	163	-1.338	0.275	-5, 5
trawl disc slope	0.063	0.003	0.01, 0.20	68	1.609	0.105	-5, 5
log_trawl disc L50	4.952	0.030	4.40, 5.20	73	1.518	0.102	-5, 5
log_srv_L50, m, bsfrf	4.395	0.045	3.59, 5.49	78	1.491	0.094	-5, 5
srv_slope, f, bsfrf	0.012	0.005	0.01, 0.435	83	1.331	0.093	-5, 5
log_srv_L50, f, bsfrf	5.334	0.508	4.09, 5.54	88	1.287	0.085	-5, 5
log_srv_L50, m, 75-81	4.351	0.011	4.09, 5.54	93	0.829	0.101	-5, 5
srv_slope, f, 75-81	0.069	0.004	0.01, 0.33	98	0.453	0.124	-5, 5
log_srv_L50, f, 75-81	4.488	0.017	4.09, 4.70	103	0.158	0.146	-5, 5
log_srv_L50, m, 82-14	4.488	0.010	4.09, 5.10	108	0.011	0.151	-5, 5
srv_slope, f, 82-14	0.061	0.002	0.01, 0.30	113	-0.246	0.178	-5, 5
log_srv_L50, f, 82-14	4.514	0.012	4.09, 4.90	118	-0.819	0.275	-5, 5
TC_slope, females	0.358	0.137	0.02, 0.40	123	-0.935	0.315	-5, 5
log_TC_L50, females	4.540	0.015	4.24, 4.90	128	-1.207	0.407	-5, 5
TC_slope, males	0.254	0.113	0.05, 0.90	133	-2.122	0.884	-5, 5
log_TC_L50, males	4.584	0.022	4.25, 5.14	138	-2.128	0.970	-5, 5
log_TC_F, males, 91	-4.159	0.087	-10.0, 1.00	143	NA	NA	
log_TC_F, males, 92	-6.130	0.088	-10.0, 1.00				
log_TC_F, males, 93	-6.860	0.090	-10.0, 1.00	Q	0.927	0.021	0.6, 1.2
log_TC_F, males, 13	-8.250	0.100	-10.0, 1.00				
log_TC_F, females, 91	-2.901	0.086	-10.0, 1.00				
log_TC_F, females, 92	-4.549	0.087	-10.0, 1.00				
log_TC_F, females, 93	-6.440	0.088	-10.0, 1.00				
log_TC_F, females, 13	-7.673	0.085	-10.0, 1.00				

Table 6(1). Annual abundance estimates (million crab), mature male biomass (MMB, 1000 t), and total survey biomass estimates (1000 t) for red king crab in Bristol Bay estimated by length-based analysis (scenario 1) from 1975-2014. Mature male biomass for year  $t$  is on Feb. 15, year  $t+1$ . Size measurements are mm CL.

Year (t)	Males				Females	Total Recruits	Trawl Survey Biomass	
	Mature (>119 mm)	Legal (>134 mm)	MMB (>119 mm)	SD MMB	Mature (>89 mm)		Model Est. (>64 mm)	Area-Swept
1975	54.578	29.101	80.681	5.522	87.096		262.081	219.637
1976	58.974	34.876	88.283	4.639	119.084	29.443	299.722	301.454
1977	60.479	36.735	90.315	3.865	146.958	49.546	309.978	380.351
1978	68.517	37.619	94.859	3.195	140.055	56.689	302.629	349.437
1979	66.356	40.385	83.250	2.662	123.888	55.590	279.160	264.248
1980	47.562	34.142	24.829	0.992	112.376	68.209	240.906	244.793
1981	14.798	8.527	8.470	0.467	48.355	39.193	98.759	122.499
1982	7.364	3.179	8.191	0.424	22.398	146.074	54.088	141.610
1983	6.455	3.041	8.450	0.398	14.595	65.238	47.165	49.322
1984	6.256	3.006	6.470	0.374	14.940	71.520	46.437	134.594
1985	8.104	2.555	11.387	0.559	12.865	10.221	38.059	34.281
1986	13.009	5.210	16.878	0.823	18.676	40.484	50.082	47.804
1987	15.798	7.403	22.775	1.011	22.461	12.070	56.678	68.935
1988	16.249	9.472	28.021	1.110	27.399	7.254	60.705	54.056
1989	17.742	11.032	31.516	1.161	25.273	7.967	63.754	61.499
1990	17.938	12.014	29.267	1.176	21.822	21.899	63.656	56.730
1991	14.447	10.744	24.089	1.148	19.801	13.550	57.759	87.499
1992	11.332	8.543	21.825	1.096	19.630	2.243	51.686	37.410
1993	11.893	7.685	19.230	1.066	17.664	10.089	49.836	53.898
1994	11.678	7.072	24.609	1.091	14.622	1.822	44.120	32.099
1995	12.098	8.862	27.272	1.059	14.251	54.545	50.377	38.116
1996	12.119	9.444	25.211	1.004	19.240	6.998	57.839	44.323
1997	11.298	8.497	23.224	0.956	28.350	3.104	62.397	84.653
1998	15.472	8.133	25.293	1.030	26.498	11.610	65.694	84.554
1999	17.017	9.680	29.628	1.129	23.230	32.164	65.324	60.878
2000	15.088	11.016	29.433	1.117	25.645	11.330	67.559	68.429
2001	14.074	10.502	28.274	1.073	29.991	10.209	70.464	52.801
2002	15.826	10.024	30.269	1.069	29.822	50.923	75.011	69.273
2003	16.626	10.917	29.109	1.056	35.181	9.733	79.957	96.781
2004	14.726	10.414	26.924	1.014	42.652	16.826	81.927	96.230
2005	16.789	9.782	26.793	1.023	41.158	50.796	86.913	106.558
2006	16.975	10.179	28.542	1.070	44.998	16.597	89.880	94.914
2007	16.429	10.677	25.731	1.090	52.003	12.364	94.881	103.801
2008	17.867	9.846	26.609	1.224	48.736	8.537	94.469	111.996
2009	18.665	10.510	29.696	1.418	44.176	9.331	91.145	91.784
2010	17.571	11.471	29.582	1.544	40.285	14.184	87.649	78.432
2011	15.095	11.014	29.676	1.591	37.683	14.637	83.116	64.555
2012	13.669	10.545	28.425	1.595	36.494	11.310	81.554	60.801
2013	13.311	9.792	27.157	1.627	35.305	7.649	79.777	61.954
2014	13.404	9.332	24.687	1.346	32.759	1.748	76.295	119.620

Table 6(1n). Annual abundance estimates (million crab), mature male biomass (MMB, 1000 t), and total survey biomass estimates (1000 t) for red king crab in Bristol Bay estimated by length-based analysis (scenario 1n) from 1975-2014. Mature male biomass for year  $t$  is on Feb. 15, year  $t+1$ . Size measurements are mm CL.

Year (t)	Males				Females	Total Recruits	Total Survey Biomass	
	Mature (>119 mm)	Legal (>134mm)	MMB (>119 mm)	SD MMB	Mature (>89 mm)		Model Est. (>64 mm)	Area-Swept (>64 mm)
1975	56.097	29.314	82.519	5.435	75.953		247.299	202.731
1976	61.567	35.659	92.039	4.622	114.509	36.764	284.349	331.868
1977	63.206	38.322	95.229	3.876	143.459	42.660	294.972	375.661
1978	69.519	39.417	98.409	3.210	137.190	50.689	287.328	349.501
1979	65.676	41.054	84.258	2.692	119.652	79.093	265.775	167.627
1980	47.207	33.973	25.266	1.024	109.443	71.558	230.635	249.322
1981	14.770	8.605	8.628	0.484	50.513	29.495	95.162	134.548
1982	7.517	3.223	8.382	0.439	23.495	139.688	53.008	144.797
1983	6.673	3.116	8.714	0.416	14.781	66.453	45.475	49.320
1984	6.283	3.094	6.563	0.388	15.003	82.710	45.288	155.311
1985	7.609	2.548	10.759	0.574	13.693	9.023	37.288	34.535
1986	12.489	4.902	15.916	0.875	20.174	42.047	49.728	48.158
1987	15.942	7.073	22.557	1.071	23.869	12.645	56.663	70.263
1988	16.726	9.478	28.535	1.174	28.724	7.625	60.893	55.372
1989	18.162	11.307	32.221	1.231	26.400	7.980	64.116	55.941
1990	18.345	12.316	30.091	1.255	22.674	22.664	64.221	60.321
1991	14.933	11.062	25.126	1.234	20.521	14.054	58.511	85.055
1992	11.834	8.921	22.950	1.182	20.237	2.233	52.536	37.687
1993	12.399	8.084	20.386	1.156	18.142	10.326	50.704	53.703
1994	12.210	7.469	25.857	1.184	14.998	1.734	45.018	32.335
1995	12.643	9.278	28.570	1.150	14.545	56.060	51.378	38.396
1996	12.622	9.874	26.470	1.093	19.689	6.919	58.710	44.649
1997	11.855	8.904	24.546	1.043	28.582	2.830	63.133	85.277
1998	16.149	8.598	26.821	1.131	26.705	11.669	66.444	85.176
1999	17.781	10.205	31.359	1.243	23.371	31.546	66.127	65.604
2000	15.809	11.601	31.171	1.229	25.628	11.220	68.129	68.342
2001	14.718	11.076	29.904	1.179	29.649	9.327	70.591	53.188
2002	16.311	10.548	31.650	1.172	29.333	52.966	74.956	69.786
2003	16.983	11.322	30.194	1.156	34.685	8.504	79.607	116.794
2004	15.119	10.720	27.964	1.112	42.046	16.005	81.324	131.910
2005	17.396	10.132	28.106	1.136	40.328	51.602	86.229	107.341
2006	17.594	10.636	29.950	1.197	44.071	17.655	89.140	95.676
2007	16.974	11.148	27.080	1.227	51.036	11.560	94.024	104.841
2008	18.466	10.307	28.026	1.384	47.895	8.630	93.723	114.430
2009	19.515	11.010	31.531	1.612	43.358	9.129	90.570	91.673
2010	18.426	12.126	31.542	1.756	39.517	14.152	87.229	81.642
2011	15.821	11.678	31.496	1.803	36.941	14.492	82.717	67.053
2012	14.305	11.136	30.095	1.800	35.736	11.149	81.002	61.248
2013	13.867	10.315	28.662	1.826	34.494	7.415	79.048	62.410
2014	13.902	9.793	25.622	1.500	31.913	1.823	75.451	114.103



Table 7(1). Comparison of projected mature male biomass (1000 t) on Feb. 15, retained catch (1000 t), their 95% limits, and mean fishing mortality with no directed fishery,  $F_{40\%}$ , and  $F_{35\%}$  harvest strategy with  $F_{35\%}$  constraint during 2014-2023. Parameter estimates with scenario 1 are used for the projection.

No Directed Fishery						
Year	MMB	95% LCI	95% UCI	Catch	95% LCI	95% UCI
2014	30.801	27.672	33.757	0.000	0.000	0.000
2015	33.972	30.520	37.231	0.000	0.000	0.000
2016	35.883	32.237	39.326	0.000	0.000	0.000
2017	35.930	32.360	39.554	0.000	0.000	0.000
2018	37.314	32.094	46.758	0.000	0.000	0.000
2019	41.061	32.066	59.383	0.000	0.000	0.000
2020	45.455	32.473	70.820	0.000	0.000	0.000
2021	49.572	32.852	77.604	0.000	0.000	0.000
2022	53.321	33.860	82.818	0.000	0.000	0.000
2023	56.632	34.915	87.299	0.000	0.000	0.000

$F_{40\%}$						
Year	MMB	95% LCI	95% UCI	Catch	95% LCI	95% UCI
2014	25.559	23.316	27.893	5.299	4.403	5.927
2015	24.392	22.516	26.361	4.793	4.031	5.515
2016	22.954	21.356	24.557	4.302	3.681	4.973
2017	20.817	19.476	22.201	3.652	3.172	4.187
2018	20.668	17.629	28.019	3.302	2.639	4.483
2019	22.816	16.692	36.168	3.500	2.244	5.618
2020	25.206	16.495	42.950	4.058	2.108	7.148
2021	26.964	16.659	45.810	4.628	2.092	8.479
2022	28.201	17.340	46.571	5.056	2.190	9.154
2023	29.027	17.619	47.882	5.349	2.377	9.270

$F_{35\%}$						
Year	MMB	95% LCI	95% UCI	Catch	95% LCI	95% UCI
2014	24.731	22.662	26.828	6.134	5.063	7.000
2015	23.192	21.512	24.829	5.222	4.429	6.055
2016	21.603	20.186	22.947	4.552	3.933	5.175
2017	19.455	18.257	20.669	3.797	3.329	4.283
2018	19.310	16.410	26.239	3.435	2.730	4.919
2019	21.369	15.529	33.960	3.707	2.298	6.198
2020	23.565	15.381	40.111	4.360	2.168	7.925
2021	25.082	15.618	42.559	4.982	2.161	9.347
2022	26.076	16.300	42.739	5.427	2.288	9.951
2023	26.696	16.442	43.681	5.707	2.491	10.061

Table 7(1n). Comparison of projected mature male biomass (1000 t) on Feb. 15, retained catch (1000 t), their 95% limits, and mean fishing mortality with no directed fishery,  $F_{40\%}$ , and  $F_{35\%}$  harvest strategy with  $F_{35\%}$  constraint during 2014-2023. Parameter estimates with scenario 1n are used for the projection.

No Directed Fishery						
Year	MMB	95% LCI	95% UCI	Catch	95% LCI	95% UCI
2014	32.174	28.888	35.277	0.000	0.000	0.000
2015	35.217	31.621	38.614	0.000	0.000	0.000
2016	36.960	33.184	40.525	0.000	0.000	0.000
2017	36.885	33.176	40.605	0.000	0.000	0.000
2018	38.255	32.725	48.903	0.000	0.000	0.000
2019	42.103	32.620	61.549	0.000	0.000	0.000
2020	46.674	32.890	72.816	0.000	0.000	0.000
2021	50.949	33.095	82.381	0.000	0.000	0.000
2022	54.825	33.948	87.659	0.000	0.000	0.000
2023	58.259	34.929	91.480	0.000	0.000	0.000

$F_{40\%}$						
Year	MMB	95% LCI	95% UCI	Catch	95% LCI	95% UCI
2014	26.605	24.215	29.095	5.633	4.727	6.253
2015	25.119	23.150	27.236	5.026	4.219	5.748
2016	23.448	21.795	25.139	4.434	3.784	5.152
2017	21.200	19.783	22.670	3.721	3.229	4.279
2018	21.064	17.867	29.103	3.350	2.658	4.601
2019	23.320	16.772	37.949	3.553	2.262	5.815
2020	25.857	16.533	44.603	4.137	2.103	7.450
2021	27.713	16.641	48.433	4.737	2.050	8.915
2022	29.002	17.164	49.163	5.186	2.123	9.651
2023	29.872	17.616	50.491	5.491	2.279	9.778

$F_{35\%}$						
Year	MMB	95% LCI	95% UCI	Catch	95% LCI	95% UCI
2014	25.694	23.519	27.973	6.552	5.430	7.385
2015	23.836	22.098	25.571	5.464	4.627	6.370
2016	22.033	20.583	23.433	4.674	4.035	5.331
2017	19.790	18.530	21.118	3.859	3.380	4.364
2018	19.663	16.584	27.296	3.482	2.744	5.086
2019	21.831	15.619	35.583	3.763	2.293	6.435
2020	24.167	15.463	41.669	4.447	2.163	8.296
2021	25.771	15.550	44.421	5.102	2.119	9.792
2022	26.807	16.099	45.252	5.569	2.212	10.394
2023	27.462	16.574	45.886	5.861	2.372	10.649




Figure 1. Current harvest rate strategy (line) for the Bristol Bay red king crab fishery and annual prohibited species catch (PSC) limits (numbers of crab) of Bristol Bay red king crab in the groundfish fisheries in zone 1 in the eastern Bering Sea. Harvest rates are based on current-year estimates of effective spawning biomass (ESB), whereas PSC limits apply to previous-year ESB.




Figure 2. Retained catch biomass and bycatch mortality biomass (t) for Bristol Bay red king crab from 1953 to 2013. Handling mortality rates were assumed to be 0.2 for the directed pot fishery 0.25 for the Tanner crab fishery and 0.8 for the trawl fisheries.




Figure 3. Comparison of survey legal male abundances and catches per unit effort for Bristol Bay red king crab from 1968 to 2013.




Figure 4. Survey abundances by 5-mm carapace length bin for male Bristol Bay red king crab from 1968 to 2014.




Figure 5. Survey abundances by 5 mm carapace length bin for female Bristol Bay red king crab from 1968 to 2014.





Figure 6. Relationship between implied effective sample sizes (section 3(a)(5)(i)) and effective sample sizes (see effective sample sizes for scenario 1) for length/sex composition data with scenario 1: trawl survey data.






Figure 7. Relationship between implied effective sample sizes (section 3(a)(5)(i)) and effective sample sizes (see effective sample sizes for scenario 1) for length/sex composition data with scenario 1: directed pot fishery data.




Figure 8a(1). Estimated trawl survey selectivities/catchability under scenario 1. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 8a(1n). Estimated trawl survey selectivities/catchability under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 8b. Estimated pot fishery selectivities and groundfish trawl bycatch selectivities under scenario 1. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 9(1). Comparison of estimated probabilities of molting of male red king crab in Bristol Bay for different periods. Molting probabilities for periods 1954-1961 and 1966-1969 were estimated by Balsiger (1974) from tagging data. Molting probabilities for 1975-2014 were estimated with a length-based model with a pot handling mortality rate of 0.2 under scenario 1.




Figure 9(1n). Comparison of estimated probabilities of molting of male red king crab in Bristol Bay for different periods. Molting probabilities for periods 1954-1961 and 1966-1969 were estimated by Balsiger (1974) from tagging data. Molting probabilities for 1975-2014 were estimated with a length-based model with pot handling mortality rate of 0.2 under scenario 1n.




Figure 10a (1 & 2). Comparisons of area-swept estimates of total survey biomass and model prediction for model estimates in 2014 under scenarios 1 and 2. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively. The error bars are plus and minus 2 standard deviations.




Figure 10a (1 & 1n). Comparisons of area-swept estimates of total survey biomass and model prediction for model estimates in 2014 under scenarios 1 and 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively. The error bars are plus and minus 2 standard deviations of the new time series.







Figure 10b (1 & 2). Comparisons of area-swept estimates of male (>119 mm) and female (>89 mm) abundance and model prediction for model estimates in 2014 under scenarios 1 and 2. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.





Figure 10b (1 & 1n). Comparisons of area-swept estimates of mature male (>119 mm) and female (>89 mm) abundance and model prediction for model estimates in 2014 under scenarios 1 and 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 10c. Comparisons of total survey biomass estimates by the BSFRF survey and the model for model estimates in 2014 (scenarios 1, 2 and 1n). The error bars are plus and minus 2 standard deviations of scenario 1.




Figure 10d(1). Estimated BSFRF survey selectivities with scenario 1. The catchability is assumed to be 1.0.




Figure 10d(1n). Estimated BSFRF survey selectivities with scenario 1n. The catchability is assumed to be 1.0.





Figure 10e(1). Comparisons of length compositions by the BSFRF survey and the model estimates in 2007 and 2008 with scenario 1.





Figure 10e(1n). Comparisons of length compositions by the BSFRF survey and the model estimates in 2007 and 2008 with scenario 1n.




Figure 11(1). Estimated recruitment time series during 1976-2014 (occurred year) with scenario 1. Mean male recruits during 1984-2014 was used to estimate  $B_{35\%}$ .




Figure 11(1n). Estimated recruitment time series during 1976-2014 (occurred year) with scenario 1n. Mean male recruits during 1984-2014 was used to estimate  $B_{35\%}$ .






Figure 11(2). Estimated recruitment time series during 1976-2014 (occurred year) with scenario 2. Mean male recruits during 1984-2014 was used to estimate  $B_{35\%}$ .




Figure 12(1). Relationships between full fishing mortalities for the directed pot fishery and mature male biomass on Feb. 15 during 1975-2013 under scenario 1. Average of recruitment from 1984 to 2014 was used to estimate  $B_{MSY}$ . Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 12(1n). Relationships between full fishing mortalities for the directed pot fishery and mature male biomass on Feb. 15 during 1975-2013 under scenario 1n. Average of recruitment from 1984 to 2014 was used to estimate  $B_{MSY}$ . Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 12(2). Relationships between full fishing mortalities for the directed pot fishery and mature male biomass on Feb. 15 during 1975-2013 under scenario 2. Average of recruitment from 1984 to 2014 was used to estimate  $B_{MSY}$ . Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 13a. Relationships between mature male biomass on Feb. 15 and total recruits at age 5 (i.e., 6-year time lag) for Bristol Bay red king crab with pot handling mortality rate to be 0.2 under scenario 1. Numerical labels are years of mating, and the vertical dotted line is the estimated  $B_{35\%}$  based on the mean recruitment level during 1984 to 2014.




Figure 13b. Relationships between log recruitment per mature male biomass and mature male biomass on Feb. 15 for Bristol Bay red king crab with pot handling mortality rate to be 0.2 under scenario 1. Numerical labels are years of mating, and the line is the regression line for data of 1978-2008.




Figure 13c. Time series of log recruitment per mature male biomass and mature male biomass on Feb. 15 for Bristol Bay red king crab with pot handling mortality rate to be 0.2 under scenario 1.




Figure 14. Average clutch fullness and proportion of empty clutches of newshell (shell conditions 1 and 2) mature female crab >89 mm CL from 1975 to 2014 from survey data. Oldshell females were excluded.






Figure 15a. Observed and predicted catch mortality biomass under scenarios 1, 1n and 2. Mortality biomass is equal to caught biomass times a handling mortality rate. Pot handling mortality rate is 0.2.





Figure 15b. Observed and predicted bycatch mortality biomass from trawl fisheries and Tanner crab fishery under scenarios 1, 1n and 2. Mortality biomass is equal to caught biomass times a handling mortality rate. Trawl handling mortality rate is 0.8, and Tanner crab pot handling mortality is 0.25. Trawl bycatch biomass was 0 before 1976.




Figure 16(1). Standardized residuals of total survey biomass under scenario 1. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 16(1n). Standardized residuals of total survey biomass under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 16(2). Standardized residuals of total survey biomass under scenario 2. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 17(1 & 2). Comparison of area-swept and model estimated survey length frequencies of Bristol Bay male red king crab by year under scenarios 1(solid black) and 2(dashed red). Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, and the first length group is 67.5 mm.




Figure 18(1n). Comparison of area-swept and model estimated survey length frequencies of Bristol Bay male red king crab by year under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.




Figure 19(1 & 2). Comparison of area-swept and model estimated survey length frequencies of Bristol Bay female red king crab by year under scenarios 1(solid black) and 2(dashed red). Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.






Figure 19(1n). Comparison of area-swept and model estimated survey length frequencies of Bristol Bay female red king crab by year under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.




Figure 20(1 & 2). Comparison of observed and model estimated retained length frequencies of Bristol Bay male red king crab by year in the directed pot fishery under scenarios 1(solid black) and 2(dashed red). Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 122.5 mm.




Figure 20(1n). Comparison of observed and model estimated retained length frequencies of Bristol Bay male red king crab by year in the directed pot fishery under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 122.5 mm.




Figure 21(1 & 2). Comparison of observer and model estimated discarded length frequencies of Bristol Bay male red king crab by year in the directed pot fishery under scenarios 1(solid black) and 2(dashed red). Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.




Figure 21(1n). Comparison of observer and model estimated discarded length frequencies of Bristol Bay male red king crab by year in the directed pot fishery under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.




Figure 22(1 & 2). Comparison of observer and model estimated discarded length frequencies of Bristol Bay female red king crab by year in the directed pot fishery under scenarios 1(solid black) and 2(dashed red). Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.




Figure 22(1n). Comparison of observer and model estimated discarded length frequencies of Bristol Bay female red king crab by year in the directed pot fishery under scenario 1n. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the first length group is 67.5 mm.




Figure 23(1 & 2). Comparison of observer and model estimated discarded length frequencies of Bristol Bay male red king crab by year in the groundfish trawl fisheries under scenarios 1(solid black) and 2(dashed red). Pot handling mortality rate is 0.2, trawl bycatch mortality rate is 0.8, and the first length group is 67.5 mm.






Figure 23(1n). Comparison of observer and model estimated discarded length frequencies of Bristol Bay male red king crab by year in the groundfish trawl fisheries under scenario 1n. Pot handling mortality rate is 0.2, trawl bycatch mortality rate is 0.8, and the first length group is 67.5 mm.




Figure 24(1 & 2). Comparison of observer and model estimated discarded length frequencies of Bristol Bay female red king crab by year in the groundfish trawl fisheries under scenarios 1(solid black) and 2(dashed red). Pot handling mortality rate is 0.2, trawl bycatch mortality rate is 0.8, and the first length group is 67.5 mm.




Figure 24(1n). Comparison of observer and model estimated discarded length frequencies of Bristol Bay female red king crab by year in the groundfish trawl fisheries under scenario 1n. Pot handling mortality rate is 0.2, trawl bycatch mortality rate is 0.8, and the first length group is 67.5 mm.




Figure 25(1). Standardized residuals of proportions of survey male red king crab under scenario 1. Solid circles are positive residuals, and open circles are negative residuals. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 25(1n). Standardized residuals of proportions of survey male red king crab under scenario 1n. Solid circles are positive residuals, and open circles are negative residuals. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 25(2). Standardized residuals of proportions of survey male red king crab under scenario 2. Solid circles are positive residuals, and open circles are negative residuals. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 26(1). Standardized residuals of proportions of survey female red king crab under scenario 1. Solid circles are positive residuals, and open circles are negative residuals. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 26(1n). Standardized residuals of proportions of survey female red king crab under scenario 1n. Solid circles are positive residuals, and open circles are negative residuals. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.






Figure 26(2). Standardized residuals of proportions of survey female red king crab under scenario 2. Solid circles are positive residuals, and open circles are negative residuals. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.

Figure 27. This figure is an empty space now and the space may be used in the future.





Figure 28a. Comparison of hindcast estimates of mature male biomass on Feb. 15 (top) and total abundance (bottom) of Bristol Bay red king crab from 1975 to 2014 made with terminal years 2008-2014 with scenario 1. These are results of the 2014 model. Legend shows the terminal year. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 28b. Comparison of hindcast estimates of total recruitment for scenario 1 of Bristol Bay red king crab from 1976 to 2014 made with terminal years 2008-2014. These are results of the 2014 model. Legend shows the terminal year. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 29. Comparison of estimates of legal male abundance (top) and mature males (bottom) of Bristol Bay red king crab from 1968 to 2014 made with terminal years 2004-2014 with the base scenarios. Scenario 1 is used for 2014. These are results of historical assessments. Legend shows the year in which the assessment was conducted. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 30. Probability distributions of estimated trawl survey catchability ( $Q$ ) under scenario 1 with the mcmc approach. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 31a. Probability distributions of estimated mature male biomass on Feb. 15, 2015 with  $F_{35\%}$  under scenario 1 with the mcmc approach. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 31b. Probability distributions of the 2014 estimated OFL with scenario 1 with the memc approach. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively.




Figure 32. Projected mature male biomass on Feb. 15 with  $F_{40\%}$  and  $F_{35\%}$  harvest strategy during 2014-2023. Input parameter estimates are based on scenario 1. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the confidence limits are for the  $F_{35\%}$  harvest strategy.






Figure 33. Projected retained catch biomass with  $F_{40\%}$  and  $F_{35\%}$  harvest strategy during 2014-2023. Input parameter estimates are based on scenario 1. Pot and trawl handling mortality rates were assumed to be 0.2 and 0.8, respectively, and the confidence limits are for the  $F_{35\%}$  harvest strategy.




Figure 34. Length frequency distributions of male (top panel) and female (bottom panel) red king crab in Bristol Bay from NMFS trawl surveys during 2010-2014. For purposes of these graphs, abundance estimates are based on area-swept methods.

## Appendix A. Description of the Bristol Bay Red King Crab Model

### a. Model Description

#### i. Population model

The original LBA model was described in detail by Zheng et al. (1995a, 1995b) and Zheng and Kruse (2002). Male crab abundances by carapace length and shell condition in any one year are modeled to result from abundances in the previous year minus catch and handling and natural mortalities, plus recruitment, and additions to or losses from each length class due to growth:

$$N_{l+1,t+1} = \sum_{l'=1}^{l'+l+1} \{P_{l',l+1} [(N_{l',t} + O_{l',t}) e^{-M_t} - (C_{l',t} + D_{l',t}) e^{(y_t-1)M_t} - T_{l',t} e^{(j_t-1)M_t}] m_{l',t}\} + R_{l+1,t+1}, \quad (1)$$

$$O_{l+1,t+1} = [(N_{l+1,t} + O_{l+1,t}) e^{-M_t} - (C_{l+1,t} + D_{l+1,t}) e^{(y_t-J)M_t} - T_{l+1,t} e^{(j_t-1)M_t}] (1 - m_{l+1,t}),$$

where

- $N_{l,t}$  is newshell crab abundance in length class  $l$  and year  $t$ ,
- $O_{l,t}$  is oldshell crab abundances in length class  $l$  and year  $t$ ,
- $M$  is the instantaneous natural mortality,
- $m_{l,t}$  is the molting probability for length class  $l$  and year  $t$ ,
- $R_{l,t}$  is recruitment into length class  $l$  in year  $t$ ,
- $y_t$  is the lag in years between the assessment survey and the mid fishery time in year  $t$ ,
- $j_t$  is the lag in years between the assessment survey and the mid Tanner crab fishery time in year  $t$ ,
- $P_{l',l}$  is the proportion of molting crab growing from length class  $l'$  to  $l$  after one molt,
- $C_{l,t}$  is the retained catch of length class  $l$  in year  $t$ , and
- $D_{l,t}$  is the discarded mortality catch of length class  $l$  in year  $t$ , including directed pot and trawl bycatch,
- $T_{l,t}$  is the discarded mortality catch of length class  $l$  in year  $t$  from the Tanner crab fishery.

The minimum carapace length for males is set at 65 mm, and crab abundance is modeled with a length-class interval of 5 mm. The last length class includes all crab  $\geq 160$ -mm CL. There are 20 length classes/groups.  $P_{l',l}$ ,  $m_l$ ,  $R_{l,t}$ ,  $C_{l,t}$ , and  $D_{l,t}$  are computed as follows:

Mean growth increment per molt is assumed to be a linear function of pre-molt length:

$$G_l = a + b l, \quad (2)$$

where  $a$  and  $b$  are constants. Growth increment per molt is assumed to follow a gamma

distribution:

$$g(x/\alpha_l, \beta) = x^{\alpha_l-1} e^{-x/\beta} / [\beta^{\alpha_l} \Gamma(\alpha_l)]. \quad (3)$$

The expected proportion of molting individuals growing from length class  $l_1$  to length class  $l_2$  after one molt is equal to the sum of probabilities within length range  $[l_1, l_2]$  of the receiving length class  $l_2$  at the beginning of the next year:

$$P_{l_1, l_2} = \int_{l_1}^{l_2} g(x/\alpha_l, \beta) dx, \quad (4)$$

where  $l$  is the mid-length of length class  $l_l$ . For the last length class  $L$ ,  $P_{L, L} = 1$ .

The molting probability for a given length class  $l$  is modeled by an inverse logistic function:

$$m_{l,t} = 1 - \frac{1}{1 + e^{-\beta(l-L_{50})}}, \quad (5)$$

where

$\beta$  and  $L_{50}$  are parameters with three sets of values for three levels of molting probabilities, and  $l$  is the mid-length of length class  $l$ .

Recruitment is defined as recruitment to the model and survey gear rather than recruitment to the fishery. Recruitment is separated into a time-dependent variable,  $R_t$ , and size-dependent variables,  $U_l$ , representing the proportion of recruits belonging to each length class.  $R_t$  is assumed to consist of crab at the recruiting age with different lengths and thus represents year class strength for year  $t$ .  $R_{l,t}$  is computed as

$$R_{l,t} = R_t U_l, \quad (6)$$

where  $U_l$  is described by a gamma distribution similar to equations (3) and (4) with a set of parameters  $\alpha_r$  and  $\beta_r$ . Because of different growth rates, recruitment was estimated separately for males and females under a constraint of approximately equal sex ratios of recruitment over time.

Before 1990, no observed bycatch data were available in the directed pot fishery; the crab that were discarded and died in those years were estimated as the product of handling mortality rate, legal harvest rates, and mean length-specific selectivities. It is difficult to estimate bycatch from the Tanner crab fishery before 1991. A reasonable index to estimate bycatch fishing mortalities is potlifts of the Tanner crab fishery within the distribution area of Bristol Bay red king crab. Thus, bycatch fishing mortalities from the Tanner crab fishery before 1991 were estimated to be proportional to the smoothing average of potlifts east of 163° W. The smoothing average is equal to  $(P_{t-2} + 2P_{t-1} + 3P_t)/6$  for the potlifts in year  $t$ . The smoothing process not only smoothes the annual number of potlifts, it also indexes the effects of lost pots during the previous years. For bycatch, all fishery catch and discard mortality bycatch are estimated as:

$$C_{l,t} \text{ or } D_{l,t} = (N_{l,t} + O_{l,t}) e^{-y_l M_t} (1 - e^{-s_l F_t}) \quad (7)$$

where

$s_l$  is selectivity for retained, pot or trawl discarded mortality catch of length class  $l$ , and

$F_t$  is full fishing mortality of retained, pot or trawl discarded mortality catch in year  $t$ .

For discarded mortality bycatch from the Tanner crab fishery,  $y_t$  is replaced by  $j_t$  in the right side of equation (7).

The female crab model is the same as the male crab model except that the retained catch equals zero, molting probability equals 1.0 to reflect annual molting (Powell 1967), and growth matrix,  $P$ , changes over time due to change in size at maturity for females. The minimum carapace length for females is set at 65 mm, and the last length class includes all crab  $\geq 140$ -mm CL, resulting in length groups 1-16. Three sets of growth increments per molt are used for females due to changes in sizes at maturity over time (Figures A2 and A3).

### ii. Fisheries Selectivities

Retained selectivity, female pot bycatch selectivity, and both male and female trawl bycatch selectivity are estimated as a function of length:

$$s_l = \frac{l}{l + e^{-\beta (l - L_{50})}}, \quad (8)$$

Different sets of parameters ( $\beta, L_{50}$ ) are estimated for retained males, female pot bycatch, male and female trawl bycatch, and discarded males and females from the Tanner crab fishery. Because some catches were from the foreign fisheries during 1968-1972, a different set of parameters ( $\beta, L_{50}$ ) are estimated for retained males for this period and a third parameter, sel\_62.5mm, is used to explain the high proportion of catches in the last length group.

Male pot bycatch selectivity is modeled by two linear functions:

$$\begin{aligned} s_l &= \varphi + \kappa l, \quad \text{if } l < 135 \text{ mm CL,} \\ s_l &= s_{l-1} + 5\gamma, \quad \text{if } l > 134 \text{ mm CL} \end{aligned} \quad (9)$$

Where

$\varphi, \kappa, \gamma$  are parameters.

During 2005-2012, a portion of legal males were also discarded in the pot fishery. The selectivity for this high grading was estimated to be the retained selectivity in each year times a high grading parameter,  $hg_t$ .

### iii. Trawl Survey Selectivities/Catchability

Trawl survey selectivities/catchability are estimated as

$$s_l = \frac{Q}{1 + e^{-\beta (t-L_{50})}}, \quad (10)$$

with different sets of parameters ( $\beta$ ,  $L_{50}$ ) estimated for males and females as well as two different periods (1975-81 and 1982-13). Survey selectivity for the first length group (67.5 mm) was assumed to be the same for both males and females, so only three parameters ( $\beta$ ,  $L_{50}$  for females and  $L_{50}$  for males) were estimated in the model for each of the four periods. Parameter  $Q$  was called the survey catchability that was estimated based on a trawl experiment by Weinberg et al. (2004, Figure A1).  $Q$  was assumed to be constant over time.

Assuming that the BSFRF survey caught all crab within the area-swept, the ratio between NMFS abundance and BSFRF abundance is a capture probability for the NMFS survey net. The Delta method was used to estimate the variance for the capture probability. A maximum likelihood method was used to estimate parameters for a logistic function as an estimated capture probability curve (Figure A1). For a given size, the estimated capture probability is smaller based on the BSFRF survey than from the trawl experiment, but the  $Q$  value is similar between the trawl experiment and the BSFRF surveys (Figure A1). Because many small-sized crab are likely in the shallow water areas that are not accessible for the trawl survey, NMFS survey catchability/selectivity consists of capture probability and crab availability.

**b. Software Used:** AD Model Builder (Otter Research Ltd. 1994).

### c. Likelihood Components

A maximum likelihood approach was used to estimate parameters. For length compositions ( $p_{l,t,s,sh}$ ), the likelihood functions are :

$$Rf = \prod_{l=1}^L \prod_{t=1}^T \prod_{s=1}^2 \prod_{sh=1}^2 \frac{\left\{ \exp \left[ -\frac{(p_{l,t,s,sh} - \hat{p}_{l,t,s,sh})^2}{2\sigma^2} \right] + 0.01 \right\}}{\sqrt{2\pi\sigma^2}}, \quad (11)$$

$$\sigma^2 = \left[ \hat{p}_{l,t,s,sh} (1 - \hat{p}_{l,t,s,sh}) + 0.1/L \right] / n,$$

where

$L$  is the number of length groups,

$T$  is the number of years, and

$n$  is the effective sample size, which was estimated for trawl survey and pot retained catch and bycatch length composition data from the directed pot fishery, and was assumed to be 50 for groundfish trawl and Tanner crab fisheries bycatch length composition data.

The weighted negative log likelihood functions are:

$$\begin{aligned}
\text{Length compositions} &: -\sum \ln(Rf_i), \\
\text{Biomasses other than survey} &: \lambda_j \sum [\ln(C_t / \hat{C}_t)^2], \\
\text{NMFS survey biomass} &: \sum [\ln(B_t / \hat{B}_t)^2 / (2\ln(CV_t^2 + 1))], \\
\text{BSFRF mature males} &: \sum [\ln(\ln(CV_t^2 + 1))^{0.5} + \ln(N_t / \hat{N}_t)^2 / (2\ln(CV_t^2 + 1))], \\
\text{R variation} &: \lambda_R \sum [\ln(R_t / \bar{R})^2], \\
\text{R sex ratio} &: \lambda_s [\ln(\bar{R}_M / \bar{R}_F)^2], \\
\text{Trawl bycatch fishing mortalities} &: \lambda_t [\ln(F_{t,t} / \bar{F}_t)^2], \\
\text{Pot female bycatch fishing mortalities} &: \lambda_p [\ln(F_{t,f} / \bar{F}_f)^2], \\
\text{Trawl survey catchability} &: (Q - \hat{Q})^2 / (2\sigma^2), \\
\text{Scenario 2, each of six growth increment parameters} &: (a - \hat{a})^2 / (2\sigma^2), \\
\text{Scenario 2, penalty for random walk at size of 50\% maturity} &: \lambda_m \delta^2.
\end{aligned} \tag{12}$$

Where

$R_t$  is the recruitment in year  $t$ ,

$\bar{R}$  is the mean recruitment,

$\bar{R}_M$  is the mean male recruitment,

$\bar{R}_F$  is the mean female recruitment,

$\bar{F}_t$  is the mean trawl bycatch fishing mortality,

$\bar{F}_f$  is the mean pot female bycatch fishing mortality,

$Q$  is summer trawl survey catchability,

$\sigma$  is the estimated standard deviation of  $Q$  (all scenarios) or each of six growth increment parameters for scenario 2.

For BSFRF mature male abundance or total survey biomass,  $CV$  is the survey  $CV$  plus  $AV$ , where  $AV$  is additional  $CV$  and estimated in the model. The mature male abundance is used for all scenarios except scenario 2. Total survey biomass is used for scenario 2.

Weights  $\lambda_j$  are assumed to be 500 for retained catch biomass, and 100 for all bycatch biomasses, 2 for recruitment variation, 10 for recruitment sex ratio, 0.2 for pot female bycatch fishing mortality, 0.1 for trawl bycatch fishing mortality, and 0.5 for female maturity (scenario 2). These  $\lambda_j$  values represent prior assumptions about the accuracy of the observed catch biomass data and about the variances of these random variables.

#### d. Population State in Year 1.

The total abundance and proportions for the first year are estimated in the model.

#### e. Parameter estimation framework:

i. Parameters estimated independently

Basic natural mortality, length-weight relationships, and mean growth increments per molt were estimated independently outside of the model. Mean length of recruits to the model depends on growth and was assumed to be 72.5 for both males and females. High grading parameters  $hg_t$  were estimated to be 0.2785 in 2005, 0.0440 in 2006, 0.0197 in 2007, 0.0198 in 2008, 0.0337 in 2009, 0.0153 in 2010, 0.0113 in 2011, and 0.0240 in 2012, based on the proportions of discarded legal males to total caught legal males. Handling mortality rates were set to 0.2 for the directed pot fishery, 0.25 for the Tanner crab fishery, and 0.8 for the trawl fisheries.

##### (1). *Natural Mortality*

Based on an assumed maximum age of 25 years and the 1% rule (Zheng 2005), basic  $M$  was estimated to be 0.18 for both males and females. Natural mortality in a given year,  $M_t$ , equals to  $M + Mm_t$  (for males) or  $M + Mf_t$  (females). One value of  $Mm_t$  during 1980-1985 was estimated and two values of  $Mf_t$  during 1980-1984 and 1976-79, 1985-93 were estimated in the model.

##### (2). *Length-weight Relationship*

Length-weight relationships for males and females were as follows:

$$\begin{aligned} \text{Immature Females: } & W = 0.000408 L^{3.127956}, \\ \text{Ovigerous Females: } & W = 0.003593 L^{2.666076}, \\ \text{Males: } & W = 0.0004031 L^{3.141334}, \end{aligned} \tag{13}$$

where

$W$  is weight in grams, and  
 $L$  is CL in mm.

##### (3). *Growth Increment per Molt*

A variety of data are available to estimate male mean growth increment per molt for Bristol Bay RKC. Tagging studies were conducted during the 1950s, 1960s and 1990s, and mean growth increment per molt data from these tagging studies in the 1950s and 1960s were analyzed by Weber and Miyahara (1962) and Balsiger (1974). Modal analyses were conducted for the data during 1957-1961 and the 1990s (Weber 1967, Loher et al. 2001). Mean growth increment per molt may be a function of body size and shell condition and vary over time (Balsiger 1974, McCaughran and Powell 1977); however, for simplicity, mean growth increment per molt was assumed to be only a function of body size in the models. Tagging data were used to estimate mean growth increment per molt as a function of pre-molt length for males (Figure A2). The results from modal analyses of 1957-1961 and the 1990s were used to estimate mean growth



increment per molt for immature females during 1975-1993 and 1994-2013, respectively, and the data presented in Gray (1963) were used to estimate those for mature females for scenarios 1 and 1n (Figure A2). To make a smooth transition of growth increment per molt from immature to mature females, weighted growth increment averages of 70% and 30% at 92.5 mm CL pre-molt length and 90% and 10% at 97.5 mm CL were used, respectively, for mature and immature females during 1983-1993. These percentages are roughly close to the composition of maturity. During 1975-1982, females matured at a smaller size, so the growth increment per molt as a function of length was shifted to smaller increments. Likewise, during 1994-2013, females matured at a slightly higher size, so the growth increment per molt was shifted to high increments for immature crab (Figure A2). Once mature, the growth increment per molt for male crab decreases slightly and annual molting probability decreases, whereas the growth increment for female crab decreases dramatically but annual molting probability remains constant at 1.0 (Powell 1967).

For females with scenario 2, some new immature female growth data from Kodiak red king crab are used to estimate initial parameter values of immature female growth increments per molt function (Figure A2(2)). Initial parameter values for three growth increments-per-molt functions are estimated using the growth increments per molt data: immature females, mature females, and males. Parameters for growth increments per molt are estimated inside the model with these initial estimates as a prior.

#### **(4). *Sizes at Maturity for Females***

NMFS collected female reproductive condition data during the summer trawl surveys. Mature females are separated from immature females by a presence of egg clutches or egg cases. Proportions of mature females at 5-mm length intervals were summarized and a logistic curve was fitted to the data each year to estimate sizes at 50% maturity. Sizes at 50% maturity are illustrated in Figure A3 with mean values for three different periods (1975-82, 1983-93 and 1994-08).

#### **(5). *Sizes at Maturity for Males***

Although size at sexual maturity for Bristol Bay red king crab males has been estimated (Paul et al. 1991), there are no data for estimating size of functional maturity collected in the natural environment. Sizes at functional maturity for Bristol Bay male RKC have been assumed to be 120 mm CL (Schmidt and Pengilly 1990). This is based on mating pair data collected off Kodiak Island (Figure A4). Sizes at maturity for Bristol Bay female RKC are about 90 mm CL, about 15 mm CL less than Kodiak female RKC (Pengilly et al. 2002). The size ratio of mature males to females is 1.3333 at sizes at maturity for Bristol Bay RKC, and since mature males grow at much larger increments than mature females, the mean size ratio of mature males to females is most likely larger than this ratio. Size ratios of the large majority of Kodiak mating pairs were less than 1.3333, and in some bays, only a small proportion of mating pairs had size ratios above 1.3333 (Figure A4).

In the laboratory, male RKC as small as 80 mm CL from Kodiak and SE Alaska can successfully mate with females (Paul and Paul 1990). But few males less than 100 mm CL were observed to mate with females in the wild. Based on the size ratios of males to

females in the Kodiak mating pair data, setting 120 mm CL as a minimum size of functional maturity for Bristol Bay male RKC is proper in terms of managing the fishery.

**(6). *Potential Reasons for High Mortality during the Early 1980s***

Bristol Bay red king crab abundance had declined sharply during the early 1980s. Many factors have been speculated for this decline: (i) completely wiped out by fishing: the directed pot fishery, the other directed pot fishery (Tanner crab fishery), and bottom trawling; and (ii) high fishing and natural mortality. With the survey abundance, harvest rates in 1980 and 1981 were among the highest, thus the directed fishing definitely had a big impact on the stock decline, especially legal and mature males. However, for the sharp decline during 1980-1984 for males, 3 out of 5 years had low mature harvest rates. During 1981-1984 for females, 3 out of 4 years had low mature harvest rates. Also pot catchability for females and immature males are generally much lower than for legal males, so the directed pot fishing alone cannot explain the sharp decline for all segments of the stock during the early 1980s.

Red king crab bycatch in the eastern Bering Sea Tanner crab fishery is another potential factor (Griffin et al. 1983). The main overlap between Tanner crab and Bristol Bay red king crab is east of 163° W. No absolute red king crab bycatch estimates are available until 1991. So there are insufficient data to fully evaluate the impact. Retained catch and potlifts from the eastern Bering Sea Tanner crab fishery are illustrated in Figure A5. The observed red king crab bycatch in the Tanner crab fishery during 1991-1993 and total potlifts east of 163° W during 1968 to 2005 were used to estimate the bycatch mortality in the current model. Because winter sea surface temperatures and air temperatures were warmer (which means a lower handling mortality rate) and there were fewer potlifts during the early 1980s than during the early 1990s, bycatch in the Tanner crab fishery is unlikely to have been a main factor for the sharp decline of Bristol Bay red king crab.

Several factors may have caused increases in natural mortality. Crab diseases in the early 1980s were documented by Sparks and Morado (1985), but inadequate data were collected to examine their effects on the stock. Stevens (1990) speculated that senescence may be a factor because many crab in the early 1980s were very old due to low temperatures in the 1960s and early 1970s. The biomass of the main crab predator, Pacific cod, increased about 10 times during the late 1970s and early 1980s. Yellowfin sole biomass also increased substantially during this period. Predation is primarily on juvenile and molting/softshell crab. But we lack stomach samples in shallow waters (juvenile habitat) and during the period when red king crab molt. Also cannibalism occurs during molting periods for red king crab. High crab abundance in the late 1970s and early 1980s may have increased the occurrence of cannibalism.

Overall, the likely causes for the sharp decline in the early 1980s are combinations of the above factors, such as pot fisheries on legal males, bycatch and predation on females and juvenile and sublegal males, senescence for older crab, and disease for all crab. In our model, we estimated one mortality parameter for males and another for females during 1980-1984. We also estimated a mortality parameter for females during 1976-1979 and 1985-1993. These three mortality parameters are additional to the basic natural mortality of 0.18, all directed fishing mortality and non-directed fishing mortality. These three

mortality parameters could be attributed to natural mortality as well as undocumented non-directed fishing mortality. The model fit the data much better with these three parameters than without them.

ii. Parameters estimated conditionally

The following model parameters were estimated for male and female crab: total recruits for each year (year class strength  $R_t$  for  $t = 1976$  to 2013), total abundance in the first year (1975), growth parameter  $\beta$  and recruitment parameter  $\beta_r$  for males and females separately. Molting probability parameters  $\beta$  and  $L_{50}$  were also estimated for male crab. Estimated parameters also include  $\beta$  and  $L_{50}$  for retained selectivity,  $\beta$  and  $L_{50}$  for pot-discarded female selectivity,  $\beta$  and  $L_{50}$  for pot-discarded male and female selectivities from the eastern Bering Sea Tanner crab fishery,  $\beta$  and  $L_{50}$  for groundfish trawl discarded selectivity,  $\phi$ ,  $\kappa$  and  $\gamma$  for pot-discarded male selectivity, and  $\beta$  for trawl survey selectivity and  $L_{50}$  for trawl survey male and females separately. NMFS survey catchabilities  $Q$  for some scenarios were also estimated. Three selectivity parameters are estimated for the survey data from the Bering Fisheries Research Foundation. Annual fishing mortalities were also estimated for the directed pot fishery for males (1975-2012), pot-discarded females from the directed fishery (1990-2012), pot-discarded males and females from the eastern Bering Sea Tanner crab fishery (1991-93), and groundfish trawl discarded males and females (1976-2013). Three additional mortality parameters for  $Mm_t$  and  $Mf_t$  were also estimated. Some estimated parameters were constrained in the model. For example, male and female recruitment estimates were forced to be close to each other for a given year.

**f. Definition of model outputs.**

- i. Biomass: two population biomass measurements are used in this report: total survey biomass (crab >64 mm CL) and mature male biomass (males >119 mm CL). Mating time is assumed to Feb. 15.
- ii. Recruitment: new number of males in the 1<sup>st</sup> seven length classes (65- 99 mm CL) and new number of females in the 1<sup>st</sup> five length classes (65-89 mm CL).
- iii. Fishing mortality: full-selected instantaneous fishing mortality rate at the time of fishery.




Figure A1. Estimated capture probabilities for NMFS Bristol Bay red king crab trawl surveys by Weinberg et al. (2004) and the Bering Sea Fisheries Research Foundation surveys.





Figure A2. Mean growth increments per molt for Bristol Bay red king crab. Note: “tagging”---based on tagging data; “mode”---based on modal analysis. The female growth increments per molt are for scenarios 1 and 1n.




Figure A2(2). Mean growth increments per molt for female Bristol Bay red king crab for scenario 2. The slope parameter of the Bristol Bay immature female function is assumed to be the same as that of Kodiak red king crab; Estimated growth increments per molt for mature females <90 mm CL are assumed as the same as that of 90 mm CL.




Figure A3. Estimated sizes at 50% maturity for Bristol Bay female red king crab from 1975 to 2008. Averages for three periods (1975-82, 1983-93, and 1994-08) are plotted with a line.




Figure A4. Histograms of carapace lengths (CL) and CL ratios of males to females for male shell ages  $\leq 13$  months of red king crab males in grasping pairs; Powell's Kodiak data. Upper plot: all locations and years pooled; middle plot: location 11; lower plot: locations 4 and 13. Sizes at maturity for Kodiak red king crab are about 15 mm larger than those for Bristol Bay red king crab. (Source: Doug Pengilly, ADF&G).







Figure A5. Retained catch and potlifts for total eastern Bering Sea Tanner crab fishery (upper plot) and the Tanner crab fishery east of 163° W (bottom).

**Appendix B. Spatial distributions of mature and juvenile male and female red king crab in Bristol Bay from 2013-2014 summer standard trawl surveys.**

