



Methods and criteria to evaluate the effects of fishing on EFH

Proposal from the SSC subcommittee

Background

MSA requires Councils to minimize adverse effects of fishing on EFH
Council is currently considering new definitions of EFH

Model-based definitions of EFH developed at AFSC

- GAM
- Hurdle GAM
- MaxEnt

Fishing Effects model for Alaska fisheries built on LEI and NEFMC SASI model

SSC requested new criteria and methods to evaluate effects of fishing

Questions for Plan Teams

Are the assessment cutoffs correct?

- Core area = upper 50th percentile of predicted abundance or suitable habitat
- Impact threshold for further impact assessment: 10% reduction in habitat
- P-value of 0.1 for significance of correlation with time trend in habitat disturbance in core area
- Evaluating cumulative habitat disturbance across EBS, AI, and GOA
 - Should assessments be based on regional boundaries for the stock or species?
 - E.g., Bristol Bay red king crab

Questions for Plan Teams

- What seasons should be used for the analysis?
 - Summer only – best data, broadest distribution
 - Seasonal based on MaxEnt for non-summer + GAMs for summer
 - Average cumulative impacts over seasons by converting GAMs to MaxEnt for summer

Is SST below
18°C?

Yes:
Recommend
Mitigation

No:
Is CEA reduction
10%?

No:
No Further
Action

Yes:
Significant ($p < 0.1$)
correlation with
life history
parameters?


No:
No Further
Action


Yes:
Elevate for
possible
mitigation

EFH and FE Model Descriptions

The background of the slide features a dark blue sky with the silhouettes of trees and foliage. The silhouettes are black and appear to be the outlines of various tree species, including what looks like a large, spreading tree on the left and some more vertical, leafy branches on the right. The overall aesthetic is clean and professional, with a focus on the text.


Fishing Effects Model






Percentiles of abundance


Core EFH area defined as 50% cumulative distribution




Proportion of habitat reduction




Example map for December 2014



Monthly proportion of habitat reduction (2003-2014)



No area exceeds 5% habitat reduction



Further guidance on what to do with this:

“Any stocks for which the proportion of habitat disturbed by fishing in the CEA is $\geq 10\%$ will be subject to additional analyses.”

“Stock assessment authors will next examine indices of growth-to-maturity, spawning success, breeding success, and feeding success (e.g., time trends in size-at-age, recruitment, spawning distributions and feeding distributions.) to determine whether there are correlations between those parameters and the trends in the proportion of the CEA impacted by fishing. If a correlation exists (negative or positive), the authors will next determine whether the correlation is significant at a p-value of 0.1. A p-value of 0.1 has been recommended to minimize the likelihood of Type II error.”

- It was requested of Martin and I that we do the correlation analysis for GOA pollock and POP as examples

Correlations:

- Proportion of habitat disturbed: Annual values calc'd as average across months (Jan-Dec)
 - pollock: 610-630 (W/CGOA)
 - POP: GOA wide
- Stock indices:
 - Growth-to-maturity: time trends in growth/maturity
 - Spawning success: recruitment
 - Breeding success: spawning distributions
 - Feeding success: feeding distributions

Correlations: pollock


- Growth-to-maturity
 - Growth: weight-at-age anomalies from Shelikof straight acoustic survey, lagged 1 year (habitat impact year prior influences weight the beginning of following year observed in survey)
 - $\rho = 0.12$, p-value > 0.1
 - Maturity: length at age at 50% maturity from Shelikof acoustic survey, lagged 1 year
 - $\rho = 0.61$, p-value > 0.1
- Spawning success: log-recruitment, lagged 1 year
 - $\rho = 0.99$, p-value > 0.1

Correlations: POP

- Growth-to-maturity
 - Growth: mean size-at-age from AFSC bottom trawl survey for most frequent ages (3-15), annual estimates of LVB parameters from bottom trawl survey
 - Maturity: only 2 years of data...
- Spawning success: recruitment, not lagged
- Breeding success/spawning distribution: assume spawning biomass proportional to distribution
- Feeding success/feeding distribution: assume total biomass proportional to distribution

Correlations: POP

- No p-values > 0.1



		ρ	p-value	
Average size-at-age	age-3	-0.49	0.33	
	age-4	-0.25	0.63	
	age-5	-0.56	0.24	
	age-6	-0.58	0.23	
	age-7	-0.20	0.71	
	age-8	-0.71	0.11	
	age-9	-0.25	0.63	
	age-10	-0.60	0.21	
	age-11	0.02	0.97	
	age-12	-0.40	0.43	
	age-13	-0.38	0.46	
	age-14	0.42	0.41	
	age-15	-0.14	0.79	
	LVB params	L_{∞}	0.56	0.33
		κ	-0.64	0.24
t_0		-0.64	0.24	
SAFE output	Spawning biomass	0.43	0.17	
	Total biomass	0.37	0.24	
	Recruitment	0.33	0.30	

Correlations: overall

“The purpose of this criterion is not to determine whether any correlation is statistically significant, but rather to provide an objective threshold to ensure that a “hard look” has been taken for each species, as appropriate. Because multiple parameters will be examined for correlation to habitat reduction, it is possible that spurious significant ($p > 0.1$) correlations will be found.

Whenever significant correlations are found, the expert judgement and opinion of the stock assessment authors will be important to determine whether there is a plausible connection to reductions in EFH as the cause, or if the result is spurious. If stock assessment authors determine that the correlation between the impacts to the CEA and life history parameter(s) suggest a stock effect, then they will raise that potential impact to the attention of the Plan Teams, SSC, and Council.”

- Martin and I took a “hard look”, no significant correlations found, no concerns at this time

Is SST below
18°C?

Yes:
Recommend
Mitigation


No:
No Further Action

No:
Is CEA reduction
10%?

Yes:
Significant ($p < 0.1$)
correlation with
life history
parameters?

No:
No Further
Action

Yes:
Elevate for
possible
mitigation



Hierarchical Impact Assessment

1. MSST – If stock is below MSST, recommend mitigation
2. Core EFH Area – CEA is predicted 50% population quantile from EFH models
 - Evaluating cumulative habitat disturbance across EBS, AI, and GOA
 - **Should assessments be based on regional boundaries for the stock or species?**
 - 50% chosen to avoid missing some important areas without washing out potential impacts if using larger area
 - **Is the 50% threshold the right one?**
 - Continue if CEA habitat reduction is 10% or more
 - **Is 10% habitat reduction threshold reasonable?**

Hierarchical Impact Assessment

3. Correlation of CEA reduction to life history parameters
 - Size at age, recruitment, spawning distribution, feeding distribution, etc.
 - Summer and non-summer

- What seasons should be used for the analysis?
 - Summer only – best data, broadest distribution
 - Seasonal based on MaxEnt for non-summer + GAMs for summer
 - Average cumulative impacts over seasons by converting GAMs to MaxEnt for summer

Hierarchical Impact Assessment

3. Correlation of CEA reduction to life history parameters
 - Are correlations significant at $p < 0.1$
 - Chosen to address possibility of Type II error not to determine significance of correlation
 - Is p-value of 0.1 reasonable?