

Norton Sound Red King Crab Stock Assessment for the fishing year 2023

Toshihide Hamazaki¹
Alaska Department of Fish and Game Commercial Fisheries Division
¹333 Raspberry Rd., Anchorage, AK 99518-1565
Phone: 907-267-2158
Email: Toshihide.Hamazaki@alaska.gov

Executive Summary

1. Stock. Red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska.
2. Catches. This stock supports three important fisheries: summer commercial, winter commercial, and winter subsistence. Of those, the summer commercial fishery accounts for 85% of total harvest. The summer commercial fishery started in 1977. Catch peaked in the late 1970s with retained catch of over 2.9 million pounds. Since 1994, the Norton Sound crab fishery has operated as superexclusive. During the 2022 fishery season, **2,708** crab (**7,6834**lb) were harvested in the winter and **125,042** crab (**317,173** lb) were harvested in the summer commercial fishery. In the winter subsistence fishery **7,565** crab (**15,130** lb) were caught and retained and **2,476** crab (**2,476** lb) were unretained. In total, **135,315** crab (**338,989** lb) were harvested during the 2022 season. Estimated discard mortality ranged from **-272** lb to **199,797** lb. Total fishing mortality ranged from **0.34** to **0.54** million lb where the exceedance of the total catch ABC is dependent on the method used to estimate discard mortality. All the estimates were lower than total catch OFL of 0.67 million lb. Thus, overfishing did not occur during the 2022 season.
3. Stock Biomass. Norton Sound red king crab is monitored not in biomass but in abundance. For the assessment model, biomass is calculated by multiplying the average weight of each length class. Abundance of the Norton Sound red king crab stock has been monitored by trawl surveys since 1976 by NMFS (1976-1991), NOAA NBS (2010-2022), and ADF&G (1996-2021). Historical survey abundance of Norton Sound red king crab of carapace length greater than 63 mm (CL > 63 mm) ranged from 1.41 million to 5.90 million crab. In 2022 abundance of crab estimated from the NOAA NBS 2022 survey was 2.103 million crab with CV 0.37 (Table 3).
4. Recruitment. Recruitment is not monitored directly. It is inferred by the assessment model. Model-estimated recruitment since the 1980s has averaged ~0.70 million, ranging from 0.20 to 1.60 million.

5. Management performance.

Status and catch specifications (million lb)

Year	MSST	Biomass (MMB)	GHL	Retained Catch	Total Catch	OFL	ABC
2018	2.41	4.08	0.30	0.31	0.34	0.43	0.35
2019	2.24	3.12	0.15	0.08	0.08	0.24	0.19
2020	2.28	3.67	0.17	Conf.	Conf.	0.29	0.21
2021	2.25	5.05	0.31	0.007	0.007	0.59	0.35
2022	2.08	5.33	0.34	0.34	TBD	0.67	0.40
2023	2.65	5.29	TBD	TBD	TBD	0.68 1.00	0.41 0.60

Notes:

MSST was calculated as $B_{MSY}/2$

OFL-ABC 2018-2020 are retained only

2019, 2020: Total catch equals retained catch. Discarded catch was estimated only for the summer commercial fishery, but the summer commercial fishery did not occur.

2023 OFL based on $F_{OFL} = 0.18$ / length dependent F_{OFL}

Status and catch specifications (k t)

Year	MSST	Biomass (MMB)	GHL	Retained Catch	Total Catch	OFL	ABC
2018	1.09	1.85	0.13	0.14	0.15	0.20	0.16
2019	1.03	1.41	0.07	0.04	0.04	0.11	0.09
2020	1.04	1.66	0.08	Conf.	Conf.	0.13	0.09
2021	1.02	2.29	0.14	0.003	0.003	0.20	0.16
2022	0.95	2.42	0.15	0.15	TBD	0.31	0.18
2023	1.20	2.40	TBD	TBD	TBD	0.31 0.45	0.19 0.27

Conversion to Metric ton: 1 Metric ton (t) = 2.2046×1000 lb


Biomass (million lb)

Year	Tier	B_{MSY}	Current MMB	B/B_{MSY} (MMB)	F_{OFL}	Years to define B_{MSY}	M	ABC Buffer	ABC
2018	4b	4.82	4.08	0.9	0.15	1980-2018	0.18	0.2	0.35
2019	4b	4.57	3.12	0.7	0.12	1980-2019	0.18	0.2	0.19
2020	4b	4.56	3.66	0.8	0.14	1980-2020	0.18	0.3	0.21
2021	4a	4.53	5.05	1.1	0.18	1980-2021	0.18	0.4	0.35
2022	4a	4.17	5.33	1.3	0.18	1980-2022	0.18	0.4	0.40
2023	4a	4.37	5.29	1.2	0.18 0.18, 0.62	1980-2023	0.18 0.18, 0.62	0.4	0.41 0.60

Biomass in kt

Year	Tier	B _{MSY}	Current MMB	B/B _{MSY} (MMB)	F _{OFL}	Years to define B _{MSY}	M	ABC Buffer	ABC
2018	4b	2.07	1.85	0.9	0.15	1980-2018	0.18	0.2	0.16
2019	4b	2.06	1.41	0.7	0.12	1980-2019	0.18	0.2	0.09
2020	4b	2.07	1.66	0.8	0.14	1980-2020	0.18	0.3	0.09
2021	4a	2.05	2.29	1.1	0.18	1980-2021	0.18	0.4	0.16
2022	4a	1.90	2.42	1.3	0.18	1980-2022	0.18	0.4	0.18
2023	4a	1.98	2.40	1.2	0.18, 0.62	1980-2023	0.18, 0.62	0.4	0.19, 0.27

6. Probability Density Function of the OFL and mcmc estimates



7. The basis for the ABC recommendation.

For Tier 4 stocks, the default maximum ABC is based on $P^*=49\%$ that is essentially identical to the OFL. Annual ABC buffer is determined by accounting for uncertainties in assessment and model results. **However, criteria for determining the level of ABC buffer are undefined.** The buffer was 10% from 2011 to 2014 (ABC = 90% OFL) that was increased to 20% (ABC = 80% OFL) in 2015, to 30% (ABC = 70% OFL) in 2020, and to 40% (ABC = 60% OFL) in 2021.

Year	ABC Buffer
2011-2014	10%
2015-2019	20%
2020	30%
2021-2022	40%

8. Summary of the results of any rebuilding analysis

NA: NSRKC is not overfished.

A. Summary of Major Changes in 2023 assessment model

1. Changes to the management of the fishery.

None.

2. Changes to the input data.

Input data update through 2022:

Winter subsistence, winter and summer commercial crab fishery harvest.

Trawl surveys: abundance, length-shell compositions: NOAA NBS survey in Aug 2022.

3. Changes to the assessment methodology.

Four assessment models are compared for September 2022 draft report, and the CPT-SSC selected the Model 21.0 for the final assessment for Jan 2023.

- a. **Model 21.0: Baseline model adopted in 2022 and for the final assessment 2023**
- b. Model 22.0: Model 21.0 + Shell based retention probability: work on earlier draft
- c. Model 22.1: Model 21.0 + individual M estimate: work on earlier draft
- d. Model 22.2: Model 22.0 + individual M estimate: work on earlier draft

4. Changes to the assessment results.

None.

B. Response to SSC and CPT Comments

Following are SSC, CPT-SSC's requests/review (received in Jan 2022, Oct 2022) and authors' responses, arranged by topic. Requests are italicized.

I. NSRKC Biology-Ecology

Natural Mortality

SSC (Jan 2022): The biological context for natural mortality used in the model is still not well understood.

CPT (Jan 2022): Evaluate the appropriateness of the use of $M=0.18 \text{ yr}^{-1}$ for all red king crab stocks.

Author reply:

Whether or not biological context and justification of $M=0.18$ specification is appropriate for NSRKC as expressed by SSC, this M specification was originally set to be 0.2 for BSAI red king crab stock (NPFMC 1998) and was changed to 0.18 with Amendment 24 (NPFMC 2011). NSRKC assessment model is **REQUIRED** to use $M = 0.18$ unless CPT-SSC grants an exception, such as the CPT-SSC-accepted higher M for crabs of $CL > 123\text{mm}$. Previous model alternatives exploring alternative M specifications (See section E, of history of model progression) and Model 22.1 and 22.2 of this year, suggest higher M could be more appropriate for NSRKC. CPT-SSC have been rejecting higher M specification for NSRKC, except for requesting more of the same analyses. Authors agree with CPT-SSC that biological context and appropriateness of the use of $M=0.18$ needs to be reevaluated not only for NSRKC but also for all RKC stocks. However, we believe that this should be conducted by CPT-SSC, not the NSRKC assessment authors who are non-member of CPT-SSC.

SSC (Jan 2022): Allowing for the estimation of M across all size classes

Author reply:

Alternative models 22.1 and 22.2 were provided, in earlier draft work. As predicted both models showed length-dependent increasing natural mortality with M of near 0 for $CL < 84\text{mm}$ (Figure 3). However, neither model showed any visual improvements of model fits for trawl and commercial length distribution (Figures 8, 9).

SSC (Jan 2022): The authors should consult with regional managers to identify the potential for additional large crab mortality sources.

Author reply:

Contrary to the SSC's assertion, the ADF&G NSRKC stock assessment author works closely with Norton Sound crab research and management biologists. The assessment author is responsible for providing biometric support for design, analyses, and interpretation of ADF&G surveys, researches, and management, including NSRKC trawl survey and tagging studies. The ADF&G NSRKC crab biologists are members of Nome community and are acquainted with many local fishermen and staff of community organizations such as Norton Sound Economic Development Corporation (NSEDC), and Kawerak, exchanging information and research ideas about crab biology and fisheries, including the model assumption of the additional large crab mortality and its source and validity. Thus far, neither the ADF&G nor the NSEDC biologists have identified any large crab mortality sources that the author has examined. NSRKC stock assessment author, research biologists, and managers welcome SSC bringing forward potential additional sources of mortality that can be evaluated.

Growth and Molting Probability

SSC (Jan 2022): Work directly with the scientists conducting the laboratory growth studies so that those data are collected in a way that informs the molting probability function in the stock assessment.

Authors reply:

The authors contacted Dr. Leah Zacher of NOAA AFSC Kodiak for consultation. Dr. Zacher expressed that directly applying lab obtained molting and growth data for assessment model may not be appropriate. NSRKC at the Kodiak lab are kept in water that is warmer than Norton Sound. Diet provided at the lab could be over or less nutritious than that of NSRKC *in situ*. NSRKC spend several months under ice (no light penetration) during winter, while they are not kept under dark condition in the lab. Those environmental difference could affect growth and molt timing that will differ from NSRKC *in situ*.

As for logistics of estimating molting probability and growth increments of the molted, at minimum 20 crab per size category or total of minimum 240 crabs need to be captured and shipped to the lab. The lab can spare a maximum of 2 holding tanks and each can hold a maximum of 20 crabs (total of 40 crabs). Catching 20-30 crabs especially for large crab (e.g., > 123 mm CL) could be challenging. In ideal conditions where all crabs are caught, shipped, healthy, and alive and no tank malfunctions occur, it would take 6 years to investigate.

II. NSRKC Assessment Surveys and Data

Discards

CPT (Jan 2022) Provide a table comparing discard estimates for all three current options (LNR2, subtraction, proportional), any new methods (survey-based), and estimates from the assessment model to better compare methods and identify years of concern.

CPT (Jan 2022) Evaluate how the spatial distribution of catch impacts the ability to estimate discards using Option 2 (survey length-compositions);

CPT-SSC (Oct 2022). Bring forward different methods for estimating discards (without survey data).

Authors reply:

As noted in the Appendix C, estimating of discards for NSRKC has TWO major issues.

1. Estimating discards (2012-2019) from **opportunistic observer surveys that were not intended to estimate discards**
2. Estimating discards (2020- present) **without observer surveys for evaluation of Annual Catch Limit overages.**

OFL and ABC for NSRKC stock has been **retained catch OFL and ABC** until 2020 because of **the lack of observer survey and thus no discards estimates**. In 2020 they were changed to **total catch OFL and ABC** based on 2012-2019 *ad hoc* observer surveys that was **not intended to estimate discards**. In the same year of 2020, **the ad hoc observer survey was terminated, and it is unlikely that observer survey will be resumed in the near future**. Given that OFL and ABC are total catch, CPT-SSC requested to bring forward methods for estimating discards **without observer survey for the purpose of determining annual catch limit (ACL) overage.**

Table: history of OFL-ABC designation for NSRKC.

Year	Discards survey	Discards/Total (retained + discards) Catch	OFL/ABC	Rationale
1997-2011	No survey	NA	Retained	No total catch = retained OFL
2012-2019	Opportunistic	<i>ad hoc</i>	Retained	No total catch = retained OFL
2020	No survey	NA	Total	Set total catch OFL given that 2012-2019 are available
2021- 2022	No survey	NA	Total	Given that total catch OFL is set, discards-total catch needs to be estimated without survey data for determination of ACL overage.

Author has provided various discards estimates (with survey data) since 2018 (Table 4, 5 Appendix C), and discards estimates without observer survey data since 2020 as requested by CPT-SSC (See Appendix C). However, CPT-SSC has not discussed or presented any **objective criteria** for selecting a method, which led no decisions for selecting the most appropriate method for NSRKC.


Regarding method for estimating discards without observer survey data, the author provided 21 methods and their estimates (Appendix C Table 12). Regarding the trawl survey method (Option 2), the majority of trawl length-composition sample came from area outside of commercial fishery grounds (See Figures A and B, Section D). This raises questions about representativeness of trawl survey length-shell composition, not only in this context of estimating discards, but also for using the data in the assessment model.

Regardless of a method selected for NSRKC, the fundamental question of jurisprudence needs to be addressed: **Should overage of the ACL be determined without data?** The discards mortality biomass estimated here is used **solely for determination of overage of the annual catch limit (ACL) that significant regulatory consequences.** The opportunistic NSRKC observer survey was temporary and would not return for foreseeable future. The jurisprudence applied for NSRKC will be a precedence that can be applied to other BSAI crab fisheries that termination of observer survey will be in consideration.

CPT(Jan 2022) Re-examine the evidence for shell condition-specific discards rates and evaluate their implications for the assessment model (e.g., would this affect the overestimation of large crab).

Authors reply:

The following figure is an observed proportion of crab retained (retention probability) by shell condition from 2012-2019 observer data. The figure shows that oldshell crabs are less likely to be retained than newshell crab. An alternative model 22.0 includes separate retention probability for newshell and oldshell crab.



VAST

CPT (Jan 2022) Updates on the use of VAST for survey abundance estimation

Authors reply:

Dr. Jon Richar of NOAA received an approval from his supervisor to work on a VAST model for the Norton Sound trawl surveys. For the 2022 assessment, NOAA assigned VAST estimate of Norton Sound trawl survey to medium priority (per Dr. William Stockhausen NOAA). We look forward to his progress.

GMACS

CPT (Jan 2022) Develop a GMACS version of the NSRKC assessment model

SSC (Oct 2022) Provide an update on GMACS development for NSRKC stock for October 2023.

Authors reply:

Dr. Andre Punt and Matthieu Veron of the University of Washington are working towards developing GMACS version of the NSRKC assessment model. The author looks forward to their further development.

Standardized CPUE

CPT (Jan 2022) Calculate the arithmetic scale CV for lognormally-distributed data

Authors reply:

Table 1 includes arithmetic scale CV calculated as $CV = \sqrt{\exp(SE^2 - 1)}$

III. NSRKC Assessment model

Include a table (if necessary) identifying any parameters estimated at a bound.

Authors reply:

The NSRKC Assessment model has two parameters that are estimated at a bound.: Trawl survey selectivity ($\log_{\phi_{stl}}$), and parameters (r1, r2) that estimate recruit size proportion.

1. Trawl Survey selectivity parameter

Trawl survey selectivity model is a one parameter logistic curve that reaches 1.0 at L_{max} (143.5 mm)

$$S_l = \frac{1}{1 + e^{(\alpha(L_{max}-L)+\ln(1/0.999-1))}}$$

where $\alpha = \exp(\log_{\phi_{stl}})$, $L_{max} = 143.5$ mm L (63.5-143.5mm)

Model estimated trawl survey selectivity is 1.0 across all size classes, or $e^{(\alpha(L_{max}-L)+\ln(1/0.999-1))} \approx 0$ for all lengths, or $\alpha(L_{max}-L)+\ln(1/0.999-1) = -\infty$ and $\alpha \approx 0$ or $\log_{\phi_{stl}} = -\infty$. Thus, parameter will hit the boundary.

Alternative option is specifying $S_l = 1.0$ for all length classes; however, this also removes the model's ability to estimate S_l when all length classes are *not* 1.0.

2. The proportion of recruits

The proportion of recruits is a multinomial formula of $n = 3$

$$p_l = \frac{\exp(r_l)}{1 + \sum_{l=1}^{n-1} \exp(r_l)} \text{ for } l = 1, \dots, n-1$$

$$p_n = 1 - \frac{\sum_{l=1}^{n-1} \exp(r_l)}{1 + \sum_{l=1}^{n-1} \exp(r_l)}$$

Model estimated recruit size proportions P_1 , P_2 , and P_3 are 0.5905, 0.4055, and 0.0040, respectively. This makes it extremely difficult for the model to estimate P_1 and P_2 . Alternative option is assuming $P_3 = 0$; however, this also removes the model's ability to estimate P_3 when P_3 is far greater than 0.

Those parameters hitting boundaries would have little influence on the model. **We welcome CPT-SSC's suggestion for an alternative parameterization form that could avoid parameters hitting the boundaries.**

SSC(Oct 2022), Explore an inverse logistic (or other descending right limb pattern) selectivity to assess whether we may be missing larger crabs, due to either movement or gear selectivity.

Authors reply:

Author notes that this is a recurring request made by the SSC that have been explored since 2017. Please refer to the section E-1 for details.

CPT (Sept 2022) Place a prior on M in the smaller size bin in future explorations of models 22.1 and 22.2 to keep the M estimate above 0.

Authors reply:

Author did explore this but ultimately did not bring forward for review. When minimum M is set to 0.18, M of the small crab sizes hit the boundary of 0.18 with no discernible difference in likelihoods and projections from models 22.1 and 22.2. Those two models were presented not as alternative assessment model, but to evaluate: 1) whether model suggest length-dependent M,

and 2) whether the length-dependent M would improve model fit to observed length-shell composition. The model exercise answered Yes to the first, and No to the second questions.

The assessment model creates abundance of smaller size classes (1, 2, 3) by mortality and growth and recruits with time invariant size proportions. Contribution of recruit is 100% for the smallest size class (64-73mm) and 80-95% for the second smallest size classes (74-83mm). The assessment model is not able to separate contribution from growth and mortality and that from recruit without setting a strict limit to either minimum M or the maximum recruit. The author see no value for setting minimum M to 0.18.

CPT (Jan 2022) Provide a brief description and discussion of the convergence criteria and other methods used to evaluate model convergence to the global minimum. Use jittering to evaluate convergence

Authors reply:

We set ADMB convergence criteria to $1e-6$, though this does not necessarily guarantee model convergence to the global minimum. Of 100 jittering, 7 did not reach to convergence. Median likelihood and gradient were 354.2 and $2.80 e-6$.

Results of Jittering are provided in Table 14.

IV. NSRKC Management

OFL

CPT (Jan 2022) Present an evaluation of Tier 4 OFL calculations for NSRKC using the standard (single M) and length-dependent M approaches, and the associated assumptions and tradeoffs, for consideration by the CPT and SSC in the fall prior to the 2023 assessment.

SSC (Oct 2022) The SSC requests that the methodology for calculating FOFL based on a length-based M as requested by the CPT be clearly documented with supporting rationale.

Authors reply:

Standard calculation of OFL for Tier 4 crab is $F_{OFL,l} = \gamma M_l = \gamma 0.18$ for all lengths. However, this is not strictly applicable for NSRKC that has length-dependent M . The length-dependent M and F_{OFL} was recommended by CPT-SSC in January 2017 but was rejected primarily because it would produce higher OFL. As a compromise, OFL for NSRKC stock in the above equations

was specified to length-independent $F_{OFL,l} = \gamma M_l = \gamma 0.18$ and length-dependent M_l for allocation of winter and summer fisheries.

This compromise works only when M is prespecified (e.g., $M = 0.18$) for the assessment model. However, when the biological justification of prespecified M is in question (this has been raised by CPT-SSC since 2019) and is estimated inside an assessment model, biological-scientific justification for selecting M for F_{OFL} needs to be clarified and specified. For instance, in 2021 Model 21.5 estimated two M s for lengths 64-123mm, and 124mm and above. In this case, should M for lengths 64-123mm be used for F_{OFL} specification? This is further complicated when M of each length is estimated within the model, such as Models 22.1 and 22.2 in which estimated M ranged from 0 to 0.7. Out of the 8 length-dependent M s, which M should be selected for F_{OFL} and what are the biological and scientific justifications? Regardless of which model is selected, a protocol for selecting a single M should be discussed and determined if size-independent M for F_{OFL} is preferred.

Length-dependent M and F_{OFL} would be more logical as this is a simple extension of the concept of the Tier 4 OFL specification. The biggest concerns raised by CPT-SSC were that length-dependent M for NSRKC would increase OFL from standard OFL , though the opposite could occur if estimated M were LOWER than 0.18. Recognizing scientific uncertainties in the determination of OFL , the FMP included the ABC Control Rule *“for setting the maximum permissible ABC for each stock as a function of the scientific uncertainty in the estimate of OFL and any other specified scientific uncertainty.”* The ABC-control rule allows CPT-SSC setting ABC-buffer they deem to be appropriate for each stock based on scientific uncertainties of OFL . If it is a concern that a length-dependent M increases the OFL , CPT-SSC should increase the ABC-buffer to a level that deems the most appropriate for NSRKC stock as CPT-SSC has increased ABC buffer of NSRKC from 10% in 2011 to 40% in 2021-2022. As an example, increasing ABC buffer to 60% for length-dependent OFL ($OFL-l$) is similar to ABC catch level as standard OFL .

	OFLst	ABCst- buffer	ABCst	OFL-l	ABC-l- buffer	ABC-l
21.0	0.70	40%	0.42	1.03	60%	0.41
22.0	0.67	40%	0.40	1.00	60%	0.40
22.1	0.63	40%	0.38	1.10	60%	0.44
22.2	0.63	40%	0.38	1.09	60%	0.44

The author recommends that CPT-SSC adopt length-dependent OFL and apply ABC control rule (increase ABC-buffer) that it deems appropriate for NSRKC stock.

The formula for calculation of a length-based M OFL has been noted in the section F.

C. Introduction

1. Species:

Red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska.

2. General Distribution:

Norton Sound red king crab (NSRKC) is one of the northernmost red king crab populations that can support a commercial fishery (Powell et al. 1983). It is distributed throughout Norton Sound with a westward limit of 167-168° W. longitude, depths less than 30 m, and summer bottom temperatures above 4° C. The Norton Sound red king crab management area consists of two units: Norton Sound Section (Q3) and Kotzebue Section (Q4) (Menard et al. 2011). The Norton Sound Section (Q3) consists of all waters in Registration Area Q north of the latitude of Cape Romanzof, east of the International Dateline, and south of 66°N latitude (Figure 1). The Kotzebue Section (Q4) lies immediately north of the Norton Sound Section and includes Kotzebue Sound. Commercial fisheries have not occurred regularly in the Kotzebue Section. This report deals with the Norton Sound Section of the NSRKC management area.

3. Evidence of stock structure:

Based on variability at 15 SNP loci and in mtDNA sequences (COI, 665 bp), NSRK stock belongs to the Okhotsk Sea–Norton Sound–Aleutian Islands evolutionary lineage (SNPs, FCT = 0.054; mtDNA FCT = 0.222) (Grant and Chen 2012). However, this does not indicate that NSRKC is a single stock. The study indicates it was incapable of detecting possible evolutionary stock differences within the NSRKC stock. No studies have investigated possible stock separation within the Norton Sound management area (Figure 1).

4. Life history characteristics relevant to management:

One of the unique life-history traits of NSRKC is that they spend their entire lives in shallow water since Norton Sound is generally less than 40 m in depth. Based on the 1976-2021 trawl surveys, NSRKC are found in areas with a mean depth range of 19 ± 6 (SD) m and bottom temperatures of $7.4^\circ \pm 2.5$ (SD) C during summer. NSRKC are consistently abundant offshore of Nome.

NSRKC migrate between deep offshore and shallow inshore waters within Norton Sound. Timing of the inshore mating migration is unknown but is assumed to be during late fall to winter (Powell et al. 1983). Offshore migration occurs in late May - July (Bell et al. 2016). The results from a study funded by North Pacific Research Board (NPRB) during 2012-2014 suggest that older/large crab (> 104 mm CL) may stay offshore in the winter, based on findings that large crab are not found nearshore during spring offshore migration periods (Jenefer Bell, ADF&G, personal communication). Molting is thought to occur in fall: late August – November for male and Jan-March for female based on laboratory observation (Leah Zacher and Jennifer Gardner NOAA-AFSC *personal comm*). Trawl surveys show that crab distributions are patchy and dynamic. Functional maturity of NSRKC male crab is as small as 79.4 mm CL (Leah Zacher NOAA Kodiak *personal comm*). Those small males could also fertilize eggs of ~ 4 females, which was comparable to the number of females larger (94-116 mm CL) crabs could fertilize. More interestingly, the largest crab (> 123 mm) was able to fertilize eggs of ~ 2 females.

5. Brief management history:

NSRKC fisheries consist of commercial and subsistence fisheries. The commercial red king crab occurs in summer (June – August) and winter (December – May), while subsistence is open year-round. The majority of NSRKC is harvested during the offshore summer commercial fishery, whereas the winter commercial and subsistence fisheries occur nearshore through ice and take a much smaller harvest.

The distinguishing characteristic of the NSRKC fisheries is that all fisheries, researches, and management are conducted by local residents of Norton Sound. Commercial fisheries is designated as super-exclusive: a vessel registered for the Norton Sound crab fishery may not be used to take king crabs in any other registration areas. The ADF&G NSRKC crab research and management biologists are members of Nome community and are acquainted with many local fishermen (commercial and subsistence) and staff of community organizations such as Norton Sound Economic Development Corporation (NSED) and Kawerak Inc, exchanging information and research ideas about crab biology and fisheries management.

Summer Commercial Fishery

A large-vessel summer commercial crab fishery started in 1977 in the Norton Sound Section (Table 1) and continued from 1977 through 1990. No summer commercial fishery occurred in 1991 because there were no staff to manage the fishery. In March 1993, the Alaska Board of Fisheries (BOF) limited participation in the fishery to small boats. Then on June 27, 1994, a super-exclusive designation went into effect for the fishery. This designation states that a vessel registered for the Norton Sound crab fishery may not be used to take king crabs in any other registration areas during that registration year. A vessel moratorium was put into place before the 1996 season. This was intended to precede a license limitation program. In 1998, Community Development Quota (CDQ) groups were allocated a portion of the summer harvest; however, no CDQ harvest occurred until the 2000 season. On January 1, 2000, the North Pacific License Limitation Program (LLP) went into effect for the Norton Sound crab fishery. The program dictates that a vessel which exceeds 32 feet in length overall must hold a valid crab license issued under the LLP by the National Marine Fisheries Service. Changes in regulations and the location of buyers resulted in eastward movement of the harvest distribution in Norton Sound in mid-1990s. In Norton Sound, a legal crab is defined as $\geq 4 \frac{3}{4}$ -inch carapace width (CW, Menard et al. 2011), which is approximately equivalent to ≥ 104 mm carapace length (CL). In 2005 and 2006, commercial buyers, specifically Norton Sound Economic Development Corporation (NSED) accepted only legal crab of ≥ 5 inch CW. This preference became permanent in 2008.

Portions of the Norton Sound area are closed to commercial fishing for red king crab. Since the beginning of the commercial fisheries in 1977, waters approximately 5-10 miles offshore of southern Seward Peninsula from Port Clarence to St. Michael have been closed to protect nearshore subsistence fisheries and to act as a refuge for crab during the summer commercial crab fishery (Figure 2). The spatial extent of closed waters has varied historically, with the

closure line being moved in to provide additional area to achieve harvest goals. In 2020 the BOF closed Norton Sound area east of 167 degrees W. longitude for commercial summer crab fishery. In 2020 and 2021 NSEDC did not purchase NSRKC resulting in small or no harvest. In 2022, the summer commercial fishery resumed.

CDQ Fishery

The Norton Sound and Lower Yukon CDQ groups divide the NSRKC CDQ allocation. Only fishers designated by the Norton Sound and Lower Yukon CDQ groups are allowed to participate in this portion of the king crab fishery. Fishers are required to have a CDQ fishing permit from the Commercial Fisheries Entry Commission (CFEC) and register their vessel with the Alaska Department of Fish and Game (ADF&G) before beginning fishing. Fishers operate under the authority of each CDQ group. CDQ harvest share is 7.5% of the guideline harvest level (GHL), and can be prosecuted in both summer and winter fisheries season.

Winter Commercial Fishery

The winter commercial crab fishery uses hand lines and pots through the nearshore ice. On average 10 permit holders harvested 2,500 crab during 1978-2009. From 2007 to 2015 the winter commercial catch increased from 3,000 crab to over 40,000 (Table 2). In 2015 the winter commercial catch reached 20% of total crab catch. The BOF responded in May 2015 by amending regulations to allocate 8% of the total commercial GHL to the winter commercial fishery, which has been in effect since the 2017 season. The timing of the winter red king crab commercial fishing season has changed over time to address ice stability. It was originally from January 1 to April 30, amended in 1985 to from November 15 to May 15. In 2015 the period was changed to from January 15 to April 30 after fisheries opened in Nov 15 in 2014, so that January 15 starting date was into effect in 2016. In 2021 it was further amended to from February 1 to April 30.

Year	Opening period
1977-1984	Jan 01 – Apr 30
1985- 2014	Nov 15 – May 15
2015	Nov 15 – Apr 30
2016-2020	Jan 15 – Apr 30
2021 -	Feb 01 – Apr 30

Subsistence Fishery

The winter subsistence fishery has a long history; however, harvest information is available only since the 1977/78 season. The majority of subsistence crab harvest occurs in winter using hand lines and pots through nearshore ice. Average annual winter subsistence harvest is 5,281 crab (1977-2021). Subsistence harvesters need to obtain a permit before fishing and record daily effort and catch. There are no size or sex specific harvest limits; however, the majority of retained catch is males of near legal size.

Summer subsistence crab fishery harvest has been monitored since 2004 with an average harvest of 1145 crab (2004-2020). The summer subsistence fishery was not included in the assessment model.

Harvest of both winter commercial and subsistence fisheries is also influenced by availability of stable ice conditions. Small harvests can occur due to poor ice condition, regardless of crab abundance.

1. Brief description of the annual ADF&G harvest strategy

Since 1997 NSRKC has been managed based on a GHL. From 1999 to 2011 the GHL for the summer commercial fishery was determined using model estimated predicted biomass: (1) 0% harvest rate of legal crab when estimated legal biomass < 1.5 million lb; (2) $\leq 5\%$ of legal male biomass when the estimated legal biomass falls within the range 1.5-2.5 million lb; and (3) $\leq 10\%$ of legal male biomass when estimated legal biomass >2.5 million lb. In 2012 the summer commercial fishery GHL was revised to (1) 0% harvest rate of legal crab when estimated legal biomass < 1.25 million lb; (2) $\leq 7\%$ of legal male biomass when the estimated legal biomass falls within the range 1.25-2.0 million lb; (3) $\leq 13\%$ of legal male biomass when the estimated legal biomass falls within the range 2.0-3.0 million lb; and (3) $\leq 15\%$ of legal male biomass when estimated legal biomass >3.0 million lb.

In 2015 the BOF passed the following regulations regarding the winter commercial fisheries:

- 1) Revise GHL to include summer and winter commercial fisheries.
- 2) Set GHL for the winter commercial fishery (GHL_w) at 8% of the total GHL

NSRKC GHL is determined in early February after the final determination of ABC. GHL is determined not to exceed (e.g., 5-10% less) retained portion of the total catch ABC.

Table: Brief NSRK fishery management history

Year	Notable historical management changes
1976	The abundance survey started
1977	Large vessel commercial fisheries began. Legal size was set to ≥ 5 inch CW
1978	Legal size was changed to ≥ 4.75 inch CW
1991	Fishery closed due to staff constraints
1993	Fishery is restricted to small boat. The end of large vessel commercial fishery operation.
1994	Super exclusive designation went into effect.
1998	Community Development Quota (CDQ) allocation went into effect
1999	Guideline Harvest Level (GHL) went into effect
2000	North Pacific License Limitation Program (LLP) went into effect.
2002	Change in closed water boundaries (Figure 2)
2006	The Statistical area Q3 section expanded (Figure 1)
2008	Start date of the open access fishery changed from July 1 to after June 15 by emergency order. Pot configuration requirement: at least 4 escape rings (> 4.5 inch diameter) per pot located within one mesh of the bottom of the pot, or at least $\frac{1}{2}$ of the vertical surface of a square pot or sloping side-wall surface of a conical or pyramid pot with mesh size > 6.5 inches.
2008	Market preferred size of ≥ 5 inch CW became a standard commercial retained size.
2012	The BOF adopted a revised GHL for summer fishery.
2016	Winter GHL for commercial fisheries was established and modified winter fishing season dates were implemented.
2020	The BOF closed summer commercial fishery E of 167 longitude
2021	Change winter fishery open date to February 1

2. Summary of the history of the B_{MSY} .

NSRKC is a Tier 4 crab stock. Direct estimation of the B_{MSY} is not possible. The B_{MSY} proxy is calculated as mean model estimated mature male biomass (MMB) from 1980 to present. Choice of this period was based on a hypothesized shift in stock productivity due to a climatic regime shift indexed by the Pacific Decadal Oscillation (PDO) in 1976-77.

D. Data

1. Summary of new information:

Winter commercial and subsistence fisheries:

The winter commercial fishery retained catch in 2022 was 2,708 crab (7,683.7 lb). Winter subsistence retained crab catch was 6,743 and unretained was 2,403 crab or 26 % of total catch (Table 2).

Summer commercial fishery:

The summer commercial fishery opened on 6/15/2022 and closed on 7/24/2022. Total of 125,042 crab (317,173 lb) were harvested (Table 1).

Standardized CPUE (Appendix B).

Standardized CPUE for the years of 1991, 2020, 2021 were not calculated because commercial fishery did not occur (1991) or no crab was harvested (2020, 2021).

Recalculate standardized CPUE: 3 periods:

1977-1992: Large Scale commercial fishery

1993-2007: Small boat commercial fishery

2008-2019: Small boat commercial fishery with high grading.

Discards

Estimate of discards is based on ratio method (Appendix C).

Summer Trawl Survey

Norton Sound portion of the NOAA NBS trawl survey was conducted in 2022. Total male crab abundance estimate (CL > 63mm) is 2103 thousand with CV 0.37 (Table 3).

Available survey, catch, and tagging data

Available NSRKC data consist of the following: trawl survey that informs abundance and size composition, catch that informs size composition, and standardized CPUE that informs an index of abundance, and tag recovery that informs growth-transition.

Trawl survey

Trawl survey consists of 3 surveys: NMFS triennial survey: 1976-1992, ADF&G survey: 1996-2021, and NOAA NBS survey: 2010, 2017-2021.

NMFS triennial survey:

Norton Sound trawl survey was initiated by NMFS in 1976 to assess stock status of crab and ground fish in Norton Sound and Kotzebue Sound. The survey established 10 nautical mile (nm) grid survey stations throughout the entire Norton Sound and 15 nm grids outside the Norton Sound area. The initial Norton Sound survey became the standard stations moving forward. The survey was conducted from mid-late August to September-October, except for 1979, which was in late July/early August. The survey used 83-112 Eastern Otter trawl gear, with tow distance of 1.3 – 1.7 nm (30 minutes tow). The survey was terminated in 1992.

ADF&G triennial -annual survey:

After the termination of the NMFS trawl survey, ADF&G began trawl surveys in 1996 using the same survey stations, but smaller boat and survey coverage. The survey started as triennially but became an annual survey in 2017. The survey usually occurs in late July – mid August, using 400 Eastern Otter trawl gear with tow distance of 1.0 nm. The survey used to have a re-tow protocol: when the first tow caught more than 5 legal red king crab, the station was re-towed. This protocol was dropped in 2012 in favor of more coverage.

NOAA biennial-annual NBS survey:

NOAA NBS trawl survey started in 2010, and biennially since 2017. The survey occurs in late July-mid August, similar to the ADF&G survey. The survey has 20 nm grid using 83-112 Eastern Otter trawl gear, with tow distance of 1.3 – 2.5 nm (30 min tow).

Abundance estimation method

Methods of estimating red king crab abundance differed among the three surveys and throughout time periods. Abundance estimates have been revised many times.

Abundance and CV of the NMFS 1976-1991 surveys were provided by NOAA (Jon Richer NOAA *personal communication*). The abundance was estimated by averaging catch CPUE ($\#/nm^2$) of all stations (including survey stations out of Norton Sound) that was multiplied by standard Norton Sound Area ($7600 nm^2$) (i.e., $N = 7600 * \text{mean CPUE}$). The ADF&G survey abundance is calculated at each station (i.e., $n = \text{CPUE} * 100 nm^2$) and summed across all surveyed stations (i.e., $N = \text{sum of } 100 * \text{CPUEs}$) (Bell and Hamazaki 2019). Extent of the ADF&G survey coverage differed among years due to survey conditions, and survey abundance has not been standardized. NOAA NBS survey abundance is estimated by the author in similar manner as ADF&G survey with the data limited to the Norton Sound survey area that overlaps the ADF&G survey area ($5841 nm^2$) (Figure A).




Figure A. ADF&G trawl survey coverage (yellow shade) and NOAA NBS trawl survey coverage where abundance estimates were made (red hashed line),

Catchability of NSRKC appears to differ between ADF&G and NMFS-NOAA NBS trawl surveys. ADF&G trawl survey abundance tends to be higher than NMFS-NOAA NBS trawl survey even though NMFS-NOAA NBS survey coverages are greater than ADF&G. The assessment model assumes (recommendation by CPT-SSC) that survey q of ADF&G trawl survey be 1.0, which resulted in $q = 0.7-0.8$ for NMFS and $0.7-0.96$ for NOAA NBS survey.

Trawl survey catches are highly patchy. The majority of catches occurred at 1 to 4 stations that caught 20% to 80% of crabs caught during the entire survey (Figure B). The most consistently abundant survey stations are near Nome (blue dots) outside of the summer commercial fishery area (red rectangular). Some offshore stations had high catches for a few years (orange dots: 1990s, yellow dots: 2020s) but they did not persist.




Figure B. Trawl survey stations where majority of catch occurred. Red rectangular indicates where the majority of summer commercial fishery occurs. Blue dots indicate the stations that had high catch consistently since 1976. Orange and yellow dots indicate high crab catch stations that occurred during the 1990s (orange) and 2020s (yellow).

Standardized CPUE

Standardized summer commercial fishery CPUE is included in the NSRKC assessment model as an index of NSRKC abundance that could supplement triennial trawl survey. In 2013, the CPUE standardization model was developed by Gretchen Bishop (ADF&G) (NPFMC 2013). Since then, the same model has been applied with updated data (Appendix B).

Tagging-recovery data

Norton Sound red king crab tagging was initially conducted in 1980 as a part of mark-recapture abundance survey (Brannian 1987). The study was conducted in 1980-1982 and 1985. From 1986 to 2012 crabs were tagged during the winter pot survey. The winter pot surveys tagged more smaller (sublegal) crabs; however, very few were recovered. Tagging resumed from 2012-2015 for a spring migration movement survey. In all the above studies, most of the tagged crabs were recovered by commercial fishermen, but subsistence fishermen also recovered a small number of tags.

Length-Shell proportion data

Length-shell data have been collected in every research and harvest monitoring survey. Of those, summer commercial harvest sampling, winter pot survey (terminated in 2012), and the trawl survey have been consistent.

Time series of the data used for the NSRKC assessment model are summarized in the following figure and table.




Table: a list of survey data

	Years	Data Types	Tables
Summer trawl survey	76,79,82,85,88,91,96, 99, 02,06,08,10,11,14,17-22	Abundance Length-shell comp	3 6
Winter pot survey	81-87, 89-91,93,95-00,02-12	Length-shell comp	7
Summer commercial fishery	77-90,92-22	Retained catch Standardized CPUE, Length-shell comp	1 1, Appendix B 4
Summer Com total catch	12-19	Length-shell comp	9
Summer Com Discards	87-90,92,94	Length-shell comp	8
Winter subsistence fishery	76-22	Total & Retained catch	2
Winter commercial fishery	78-22	Retained catch	2
	15-18	Retained Length-Shell	5
Tag recovery	80-19	Recovered tagged crab	10

Table: A list of data available but not used for assessment

Data	Years	Data Types	Reason for not used
Summer pot survey	80-82,85	Abundance Length proportion	Uncertainties on how estimates were made.
Summer preseason survey	95	Length proportion	Just one year of data
Summer subsistence fishery	2005-2019	retained catch	Too few catches, ignored.
Winter Pot survey	87, 89-91,93,95- 00,02-12	CPUE	CPUE data unreliable.
Preseason Spring pot survey	2011-15	CPUE, Length proportion	Years of data too short
Postseason Fall pot survey	2013-15	CPUE, Length proportion	Years of data too short

Catches in other fisheries

In Norton Sound, the directed Pacific cod pot fishery was issued in 2018 under the CDQ permit. In 2018 and 2019 fishery seasons, a total of 8 and 13 kg (mortality applied) of NSRKC were taken in the groundfish fisheries (CPT 2020). However, all bycatch occurred to the west of 168.0 longitude where NSRKC survey has not been conducted. Norton Sound Fishery Management Area (Q3) extends to St. Lawrence Island and US-Russia border (Figure 1). In the absence of survey abundance extended to those area, it is questionable whether those bycatch mortalities should be included in the NSRKC population.

	Fishery	Data availability
Other crab fisheries	Does not exist	NA
Groundfish pot	Pacific cod	Y
Groundfish trawl	Does not exist	NA
Scallop fishery	Does not exist	NA

Other miscellaneous data:

- Tagging-recovery locations (2012-2019)
- Satellite tag migration tracking (NOAA 2016, ADF&G 2020-21)
- Spring offshore migration distance and direction (2012-2015)
- Monthly blood hormone level (indication of molting timing) (2014-2015)
- Functional maturity and mating success of captured crab (2021-)

Data aggregated:

- Length data were aggregated by 10 mm range, starting from 64-73 mm. Crab length greater than 133 mm were aggregated in >133 mm class.
- Shell condition data were aggregated from very new, and new to newshell, and from old, very old, and very very old to oldshell.
- Tag-recovery data were aggregated regardless of tagging years.

Data estimated outside the model:

- Summer commercial catch standardized CPUE (Table 1, Appendix B)
- Proportion of legal-size crab, estimated from trawl survey and observer data. (Table 13)
- Average weight of crab by length class (Table 13)

E. Analytic Approach

1. History of the modeling approach and issues:

The Norton Sound red king crab stock was assessed using a length-based synthesis model (Zheng et al. 1998). Since adoption of the model, the model had the following model mismatches:

1. Model projects higher abundance-proportions of large size class (> 123mm CL) of crab than observed. This problem was further exasperated when natural mortality M was set to 0.18 from previous $M = 0.3$ in 2011 (NPFMC 2011).

2. Poor model fit to trawl survey abundance. This was further exasperated when $M = 0.18$ was applied to all lengths.
3. Some model parameters hit boundaries

Those issues resulted in the model overestimating projected abundance. The following describes historical model adjustments attempted.

1. Model projects higher abundance-proportions of large size class ($> 123\text{mm CL}$) of crab.

This issue has been solved by assuming (3-4 times) higher M for the large crab (i.e., $M = 0.18$ for length classes ≤ 123 mm, and higher M for > 123 mm) (NPFMC 2012, 2013, 2014, 2015, 2016, 2017, 2018). However, because this solution is biologically suspect, several alternative assumptions have been considered in the past.

- a. Large crabs move out of the survey and fishing area

In modeling, this was dealt with by setting dome shaped survey and commercial catch selectivity (i.e., lower catchability for large crab). This modeling configuration resulted in estimating MMB two times higher than the default model, which indicates that true NSRKC abundance is twice larger than current trawl survey and commercial crab fishery indicate (NPFMC 2017). The NOAA NBS surveys (2010, 2017, 2019, 2021) did not find high numbers of red king crab outside Norton Sound. The large crab could also be nearshore where the commercial fishery is closed, and trawl survey is not conducted due to rocky bottom. However, spring tagging studies showed that most crab migrated from near shore to offshore (fishing) area (Jenefer Bell, ADF&G personal comm.). There was little evidence that large crabs stay in nearshore waters during summer.

- b. Molting and growth of NSRKC are slower. (i.e., model overestimating molting and growth probability: transition matrix)

The model originally estimated the transition outside of the model. In 2014 the model was configured to estimate the transition matrix inside of the assessment model (NPFMC 2014). The transition matrix estimated inside of the model was similar to that estimated outside of the model. When length-specific molting probability was estimated individually, the shape of the probability curve was also similar to the default inverse logistic molting function (NPFMC 2016). A time varying molting function (random walk) process did not improve model fit. Laboratory studies showed that observed growth after molting was comparable to those from tag-recovery data, though sample size was limited and comparable tank-natural condition factors such as water temperature and food availability, were questionable (Leah Zacher of NOAA AFSC Kodiak personal comm),

- c. Higher natural mortality (M) than assumed $M = 0.18$

Profile analyses and estimating M across all length classes resulted in higher M (0.3-0.45) than default $M=0.18$ (NPFMC 2013, 2017). However, the model fit is slightly worse than the default model.

- d. Higher natural mortality ($M > 0.18$) for small crab and large crab having higher mortality than small crab.

This model configuration had the best fit to data (NPFMC 2017).

- e. Gradual size dependent natural mortality.

The default assessment model assumes abrupt M increase at size CL 124mm or greater. An alternative model suggested that M gradually increasing from size as low as 94 mm CL; however, the overall model fit did not greatly improve from the default model (NPFMC 2017). In 2022, CPT requested estimating M for each length class, which also suggested length-dependent natural mortality.

Natural mortality M specification was originally set to be 0.2 for BSAI red king crab stock, including NSRKC (NPFMC 1998) and was changed to 0.18 with Amendment 24 (NPFMC 2011), **which means that the NSRKC stock assessment model is REQUIRED to use $M=0.18$** . Since adoption of the model, the CPT-SSC has required reexamining the appropriateness of M specification for NSRKC; however, the CPT-SSC have been rejecting NSRKC having higher M , except for current model configuration.

- 2. Poor model fit to trawl survey abundance, especially NMFS survey (1976-1992) data

The NSRKC assessment model suggest higher crab abundance during the 1976-1990s period. The model deals with this issue by including survey q ($q < 1$), or the model assumes the NMFS trawl surveys **underestimated** NSRKC abundance. However, this assumption is arbitrary, which is also affected by other model configurations. For instance, when $M = 0.18$ is assumed for all length classes, the model suggests that survey q for NMFS is greater than 1.0, or that the NMFS trawl surveys **overestimated** NSRKC abundance (Model 21.1). This indicates the trawl surveys overestimated NSRKC abundance (NPFMC 2022). Alternatively, assuming the NMFS survey q to be 1.0 resulted in ADF&G trawl survey q greater than 1.0 (i.e., trawl survey overestimates abundance), even though ADF&G trawl survey area is generally smaller than NMFS and NBS survey areas.

This model fitting issue was also influenced by input sample sizes for size-shell compositions. Increasing the input sample size resulted in the model estimating lower abundance. Reducing the input sample sizes improved model fit to the trawl survey data but caused lower fit to size-shell composition data (NPFMC 2012, 2013, 2015). Alternative model weighting methods (e.g., Francis 2012) have been tried, but those did not improve model fit.

3. Some model parameters hit boundaries.

There are two model parameters that hits boundaries: Trawl survey selectivity ($\log_{\phi_{stl}}$), and the proportion of recruits (r_1, r_2).

3. Trawl Survey selectivity parameter

Trawl survey selectivity model is a one parameter logistic curve that reaches 1.0 at L_{max} (143.5 mm)

$$S_l = \frac{1}{1 + e^{(\alpha(L_{max} - L) + \ln(1/0.999 - 1))}}$$

where $\alpha = \exp(\log_{\phi_{stl}})$, $L_{max} = 143.5$ mm L (63.5-143.5 mm)

Model estimated trawl survey selectivity is 1.0 across all size classes. This means that $e^{(\alpha(L_{max} - L) + \ln(1/0.999 - 1))} \approx 0$, $\alpha(L_{max} - L) + \ln(1/0.999 - 1) = -\infty$, $\alpha \approx 0$, and $\log_{\phi_{stl}} = -\infty$. Hence, the parameter will hit the boundary.

Alternative option is assuming $S_l = 1.0$ for all length classes; however, this also removes the model's ability to estimate S_l when all length classes are NOT 1.0.

4. The proportion of recruits

The proportion of recruits is a multinomial formula of $n=3$

$$p_l = \frac{\exp(r_l)}{1 + \sum_{l=1}^{n-1} \exp(r_l)} \text{ for } l = 1, \dots, n-1$$

$$p_n = 1 - \frac{\sum_{l=1}^{n-1} \exp(r_l)}{1 + \sum_{l=1}^{n-1} \exp(r_l)}$$

Model estimated recruit proportions for length classes 1, 2, 3 (P_1, P_2, P_3) are 0.592, 0.403, and 0.003. $P_3 \approx 0$ makes it extremely difficult for the model to estimate P_1 and P_2 , and **thus**

model parameters r_1 and r_2 (Tables 11, 12). Increasing the upper bound of the r parameters would still make r_1 hit the boundary and make estimates of P_1, P_2, P_3 to be closer to $P_1 = 0.60, P_2 = 0.40, \text{ and } P_3 = 0$. Alternative option is assuming $P_3 = 0$; however, this also removes the model's ability to estimate P_3 when P_3 is far greater than 0.

Historical Model configuration progression:

2011 (NPFMC 2011)

- 1). $M = 0.18$.
- 2). M of the last length class = 0.288.
- 3). Include summer commercial discards mortality = 0.2.
- 4). Weight of fishing effort = 20.
- 5). The maximum effective sample size for commercial catch and winter surveys = 100.

2012 (NPFMC 2012)

- 1) M of the last length class = $3.6 \times M$.
- 2) The maximum effective sample size for commercial catch and winter surveys = 50.
- 3) Weight of fishing effort = 50.

2013 (NPFMC 2013)

- 3) Standardize commercial catch CPUE and replace likelihood of commercial catch efforts to standardized commercial catch CPUE with weight = 1.0.
- 4) Eliminate summer pot survey data from likelihood.
- 5) Estimate survey q of 1976-1991 NMFS survey with maximum of 1.0.
- 6) The maximum effective sample size for commercial catch and winter surveys = 20.

2014 (NPFMC 2014)

- 1) Modify functional form of selectivity and molting probability to improve parameter estimates (2 parameter logistic to 1 parameter logistic).
- 2) Include additional variance for the standardized CPUE.
- 3) Include winter pot survey CPUE (was removed from the final model due to lack of fit).
- 4) Estimate growth transition matrix from tagged crab recovery data.

2015 (NPFMC 2015)

- 1) Winter pot survey selectivity is an inverse logistic, estimating selectivity of the smallest length group independently.
- 2) Reduce weight of tag-recovery: $W = 0.5$.
- 3) Model parsimony: one trawl survey selectivity and one commercial pot selectivity.

2016 (NPFMC 2016)

- 1) Length range extended from 74 mm – 124 mm above to 64 mm – 134 mm above.
- 2) Estimate multiplier for the largest ($> 123\text{mm}$) length classes.

2017 (NPFMC 2017)

- 1) Change molting probability function from 1 to 2 parameter logistic. Assume molting probability not reaching 1 for the smallest length class.

2018 (NPFMC 2017) CPT-SSC suggested no model alternatives

2019 (NPFMC 2019)

- 1) Fit total catch length composition and estimate retention probability for summer and winter commercial fishery.
- 2) Include winter commercial retained length data.

2020 (NPFMC 2020) The CPT and SSC suggested no model alternatives

2021 (NPFMC 2021) Include discards data at the request of CPT and SSC

- 1) Models that bridge from the Model 19.0e to 21.0
- 2) Model 21.0 with natural mortality estimated by model.
- 3) Estimate size specific natural mortality.

2022 (NPFMC 2022)

- 1) Shell based retention probability
- 2) Estimate individual length class M

2. Model Description

- a. Description of overall modeling approach:

The model is a male-only size structured model based on **abundance** that combines multiple sources of surveys, fishery catches and discards, and mark-recovery data using a maximum likelihood modeling framework to estimate population dynamics under fisheries. The model is an extension of the length-based model developed by Zheng et al. (1998) for NSRKC. A detailed description of the model is in Appendix A.

The model estimates abundances of male crab with $CL \geq 64$ mm and with 10 mm length intervals (8 length classes, ≥ 134 mm) because few crab measuring less than 64 mm CL were caught during surveys or fisheries.

The model assumes newshell crab as molted and oldshell crab as unmolted.

One critical characteristic of the model is that it does not estimate fishing mortality (F). Observed harvests were considered accurate and thus directly subtracted from the model estimated abundance.

The modeling scheme and data is described in the following figure.




Figure C. Norton Sound red king crab model and data scheme. Bold type indicate data that were fitted to the model. Boxes in dotted line indicate model estimated parameters and quantities. Natural mortality, M was set to 0.18 except for CL greater than 123mm that was estimated in the model.

Timeline of calendar events and crab modeling events:

- **Model year starts February 1st to January 31st of the following year.**
- **Initial Population Date: February 1st, 1976, consisting of only newshell crab.**
- **Instantaneous fishing mortality: winter (February 1st) and summer (July 1st) fisheries**
- **Instantaneous molting and recruitment occur on July 1st**

Critical model assumptions

NSRKC Crab Biology

1. Instantaneous annual natural mortality (M) is 0.18 and increases at the size greater than 123 mm CL. M is constant over time.

See History of the modeling approach and issues section for detailed discussion regarding this assumption

2. Male crab size at maturity is 94mm CL.

Size at maturity of NSRKC is highly uncertain (NPFMC 2018, 2019, 2020, 2021). First, maturity has two categories (biological and functional). Biological maturity indicates that male red king crab can produce viable sperm, whereas functional maturity indicates that male red king crab are large enough to mate. The former can be determined using the presence/absence of spermatophores in the vas deferens, whereas the latter can be inferred by measuring mating pairs *in situ* or in lab experiments. The current NSRKC functional maturity size (>94 mm) was inferred from Bristol Bay red king crab by incorporating the fact that Norton Sound red king crab are smaller.

SSC suggested investigating size at functional maturity of other stocks, such as of Barents Sea red king crab (NPFMC 2021). However, it is unlikely that those metadata analyses would provide insights about size at functional maturity of Norton Sound red king crab because NSRKC is the smallest among red king crab stocks. Author was not able to find any other red king crab stocks that are comparable to the size of NSRKC.

Recent laboratory studies reported that NSRKC male crab as small as 79.4 mm CL can fertilize females. Those small males could also fertilize eggs of ~ 4 females, which was comparable to the number of females larger (94-116 mm CL) crabs could fertilize. More interestingly, the largest crab (> 123 mm) was able to fertilize eggs of ~ 2 females (Leah Zacher NOAA Kodiak *personal comm*). This functional maturity/mating success study is ongoing but does suggest that functional maturity of NSRKC is smaller than current model assumption of 94 mm CL.

Although determining size at functional maturity is important biologically, there is limited utility of this information for Tier 4 crab stock assessment. In Tier 4 stock assessment, size at maturity is used only for calculation of mature male biomass (MMB) and B_{MSY} (average MMB). Harvest control (F_{OFL}) is based on the ratio of projected MMB and B_{MSY} (projected MMB/ B_{MSY}).

Level	Criteria	F_{OFL}
a	$B / B_{MSY\ proxy} > 1$	$F_{OFL} = \gamma M$
b	$\beta < B / B_{MSY\ proxy} \leq 1$	$F_{OFL} = \gamma M (B / B_{MSY\ proxy} - \alpha) / (1 - \alpha)$
c	$B / B_{MSY\ proxy} \leq \beta$	$F_{OFL} = \text{bycatch mortality \& directed fishery } F = 0$

where B is a mature male biomass (MMB), $B_{MSY\ proxy}$ is average mature male biomass over a specified time period, $M = 0.18$, $\gamma = 1$, $\alpha = 0.1$, and $\beta = 0.25$.

The MMB/B_{MSY} ratio is affected very little by changes of maturity size unless the ratio is very close to 1.0 (Tier 4a vs Tier 4b borderline). To illustrate this, we present 2022 assessment model results with various minimum size at maturity cutoffs, as follows.

Maturity size	94mm (default)	74mm	84mm	104mm	114mm	124mm	>134mm
B_{MSY} mil. lb	4.88	5.21	4.92	3.76	2.71	1.33	0.39
MMB(2022) mil. lb	5.21	5.91	5.61	4.42	2.86	1.03	0.27
MMB/ B_{MSY}	1.16	1.13	1.14	1.18	1.06	0.77	0.70
Tier 4 level	a	a	a	a	a	b	b
F_{OFL}	0.18	0.18	0.18	0.18	0.18	0.13	0.12

As illustrated in the above table, changing minimum maturity size has little effect on MMB/B_{MSY} ratio and Tier 4 level designation. OFL and ABC are based on retained and unretained catch by size applied by F_{OFL} .

3. Molting occurs right after the summer fishery.

Molt timing of NSRKC was verified by field and laboratory survey. Double shelled crabs are often observed in September (Joyce Soong ADFG personal comm.), and crabs sent to Kodiak Lab molted in September-October (Leah Zacher NOAA personal comm).

4. Recruitment occurs in fall at the same time as molting.

In NSRKC assessment modeling, recruitment is not a function of mature males, but estimated model parameters entering to the immature length classes 64 mm - 93 mm. In modeling, this adjustment is done at the same time as molting-growth.

5. Molting probability is a descending logistic function of crab size. Molted crab become newshell and unmolted crab become oldshell crab.

Tag recovery data during the 2012-2014 study suggest lower molting probability for larger crabs. The table below shows the number of newshell crab tagged, released, and recaptured at 1 year of liberty. Crabs recaptured newshell is considered as molted and oldshell is considered as unmolted.

Length Class	Newshell	Oldshell	% molted
1 (64-73mm)	3	0	100
2 (74-83mm)	30	0	100
3 (84-93mm)	64	5	93
4 (94-103mm)	113	9	93
5 (104-113mm)	44	36	56
6 (114-123mm)	22	21	51
7 (124-133mm)	5	10	22
8 (>133mm)	0	4	0

This assumes that shell condition observations are correct, which is difficult to verify objectively. For instance, in tag-recovery data (2012-2016) below, 125 crabs had no growth (+/- 3 mm) in one year of liberty. Of those, 100 crabs were released as newshell and 25 crabs were released as oldshell. If no growth is considered unmolted, all those crabs should be recaptured as oldshell. However, 29% of crabs released as newshell were recaptured as newshell crab and 48% of crabs released as oldshell were recaptured as newshell.

Released/Recovered	Newshell	Oldshell
Newshell	28	70
Oldshell	12	13

This could be caused by (1) inaccurate length measurement, (2) inaccurate shell condition assessment, or (3) no growth after molting.

6. Growth increment is a function of length, constant over time. Molted crab does not shrink.

Tag recovery data showed that growth increment of large crab tend to be smaller than that of small crab (Figure D). The data also showed negative growth increment, at the largest length class.




Figure D. Growth increment by tagged length class of molted (newshell recovered) crab with 1 year at liberty.

NSRKC Surveys

1. ADF&G trawl survey (1996-2021) abundance has the same scale as the population (i.e., catchability $q = 1.0$). Abundances by historical NMFS (1976-1991) and NOAA NBS (2010-present) survey are biased low (i.e., $q < 1.0$).

Survey $q = 1.0$ for ADF&G trawl survey and lower survey q for NOAA survey was adopted in 2013 assessment (NPFMC 2013). However, it is possible that ADF&G surveys are overestimates of abundance. Model estimated survey q for ADF&G trawl survey was greater than 1.0 (NPFMC 2013, 2019).

2. Size selectivity is an asymptotic one parameter logistic function of 1.0 at the length class 134 mm CL and the same across years and survey agencies.

$$S_l = \frac{1}{1 + e^{(\alpha(L_{\max} - L) + \ln(1/0.999 - 1))}}$$

This logistic function form was adopted during the crab workshop in 2005 to reduce model parameters and increase parameter estimation stability.

Although the surveys differ among NOAA (1976-1991), ADF&G (1996-2021), and NOAA NBS (2010-present) in terms of survey vessel and trawl net structure, selectivity of all surveys were assumed to be identical. Model fits separating and combining the surveys were examined in 2015; however, selectivity was essentially identical (1.0 across all size classes) (NPFMC 2015). For model parsimony, the SSC recommended using only one selectivity.

3. Winter pot survey selectivity is a dome shaped function: a combination of a reverse logistic function starting from length class 84 mm CL and model estimate for CL < 84 mm length classes. The selectivity is constant over time.

$$S_{w,l} = \frac{I}{1 + e^{\alpha(L-\beta)}}$$

This assumption is based on the low proportion of large crab that are caught in the nearshore area where winter surveys occur. This does not necessarily imply that the crab pots are less selective to large crabs. Alternatively, this may imply that fewer large crab migrate into nearshore waters in winter.

NSRKC Fisheries

1. Fisheries occur twice on July 01 and Feb 01 and are instantaneous.
2. Summer commercial fishery size selectivity is an asymptotic one parameter logistic function of length, with the selectivity in length class 134 mm CL set to 1. Selectivity is constant over time.

$$S_l = \frac{I}{1 + e^{(\alpha(L_{\max}-L)+\ln(1/0.999-1))}}$$

This logistic function form was adopted during the crab workshop in 2005 to reduce model parameters and parameter estimation stability. Although summer commercial fishery changed greatly between the periods (1977-1992, 1993-present) in terms of fishing vessel composition, and pot configuration, the selectivity of each period is assumed to be identical. Model fits of separating and combining the two periods were examined in 2015 and showed no difference between the two (NPFMC 2015). For model parsimony, the SSC recommended using only one selectivity.

3. Not all legal sized crabs are retained. Retention probability is an asymptotic logistic function.

Legal size of NSRKC is defined as carapace width (CW) greater than 4.75 inches that was conventionally equated as greater than 104 mm CL. Since 1996 ADF&G has started noting legal size crab based on carapace width in trawl, commercial fishery observer, and other miscellaneous surveys to complement the carapace length measurement. Originally, the proportion was based solely from the trawl survey. As more data are collected from commercial observer surveys, recent proportions are based on more observer data.

Proportion of legal (CW>4.75 inch) crab in Trawl survey

size class	64	74	84	94	104	114	124	134
1996	0.00	0.00	0.00	0.18	0.93	1.00	1.00	1.00
1999	0.00	0.00	0.00	0.40	0.98	0.98	1.00	1.00
2002	0.00	0.00	0.00	0.28	0.97	1.00	1.00	1.00
2006	0.00	0.00	0.00	0.18	1.00	1.00	1.00	1.00
2008	0.00	0.00	0.00	0.19	0.96	1.00	1.00	1.00
2011	0.00	0.00	0.00	0.24	0.99	1.00	1.00	1.00
2014	0.00	0.00	0.00	0.21	0.98	1.00	1.00	1.00
2017	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00
2018	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00
2019	0.00	0.00	0.00	0.33	1.00	1.00	1.00	1.00
2020	0.00	0.00	0.00	0.22	1.00	1.00	1.00	1.00
Average	0.00	0.00	0.00	0.25	0.98	1.00	1.00	1.00

Proportion of legal (CW>4.75 inch) crab in Observer survey

size class	64	74	84	94	104	114	124	134
2012	0.00	0.01	0.02	0.22	0.90	1.00	1.00	1.00
2013	0.00	0.00	0.00	0.44	0.98	1.00	1.00	1.00
2014	0.00	0.00	0.00	0.22	0.91	1.00	1.00	1.00
2015	0.00	0.00	0.00	0.38	0.98	1.00	1.00	1.00
2016	0.00	0.00	0.00	0.46	1.00	1.00	1.00	1.00
2017	0.00	0.00	0.00	0.13	0.91	1.00	1.00	1.00
2018	0.00	0.00	0.00	0.16	0.95	0.99	1.00	1.00
2019	0.00	0.00	0.00	0.18	0.93	1.00	1.00	1.00
Average	0.00	0.00	0.00	0.30	0.95	1.00	1.00	1.00

The proportion of legal crab used in the assessment model is an average proportion based on observer survey data. In the assessment model, this proportion is used to estimate the number of retained crab in winter and summer commercial fisheries prior to 2008. It is assumed prior to 2008, all legal sized crab were retained.

Since 2008 commercially retained crab size is $CW > 5.0$ inches and retention probability is estimated from the observer survey.

The table below shows the proportion of legal vs. retained crab during the 2012-2019 observer survey, in response to request from the public.

Year		64	74	84	94	104	114	124	134
2012	Legal	0	0.01	0.02	0.22	0.9	1	1	1
	Retained	0	0	0	0.05	0.46	0.63	0.64	0.85
2013	Legal	0	0	0	0.44	0.98	1	1	1
	Retained	0	0	0	0.14	0.86	0.99	1	1
2014	Legal	0	0	0	0.22	0.91	1	1	1
	Retained	0	0	0	0.04	0.74	0.97	0.99	1
2015	Legal	0	0	0	0.38	0.98	1	1	1
	Retained	0	0	0	0.11	0.74	0.91	0.94	0.89
2016	Legal	0	0	0	0.46	1	1	1	1
	Retained	0	0	0	0.13	0.89	0.99	1	1
2017	Legal	0	0	0	0.12	0.91	1	1	1
	Retained	0	0	0	0.02	0.75	0.99	1	1
2018	Legal	0	0	0	0.16	0.95	0.99	1	1
	Retained	0	0	0	0.14	0.92	0.99	1	0.99
2019	Legal	0	0	0	0.18	0.93	1	1	1
	Retained	0	0	0	0.15	0.93	1	1	1

The proportion of legal sized crab retained from observer survey 2012-2019

Year	64	74	84	94	104	114	124	134
2012	0	0	0	0.23	0.51	0.63	0.64	0.85
2013	0	0	0	0.31	0.88	0.99	1	1
2014	0	0	0	0.19	0.82	0.97	0.99	1
2015	0	0	0	0.28	0.76	0.91	0.94	0.89
2016	0	0	0	0.28	0.89	0.99	1	1
2017	0	0	0	0.14	0.82	0.99	1	1
2018	0	0	0	0.87	0.98	1	1	0.99
2019	0	0	0	0.86	1	1	1	1

The above data justifies using logistic function as selection criteria.

Fishery	Model retention	Data
Summer:1977-2007	Logistic retention prob	Discard, retained size prop
Summer: 2008-2022	Logistic retention prob	Total, retained size prop
Winter: 1977-2007	Mean legal crab proportion	No data
Winter: 2008-2022	Logistic retention prob	Retained size prop
Winter: Subsistence	All crab > 94mm retained	No data (No legal size limit)

3. Winter commercial pot selectivity is the same as the selectivity of the winter pot survey.

This assumption is based on the survey pot being similar to the one used for subsistence, and that many commercial fishermen are also subsistence harvesters. However, by regulation winter commercial king crab pots can be any dimension (5AAC 34.925(d)) and recent popularity of winter commercial fishery may have deviated this assumption.

4. Winter subsistence fishery retains crab size greater than 94 mm CL.

This was based on the assumption that subsistence fishermen would keep crab smaller than legal crab size. By regulation, subsistence fishery had no size limit for retention. Size of crab caught by subsistence fishery has never been monitored.

5. Discards handling mortality rate for all fisheries is 20%.

Discards mortality rate was specified by CPT. No empirical estimates are available.

Data quality assumptions

1. All size-shell composition data are collected accurately without systematic bias.

Although this assumption is reasonable, it is difficult to verify the assumption objectively. For instance, in tag-recovery data (2012-2016), 125 crabs had no growth (+/- 3mm) in one year of liberty. Of those, 100 crabs were released as newshell and 25 crabs were released as oldshell. If no growth is considered unmolted, all those crabs should be recaptured as oldshell. However, 29% of crabs released as newshell were recaptured as newshell crab and 48% of crabs released as oldshell were recaptured as newshell.

Released\Recovered	Newshell	Oldshell
Newshell	29	71
Oldshell	12	13

This could be caused by (1) inaccurate length measurement, (2) inaccurate shell condition assessment, or (3) no growth after molting.

- Annual retained catch is accurate without error.

In Norton Sound, almost all crabs caught by commercial fisheries are sold to NSEDC. Subsistence fishery harvest are self-reporting. Accuracy of self-reporting has never been evaluated.

Model data weighting

Survey data	Input sample size
Summer commercial, winter pot, and summer observer	minimum of $0.1 \times$ actual sample size or 10
Summer trawl and pot survey	minimum of $0.5 \times$ actual sample size or 20
Tag recovery	$0.5 \times$ actual sample size

Recruitment SD: 0.5.

Discards CV: 0.3

“Implied” effective sample sizes were calculated as

$$n = \sum_l \hat{P}_{y,l} (1 - \hat{P}_{y,l}) / \sum_l (P_{y,l} - \hat{P}_{y,l})^2$$

Where $P_{y,l}$ and $\hat{P}_{y,l}$ are observed and estimated length compositions in year y and length group l , respectively. Estimated implied effective sample sizes vary greatly over time.

Data-weighting for NSRKC model is aimed at achieving a balance between various data sets. The current model data weighting schemes, although arbitrary, were deemed appropriate by the CPT-SSC (NPFMC 2011, 2012, See Section E. 1. *Historical Model configuration progression* section). As illustrated in the figure below, increasing weight of size composition data (input sample size: from minimum) would lower model fit to the trawl survey abundance data.

Trawl survey crab abundance




Figure E. Model 21.0 default input sample size (20: trawl, 10: others) (black) vs. increased input sample size (200, 100) (blue dash line), and reduce input size (10, 5) (red dash line).

Thus far, there is no objective criteria for determining the balance (i.e., how much a model should fit observed trawl abundance data vs. size composition data). The author has tried alternative weighting schemes (NPFMC 2019, 2020, 2021) and found current ones are most appropriate.

Changes of assumptions since last assessment:

None

3. Model Selection and Evaluation

- a. Description of alternative model configurations.

For the 2023 draft assessment, the following alternative models are presented.

Model 21.0: Adopted by CPT-SSC for the 2022 final assessment.

Model 22.0: Model 21.0 with shell-dependent retention probability

Model 22.1: Model 21.0 with individual M estimated for all length classes

Model 22.2: Model 21.0 with individual M estimated for all length classes

Model	Retention probability	M	Parameters
21.0	1	0.18+est (L)	
22.0	2	0.18+est (L)	+4
22.1	1	Est	+8
22.2	2	Est	+12

Model 22.0 was proposed to compare efficacy of shell-dependent retention probability on model fit. The shell-dependent retention probability is expected to improve model fit to commercial retention length-shell proportion. This would improve model fit to trawl and total catch length-shell proportion as this allows more oldshell crabs caught in trawl survey. Models 22.1 and 22.2 are not alternatives but simply exploration if full length-dependent M would improve model fit to trawl survey length composition.

4. Results

As predicted, retention probability of oldshell crabs was lower than newshell crabs and individual estimates of M resulted in size-dependent increasing M . M was estimated to be nearly zero for length sizes 64-84 classes (Figure 3). Abundance of the first two length classes is a sum of crab that grew and survived from smaller length classes and independently added recruit (Appendix A equations 6,7). The model was not able to estimate M under this condition. Individual M estimates (Models 22.1, 22.2) also increased the abundance of recruits and thus total crab abundance and mature male biomass during the 1970-2010s (Figures 5, 6, Table 12), but had little influence on model fit to trawl and st. CPUE (Figure 16). This is because model would estimate higher q (Table 11).

Adding shell dependent retention probability (Model 21.0 vs. 22.0) did not reduce or improve retention or discards and total catch size composition (CLP, OBS), but improved trawl survey likelihood (TLP) by 3 points. Similarly, estimating individual M (Model 21.0 vs. 22.1) also lowered trawl survey likelihood by 3 points. Combined together, those reduced trawl survey

likelihood by 5 points (Model 21.0 vs. 22.1). However, there was no observable difference in fits among all four models (Figures 8, 9).

In conclusion, size-dependent M and shell-dependent retention probability adds yet another failure of improving model fits since adoption of the NSRKC assessment model (Section E.1). The NSRKC assessment model has been failing to produce more oldshell crabs and fewer large-sized crabs to match observed size-shell proportions of the trawl survey and commercial retention. Those discrepancies cannot be explained by current understanding of NSRKC biology and fishery.

For selection of an assessment model for the final draft, the author recommends model 21.0 over 22.0. Although inclusion of shell-dependent retention probability is closer to the reality of the fishery, no change in model fit at the cost of additional 4 parameters would be against model parsimony.

In 2022 CPT and SSC recommended the model 21.0 for the final assessment model for 2023 OFL calculation.

Evaluation of negative log-likelihood values.

	Final	Sept 2022			
Model	21.0	21.0	22.0	22.1	22.2
Additional Parameters			+4	+8	+12
AIC change			+6	+5.4	+24
Total	354.1	347.9	346.1	342.6	341.1
TSA	11.0	11.0	10.8	10.5	10.5
DIS	3.4	3.5	4.5	3.3	3.6
St.CPUE	-14.8	-14.8	-14.9	-15.1	-15.0
TLP	134.0	129.0	126.4	125.5	123.7
WLP	39.6	39.5	39.3	39.3	39.1
CLP	49.5	49.3	48.5	48.7	48.9
OBS	24.3	24.3	25.0	24.9	25.1
WCLP	2.8	2.7	2.9	2.5	2.7
REC	19.4	19.5	19.6	20.1	20.1
TAG	85.0	83.9	83.9	82.9	83.4
Max gradient (e-6)	7.6	4.9	2009	14.7	4.55
RMSE Trawl	0.34	0.34	0.34	0.33	0.33
RMSE CPUE	0.44	0.44	0.44	0.44	0.44

TSA: Trawl Survey Abundance

St. CPUE: Summer commercial catch standardized CPUE

TLP: Trawl survey length composition:

WLP: Winter pot survey length composition

CLP: Summer commercial retention catch length composition

REC: Recruitment deviation

OBS: Summer commercial catch observer discards and total catch length composition

TAG: Tagging recovery data composition

WCLP: Winter commercial length-shell composition

DIS: Summer commercial discards abundance

F. Calculation of the OFL

1. Specification of the Tier level and stock status.

NSRKC stock is placed in Tier 4. It is not possible to estimate the spawner-recruit relationship, but some abundance and harvest estimates are available to build a computer simulation model that captures the essential population dynamics. Tier 4 stocks are assumed to have reliable estimates of current survey biomass and instantaneous M ; however, the estimates of M for NSRKC stock are uncertain.

At the Tier 4 level the OFL is determined by the F_{MSY} proxy, B_{MSY} proxy, and estimated legal male abundance and biomass:

Level	Criteria	F_{OFL}
a	$B / B_{MSY\ proxy} > 1$	$F_{OFL} = \gamma M$
b	$\beta < B / B_{MSY\ proxy} \leq 1$	$F_{OFL} = \gamma M (B / B_{MSY\ proxy} - \alpha) / (1 - \alpha)$
c	$B / B_{MSY\ proxy} \leq \beta$	$F_{OFL} = \text{bycatch mortality \& directed fishery } F = 0$

where B is a mature male biomass (MMB), B_{MSY} proxy is average mature male biomass over a specified time period, $M = 0.18$, $\gamma = 1$, $\alpha = 0.1$, and $\beta = 0.25$.

For NSRKC, MMB is defined as the biomass of males > 94 mm CL on February 01 (Appendix A). B_{MSY} proxy is

B_{MSY} proxy = average model estimated MMB from 1980-2023.

Estimated B_{MSY} proxy: 4.3726 million lb

Predicted mature male biomass in 2023 on February 01

Mature male biomass: 5.2872 million lb

Since the projected MMB is above B_{MSY} proxy,

The NSRKC status is Tire 4a

And F_{OFL} for calculation of the OFL is

$F_{OFL} = \gamma \cdot M$ for M is length invariant of 0.18 and $F_{OFL,l} = \gamma \cdot M_l$ for length-dependent M

2. Calculation formula of NSRKC OFL.

OFL of NSRKC is total OFL (OFL_T) that is a sum of the retained and unretained OFL (OFL_r , OFL_{nr}).

$$OFL_T = OFL_r + OFL_{ur}$$

where

$$OFL_r = \text{retained_}B \cdot F_{OFL} \text{ and } OFL_{ur} = \text{unretained_}B \cdot F_{OFL} \cdot hm$$

retained_B is a biomass of crab subject to fisheries that is a sum of the products of crab abundance ($N_{w,l} + O_{w,l}$), fishery selectivity ($S_{s,l}$), retention probability ($S_{r,l}$), and average weight lb (wm_l) by length class (l).

$$\text{retained_}B = \sum_l (N_{w,l} + O_{w,l}) S_{s,l} S_{r,l} wm_l$$

unretained_B is a biomass of crab subject to fisheries and is a sum of the products of crab abundance ($N_{w,l} + O_{w,l}$), fishery selectivity ($S_{s,l}$), 1 minus retention probability ($S_{r,l}$), and average weight lb (wm_l) by length class (l).

$$\text{unretained_}B = \sum_l (N_{w,l} + O_{w,l}) S_{s,l} (1 - S_{r,l}) wm_l$$

hm is handling mortality, default 0.2

The NSRKC fishery consists of two distinct fisheries: winter and summer. The two fisheries are discontinuous with 5 months (0.42 year) between the two fisheries during which natural mortality occurs. To estimate the OFL for the two fisheries, the CPT in 2016 recommended the following formula that the sum of winter and summer catch (H_w, H_s) equals total OFL ($OFL = H_w + H_s$) and that winter catch is a fraction (p) of total OFL: $H_w = p \cdot OFL$, where ***p* is predetermined fraction of the winter fishery to total fishery**. In NSRKC fishery $p = 0.16$ is used.

$$H_w = B_w (1 - \exp(-x \cdot F_{OFL})),$$

$$H_s = B_s (1 - \exp((1 - x) \cdot F_{OFL})), \text{ and}$$

$$B_s = (B_w - H_w) e^{-0.42 \cdot M}$$

where

B_w is the winter NSRKC biomass, B_s is the summer NSRKC biomass, and x is a fraction parameter,

Solving x of the above (see Appendix A for derivation), retained and unretained OFL is calculated as:

(1)

and

$$OFL_{nr} = \text{unretained_} B_w \cdot FOFL_a \cdot hm$$

$$\text{where } FOFL_a = \left(1 - e^{-(F_{OFL} + 0.42M)} - (1 - e^{-0.42M}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL} + 0.42M)})}{1 - p \cdot (1 - e^{-0.42M})} \right) \right) \quad (2)$$

Because M of NSRKC is length-dependent, the proper calculation of NSRKC OFL should account for length-dependent M as:

$$OFL_r = \sum_l \left[\text{retained_} B_{w,l} \cdot FOFL_{a,l} \right]$$

$$\text{where } FOFL_{a,l} = \left(1 - e^{-(F_{OFL,l} + 0.42M_l)} - (1 - e^{-0.42M_l}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL,l} + 0.42M_l)})}{1 - p \cdot (1 - e^{-0.42M_l})} \right) \right) \quad (3)$$

and

$$OFL_{ur} = \sum_l \left[\text{unretained_} B_{w,l} \cdot FOFL_{a,l} \right] \cdot hm$$

$$\text{where } FOFL_{a,l} = \left(1 - e^{-(F_{OFL,l} + 0.42M_l)} - (1 - e^{-0.42M_l}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL,l} + 0.42M_l)})}{1 - p \cdot (1 - e^{-0.42M_l})} \right) \right) \quad (4)$$

where M_l is a size specific natural mortality,

Determination of Total catch OFL

Total catch OFL is calculable by adding retained and unretained portion of the OFL (i.e., Total OFL = $OFL_r + OFL_{nr}$). Standard calculation of OFL for Tier 4 crab is $F_{OFL} = \gamma M = \gamma 0.18$, which assumes that **natural mortality of a stock is F_{MSY} proxy or that OFL is proxy of a stock harvested at F_{MSY} (See Report of the Alaska Crab Stock Assessment Workshop May 13-14 2009)**. However, this assumption is not strictly applicable for NSRKC that has length-dependent M because **F_{OFL} is NOT equal to natural mortality of the stock and OFL is not proxy of a stock harvested at F_{MSY}** .

The length-dependent M and F_{OFL} was recommended to CPT-SSC in January 2017 but was rejected primarily because it would produce higher OFL. As a compromise, OFL for NSRKC stock in the above equations was specified to length-independent $F_{OFL,l} = \gamma M_l = \gamma 0.18$ and length-dependent M_l for allocation of winter and summer fisheries (see equation 2, 3, 4, $FOFL_{a,l}$). This compromise works only when M is prespecified (e.g., $M = 0.18$) for the assessment model. However, when the biological justification of prespecified M is in question (this has been raised by CPT-SSC since 2019) and is estimated inside an assessment model, biological-scientific justification for selecting M for F_{OFL} needs to be clarified and specified. For instance, in 2021

Model 21.5 estimated two M s for lengths 64-123mm, and 124mm and above. In this case, should M for lengths 64-123mm be used for F_{OFL} specification? This is further complicated when M of each length is estimated within the model, such as Models 22.1 and 22.2 in which estimated M ranged from 0 to 0.7. Out of the 8 length-dependent M s, which M should be selected for F_{OFL} and what are biological and scientific justifications? **If length-independent F_{OFL} is preferred, a protocol for selecting a single M should be established when M is length-dependent.**

Length-dependent M and F_{OFL} on the other hand, would be more logical and in line with the assumption of the Tier 4 OFL as **proxy of a stock harvested at Fmsy**. The biggest concerns raised by CPT-SSC were that length-dependent M for NSRKC would increase OFL from standard OFL , though the opposite could occur if estimated M were LOWER than 0.18. Recognizing scientific uncertainties in the determination of OFL , the FMP included ABC Control Rule “for setting the maximum permissible ABC for each stock as a function of the scientific uncertainty in the estimate of OFL and any other specified scientific uncertainty.” The ABC-control rule allows CPT-SSC setting ABC-buffer that they deem to be appropriate for each stock based on scientific uncertainties of OFL . Therefore, if a length-dependent M increases the OFL , the CPT-SSC would simply increase the ABC-buffer to a level that it deems the most appropriate for NSRKC stock. In fact, ABC buffer of NSRKC has increased from 10% in 2011 to 40% in 2021-2022. For instance, increasing ABC buffer to 60% for length-dependent OFL is similar to ABC catch level as standard OFL . The author recommends that CPT-SSC adopt length-dependent OFL and apply ABC control rule (increase ABC-buffer) that it deems appropriate for NSRKC stock.

3. Determination of NSRKC OFL for the 2023 fishery season.

Projected NSRKC biomass catchable to fishery in 2023 is 0.64 million lb. Applying the equations 1-4

OFL s are as follows:

Length independent M .

Retained catch OFL (OFL_r) = 0.643 million lb or 0.292 kt
Unretained catch OFL (OFL_{nr}) = 0.037 million lb or 0.017 kt
Total catch OFL (OFL_T) = 0.680 million lb or 0.31 kt

Length dependent M .

Retained catch OFL (OFL_r) = 0.965 million lb or 0.438 kt
Unretained catch OFL (OFL_{nr}) = 0.037million lb or 0.017 kt
Total catch OFL (OFL_T) = 1.00 million lb or 0.45 kt

Table: Derivation of length-dependent and length-independent OFL

Length class	64	74	84	94	104	114	124	134	Total
Newshell (abundance 10 ³)	515.52	397.95	392.17	371.76	404.69	473.50	231.98	49.17	
Oldshell (abundance 10 ³)	9.85	14.10	24.86	56.62	132.51	176.87	53.24	21.07	
S_l	0.12	0.33	0.63	0.86	0.99	1.00	1.00	1.00	
S_{lr}	0.00	0.00	0.00	0.02	0.90	1.00	1.00	1.00	
wm_l (lb)	0.52	0.82	1.20	1.70	2.32	2.99	3.69	4.37	
Retained (10 ³ lb)	0	0	0	12	1072	1922	1048	307	4361
Unretained (10 ³ lb)	33	111	317	620	121	0	0	0	1202
Length dependent M									
M_l	0.18	0.18	0.18	0.18	0.18	0.18	0.62	0.62	
FOFL _{<i>l</i>}	0.15	0.15	0.15	0.15	0.15	0.15	0.37	0.37	
OFL _{<i>r,l</i>} (x1000)	0	0	0	2	166	297	387	113	965
OFL _{<i>nr,l</i>} (x1000)	1	3	10	19	4	0	0	0	37
Length independent M									
M	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	
FOFL	0.15	0.15	0.15	0.15	0.15	0.15	0.13	0.13	
OFL _{<i>r</i>} (x1000)	0	0	0	2	166	297	138	40	643
OFL _{<i>nr</i>} (x1000)	1	3	10	19	4	0	0	0	37

G. Calculation of the ABC

1. Specification of the probability distribution of the OFL.

ABC is calculated as (1-ABC buffer)·OFL

In 2015 ABC buffer of Norton Sound Red King Crab was set to 20%, which was increased to 30% in 2020 and to 40% in 2021.

Applying the 40% buffer, NSRKC ABC for the 2023 fishery is

ABC = 0.408 million lb or 0.185 kt for length-independent and 0.60 million lb or 0.272 kt for length-dependent.

H. Rebuilding Analyses

Not applicable

I. Data Gaps and Research Priorities

The major data gap of NSRKC is an incomplete understanding of NSRKC biology, including natural mortality, and fate of oldshell and large crabs. Additionally, research should focus on female abundance and fecundity as well as their reproductive potential. As for management, the number and length-shell composition of unretained crab in fisheries are needed for calculation of total catch. Viability of unretained crab need to be researched to examine current default 20% mortality. Incorporation of local and traditional knowledge (LK/TK) and socio-economic impacts of NSRKC fisheries on the region, could bring further insights about NSRKC biology and management.

Acknowledgments

We thank many ADF&G, CPT, SSC for review of the assessment model and suggestions for improvements and diagnoses.

J. References

- Bell, J., J. M. Leon, T. Hamazaki, S. Kent, and W. W. Jones. 2016. Red king crab movement, growth, and size composition within eastern Norton Sound, Alaska, 2012-2014. Alaska Department of Fish and Game, Fishery Data Series No. 16-37, Anchorage.
- Bell, J. and T. Hamazaki 2019. Summary of 2017 and 2018 Norton Sound red king crab bottom trawl survey. Fishery Data Series No. 19-33. Alaska Department of Fish and Game Anchorage
- Fournier, D., and C.P. Archibald. 1982. A general theory for analyzing catch at age data. *Can. J. Fish. Aquat. Sci.* 39:1195-1207.
- Fournier, D.A., H.J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J. Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim. Methods Softw.* 27:233-249.
- Grant, S.W. and W. Chen. 2012. Incorporating deep and shallow components of genetic structure into the management of Alaskan red king crab. *Evolutionary Applications.* 5:820-837.
- Menard, J., J. Soong, and S. Kent 2011. 2009 Annual Management Report Norton Sound, Port Clarence, and Kotzebue. Fishery Management Report No. 11-46. Alaska Department of Fish and Game Anchorage
- Method, R.D. 1989. Synthetic estimates of historical abundance and mortality for northern anchovy. *Amer. Fish. Soc. Sym.* 6:66-82.
- Mohn, R. 1999. The retrospective problem in sequential population analysis: An investigation using

cod fishery and simulated data. ICES Journal of Marine Science, 56:473-488.

NPFMC 1998. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 1998 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2010. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2010 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2011. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2011 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2012. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2012 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2013. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2013 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2014. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2014 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2015. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2015 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2016. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2016 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2017. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2017 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2018. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2018 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2019. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2019 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

NPFMC 2020. Stock assessment and fishery evaluation report for the King and Tanner crab

fisheries of the Bering Sea and Aleutian Islands regions. 2020 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.

Powell, G.C., R. Peterson, and L. Schwarz. 1983. The red king crab, *Paralithodes camtschatica* (Tilesius), in Norton Sound, Alaska: History of biological research and resource utilization through 1982. Alaska Dept. Fish and Game, Inf. Leaflet. 222. 103 pp.

Zheng, J., G.H. Kruse, and L. Fair. 1998. Use of multiple data sets to assess red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska: A length-based stock synthesis approach. Pages 591-612 *In* Fishery Stock Assessment Models, edited by F. Funk, T.J. Quinn II, J. Heifetz, J.N. Ianelli, J.E. Powers, J.F. Schweigert, P.J. Sullivan, and C.-I. Zhang, Alaska Sea Grant College Program Report No. AK-SG-98-01, University of Alaska Fairbanks.

Table 1. Historical summer commercial red king crab fishery harvest and economic performance, Norton Sound Section, eastern Bering Sea. Bold type shows data that are used for the assessment model.

Year	Guideline Harvest Level (lb) ^b	Commercial Harvest (lb) ^{a, b}		Number Harvest	Total Number (Open Access)			Total Pots		ST CPUE		Season Length		Mid-day from July
		Open Access	CDQ		Vessels	Permits	Landings	Registered	Pulls	CPUE	CV	Days	Dates	
1977	^c	517.787		195,877	7	7	13		5,457	2.03	0.32	60	^c	0.049
1978	3,000.000	2,091.961		660,829	8	8	54		10,817	3.87	0.16	60	6/07-8/15	0.142
1979	3,000.000	2,931.672		970,962	34	34	76		34,773	1.30	0.23	16	7/15-7/31	0.088
1980	1,000.000	1,186.596		329,778	9	9	50		11,199	1.64	0.27	16	7/15-7/31	0.066
1981	2,500.000	1,379.014		376,313	36	36	108		33,745	0.57	0.19	38	7/15-8/22	0.096
1982	500.000	228.921		63,949	11	11	33		11,230	0.25	0.15	23	8/09-9/01	0.151
1983	300.000	368.032		132,205	23	23	26	3,583	11,195	0.50	0.18	3.8	8/01-8/05	0.096
1984	400.000	387.427		139,759	8	8	21	1,245	9,706	1.13	0.19	13.6	8/01-8/15	0.110
1985	450.000	427.011		146,669	6	6	72	1,116	13,209	0.69	0.17	21.7	8/01-8/23	0.118
1986	420.000	479.463		162,438	3	3		578	4,284	2.24	0.47	13	8/01-8/25	0.153
1987	400.000	327.121		103,338	9	9		1,430	10,258	0.88	0.33	11	8/01-8/12	0.107
1988	200.000	236.688		76,148	2	2		360	2,350	2.16	0.41	9.9	8/01-8/11	0.110
1989	200.000	246.487		79,116	10	10		2,555	5,149	0.99	0.29	3	8/01-8/04	0.096
1990	200.000	192.831		59,132	4	4		1,388	3,172	2.03	0.32	4	8/01-8/05	0.099
1991	340.000			0	No Summer Fishery									
1992	340.000	74.029		24,902	27	27		2,635	5,746	1.47	0.47	2	8/01-8/03	0.093
1993	340.000	335.790		115,913	14	20	208	560	7,063	0.17	0.22	52	7/01-8/28	0.093
1994	340.000	327.858		108,824	34	52	407	1,360	11,729	1.02	0.09	31	7/01-7/31	0.044
1995	340.000	322.676		105,967	48	81	665	1,900	18,782	0.43	0.17	67	7/01-9/05	0.093
1996	340.000	224.231		74,752	41	50	264	1,640	10,453	1.08	0.13	57	7/01-9/03	0.101
1997	80.000	92.988		32,606	13	15	100	520	2,982	1.01	0.09	44	7/01-8/13	0.074
1998	80.000	29.684	0.00	10,661	8	11	50	360	1,639	1.14	0.09	65	7/01-9/03	0.110
1999	80.000	23.553	0.00	8,734	10	9	53	360	1,630	1.30	0.13	66	7/01-9/04	0.104
2000	336.000	297.654	14.87	111,728	15	22	201	560	6,345	0.97	0.10	91	7/01- 9/29	0.126
2001	303.000	288.199	0	98,321	30	37	319	1,200	11,918	2.08	0.11	97	7/01- 9/09	0.104
2002	248.000	244.376	15.226	86,666	32	49	201	1,120	6,491	0.76	0.26	77	6/15-9/03	0.060
2003	253.000	253.284	13.923	93,638	25	43	236	960	8,494	0.76	0.10	68	6/15-8/24	0.058
2004	326.500	314.472	26.274	120,289	26	39	227	1,120	8,066	1.65	0.09	51	6/15-8/08	0.033
2005	370.000	370.744	30.06	138,926	31	42	255	1,320	8,867	1.36	0.07	73	6/15-8/27	0.058
2006	454.000	419.191	32.557	150,358	28	40	249	1,120	8,867	0.64	0.12	68	6/15-8/22	0.052
2007	315.000	289.264	23.611	110,344	38	30	251	1,200	9,118	0.93	0.10	52	6/15-8/17	0.036
2008	412.000	364.235	30.9	143,337	23	30	248	920	8,721	0.89	0.23	73	6/23-9/03	0.079
2009	375.000	369.462	28.125	143,485	22	27	359	920	11,934	1.35	0.05	98	6/15-9/20	0.090
2010	400.000	387.304	30	149,822	23	32	286	1,040	9,698	0.92	0.04	58	6/28-8/24	0.074
2011	358.000	373.990	26.851	141,626	24	25	173	1,040	6,808	1.35	0.04	33	6/28-7/30	0.038
2012	465.450	441.080	34.91	161,113	40	29	312	1,200	10,041	1.54	0.05	72	6/29-9/08	0.093
2013	495.600	373.278	18.585	130,603	37	33	460	1,420	15,058	1.36	0.04	74	7/3-9/14	0.110
2014	382.800	360.860	28.148	129,657	52	33	309	1,560	10,127	0.71	0.04	52	6/25-8/15	0.052
2015	394.600	371.520	29.595	144,255	42	36	251	1,480	8,356	1.08	0.04	26	6/29-7/24	0.033
2016	517.200	416.576	3,583	138,997	36	37	220	1,520	8,009	1.33	0.05	25	6/27-7/21	0.025
2017	496.800	411,736	0	135,322	36	36	270	1,640	9,401	1.17	0.05	30	6/26-7/25	0.027
2018	319,400	298,396	0	89,613	34	34	256	1,400	8,797	1.00	0.05	35	6/24-7/29	0.030
2019	150,600	73,784	1,239	24,506	24	26	146	1,096	5,438	0.58	0.05	62	6/25-9/03	0.068
2020	170,000	0	0	0	0	0	0	0	0			0	6/25-9/03	NA
2021	290,000	0	0	0	0	0	0	0	0			0	6/15-9/03	NA
2022	341.600	291,553	25,620	121,323	26	26	138		5,154	1.51	0.07	40	6/15-7/24	0.014

^a Deadloss included in total. ^b Millions of pounds. ^c Information not available.

Table 2. Historical winter commercial and subsistence red king crab fisheries, Norton Sound Section, eastern Bering Sea. Bold typed data are used for the assessment model.

Model Year	Year ^a	Commercial		Winter ^b	Subsistence			Total Crab	
		# of Fishers	# of Crab Harvested		Issued	Permits Returned	Fished	Caught ^c	Retained ^d
1978	1978	37	9,625	1977/78	290	206	149	NA	12,506
1979	1979	1 ^f	221^f	1978/79	48	43	38	NA	224
1980	1980	1 ^f	22^f	1979/80	22	14	9	NA	213
1981	1981	0	0	1980/81	51	39	23	NA	360
1982	1982	1 ^f	17^f	1981/82	101	76	54	NA	1,288
1983	1983	5	549	1982/83	172	106	85	NA	10,432
1984	1984	8	856	1983/84	222	183	143	15,923	11,220
1985	1985	9	1,168	1984/85	203	166	132	10,757	8,377
1986	1985/86	5	2,168	1985/86	136	133	107	10,751	7,052
1987	1986/87	7	1,040	1986/87	138	134	98	7,406	5,772
1988	1987/88	10	425	1987/88	71	58	40	3,573	2,724
1989	1988/89	5	403	1988/89	139	115	94	7,945	6,126
1990	1989/90	13	3,626	1989/90	136	118	107	16,635	12,152
1991	1990/91	11	3,800	1990/91	119	104	79	9,295	7,366
1992	1991/92	13	7,478	1991/92	158	105	105	15,051	11,736
1993	1992/93	8	1,788	1992/93	88	79	37	1,193	1,097
1994	1993/94	25	5,753	1993/94	118	95	71	4,894	4,113
1995	1994/95	42	7,538	1994/95	166	131	97	7,777	5,426
1996	1995/96	9	1,778	1995/96	84	44	35	2,936	1,679
1997	1996/97	2 ^f	83^f	1996/97	38	22	13	1,617	745
1998	1997/98	5	984	1997/98	94	73	64	20,327	8,622
1999	1998/99	5	2,714	1998/99	95	80	71	10,651	7,533
2000	1999/00	10	3,045	1999/00	98	64	52	9,816	5,723
2001	2000/01	3	1,098	2000/01	50	27	12	366	256
2002	2001/02	11	2,591	2001/02	114	61	45	5,119	2,177
2003	2002/03	13	6,853	2002/03	107	70	61	9,052	4,140
2004	2003/04	2 ^f	522^f	2003/04 ^h	96	77	41	1,775	1,181
2005	2004/05	4	2,091	2004/05	170	98	58	6,484	3,973
2006	2005/06	1 ^f	75^f	2005/06	98	97	67	2,083	1,239
2007	2006/07	8	3,313	2006/07	129	127	116	21,444	10,690
2008	2007/08	9	5,796	2007/08	139	137	108	18,621	9,485
2009	2008/09	7	4,951	2008/09	105	105	70	6,971	4,752
2010	2009/10	10	4,834	2009/10	125	123	85	9,004	7,044
2011	2010/11	5	3,365	2010/11	148	148	95	9,183	6,640
2012	2011/12	35	9,157	2011/12	204	204	138	11,341	7,311
2013	2012/13	26	22,639	2012/13	149	148	104	21,524	7,622
2014	2013/14	21	14,986	2013/14	103	103	75	5,421	3,252
2015	2014/15	44	41,062	2014/15	155	153	107	9,840	7,651
2016	2015/16	25	29,792	2015/16	139	97	64	6,468	5,340
2017	2017	43	26,008	2017	163	163	109	7,185	6,039
2018	2018	28	9,180	2018	123	120	82	5,767	4,424
2019	2019	6	1,050	2019	101	101	60	2,080	1,545
2020	2020	1	conf	2020	79	79	50	813	548
2021	2021	5	320	2021	103	103	76	4,655	2,892
2022	2022	8	2,424	2022				9,146	6,743

a Prior to 1985 the winter commercial fishery occurred from January 1 - April 30. As of March 1985, fishing may occur from November 15 - May 15.

b The winter subsistence fishery occurs during months of two calendar years (as early as December, through May).

c The number of crab actually caught, **including females**

d The number of crab retained is the number of crab caught and kept, **including females**

f Confidentiality was waived by the fishers.9963+

h Prior to 2005, permits were only given out of the Nome ADF&G office. Starting with the 2004-5 season, permits were given out in Elim, Golovin, Shaktoolik, and White Mountain.

Table 3. Summary of Norton Sound red king crab trawl survey abundance estimates (x 1000) (CL ≥ 64mm). NMFS and ADF&G trawl survey abundance estimate is based on 10×10 nm² grids, and NBS trawl survey is based on 20×20 nm² grids. Bold typed data are used for the assessment model.

Year	Dates	Survey Agency	Survey method	Abundance ≥64 mm		Female			
				CV	N	% barren	% clutch full	% clutch full 95% CI	
1976	9/02 – 9/25	NMFS	Trawl	4301.8	0.31	181	2.6	66.7	62.4-71.0
1979	7/26 - 8/05	NMFS	Trawl	1457.4	0.22	42	25.0	79.9	64.8-94.8
1980	7/04 - 7/14	ADF&G	Pots	2092.3	N/A				
1981	6/28 - 7/14	ADF&G	Pots	2153.4	N/A				
1982	7/06 - 7/20	ADF&G	Pots	1140.5	N/A				
1982	9/05 - 9/11	NMFS	Trawl	3548.9	0.25	269	0	84.3	81.5-87.2
1985	7/01 - 7/14	ADF&G	Pots	2320.4	0.08				
1985	9/16 -10/01	NMFS	Trawl	2424.9	0.26	151	0	87.5	NA
1988	8/16 - 8/30	NMFS	Trawl	2702.3	0.29	219	1.0	80.7	77.3-84.2
1991	8/22- 8/30	NMFS	Trawl	3132.5	0.43	105	0	69.3	57.7-80.8
1996	8/07 - 8/18	ADF&G	Trawl	1283.0	0.25	168	30.8	71.9	65.9-77.9
1999	7/28 - 8/07	ADF&G	Trawl	2608.0	0.24	81	4.7	80.4	76.0-84.7
2002	7/27 - 8/06	ADF&G	Trawl	2056.0	0.36	168	4.7	76.8	73.4-80.2
2006	7/25 - 8/08	ADF&G	Trawl	3336.0	0.39	194	3.6	67.3	63.2-71.5
2008	7/24 - 8/11	ADF&G	Trawl	2894.2	0.31	116	3.3	56.1	48.5-61.7
2010	7/27 - 8/09	NBS	Trawl	1980.1	0.44	28	0	70.2	63.8-78.5
2011	7/18 - 8/15	ADF&G	Trawl	3209.3	0.29	135	9.8	67.2	61.7-72.6
2014	7/18 - 7/30	ADF&G	Trawl	5934.6	0.47	60	0	60.4	54.3-66.6
2017	7/28 - 8/08	ADF&G	Trawl	1762.1	0.22	43	21.4	71.6	60.0-82.7
2017	8/18 - 8/29	NBS	Trawl	1035.8	0.40	58	0	80.0	72.5-87.5
2018	7/22 - 7/29	ADF&G	Trawl	1108.9	0.25	424	15.8	76.3	59.7-83.5
2019	7/17-7/29	ADF&G	Trawl	4660.8	0.60	386	47.8	50.6	43.1-56.4
2019	8/04-8/07	NBS	Trawl	2532.4	0.26	94	17.6	47.9	36.8-58.9
2020	7/31-8/14	ADF&G	Trawl	1716.5	0.27	186	4.5	66.2	61.6-70.8
2021	7/19-8/03	ADF&G	Trawl	2400.0	0.60	90	3.4	59.8	54.9-64.6
2021	7/29-8/07	NBS	Trawl	2370.0	0.43	139	2.6	61.1	58.8-63.4
2022	8/03-8/12	NBS	Trawl	2103.0	0.37	387	3.5	66.5	64.2-68.7

Abundance of NMFS survey was estimated by NMFS, by multiplying the mean CPUE (# NRKC/nm²) across all hauls (including re-tows) to a standard survey area (7600nm²). Abundance of ADF&G and NBS survey was estimated by ADF&G by multiplying CPUE (# NRKC/nm²) of each station to the grid represented by the station and summing across all surveyed station (ADF&G: 4700 – 5200nm². NBS 5841 nm²).

%barren is calculated by dividing the number of mature females with no eggs by total number of mature females.

Mean and 95% CI of % clutch full is calculated among non-barren mature females. Clutch fullness of each non-barren female was assigned by fullness index that was converted to percentage in the table below.

Clutch fullness index of both NMFS-NBS and ADF&G were converted as follows

NMFS and NBS Code	NMFS and NBS Fullness	Assigned %	ADF&G code	ADF&G Fullness	Assigned %
2	0-1/8	6.25	3	1-29%	15
3	1/8-1/4	18.75	4	30-59%	45
4	1/4 – 1/2	27.5	5	60-89%	75
5	1/2 – 3/4	62.5	6	90-100%	95
6	3/4 – 1	87.5			
7	>1	100			

Table 4. Summer commercial retained catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1977	1549	0	0	0	0.00	0.42	0.34	0.08	0.05	0	0	0	0.00	0.06	0.04	0.01	0.00
1978	389	0	0	0	0.01	0.19	0.47	0.26	0.04	0	0	0	0.00	0.01	0.01	0.01	0.00
1979	1660	0	0	0	0.03	0.23	0.38	0.26	0.07	0	0	0	0.00	0.03	0.00	0.00	0.01
1980	1068	0	0	0	0.00	0.10	0.31	0.37	0.18	0	0	0	0.00	0.00	0.01	0.02	0.01
1981	1784	0	0	0	0.00	0.07	0.15	0.28	0.23	0	0	0	0.00	0.00	0.05	0.12	0.09
1982	1093	0	0	0	0.04	0.19	0.16	0.22	0.29	0	0	0	0.00	0.01	0.02	0.03	0.03
1983	802	0	0	0	0.04	0.41	0.36	0.06	0.03	0	0	0	0.00	0.04	0.01	0.02	0.02
1984	963	0	0	0	0.10	0.42	0.28	0.06	0.01	0	0	0	0.01	0.07	0.05	0.01	0.00
1985	2691	0	0	0.00	0.06	0.31	0.37	0.15	0.02	0	0	0	0.00	0.03	0.03	0.01	0.00
1986	1138	0	0	0	0.03	0.36	0.39	0.12	0.02	0	0	0	0.00	0.02	0.04	0.02	0.00
1987	1985	0	0	0	0.02	0.18	0.29	0.27	0.11	0	0	0	0.00	0.03	0.06	0.03	0.01
1988	1522	0	0.00	0	0.02	0.20	0.30	0.18	0.04	0	0	0	0.01	0.06	0.10	0.07	0.02
1989	2595	0	0	0	0.01	0.16	0.32	0.17	0.05	0	0	0	0.00	0.06	0.12	0.09	0.02
1990	1289	0	0	0	0.01	0.14	0.35	0.26	0.07	0	0	0	0.00	0.04	0.07	0.05	0.01
1991																	
1992	2566	0	0	0	0.02	0.20	0.27	0.14	0.09	0	0	0	0.00	0.08	0.13	0.06	0.02
1993	17804	0	0	0	0.01	0.23	0.39	0.23	0.03	0	0	0	0.00	0.02	0.04	0.03	0.01
1994	404	0	0	0	0.02	0.09	0.08	0.07	0.02	0	0	0	0.02	0.19	0.25	0.20	0.05
1995	1167	0	0	0	0.04	0.26	0.29	0.15	0.05	0	0	0	0.01	0.05	0.07	0.06	0.01
1996	787	0	0	0	0.03	0.22	0.24	0.09	0.05	0	0	0	0.01	0.12	0.14	0.08	0.02
1997	1198	0	0	0	0.03	0.37	0.34	0.10	0.03	0	0	0	0.00	0.06	0.04	0.03	0.01
1998	1055	0	0	0	0.03	0.23	0.24	0.08	0.03	0	0	0	0.02	0.11	0.14	0.08	0.03
1999	562	0	0	0	0.06	0.29	0.24	0.18	0.09	0	0	0	0.00	0.02	0.05	0.04	0.00
2000	17213	0	0	0	0.02	0.30	0.39	0.11	0.02	0	0	0	0.00	0.05	0.07	0.04	0.01
2001	20030	0	0	0	0.02	0.22	0.37	0.21	0.07	0	0	0	0.00	0.02	0.05	0.02	0.01
2002	5219	0	0	0	0.04	0.23	0.28	0.25	0.07	0	0	0	0.00	0.03	0.04	0.03	0.01
2003	5226	0	0	0	0.02	0.37	0.32	0.12	0.03	0	0	0	0.00	0.02	0.05	0.05	0.01
2004	9606	0	0	0	0.01	0.38	0.39	0.11	0.03	0	0	0	0.00	0.03	0.03	0.01	0.01
2005	5360	0	0	0	0.00	0.25	0.47	0.16	0.02	0	0	0	0.00	0.02	0.05	0.02	0.01
2006	6707	0	0	0	0.00	0.18	0.35	0.17	0.02	0	0	0	0.00	0.05	0.14	0.07	0.01
2007	6125	0	0	0	0.01	0.36	0.34	0.14	0.03	0	0	0	0.00	0.02	0.06	0.03	0.01
2008	5766	0	0	0	0.00	0.35	0.35	0.06	0.01	0	0	0	0.00	0.09	0.09	0.04	0.01
2009	6026	0	0	0	0.01	0.34	0.33	0.11	0.02	0	0	0	0.00	0.08	0.08	0.02	0.01
2010	5902	0	0	0	0.01	0.39	0.36	0.10	0.01	0	0	0	0.00	0.05	0.05	0.02	0.00
2011	2552	0	0	0	0.00	0.32	0.40	0.12	0.02	0	0	0	0.00	0.06	0.06	0.02	0.00
2012	5056	0	0	0	0.00	0.24	0.46	0.18	0.02	0	0	0	0.00	0.03	0.04	0.02	0.00
2013	6072	0	0	0	0.00	0.24	0.37	0.24	0.06	0	0	0	0.00	0.01	0.04	0.02	0.00
2014	4682	0	0	0	0.01	0.28	0.24	0.18	0.07	0	0	0	0.00	0.04	0.09	0.07	0.02
2015	4173	0	0	0	0.01	0.48	0.28	0.10	0.03	0	0	0	0.00	0.02	0.03	0.03	0.01
2016	1543	0	0	0	0.00	0.25	0.47	0.16	0.03	0	0	0	0.00	0.02	0.02	0.03	0.01
2017	3412	0	0	0	0.00	0.18	0.39	0.21	0.03	0	0	0	0.01	0.03	0.12	0.05	0.01
2018	2609	0	0	0	0.00	0.11	0.32	0.32	0.08	0	0	0	0	0.01	0.08	0.08	0.02
2019	1136	0	0	0	0.01	0.32	0.23	0.13	0.03	0	0	0	0	0.02	0.10	0.14	0.03
2020																	
2021																	
2022	2981	0	0	0	0.02	0.46	0.30	0.03	0.00	0	0	0	0.00	0.12	0.05	0.01	0.00

Table 5. Winter commercial catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
2015	576	0	0	0	0.07	0.50	0.24	0.06	0.01	0	0	0	0.01	0.04	0.03	0.03	0.01
2016	1016	0	0	0	0.03	0.45	0.31	0.03	0.00	0	0	0	0.01	0.09	0.04	0.02	0.01
2017	540	0	0	0	0.00	0.20	0.30	0.13	0.02	0	0	0	0.00	0.08	0.19	0.06	0.02
2018	401	0	0	0	0.00	0.11	0.25	0.27	0.05	0	0	0	0.04	0.16	0.10	0.02	

Table 6. Summer Trawl Survey length-shell compositions.

Year	Survey	Sample	New Shell								Old Shell							
			64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1976	NMFS	1326	0.01	0.02	0.10	0.19	0.34	0.18	0.02	0.00	0.00	0.00	0.01	0.02	0.03	0.04	0.01	0.01
1979	NMFS	220	0.01	0.01	0.00	0.02	0.05	0.05	0.03	0.01	0.01	0.00	0.01	0.04	0.14	0.40	0.19	0.03
1982	NMFS	327	0.22	0.07	0.16	0.23	0.17	0.03	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.02	0.02	0.03
1985	NMFS	350	0.11	0.11	0.19	0.17	0.16	0.06	0.01	0.00	0.00	0.00	0.00	0.02	0.05	0.08	0.05	0.01
1988	NMFS	366	0.16	0.19	0.12	0.13	0.11	0.06	0.03	0.00	0.00	0.00	0.01	0.01	0.03	0.07	0.05	0.03
1991	NMFS	340	0.18	0.08	0.02	0.03	0.06	0.03	0.01	0.01	0.03	0.06	0.02	0.08	0.16	0.14	0.09	0.02
1996	ADF&G	269	0.29	0.21	0.13	0.09	0.05	0.00	0.00	0.01	0.00	0.00	0.03	0.03	0.04	0.04	0.04	0.03
1999	ADF&G	283	0.03	0.01	0.10	0.29	0.26	0.13	0.03	0.01	0.00	0.00	0.00	0.03	0.05	0.04	0.02	0.00
2002	ADF&G	244	0.09	0.12	0.14	0.11	0.02	0.03	0.02	0.01	0.01	0.03	0.07	0.10	0.09	0.09	0.05	0.02
2006	ADF&G	373	0.18	0.26	0.21	0.11	0.06	0.04	0.02	0.00	0.00	0.00	0.00	0.02	0.04	0.04	0.01	0.00
2008	ADF&G	275	0.12	0.15	0.21	0.11	0.10	0.03	0.02	0.01	0.00	0.01	0.04	0.06	0.08	0.01	0.04	0.00
2010	NOAA	69	0.01	0.04	0.06	0.17	0.06	0.03	0.00	0.00	0.00	0.03	0.09	0.20	0.19	0.07	0.03	0.01
2011	ADF&G	315	0.13	0.11	0.09	0.11	0.18	0.14	0.03	0.01	0.00	0.00	0.01	0.02	0.09	0.04	0.03	0.00
2014	ADF&G	387	0.08	0.15	0.24	0.18	0.09	0.02	0.01	0.01	0.00	0.00	0.03	0.10	0.05	0.04	0.01	0.00
2017	ADF&G	116	0.14	0.12	0.05	0.09	0.10	0.04	0.00	0.00	0.01	0.02	0.02	0.02	0.07	0.18	0.04	0.00
2017	NOAA	58	0.09	0.10	0.14	0.05	0.05	0.05	0.05	0.03	0.03	0.00	0.03	0.05	0.03	0.19	0.05	0.03
2018	ADF&G	73	0.37	0.10	0.11	0.03	0.01	0.03	0.04	0.01	0	0.07	0.01	0.04	0.03	0.03	0.10	0.03
2019	ADF&G	307	0.55	0.30	0.03	0	0.00	0.00	0.00	0	0.00	0.00	0.01	0.02	0.01	0.02	0.03	0.01
2019	NOAA	135	0.36	0.30	0.08	0.04	0.01	0	0.01	0.01	0.04	0.01	0.04	0.02	0.01	0.01	0.04	0.01
2020	ADF&G	111	0.13	0.22	0.30	0.06	0.05	0.01	0	0	0.03	0.08	0.05	0.02	0.02	0.02	0	0.01
2021	ADF&G	158	0.06	0.17	0.22	0.22	0.22	0.04	0.01	0.01	0	0	0.01	0	0.02	0.01	0.01	0.01
2021	NOAA	82	0.05	0.16	0.21	0.16	0.10	0.02	0	0	0.01	0.05	0.11	0.06	0.06	0.01	0	0
2022	NOAA	378	0.16	0.17	0.11	0.10	0.07	0.03	0.01	0.01	0.02	0.02	0.07	0.08	0.087	0.05	0.02	0.01

Table 7. Winter pot survey length-shell compositions.

Year	CPUE	Sample	New Shell								Old Shell							
			64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1981/82	NA	719	0.00	0.10	0.23	0.21	0.07	0.02	0.02	0.00	0.00	0.05	0.11	0.11	0.04	0.02	0.02	0.00
1982/83	24.2	2583	0.03	0.08	0.28	0.28	0.21	0.07	0.01	0.00	0.00	0.00	0.00	0.00	0.02	0.01	0.01	0.01
1983/84	24.0	1677	0.01	0.16	0.26	0.23	0.15	0.06	0.01	0.00	0.00	0.00	0.00	0.02	0.06	0.03	0.01	0.01
1984/85	24.5	789	0.02	0.09	0.25	0.35	0.16	0.06	0.01	0.00	0.00	0.00	0.00	0.01	0.03	0.02	0.00	0.00
1985/86	19.2	594	0.04	0.12	0.17	0.24	0.19	0.08	0.01	0.00	0.00	0.00	0.00	0.01	0.06	0.04	0.01	0.00
1986/87	5.8	144	0.00	0.06	0.15	0.19	0.07	0.04	0.00	0.00	0.00	0.00	0.01	0.04	0.30	0.11	0.03	0.00
1987/88																		
1988/89	13.0	500	0.02	0.13	0.15	0.13	0.19	0.17	0.03	0.00	0.00	0.00	0.00	0.05	0.08	0.03	0.00	
1989/90	21.0	2076	0.00	0.05	0.21	0.26	0.18	0.12	0.06	0.01	0.00	0.00	0.00	0.00	0.03	0.06	0.02	0.00
1990/91	22.9	1283	0.00	0.01	0.09	0.29	0.27	0.10	0.01	0.00	0.00	0.00	0.00	0.00	0.03	0.12	0.07	0.02
1992/93	5.5	181	0.00	0.01	0.03	0.06	0.13	0.12	0.03	0.00	0.00	0.00	0.00	0.02	0.19	0.27	0.10	0.05
1993/94																		
1994/95	6.2	858	0.01	0.06	0.08	0.10	0.26	0.23	0.07	0.01	0.00	0.00	0.00	0.00	0.03	0.07	0.06	0.02
1995/96	9.9	1580	0.06	0.14	0.20	0.19	0.11	0.07	0.03	0.00	0.00	0.00	0.00	0.01	0.06	0.07	0.03	0.01
1996/97	2.9	398	0.07	0.21	0.22	0.11	0.15	0.11	0.05	0.01	0.00	0.00	0.00	0.00	0.02	0.03	0.01	0.01
1997/98	10.9	881	0.00	0.14	0.41	0.27	0.05	0.02	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.02	0.02	0.01
1998/99	10.7	1307	0.00	0.02	0.12	0.36	0.36	0.08	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.01	0.01	0.00
1999/00	6.2	575	0.02	0.09	0.10	0.16	0.33	0.18	0.03	0.00	0.00	0.00	0.00	0.00	0.05	0.02	0.01	0.00
2000/01	3.1	44																
2001/02	13.0	828	0.05	0.29	0.26	0.17	0.06	0.06	0.04	0.01	0.01	0.00	0.01	0.01	0.00	0.01	0.00	0.00
2002/03	9.6	824	0.02	0.10	0.22	0.28	0.18	0.06	0.02	0.00	0.00	0.01	0.01	0.02	0.02	0.03	0.02	0.01
2003/04	3.7	296	0.00	0.02	0.16	0.26	0.32	0.14	0.01	0.00	0.00	0.00	0.01	0.02	0.02	0.01	0.02	0.01
2004/05	4.4	405	0.00	0.07	0.14	0.18	0.22	0.19	0.07	0.00	0.00	0.00	0.00	0.00	0.04	0.06	0.01	0.00
2005/06	6.0	512	0.00	0.14	0.23	0.21	0.16	0.05	0.02	0.00	0.00	0.01	0.01	0.02	0.04	0.07	0.03	0.01
2006/07	7.3	159	0.07	0.14	0.19	0.35	0.13	0.04	0.00	0.00	0.00	0.00	0.01	0.01	0.02	0.04	0.00	0.00
2007/08	25.0	3552	0.01	0.14	0.25	0.17	0.14	0.07	0.01	0.00	0.01	0.04	0.07	0.03	0.03	0.01	0.01	0.00
2008/09	21.9	525	0.00	0.07	0.13	0.35	0.20	0.08	0.01	0.00	0.00	0.00	0.00	0.00	0.04	0.10	0.00	0.00
2009/10	25.3	578	0.01	0.05	0.13	0.21	0.24	0.11	0.02	0.00	0.00	0.00	0.01	0.06	0.10	0.05	0.01	0.00
2010/11	22.1	596	0.02	0.08	0.13	0.20	0.17	0.13	0.05	0.00	0.00	0.00	0.01	0.03	0.11	0.05	0.01	0.00
2011/12	29.4	675	0.03	0.11	0.23	0.19	0.12	0.13	0.04	0.00	0.00	0.00	0.00	0.01	0.05	0.05	0.03	0.00

Table 8. Summer commercial 1987-1994 observer discards length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1987	1146	0.06	0.19	0.32	0.33	0.03	0.00	0.00	0.00	0.00	0.00	0.02	0.04	0.00	0.00	0.00	0.00
1988	722	0.01	0.04	0.15	0.48	0.14	0.00	0.00	0.00	0.00	0.01	0.03	0.10	0.04	0.00	0.00	0.00
1989	1000	0.07	0.19	0.24	0.22	0.03	0.00	0.00	0.00	0.02	0.03	0.07	0.11	0.03	0.00	0.00	0.00
1990	507	0.08	0.23	0.27	0.27	0.04	0.00	0.00	0.00	0.02	0.02	0.02	0.05	0.01	0.00	0.00	0.00
1992	580	0.11	0.17	0.30	0.29	0.03	0.00	0.00	0.00	0.01	0.02	0.02	0.04	0.01	0.00	0.00	0.00
1994	850	0.07	0.06	0.11	0.15	0.02	0.00	0.00	0.00	0.07	0.07	0.15	0.24	0.05	0.00	0.00	0.00

Table 9. Summer commercial observer total catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
2012	3055	0.10	0.05	0.08	0.15	0.15	0.17	0.06	0.01	0.00	0.00	0.00	0.03	0.08	0.09	0.03	0.00
2013	4762	0.19	0.16	0.09	0.10	0.16	0.16	0.09	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.01	0.00
2014	3506	0.02	0.05	0.13	0.22	0.22	0.12	0.08	0.03	0.00	0.00	0.00	0.02	0.03	0.03	0.02	0.01
2015	1671	0.01	0.04	0.09	0.23	0.37	0.14	0.05	0.01	0.00	0.00	0.00	0.01	0.02	0.02	0.01	0.00
2016	2114	0.01	0.01	0.03	0.12	0.29	0.36	0.08	0.02	0.00	0.00	0.00	0.01	0.03	0.03	0.02	0.00
2017	2748	0.02	0.03	0.03	0.06	0.19	0.33	0.18	0.02	0.00	0.00	0.00	0.00	0.02	0.07	0.03	0.01
2018	1628	0.03	0.06	0.12	0.11	0.09	0.17	0.18	0.04	0.00	0.00	0.01	0.01	0.15	0.07	0.08	0.02
2019	236	0.13	0.06	0.06	0.13	0.08	0.05	0.01	0.01	0	0	0.00	0.04	0.11	0.14	0.14	0.05

Table 10. The observed proportion of tagged crab by each size class released and recovered after 1 -3 year of liberty 1980-2019 periods.

Year at liberty 1

	64-73	74-83	84-93	94-103	104-113	114-123	124-33	> 134	n
64-73	0	0.2	0.8	0	0	0	0	0	5
74-83		0	0.44	0.47	0.09	0	0	0	47
84-93			0	0.32	0.62	0.05	0.01	0	146
94-103				0.03	0.62	0.34	0.01	0.00	317
104-113					0.31	0.59	0.09	0	241
114-123						0.42	0.47	0.11	210
124-133							0.69	0.31	81
>134								1	26

Year at liberty 2

	64-73	74-83	84-93	94-103	104-113	114-123	124-33	> 134	n
64-73	0	0	0.09	0.55	0.36	0	0	0	11
74-83		0	0	0.11	0.85	0.04	0	0	113
84-93			0	0.04	0.32	0.61	0.03	0	114
94-103				0.02	0.36	0.41	0.20	0	94
104-113					0.06	0.71	0.22	0	108
114-123						0.17	0.72	0.11	65
124-133							0.36	0.64	25
>134								1	8

Year at liberty 3

	64-73	74-83	84-93	94-103	104-113	114-123	124-33	> 134	n
64-73	0	0	0	0	0.5	0.5	0	0	22
74-83	0	0	0	0	0.26	0.66	0.082	0	73
84-93	0	0	0	0.04	0.26	0.53	0.17	0	53
94-103	0	0	0	0	0.06	0.67	0.27	0	52
104-113	0	0	0	0	0	0.26	0.62	0.12	34
114-123	0	0	0	0	0	0	0.79	0.21	14
124-133	0	0	0	0	0	0	0.1	0.9	10
>134	0	0	0	0	0	0	0	1	1

Table 11. Summary of bounds and model estimated parameters for a length-based population model of Norton Sound red king crab. Parameters with “log_” indicate log scaled parameters

Parameter	Parameter description	Lower	Upper
log q_1	Commercial fishery catchability (1977-93)	-20.5	20
log q_2	Commercial fishery catchability (1994-2007)	-20.5	20
log q_3	Commercial fishery catchability (2008-2019)	-20.5	20
log N_{76}	Initial abundance	2.0	15.0
R_0	Mean Recruit	2.0	12.0
log σ_R^2	Recruit standard deviation	-40.0	40.0
a_{1-7}	Intimal length proportion	0	10.0
$r_{1,2}$	Proportion of length class 1 for recruit	0	5.0
log α	Inverse logistic molting parameter	-5.0	-1.0
log β	Inverse logistic molting parameter	1.0	5.5
log ϕ_{st1}	Logistic trawl selectivity parameter	-5.0	1.0
log ϕ_{wa}	Inverse logistic winter pot selectivity parameter	-5.0	1.0
log ϕ_{wb}	Inverse logistic winter pot selectivity parameter	0.0	6.0
$Sw_{1,2}$	Winter pot selectivity of length class 1,2	0.1	1.0
log ϕ_l	Logistic commercial catch selectivity parameter	-5.0	1.0
log ϕ_{ra}	Logistic summer commercial retention selectivity Newshell (1976-2007, 2008-2022)	-5.0	1.0
log ϕ_{rb}	Logistic summer commercial retention selectivity Newshell (1976-2007, 2008-2022)	0.0	6.0
log ϕ_{rao}	Logistic summer commercial retention selectivity Oldshell (1976-2007, 2008-2022)	-5.0	1.0
log ϕ_{rbo}	Logistic summer commercial retention selectivity Oldshell (1976-2007, 2008-2022)	0.0	6.0
log ϕ_{wra}	Logistic winter commercial retention selectivity p	-5.0	1.0
log ϕ_{wrb}	Logistic winter commercial retention selectivity	0.0	6.0
w^2_t	Additional variance for standard CPUE	0.0	6.0
m_{1-8}	Natural mortality multipliers	0	5.0
$q_{.1}$	Survey q for NMFS trawl 1976-91	0.1	1.0
$q_{.2}$	Survey q for NMFS NBS trawl 2010,17,19	0.1	1.0
σ	Growth transition sigma	0.0	30.0
β_1	Growth transition mean	0.0	20.0
β_2	Growth transition increment	0.0	20.0

*: Parameter was unestimable because model estimated trawl survey selectivity was 1.0 across all size classes.

Name	21.0 Final		21.0		22.0		22.1		22.2	
	Estimate	std.dev	Estimate	std.dev	Estimate	std.dev	Estimate	std.dev	Estimate	std.dev
log q ₁	-7.315	0.195	-7.313	0.195	-7.297	0.196	-7.358	0.197	-7.343	0.198
log q ₂	-6.721	0.166	-6.721	0.166	-6.724	0.166	-6.699	0.169	-6.697	0.169
log q ₃	-6.807	0.156	-6.811	0.156	-6.790	0.153	-6.811	0.159	-6.814	0.159
log N ₇₆	9.136	0.139	9.133	0.139	9.124	0.138	9.331	0.165	9.318	0.165
R ₀	6.454	0.082	6.449	0.083	6.450	0.083	6.485	0.126	6.482	0.126
a ₁	1.016	4.460	1.038	4.456	1.068	4.474	0.983	4.420	0.982	4.439
a ₂	1.016	4.460	1.763	4.190	1.787	4.209	1.693	4.165	1.685	4.183
a ₃	1.749	4.191	3.497	3.932	3.520	3.952	3.519	3.906	3.525	3.924
a ₄	3.492	3.932	3.979	3.910	4.000	3.930	3.998	3.884	4.006	3.901
a ₅	3.978	3.910	4.246	3.901	4.287	3.922	4.227	3.875	4.259	3.893
a ₆	4.243	3.902	3.506	3.930	3.520	3.950	3.482	3.905	3.498	3.922
a ₇	3.503	3.930	2.062	4.198	2.077	4.212	1.881	4.177	1.891	4.190
r1	5.000	0.002	5.000	0.002	5.000	0.002	5.000	0.003	5.000	0.003
r2	4.624	0.163	4.617	0.167	4.610	0.168	4.651	0.190	4.648	0.191
log a	-2.750	0.090	-2.729	0.089	-2.711	0.089	-2.690	0.091	-2.685	0.090
log b	4.835	0.016	4.834	0.015	4.827	0.015	4.824	0.015	4.821	0.015
log ϕ_{st}	-5.000	0.028	-5.000	0.032	-5.000	0.033	-5.000	0.070	-5.000	0.067
log ϕ_{wa}	-2.392	0.421	-2.398	0.422	-2.410	0.427	-2.322	0.466	-2.343	0.475
log ϕ_{wb}	4.773	0.067	4.771	0.069	4.770	0.071	4.805	0.056	4.802	0.060
Sw1	0.061	0.033	0.060	0.033	0.060	0.033	0.072	0.038	0.071	0.038
Sw2	0.425	0.146	0.425	0.147	0.423	0.149	0.480	0.151	0.474	0.153
Sw3	0.731	0.234	0.727	0.236	0.718	0.238	0.778	0.216	0.769	0.222
log ϕ_l	-2.063	0.044	-2.061	0.044	-2.039	0.045	-2.036	0.048	-2.037	0.048
log ϕ_{ra1}	-0.856	0.143	-0.855	0.143	-0.809	0.157	-0.881	0.145	-0.839	0.159
log ϕ_{rb1}	4.641	0.008	4.641	0.008	4.637	0.009	4.645	0.009	4.641	0.010
log ϕ_{ra2}	-0.491	0.278	-0.497	0.276	0.906	560.740	-0.500	0.280	-0.361	0.491
log ϕ_{rb2}	4.653	0.013	4.654	0.013	4.606	8.653	4.654	0.014	4.645	0.025
log ϕ_{rao1}					-1.036	0.448			-1.099	0.469
log ϕ_{rbo1}					4.661	0.019			4.667	0.020
log ϕ_{rao2}					-1.969	0.704			0.849	524.780
log ϕ_{rbo2}					4.711	0.040			4.685	0.860
log ϕ_{wra}	-0.948	0.566	-0.949	0.564	-0.945	0.573	-0.970	0.561	-0.983	0.552
log ϕ_{wrb}	4.654	0.038	4.654	0.038	4.653	0.039	4.654	0.038	4.655	0.038
w ² _t	0.142	0.040	0.142	0.040	0.141	0.040	0.139	0.039	0.140	0.039
q.1	0.713	0.128	0.716	0.128	0.723	0.129	0.634	0.117	0.641	0.118
q.2	0.871	0.173	0.891	0.191	0.883	0.189	0.861	0.185	0.861	0.184
σ	3.815	0.208	3.824	0.208	3.792	0.211	3.809	0.214	3.782	0.216
β_1	11.788	0.696	11.808	0.697	11.668	0.714	11.960	0.754	11.868	0.760
β_2	7.813	0.171	7.806	0.171	7.850	0.176	7.786	0.186	7.816	0.188
m1							0.000	0.000	0.000	0.002
m2							0.000	0.754	0.000	0.000
m3							0.928	0.593	1.005	0.754
m4							1.249	0.395	1.268	0.593
m5							1.224	0.294	1.107	0.396
m6							1.680	0.339	1.708	0.296
m7							3.033	0.574	3.041	0.340
m8	3.815	0.208	3.428	0.266	3.427	0.266	3.746	0.214	3.752	0.574

Table 12. Annual abundance estimates of legal crab (million crab) and mature male biomass (Feb 01) (MMB, million lb) for Norton Sound red king crab estimated by a length-based analysis.

MMB

Year	Model 21.0 final	Model 21.0	Model 22.0	Model 22.1	Model 22.2
1976	17.33	17.45	17.29	21.02	20.84
1977	19.15	19.24	19.06	22.27	22.11
1978	16.71	16.76	16.61	18.61	18.50
1979	11.86	11.90	11.79	12.88	12.80
1980	6.42	6.44	6.37	6.95	6.90
1981	4.36	4.38	4.34	4.71	4.67
1982	3.40	3.42	3.38	3.77	3.73
1983	4.30	4.33	4.28	4.79	4.73
1984	5.04	5.07	5.01	5.61	5.53
1985	5.70	5.73	5.65	6.32	6.22
1986	6.05	6.09	5.99	6.65	6.54
1987	5.96	5.99	5.88	6.48	6.37
1988	5.88	5.91	5.81	6.36	6.27
1989	5.62	5.65	5.57	6.02	5.96
1990	5.31	5.33	5.28	5.66	5.62
1991	5.00	5.02	4.99	5.32	5.29
1992	4.76	4.78	4.76	5.01	5.00
1993	4.45	4.47	4.47	4.65	4.64
1994	3.85	3.87	3.87	3.99	3.99
1995	3.29	3.30	3.32	3.39	3.40
1996	2.90	2.91	2.94	2.99	2.99
1997	2.88	2.90	2.93	2.97	2.98
1998	3.49	3.52	3.54	3.63	3.63
1999	4.38	4.42	4.44	4.59	4.59
2000	4.62	4.66	4.68	4.76	4.77
2001	4.18	4.21	4.23	4.23	4.24
2002	3.98	4.01	4.03	4.04	4.05
2003	4.07	4.10	4.14	4.18	4.19
2004	4.08	4.11	4.16	4.21	4.22
2005	3.81	3.83	3.89	3.92	3.93
2006	3.57	3.60	3.66	3.70	3.71
2007	3.74	3.78	3.84	3.95	3.95
2008	4.26	4.31	4.38	4.55	4.54
2009	4.72	4.78	4.87	5.08	5.07
2010	4.92	4.98	5.10	5.28	5.28
2011	4.60	4.65	4.77	4.88	4.89
2012	4.15	4.19	4.28	4.35	4.36
2013	3.96	3.99	4.05	4.17	4.16
2014	4.36	4.40	4.44	4.69	4.67
2015	4.91	4.96	5.01	5.30	5.30
2016	4.50	4.54	4.61	4.77	4.78
2017	3.67	3.70	3.76	3.78	3.80
2018	2.82	2.85	2.88	2.83	2.85
2019	2.39	2.42	2.43	2.37	2.37
2020	3.00	3.08	3.08	3.04	3.04
2021	4.45	4.64	4.65	4.57	4.57
2022	5.31	5.52	5.52	5.30	5.31
2023	5.29	5.39	5.39	5.05	5.06

Legal abundance (x million) (≥ 104 mm CL)

Year	Model 21.0 Final	Model 21.0	Model 22.0	Model 22.1	Model 22.2
1976	5.02	7.39	7.33	8.98	8.89
1977	6.15	7.14	7.08	8.32	8.25
1978	5.22	5.59	5.54	6.23	6.19
1979	3.51	3.77	3.74	4.09	4.07
1980	1.83	2.02	2.01	2.19	2.18
1981	1.15	1.46	1.45	1.59	1.58
1982	0.84	1.34	1.33	1.50	1.48
1983	1.14	1.76	1.74	1.97	1.94
1984	1.40	2.03	2.00	2.27	2.24
1985	1.60	2.25	2.21	2.51	2.47
1986	1.76	2.31	2.27	2.56	2.51
1987	1.71	2.22	2.18	2.43	2.39
1988	1.70	2.15	2.12	2.35	2.32
1989	1.62	2.01	1.99	2.18	2.16
1990	1.51	1.90	1.89	2.05	2.04
1991	1.43	1.78	1.77	1.92	1.91
1992	1.36	1.68	1.68	1.79	1.78
1993	1.27	1.57	1.57	1.65	1.65
1994	1.10	1.35	1.36	1.42	1.42
1995	0.93	1.16	1.17	1.21	1.22
1996	0.80	1.05	1.06	1.10	1.10
1997	0.78	1.09	1.10	1.13	1.13
1998	0.89	1.40	1.40	1.47	1.47
1999	1.24	1.74	1.75	1.83	1.83
2000	1.40	1.71	1.72	1.77	1.77
2001	1.22	1.49	1.50	1.52	1.52
2002	1.09	1.46	1.47	1.50	1.50
2003	1.11	1.54	1.56	1.61	1.61
2004	1.17	1.53	1.55	1.60	1.60
2005	1.11	1.40	1.42	1.45	1.46
2006	0.98	1.34	1.36	1.41	1.41
2007	0.99	1.48	1.50	1.58	1.57
2008	1.16	1.71	1.74	1.84	1.84
2009	1.32	1.88	1.92	2.04	2.03
2010	1.44	1.90	1.94	2.04	2.04
2011	1.37	1.69	1.73	1.80	1.80
2012	1.18	1.51	1.54	1.59	1.59
2013	1.07	1.51	1.52	1.60	1.59
2014	1.13	1.76	1.77	1.91	1.90
2015	1.41	1.95	1.97	2.11	2.11
2016	1.38	1.66	1.68	1.76	1.76
2017	1.11	1.27	1.29	1.31	1.32
2018	0.81	0.97	0.98	0.97	0.98
2019	0.65	0.86	0.86	0.85	0.85
2020	0.68	1.26	1.26	1.26	1.26
2021	1.19	1.92	1.93	1.92	1.92
2022	1.58	2.13	2.13	2.06	2.06
2023	1.54	1.98	1.98	1.88	1.88

Table 13. Summary of observed catch (million lb) for Norton Sound red king crab.

Year	Summer Com	Winter Com	Winter Sub	Discards Winter Sub	Total
1977	0.52	0.000	0.000	0	0.520
1978	2.09	0.024	0.025	0.008	2.147
1979	2.93	0.001	0.000	0	2.931
1980	1.19	0.000	0.000	0	1.190
1981	1.38	0.000	0.001	0	1.381
1982	0.23	0.000	0.003	0.001	0.234
1983	0.37	0.001	0.021	0.006	0.398
1984	0.39	0.002	0.022	0.005	0.419
1985	0.43	0.003	0.017	0.002	0.452
1986	0.48	0.005	0.014	0.004	0.503
1987	0.33	0.003	0.012	0.002	0.347
1988	0.24	0.001	0.005	0.001	0.247
1989	0.25	0.000	0.012	0.002	0.264
1990	0.19	0.010	0.024	0.004	0.228
1991	0	0.010	0.015	0.002	0.027
1992	0.07	0.021	0.023	0.003	0.117
1993	0.33	0.005	0.002	0	0.337
1994	0.32	0.017	0.008	0.001	0.346
1995	0.32	0.022	0.011	0.002	0.355
1996	0.22	0.005	0.003	0.001	0.229
1997	0.09	0.000	0.001	0.001	0.092
1998	0.03	0.002	0.017	0.012	0.061
1999	0.02	0.007	0.015	0.003	0.045
2000	0.3	0.008	0.011	0.004	0.323
2001	0.28	0.003	0.001	0	0.284
2002	0.25	0.007	0.004	0.003	0.264
2003	0.26	0.017	0.008	0.005	0.290
2004	0.34	0.001	0.002	0.001	0.344
2005	0.4	0.006	0.008	0.003	0.417
2006	0.45	0.000	0.002	0.001	0.453
2007	0.31	0.008	0.021	0.011	0.350
2008	0.39	0.015	0.019	0.009	0.433
2009	0.4	0.012	0.010	0.002	0.424
2010	0.42	0.012	0.014	0.002	0.448
2011	0.4	0.009	0.013	0.003	0.425
2012	0.47	0.025	0.015	0.004	0.514
2013	0.35	0.061	0.015	0.014	0.440
2014	0.39	0.035	0.007	0.002	0.434
2015	0.40	0.099	0.019	0.005	0.523
2016	0.42	0.080	0.011	0.001	0.512
2017	0.41	0.078	0.012	0.001	0.501
2018	0.30	0.029	0.008	0.001	0.338
2019	0.08	0.032	0.003	0.001	0.116
2020	0	Conf.	0.001	0.000	Conf
2021	0	0.0	0.004	0.002	0.006
2022	0.32	0.070	0.006	0.003	0.400

Table 14: Jittering

replicate	likelihood	gradient	replicate	likelihood	gradient
1	NaN	3.63E+02	51	353.4257	5.95E-07
2	NaN	1.03E+03	52	354.6813	3.06E-06
3	354.7677	4.12E-06	53	353.1181	3.04E-07
4	NaN	1.93E+02	54	353.4668	1.17E-06
5	354.2047	2.28E-06	55	352.535	5.84E-06
6	353.9893	3.86E-06	56	354.4949	7.88E-08
7	355.339	4.91E-07	57	675.3477	3.80E+04
8	404.9138	2.82E+03	58	353.4821	1.34E-07
9	355.1658	1.68E-06	59	353.2222	5.38E-08
10	354.3164	9.41E-06	60	354.1651	1.14E-06
11	353.9742	8.86E-06	61	353.7704	2.64E-05
12	354.3997	8.54E-07	62	353.5346	1.09E-06
13	354.4445	2.22E-06	63	353.5866	1.34E-05
14	353.2734	2.05E-07	64	353.8928	5.23E-05
15	354.0722	5.74E-07	65	353.1969	2.28E-05
16	353.9825	1.82E-05	66	354.2969	2.13E-05
17	354.8302	9.90E-07	67	353.6836	3.35E-08
18	354.2344	7.64E-05	68	568123.7	4.11E+05
19	353.5129	1.29E-06	69	355.4708	3.14E-08
20	353.6984	6.07E-07	70	352.9499	2.92E-06
21	354.8259	8.81E-06	71	353.9816	6.30E-08
22	355.6285	2.57E-06	72	354.4901	5.71E-07
23	355.7144	1.22E-07	73	353.725	8.31E-06
24	353.7708	5.08E-06	74	354.5325	2.47E-05
25	353.2279	1.20E-05	75	353.929	1.08E-05
26	355.2713	2.85E-07	76	410.8851	2.23E+02
27	355.3182	5.23E-06	77	354.6617	2.85E-05
28	354.4986	1.19E-07	78	353.9638	2.33E-05
29	353.0543	3.09E-07	79	353.3969	3.02E-06
30	353.7164	4.49E-07	80	353.2247	7.15E-07
31	353.2282	7.52E-07	81	NaN	1.74E+03
32	354.0614	1.56E-06	82	NaN	3.07E+02
33	353.8865	1.40E-05	83	354.6141	7.40E-07
34	354.2521	4.80E-06	84	354.9823	3.37E-07
35	355.4237	1.85E-07	85	353.144	3.64E-06
36	354.6227	1.10E-07	86	355.0196	4.33E-07
37	354.7637	3.43E-07	87	354.4193	2.35E-06
38	354.6088	6.82E-07	88	354.3445	7.34E+02
39	354.0906	9.60E-03	89	353.6029	1.83E-05
40	354.8387	4.89E-07	90	354.9923	4.74E-06
41	353.6605	7.34E-08	91	353.9155	8.37E-06
42	354.2417	3.55E-07	92	353.8707	5.32E-07
43	353.528	1.26E-06	93	393.468	3.03E-07
44	354.2095	7.95E-05	94	353.7274	1.08E-06
45	NaN	6.05E+02	95	354.3414	1.64E-05
46	354.528	5.21E-06	96	354.7069	5.12E-06
47	352.8339	7.16E-08	97	354.3749	8.77E-06
48	390.5528	4.89E-05	98	355.3819	4.66E-07
49	393.4449	8.52E-05	99	353.8777	8.60E-07
50	353.9817	2.68E-06	100	NaN	1.91E+02

Figure 1. King crab fishing districts and sections of Statistical Area Q.




Figure 2. Closed water regulations in effect for the Norton Sound commercial crab fishery. Line around the coastline delineates the 3-mil state waters zone.




Figure 3. Model estimated natural mortality, annual molting probability, selectivity for trawl survey, winter pot survey, and summer commercial fishery, and retention probability for winter commercial and summer commercial (newshell and oldshell (for models 22.0 and 22.1)). X-axis is carapace length (mm).




Figure 4. Model estimated transition probability for each size classes.




Figure 5. Model estimated abundances of total, legal (CL>104 mm) and recruit (CL 64-94 mm) males during 1976-2022.




Figure 5. Estimated MMB during 1976-2023. Horizontal line Bmsy (Average MMB of 1980-2023).




Figure 6. Observed (open circle) (White: NMFS, Red: ADF&G) and model trawl survey male abundances with 95% lognormal Confidence Intervals (crab ≥ 64 mm CL).




Figure 7. Observed (open circle) with 95% lognormal Confidence Intervals with additional variance (gray), and model estimated standardized CPUE.




Figure 8. Predicted (line) vs. observed (bar New Shell: blue, Old Shell: green) length class proportions for the summer commercial harvest 1977-2022.




Figure 9. Predicted (line) vs. observed (bar New Shell: blue, Old Shell: green) length class proportions for trawl survey 1976-2022.




Figure 10. Predicted (line) vs. observed (bar New Shell: blue, Old Shell: green) length class proportions for winter pot survey 1982-2012.




Figure 11. Predicted (line) vs. observed (bar New Shell: left blue, Old Shell: right green) length class proportions for summer commercial total and discards (1987-1994, 2012-2019) and winter commercial retained fishery 2015-2018




Figure 12. Predicted (line) vs. observed (bar) length class proportions for tag recovery data.







Figure 13. Input vs. model implied effective sample size. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the harmonic mean of implied sample size. Figures in the second column show input sample sizes (x-axis) vs. implied effective sample sizes (y-axis). Dashed line indicates the linear regression slope, and solid line is 1:1 line. Figures in the third column show years (x-axis) vs. implied effective sample sizes (y-axis). Horizontal solid line is the harmonic mean of implied sample size.




Model 21.0



Model 22.0



Model 22.1



Model 22.2









Figure 14. Bubble plots of predicted and observed length proportions. Black circles indicate model underestimates compared to observed, and white circles indicate model overestimates compared to observed. Size of circle indicates degree of deviance (larger circle = larger deviance). In ideal model fit case, distribution of sizes and colors of circles should be random (i.e., no systematic model misfits).



















Figure 15. Standardized Pearson residual plots for trawl survey, summer commercial retained catch, winter pot survey, and observer for length size classes 1-8.


















Figure 16. QQ Plot of Trawl survey and Commercial CPUE.







Retrospective Analysis Model 21.0




Figure 17. Retrospective Analyses of Norton Sound Red King Crab MMB from 2012 to 2023. Solid black line: 2023 assessment model results.