


Ecosystem Status Report: Gulf of Alaska 2023



Bridget Ferriss

With contributions from:

Anna Abelman, Grant Adams, Zoe Almeida, Mayumi Arimitsu, Kelia Axler, Kerim Aydin, Brenda Ballachey, Sonia Batten, James Bodkin, Nick Bond, Maile Branson, Brian Brettschneider, Matt W. Callahan, Rob Campbell, Reuben Cash, Louisa Castrodale, Shannon Cellan, Wei Cheng, Heather Coletti, Bryan Cormack, Deana Crouser, Daniel Cushing, Seth Danielson, Alison Deary, Alexandra Dowlin, Sherri Dressel, Brie Drummond, Kathleen Easley, Grace Ellwanger, Daniel Esler, Stacie Evans, Ed Farley, Thomas Farrugia, Will Fennie, Emily Fergusson, Teresa Fish, Christine Gabriele, Jeanette Gann, Andrew Gray, Scott Hatch, Kyle Hebert, Tyler Hennon, Kirstin Holsman, Russell Hopcroft, John Horne, Peter Hulson, Jim Ianelli, Katrin Iken, Annette Jarosz, Timothy Jones, Robert Kaler, Julie Keister, Tom Kelley, Mandy Keogh, David Kimmel, Kim Kloecker, Brenda Konar, Elizabeth Labunski, Ned Laman, Jesse Lamb, Geoffrey M. Lang, Kari Lanphier, Ben Laurel, Emily Lemagie, Mandy Lindeberg, Jackie Lindsey, Emily Mailman, Jasmine Maurer, Lauren E. McCaslin, Jessica Miller, Cole Monnahan, Daniel Monson, John Moran, Calvin Mordy, Jennifer Morella, Jens Nielsen, Cecilia O'Leary, John Olson, Clare Ostle, Janet Neilson, Emma Pate, W. Scott Pegau, Noel Pelland, John Piatt, Cody Pinger, Andrew Piston, Meredith Pochardt, André Punt, Heather Renner, Felipe Restrepo, Brian Robinson, Lauren Rogers, Sean Rohan, Natalie Rouse, Kalei Shotwell, Kevin Siwicke, Scott Smeltz, Mason Smith, Hank Statscewich, Ian Stewart, William Stockhausen, Wesley Strasburger, Janice Straley, Suzanne Strom, Robert Suryan, Hillary Thalmann, Rick Thoman, Sarah Traiger, Scott Vulstek, Andy Wall, Benjamin Weitzman, Shannon Whelan, George A. Whitehouse, Carrie Worton, Ellen Yasumiishi, Stephani Zador, Joshua Zahner, and Molly Zaleski



GOA 2023: Key Messages



1. 2023 Average productivity; Spatially variable; 4-year consistent environmental drivers



2. Variable pelagic prey base (zooplankton, forage fish); Reduced from 2022



3. Pacific cod & capelin show signs of increase (first since marine heatwave years)



4. Looking ahead to 2024 (El Niño): which groundfish are vulnerable & which might benefit?



GOA Full & Update Assessment Risk Tables: Environmental/ Ecosystem Considerations

Level 1

*(No apparent environmental/
ecosystem concerns)*

- Walleye pollock (+ESP)
- Pacific cod (+ESP)
- Sablefish (statewide) (+ESP)
- Pacific ocean perch
- Deepwater flatfish
- Rougheye/blackspotted rockfish
- Shortraker rockfish
- Other rockfish
- Skates

Environment: ocean temperatures cool to average

Prey:

- Pelagic: (Zooplankton & forage fish) reduced/variable
- Benthic & infauna: variable /unknown


Predation: relatively low (P. cod, P. halibut, arrowtooth flounder) except sablefish; no major changes in seabirds & marine mammal populations

Competition: potentially higher for zooplankton-eating groundfish due to pink salmon, P. ocean perch, pollock


2024: El Niño primarily impact larval survival & prey base for juveniles and zooplankton-eating adults

Reduced primary productivity

J. Gann, M. Callahan



- Satellite-derived chl-a (1998-2023)
- Indicated low phytoplankton biomass
- Late timing of the chl-a spring bloom
- Unique in time series




→ *2024 early, larger phytoplankton spring bloom*

Zooplankton

D. Kimmel, R. Hopcroft, E. Fergusson

WGOA




- Surveys: EcoFOCI Shelikof spring, Seward Line spring, Icy Strait (SEAK) summer
- Lower total zooplankton biomass than 2022 (below average to average)
- Higher biomass of large copepods & euphausiids (Shelikof, Seward)
- Energy density (lipid content) above average in Icy St.

→ 2024 community shift to smaller copepods

EGOA

Forage Fish Prey Base: variable

B. Drummond, D. Cushing, S. Hatch, K. Hebert, S. Pegau, E. Pochardt, W. Strasburger, C. Worton



Prey for:
W. pollock
P. cod
 Arrowtooth fl.
 Sablefish
 Flatfish

- Survey baselines from 1990's/early 2000's to present)
- Below to above average
- Incr: capelin, herring
- Decr.: sandlance, juv. salmon, age-0 pollock

Legend



- █ Surveys
- Seabird Reproductive Success
- + Above Average
- ~ Average
- Below Average

Capelin returning in core habitat


S. Whelan, D. McGowan, L. Rogers, N. Laman, Skipper Science

- Capelin are rebounding in their core habitat (at least)
- Present around Kodiak, Middleton Isl., Chowiet Isl., Sitka
- Low biomass around AK peninsula (NOAA EcoFOCI summer survey) & eastern GOA (NOAA summer acoustic survey & Bottom trawl survey)

NOAA summer acoustic survey



Black-legged kittiwake diet (June-Aug) Middleton Island (CGOA)




Capelin

- Capelin observed around Sitka;
- Not uncommon but hadn't seen since heatwave years;
- Observed in chinook salmon stomachs in the area;
- Observed large groups of seabirds (rhinoceros auklets and murrelets) feeding in these areas
- (*synthesized from Skipper Science*)

GOA Pacific Cod age 0 & 1: 2017-2023

B. Laurel et al.

Age 0 & 1 Year Class Strength of GOA Pacific Cod



Fluctuations in year class strength

- Marine heatwave: warm SST & spawning habitat/egg survival (2014-2016, 2019)
- Warm fall SST (2017, 2018)

Surveys: NOAA beach seine Kodiak (since 2006) & AK peninsula (since 2018); NOAA EcoFOCI spring (odd years) and summer (2023, 2019)

Looking ahead to 2024 (El Niño): warming SST

N. Bond

- National Multi-Model Ensemble predictions of sea surface temperatures in 2024 (baseline: 1991-2020)

VULNERABLE? (larvae favor cooler springs):

P. cod yolk-sac larvae & feeding larvae (5-6°C)

W. pollock yolk-sac larvae (3-7°C)

N. rock sole larvae




BENEFIT? (larvae favor warm springs):

Sablefish larvae & YOY (12-16°C)

S. rock sole larvae

P. ocean perch larvae

Rockfish larvae



Looking ahead to 2024 (El Niño)

VULNERABLE (?) 2024

SST (-)

Zoop (-)

P. cod: Larvae

W. pollock: Larvae, Adult

N. rock sole: Larvae, Adult

P. ocean perch: Adult

Dusky rockfish: Adult

BENEFIT (?) 2024

SST (+)

Zooplankton (-)

Transport (+)

S. rock sole: Larvae

P. ocean perch: Larvae

Rockfish: Larvae

Sablefish: Larvae

Arrowtooth flounder: Larvae

Rex sole: Larvae

P. halibut: Larvae

Looking ahead to 2024 (El Niño)

VULNERABLE (?) 2024

SST (-)

Zoop (-)

P. cod: Larvae

W. pollock: Larvae, Adult

N. rock sole: Larvae, Adult

P. ocean perch: Adult

Dusky rockfish: Adult



BENEFIT (?) 2024

SST (+)

Zooplankton (-)

Transport (+)

S. rock sole: Larvae

P. ocean perch: Larvae

Rockfish: Larvae

Sablefish: Larvae

Arrowtooth flounder: Larvae

Rex sole: Larvae

P. halibut: Larvae

Looking ahead to 2024 (El Niño)

VULNERABLE (?) 2024

P. cod: Larvae

W. pollock: Larvae, Adult

N. rock sole: Larvae, Adult

P. ocean perch: Adult

Dusky rockfish: Adult

BENEFIT (?) 2024

S. rock sole: Larvae

P. ocean perch: Larvae

Rockfish: Larvae

Sablefish: Larvae

Arrowtooth flounder: Larvae

Rex sole: Larvae

P. halibut: Larvae

SST (-)

Zoop (-)

SST (+)

Zooplankton (-)

Transport (+)



Questions:

- How would a low recruitment year in 2023 affect populations? 2 low recruitment years ('23/'24)?
- If heat persists and mixes to depth (late 2024/2025?): which adult populations are vulnerable?
- Do populations have a buffer for unknown/indirect/cumulative ecological responses?

GOA 2023: Key Messages

1. 2023 Average productivity; Spatially variable; 4-year consistent environmental drivers

- Regional variability, average productivity, 3 La Niña's

2. Variable pelagic prey base; Reduced from 2022


- Zooplankton: below average to average
- Forage fish: below average to above average

3. Pacific cod & capelin show signs of increase (first since marine heatwave years)


4. Looking ahead to 2024 (El Niño):

- Warm surface temperatures, potentially lower quality zooplankton prey, increased cross shelf transport
- Larval & juvenile groundfish most sensitive (some vulnerable, some benefit)
- Adult POP, pollock, dusky rockfish, n. rock sole are more vulnerable


Where are we headed (2024 +): El Niño



Aleutian Low: deeper and displaced SE



Warmer & wetter winter




Shallower mixed layer depth (earlier stratification)



Earlier & more intense spring Bloom


?




Jan, Feb Mar (2024)



Warmer sea surface temperature



Less lipid rich zooplankton community



Earlier phenology

Optimal thermal ranges for groundfish?