

BSAI Alaska Plaice Transition to Stock Synthesis

Lee Cronin-Fine September 2024

Outline

•Transition to Stock Synthesis (SS3)

Updating new base model



 Model: Sex specific, age and length population dynamics model coded in ADMB



 Model: Sex specific, age and length population dynamics model coded in ADMB

 Data: catch, Biomass Index from survey, age composition data from the fishery and survey and length composition data from the fishery and survey.



 Model: Sex specific, age and length population dynamics model coded in ADMB

 Data: catch, Biomass Index from survey, age composition data from the fishery and survey and length composition data from the fishery and survey.

• Sex specific: length-at-age, weight-at-length, weight-at-age, age-length transition matrix (ALK) and selectivity.

• Estimated Externally: length-at-age (2016), weight-at-age (2016), weight-at-length (2016), ALK (<2003), age-at-maturity (2014).



• Estimated Externally: length-at-age (2016), weight-at-age (2016), weight-at-length (2016), ALK (<2003), age-at-maturity (2014).

Fixed Parameters: survey catchability, natural mortality



• Estimated Externally: length-at-age (2016), weight-at-age (2016), weight-at-length (2016), ALK (<2003), age-at-maturity (2014).

 Fixed Parameters: survey catchability, natural mortality.

• Estimated Parameters: Mean recruitment, recruitment deviations, annual fishing mortality, Age-based selectivity (fishery and survey).



• Estimated Externally: length-at-age (2016), weight-at-age (2016), weight-at-length (2016), ALK (<2003), age-at-maturity (2014).

- Fixed Parameters: SUrvey catchability, natural mortality
- **Estimated Parameters**: Age-based selectivity (fishery and survey), R0, recruitment deviations, annual fishing mortality.
- Population age range 3-25, length range 10cm 60cm

Why Transition to SS3?

Estimate more parameters internally

More flexible

Better external resources

Better documented



Transitioning to SS3

- •Base-1
- •Base-2
- •Base-3
- ·Base-4




Deterministic Model

•All parameters fixed except annual fishing mortality and selectivity






Biomass Index




Model - Old - Base-1





Model - Old - Base-1





Growth in Old Model

Length-at-age


<u>ALK</u>


2016

< 2003

Length-at-age * Weightat-length = Weight-at-age Length-composition








•Updated version of Base-1 with new growth curve

 All parameters fixed except annual fishing mortality and selectivity











Model - Old - Base-1 - Base-2









Updated version of Base-2


• Estimate mean recruitment, recruitment deviations, annual fishing mortality and selectivity

Use new growth curve




Biomass Index





Model - Old - Base-1 - Base-2 - Base-3







Likelihood Components

	Base-1	Base-2	Base-3
Total	1623.974	992.067	982.7518
Survey	-58.9898	-59.209	-58.8731
Length-comp	1007.89	383.985	373.708
Age-comp	656.727	648.944	644.961
Recruitment	18.347	18.347	22.9559




Updated version of Base-3

 Estimate mean recruitment, recruitment deviations, annual fishing mortality and selectivity, growth




Total Biomass, age 3 up






Model - Old - Base-1 - Base-2 - Base-3 · Base-4









Likelihood Components

	Base-1	Base-2	Base-3	Base-4
Total	1623.974	992.067	982.7518	985.547
Survey	-58.9898	-59.209	-58.8731	-58.9021
Length- comp	1007.89	383.985	373.708	374.879
Age-comp	656.727	648.944	644.961	645.987
Recruitment	18.347	18.347	22.9559	23.5831



Update/Candidate Models

·Base-4

•Model 24.0



- Updated survey age-composition input sample size
- General bootstrap framework in "surveyISS" Rpackage (Williams and Hulson 2024)

^{*} Williams B, Hulson P (2024). *surveyISS: Survey composition input sample size*. R package version 1.0.0, https://benwilliams-noaa.github.io/surveyISS/, https://github.com/BenWilliams-NOAA/surveyISS/.



- Updated survey age-composition input sample size
- Update survey length-composition input sample size
 - Number of hauls



- Updated survey age-composition input sample size
- Update survey length-composition input sample size
- Add age-1 and age-2 to agecomposition data




- Updated survey age-composition input sample size
- Update survey length-composition input sample size
- Add age-1 and age-2 to agecomposition data
- Estimate all growth parameter except CV




- Updated survey age-composition input sample size
- Update survey length-composition input sample size
- Add age-1 and age-2 to agecomposition data
- Estimate all growth parameter except
 CV
- Update length-weight relationship parameters





- Updated survey age-composition input sample size
- Update survey length-composition input sample size
- Add age-1 and age-2 to age-composition data
- Estimate all growth parameter except CV
- Update length-weight relationship parameters
- Update old age growth CVs







•Same as Model 24.0 except weightat-age is not a manual input



Candidate Models 1,000,000 300,000 Spawning Biomass (mt) 800,000 **Biomass Index** 600,000 400,000 100,000 1980 1990 2000 2010 2020 1980 1990 2000 2010 2020 Year Year Model – Base-4 – Model 24.0 – Model 24.1




Candidate Models 600,000 -0.16 0.12 Fishing Mortality 400,000 -# of Recruits 200,000 0.04 1980 1990 2000 2010 2020 1980 2000 2010 2020 1990 Year Year





Candidate Models





Likelihood Components

	Base-4	Model 24.0	Model 24.1
Total	1010.07	670.014	673.004
Survey	-66.6602	-66.4449	-65.701
Length-comp	417.408	379.449	380.43
Age-comp	633.643	335.088	335.743
Recruitment	25.6784	21.9209	22.5315



Conclusion

Transition to SS3

Base-4 Model that estimates growth

<u>Updates for 2024</u>

- 3 options (Base-4, Model 24.0, Model 24.1)
 - Recommendation: Model 24.1

