


Climate update



Dec 2024

Kirstin Holsman

NOAA Alaska Fisheries Science Center

kirstin.holsman@noaa.gov



Today's talk

Part 1: Climate challenges continue to grow, with uncertainty about future trajectories

Part 2: Climate-informed tools are available to support future sustainable fisheries (ACLIM)

Part 3: CEFI accelerates the capacity to deliver climate-informed tools and advice

1 Climate challenges continue to grow, with uncertainty about future trajectories





Greenhouse gases reached record observed levels in 2023.

Real time data indicate that they continued to rise in 2024.





Figure 1: Annual global mean temperature anomalies (relative to 1850–1900) from 1850 to 2024 from six datasets. The 2024 average is based on data from January-September.

January-September 2024 was 1.54±0.13°C above the pre-industrial average.



Greenhouse gases reached record observed levels in 2023.

Real time data indicate that they continued to rise in 2024.

- CO₂

420.0 ppm ± 0.2

151% of pre-industrial levels
- CH₄

1934 ppb ± 2

265% of pre-industrial levels.
- N₂O

336.9 ppb ± 0.1

124% of pre-industrial levels

2.0 C of Warming : Critical Tipping point






Figure 1: Annual global mean temperature anomalies (relative to 1850–1900) from 1850 to 2024 from six datasets. The 2024 average is based on data from January–September.

**January–September 2024 was
1.54±0.13°C above the pre-industrial average.**

UNFCCC 2022 Nationally Determined Contributions (NDCs) Synthesis report

PARIS CLIMATE AGREEMENT


1.  $<2^{\circ}\text{C}$

 NET ZERO EMISSIONS
 Limit the avg. global temperature increase to $<2^{\circ}$ centigrade + achieve net zero emissions by mid-century

2. 

 Enhance resilience and adaptation to climate impacts certain to occur

3. 

 Align financial flows in the world with these objectives




ACLIM found fisheries have higher risk above this (SSP585, RCP8.5)

ACLIM found fisheries have lower risk in this scenario (SSP126)

Climate change is expected to continue to impact AK Ecosystems & Fisheries




a) Global surface temperature change relative to 1850-1900



Warming will continue and is greater in scenarios with **low CO₂ mitigation and higher warming**


b) September Arctic sea ice area



Sea Ice will continue to decline, more so under scenarios with **high global warming and low CO₂ mitigation**

Figures from the IPCC AR6 WGI Summary for Policymakers:
https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_SPM.pdf

Sea ice loss impacts fisheries



Alaska Snow Crab



Plummeting Bering Sea crab populations

Snow crab and king crab have long been mainstays of commercial harvests.



Red king crab
Paralithodes camtschaticus

Long-term decline in mature red king crab populations (for Bristol Bay District)





Source: Surveys conducted by NOAA Fisheries



Snow crab
Chionoecetes opilio

Sharp drops in snow crab populations (all districts)



MARK NOWLIN / THE SEATTLE TIMES

www.noaa.gov/stories/unprecedented-2018-bering-sea-ice-loss-repeated-in-2019

www.seattletimes.com/seattle-news/as-bering-sea-ice-melts-nature-is-changing-on-a-massive-scale-and-alaska-crab-pots-are-pulling-up-cod/


www.seattletimes.com/seattle-news/valuable-crab-populations-crash-in-a-warming-bering-sea/

2 Climate-informed tools are available to support future sustainable fisheries



New predictive tools can help fisheries prepare & plan

psl.noaa.gov/marine-heatwaves



The screenshot shows the NOAA Physical Sciences Laboratory website. The header includes the NOAA logo and the text "Physical Sciences Laboratory". Navigation links include "About", "People", "Research", "Data", "Products", "News | Events", "Learn", and a search icon. The main content area is titled "Overview" and contains a paragraph: "Marine heatwaves are periods of persistent anomalously warm ocean temperatures, which can have significant impacts on marine life as well as coastal communities and economies. Scientists at PSL are working to characterize marine heatwaves, understand how they form and dissipate, and predict them in advance. On this page, we will provide current ocean maps, forecasts of heatwaves, interactive tools for users to explore ocean heatwaves themselves, links to research results and to webpages at other institutions." Below the text is a globe showing ocean temperature anomalies, with significant red and orange areas indicating warm events, particularly in the North Pacific and North Atlantic.

Can predict the probability of a marine heatwave 1-12 mo ahead of time

Jacox et al. 2022.

www.nature.com/articles/s41586-022-04573-9

Climate information “on ramps” for EBFM

Climate informed annual* stock and ecosystem assessments & EBFM advice

Climate information in near-term ecosystem based management targets

Climate-ready Ecosystem Based Fisheries Management planning, information & design

KEY: Matching climate information & projections to the scale of decision making & advice

On-ramp 1



Tactical Near-term Advice (<2 yr)

Climate change information incorporated into stock assessment models, stock-specific indicators (ESPs), stock-specific risk tables (as appropriate).

E.g., ABC based on climate forecasts

On-ramp 2



Strategic Near-term Advice (<2 yr)

Climate change context for observed changes in social, ecological, & oceanographic conditions relevant for harvest advice and targets.

E.g., Forecasts of climate-driven distributions, tipping points, & thresholds

On-ramp 3 (new)



Strategic & Long-term Advice (>2 yr)

Climate - informed long-term strategic decision making & planning informed by IK, LK, and climate & management scenario evaluations, risk assessments, & adaptation efficacy & feasibility evaluations.

E.g., Targets based on climate projections

Legend

- Existing tools or process
- Help ID climate change gaps
- New tools or process

The Alaska Climate Integrated Modeling Project

Goal: To address climate information needs with best available science & tools

What to expect?

- Project physical and ecological conditions under levels of climate change (levels of global carbon mitigation)
- Characterize uncertainty

What can be done?

- Evaluate effectiveness of adaptation actions including those supported by fisheries management

Scenarios form the basis for comparative simulations & Management Strategy Evaluations

Global General Circulation Models (GCM)

CESM
GFDL
MIROC

Carbon Scenarios:

Hindcast (1970-2021)
(high CO2 mitigation)

- CMIP6 SSP1 2.6

(med CO2 mitigation)


- CMIP5 RCP 4.5

(low CO2 mitigation)

- CMIP5 RCP 8.5
- CMIP6 SSP5 8.5

Climate Enhanced Biological models (x 5+)


CE- single-spp assessment models
CE- multi-spp model (CEATTLE)
CE- Size spectrum model
CE- Ecopath with Ecosim
End-to-End model (FEAST)
CE- spatial MICE model
CE - IBM (crab)



Hollowed et al. 2020. <https://doi.org/10.3389/fmars.2019.00775>

www.fisheries.noaa.gov/alaska/ecosystems/alaska-climate-integrated-modeling-project

ACLIM3 Team



Supporting climate
resilience through
climate-informed
Ecosystem Based
Management advice

Kirstin Holsman, Sarah Wise, Andre Punt, Albert Hermann, Cheryl Barnes, Cody Szuwalski, Kerim Aydin, Kelly Kearney, Anne Hollowed, Alberto Rovellini, Andrea Havron, Andy Whitehouse, Anna Amalka Sulc, Carey McGilliard, Catherine Moncrieff, Darren Pilcher, Diana Stram, Ed Farley, Elizabeth McHuron, Elizabeth Siddon, Ellen Yasumiishi, Grant Adams, Ingrid Spies, Ivonne Ortiz, James Ianelli, James Thorson, Jean Lee, Jennifer Bigman, Jeremy Sterling, Jodi Pirtle, Jonathan Reum, Kalei Shotwell, Kate Haapala, Kelly Kearney, Lorenzo Ciannelli, Mabel Baldwin-Schaeffer, Maggie Mooney-Seus, Martin Dorn, Maurice Goodman, Meaghan Bryan, Melissa Haltuch, Melissa Parks, Michael Litzow, Mike Dalton, Molly Graham, Patricia Pinto da Silva, Paul Spencer, Sarah Stone, Serena Fitka, Steve Barbeaux, Trond Kristiansen, Wei Cheng, William Stockhausen, Lia Domke, Anne Beaudreau, Justin Hansen, Angela Abolhassani, Matt Callahan, Brett Holycross

www.fisheries.noaa.gov/alaska/ecosystems/alaska-climate-integrated-modeling-project



Integrated Modeling to Evaluate Climate Change Impacts on Coupled Social-Ecological Systems in Alaska

Anne Babcock Hollowed^{1*}, Kirstin Kari Holsman¹, Alan C. Haynie¹, Albert J. Hermann^{2,3}, Andre E. Punt⁴, Kerim Aydin¹, James N. Ianelli¹, Stephen Kasperski¹, Wei Cheng^{2,3}, Amanda Faig^{2,4}, Kelly A. Kearney^{1,2}, Jonathan C. P. Reum^{1,5}, Paul Spencer¹, Ingrid Spies¹, William Stockhausen¹, Cody S. Szuwalski¹, George A. Whitehouse^{2,4} and Thomas K. Wilderbuhr¹

¹ Alaska Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA, United States, ² Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, WA, United States, ³ Pacific Marine Environmental Laboratory, Oceans and Atmospheric Research, National Oceanic and Atmospheric Administration, Seattle, WA, United States, ⁴ School of Aquatic and Fishery Sciences, College of the Environment, University of Washington, Seattle, WA, United States, ⁵ Centre for Marine Socioecology, Institute for Marine and Antarctic Studies, College of Sciences and Engineering, University of Tasmania, Hobart, TAS, Australia

OPEN ACCESS

Edited by:

Jamie C. Torn,
Bedford Institute of Oceanography
(BIO), Canada

Reviewed by:

Nancy Shackal,
Bedford Institute of Oceanography
(BIO), Canada
Daniel Howald,
Norwegian Institute of Marine
Research (IMR), Norway

*Correspondence:

Anne Babcock Hollowed
Anne.Hollowed@noaa.gov

Specialty section:

This article was submitted to
Global Change and the Future Ocean,
a section of the journal
Frontiers in Marine Science

Received: 20 August 2019

Accepted: 02 December 2019

Published: 14 January 2020

Citation:

Hollowed AB, Holsman KK,
Haynie AC, Hermann AJ, Punt AE,
Aydin K, Ianelli JN, Kasperski S,
Cheng W, Faig A, Kearney KA,
Reum JCP, Spencer P, Spies I,
Stockhausen W, Szuwalski CS,
Whitehouse GA and Wilderbuhr TK
(2020) Integrated Modeling
to Evaluate Climate Change Impacts
on Coupled Social-Ecological
Systems in Alaska.
Front. Mar. Sci. 6:775.
doi: 10.3389/fmars.2019.00775

The Alaska Climate Integrated Modeling (ACLIM) project represents a comprehensive, multi-year, interdisciplinary effort to characterize and project climate-driven changes to the eastern Bering Sea (EBS) ecosystem, from physics to fishing communities. Results from the ACLIM project are being used to understand how different regional fisheries management approaches can help promote adaptation to climate-driven changes to sustain fish and shellfish populations and to inform managers and fishery dependent communities of the risks associated with different future climate scenarios. The project relies on iterative communications and outreaches with managers and fishery-dependent communities that have informed the selection of fishing scenarios. This iterative approach ensures that the research team focuses on policy relevant scenarios that explore realistic adaptation options for managers and communities. Within each iterative cycle, the interdisciplinary research team continues to improve: methods for downscaling climate models, climate-enhanced biological models, socio-economic modeling, and management strategy evaluation (MSE) within a common analytical framework. The evolving nature of the ACLIM framework ensures improved understanding of system responses and feedbacks are considered within the projections and that the fishing scenarios continue to reflect the management objectives of the regional fisheries management bodies. The multi-model approach used for projection of biological responses, facilitates the quantification of the relative contributions of climate forcing scenario, fishing scenario, parameter, and structural uncertainty within and between models. Ensemble means and variance within and between models inform risk assessments under different future scenarios. The first phase of projections of climate conditions to the end of the 21st century is complete,

Hollowed et al. 2020 ACLIM overview paper



Also see list of 20+ publications at end of ppt

Key Takeaways from ACLIM to date


Climate Informed EBM advice




Climate impacts ecosystems & food webs



Climate impacts on growth, survival & biomass



Changes to fish distributions (& fishing grounds)



Climate change (oceanography)




1. Need to account for ecosystem-wide productivity changes
2. Manage predator-prey Interactions using new approaches
3. Need to (and can) adapt to climate-driven species range shifts
4. Expand Ecosystem-Based Fisheries Management through coordination and collaboration
5. Address uncertainty with ensembles and participatory scenario planning
6. Incorporate socio-economic resilience into planning and response
7. Strengthen research and monitoring infrastructure

Key Outcomes of ACLIM to date

1. **Provided baseline engineering capacity for ocean and ecosystems predictions and forecasts**
2. Identified key couplings, and gaps and needs in linked climate- oceanographic- biological- social-economic models
3. Evaluated trajectories under future scenarios, alternative management strategies, alternative harvest control rules → emergent understanding of common pitfalls and best practices
4. Sustained community of practice for climate-informed advice, including support for partnership building around adaptation planning

High-res model reproduces the Bering Sea environment




Kearney K (2021). U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-415, 40 p. [link](#).

Supporting Publications

Szuwalski et al. (2022, 2023), Pilcher et al. (2022), Reum et al. (2020), Whitehouse et al. (2021), Cheng et al. (2021,2023), Hermann et al. (2021,2023), Hollowed et al. (2022), Thorson et al. (2021)

Key Outcomes of ACLIM to date

1. Provided baseline engineering capacity for ocean and ecosystems predictions and forecasts
2. **Identified key couplings, and gaps and needs in linked climate- oceanographic- biological- social-economic models**
3. Evaluated trajectories under future scenarios, alternative management strategies, alternative harvest control rules → emergent understanding of common pitfalls and best practices
4. Sustained community of practice for climate-informed advice, including support for partnership building around adaptation planning



Supporting Publications

Goodman et al. (2024), Punt et al. (2023), Szuwalski et al. (2023), Olmos et al. (2023), McHuron et al. (2024), Barnes et al. (2022), Thorson et al. (2021), Whitehouse et al. (2021), Kearney et al (2020), Pilcher et al. (2022), Hollowed et al. (2020).

Key Outcomes of ACLIM to date

1. Provided baseline engineering capacity for ocean and ecosystems predictions and forecasts
2. Identified key couplings, and gaps and needs in linked climate- oceanographic- biological- social-economic models
3. Evaluated trajectories under future scenarios, alternative management strategies, alternative harvest control rules → emergent understanding of common pitfalls and best practices


SSB targets under Climate Change; HCRs; 2 mt cap effects

Supporting Publications

Szuwalski et al. (2022, 2023), Pilcher et al. (2022), Reum et al. (2020), Whitehouse et al. (2021), Holsman et al. (2020), Hollowed et al. (2024), Barnes et al., (2022), Hermann et al. (2021,2023), Cheng et al. (2021,2023), Punt et al. (2023), Goodman et al. (2024), McHuron et al. (2024), Punt et al. (2023)



Adapting reference points to reflect changes in productivity

- MSA directs reference points to reflect current and probable future environmental conditions
- Changing reference points for stocks undergoing climate-related productivity shifts can result in counter-intuitive management actions:
 - Declining stocks could be fished harder
 - Flourishing stocks could be fished more conservatively



Szuwalski et al. 2023


A) Biological reference points



Holsman et al. 2020. <https://www.nature.com/articles/s41467-020-18300-3>


June 2024 NPFMC Climate Scenarios Workshop

Based on ACLIM scenarios, updated by CCTF and refined by CSW planning group



Future Climate Change Scenarios

Discussion Roadmap



Reflect back: what is the set of tools or processes that work in all scenarios?

Key Outcomes of ACLIM to date

1. Provided baseline engineering capacity for ocean and ecosystems predictions and forecasts
2. Identified key couplings, and gaps and needs in linked climate- oceanographic- biological- social-economic models
3. Evaluated trajectories under future scenarios, alternative management strategies, alternative harvest control rules → emergent understanding of common pitfalls and best practices
4. **Sustained community of practice for climate-informed advice, including support for partnership building around adaptation planning**



Salmon & Communities

Identify candidate ROMS/NPZ indicators for Yukon River Chinook salmon survival based on scientific and traditional knowledge.

H1: Ocean temperatures during the 1st and 2nd year at sea impacts growth & survival.

Spring 2023 trip to Lower Yukon LTK:

Good for salmon returns: Strong north winds, high river water, ice break up but not thaw, & yellow butterflies. (wish list indicators)

Produce recruitment projections under different climate & emission scenarios at various lags



*Slide Courtesy of S. Wise, E. Yasumiishi, J. Reynolds (AFSC-NOAA)
Draft results, please do not copy or distribute without permission of the author*

- Conducted preliminary community meetings in lower Yukon to inform household survey. Planned final phase fieldwork for 2025.
- Coordinated Multiple Knowledge systems on Yukon River Chinook marine survival. Continue to coordinate with upriver findings.
- Explored food security as driver in management decision-making






ACLIM3 Decision Support System

Partnership building & scenario discussions


Multiple Knowledge systems

Climate Scenarios
Multiple ESMS



Fishing & harvest Scenarios
Climate Informed Policies

Alternative Food Web & Climate Coupling



Species Distribution
Biomass



Climate Integrated Assessments
Climate Smart HCRs
Climate Informed BRPs

ABC



Catch

TAC

Bycatch

Economic benefits

Wellbeing

Livelihoods

Food Security

- ▲ Carrying Capacity
- ▲ Distribution
- ▲ Biomass
- ▲ Fish Condition

- ▲ Catch
- ▲ Value
- ▲ Cost
- ▲ Wellbeing

PROJECTIONS
Species distribution & biomass

ADVICE
CI harvest recommendation

TOOL BOX
CI smart tools

RISK ASSESSMENT
Risk & Adaptation

DISASTER RESPONSE
Effective adaptation

Decision Support & Adaptation Planning
Community-determined decision-making support



Open Science: interactive tools

roms-bering-sea

Posts About Literature



The Bering10K ROMS configuration

The Bering10K ROMS configuration, including associated biological modules (research conducted through the University of Washington, CICOES)

GitHub

The Bering10K dataset

3 minute read

Numerous Bering 10K ROMS model simulations have been run to date, including hindcasts of the past few decades, long-term forecasts under CMIP5 and CMIP6 emissions scenarios, and seasonal retrospective forecasts. Data and metadata related to these simulations are held in a number of locations. This page serves as a centralized hub for this data and metadata.

The model

Model source code is available on GitHub: [beringnpz/roms-bering-sea](https://beringnpz.github.io/roms-bering-sea)

The documentation

A few guides for working with the Bering10K output dataset can be found

- [The Bering10K Dataset documentation](#): A pdf describing the dataset, including:



1. A description of the various simulations (base model versions, parent model forcing datasets, and biological modules) and the output naming scheme for each



<https://beringnpz.github.io/roms-bering-sea/B10K-dataset-docs>

This screenshot shows the GitHub repository for 'kholman / ACLIM2'. It displays the repository name, public status, and navigation options like Code, Issues, Pull requests, Actions, Projects, Security, and Insights. Below these are statistics for branches (3), tags (0), and commits (218). A list of recent commits is visible, including one by 'kholman' to 'fix readme' and another to 'updates'.

This screenshot shows the 'Alaska Climate Integrated Modeling Project' catalog. It lists several datasets with icons representing different simulation scenarios, such as 'B10K-H16_CMIP5_CESM_B10_rcp85', 'B10K-H16_CMIP5_CESM_rcp85', 'B10K-H16_CMIP5_CESM_rcp45', 'B10K-H16_CMIP5_GFDL_B10_rcp85', 'B10K-H16_CMIP5_GFDL_rcp45', 'B10K-H16_CMIP5_GFDL_rcp85', 'B10K-H16_CMIP5_MIROC_rcp45', and 'B10K-H16_CMIP5_MIROC_rcp85'.



Open Science: interactive tools

ACLIM2 SDMs: Species range and overlap forecasts

Species 1: arrowtooth flounder
Species 2: walleye pollock
Species 1 size bin: adult
Species 2 size bin: juvenile

Compute overlap indices using:
Estimated biomass (selected) | Probability of occurrence

Climate scenarios:
SSP1-2.6 (CMIP4) | RCP4.7 (CMIP3) | SSP3-6.5 (CMIP4)

Climate models:
Ensemble (selected) | CLSM | GFDL | MIROC

Download output:
Contact Maurice Goodman (goodman3@stanford.edu) for other outputs
Overlap | Species 1 | Species 2

Background

This interface displays fitted means and confidence intervals from SDMs built for a suite of eastern Bering Sea groundfish and crab species as part of the ACLIM2 project. Generalized additive models (GAMs) were built for each species with environmental covariates selected using time-series cross validation (i.e., by optimizing overall year-ahead predictive performance). Terms considered in the models include temperature, depth, oxygen, pH, and spatially-varying effects of the cold pool. Models were fit separately for juveniles and adults for most species, but not for snow crabs, and red king crab - when plotting these species, choose 'all' for the size bin. Models were fit as delta-GAMs, with a binomial component for estimating probability of occurrence, and lognormal component for estimating positive (non-zero) biomass. Overlap projections are plotted below, with model summaries and range projections for individual species available in tabs to the right.

Overlap metrics can be computed from either probability of occurrence, in which case only the binomial component of each delta model is used, or expected biomass (i.e., the product of the expected values from the binomial and Lognormal model component). It is generally much better for the binomial component (i.e., the models tend to do a better job describing the distribution of encounters / non-encounters for each species than they do for species biomass distributions), so overlap estimates based on probability of occurrence only may be more reliable, possibly at the cost of ecological interpretability.

Spatial overlap

Each of the below overlap metrics varies between 0 and 1, with 0 indicating no overlap and 1 indicating complete overlap, and each is invariant under linear transformation, i.e., impacted only by the relative biomass distributions of each species, not by the aggregate sum biomass in a given year. The years containing groundfish survey data (1982-2019) are shaded.

Local index of collocation

$$\frac{\sum_i p_i p_j}{\sum_i p_i^2 \sum_j p_j^2}$$



The local index of collocation is (loosely) a relative metric of interspecies encounter, which measures co-occurrence as a non-centered correlation among estimated species biomass:




Juv. pollock





Shifting distributions: Habitat novelty



Open Science: interactive tools



Changes in predator & prey overlap




Climate information on-ramps in 2024



ESPs, ESRs, Stock assessments


Future ABC



New this year
In the
multispecies
assessment

Climate scenario

- Persistence
- High mitigation
- Med mitigation
- Low mitigation



[Link to 2024 Climate-enhanced multispecies assessment \(EBS\)](#)

Community of practice is needed. In particular, need capacity to: co-develop, deliver, & refine



3 CEFI accelerates the capacity to deliver climate-informed tools and advice






**Climate, Ecosystems,
& Fisheries Initiative**

The Alaska Climate/CEFI Team (ACT)

Kirstin Holsman
kirstin.holsman@noaa.gov



CEFI Decision Support System



Alaska's Climate Ecosystem and Fisheries Initiative




CEFI High resolution oceanographic model (MOM6) grids

Pacific

Arctic

Atlantic



CEFI-OAR High resolution models (operational delivery)

hindcasts


seasonal forecasts

decadal predictions

multidecadal climate change projections

(e.g., sea ice, water temp, pH, winds, currents, zooplankton)


NOAA's Climate Ecosystem and Fisheries Initiative (CEFI)






Current Bering10K high resolution oceanographic seasonal forecasts

Spring

Predicted 2024 July 1
bottom temperature




Current Bering10K high resolution oceanographic seasonal forecasts




Current Bering10K high resolution oceanographic seasonal forecasts



Updated in fall



model




Obs.




Spring

Predicted 2024 July 1
bottom temperature




Similar based on...




Current Bering10K high resolution oceanographic seasonal forecasts


Updated in fall



model





Obs.




Spring

Predicted 2024 July 1
bottom temperature





CEFI synergies at work in the Bering Sea



fraction of the trawl survey area

MOM6 Bering Sea Cold Pool



Year	West Coast and Arctic
FY23	Initial Configuration ✓
FY24	Initial hindcast ✓
FY25	Hindcast update, retrospective seasonal predictions , initial climate change projections
FY26	Hindcast update, retrospective decadal predictions , initial climate change projections
FY27	Hindcast update, expanded projections , seasonal outlooks reliably delivered
FY28	All products reliably delivered
FY29	All products reliably delivered

Thanks to Wei Cheng, Vivek Seelanki, Liz Drenkard, Kelly Kearney, Al Hermann, Darren Pilcher, Theresa Morrison, Bob Hallberg and others in the regional MOM6 development forum...




NOAA Climate, Ecosystems, & Fisheries Initiative

Decision Support Teams




Co-generate regionally, locally, and community tailored tools & advice





Species, stock & ecosystem status

Climate-informed advice




Strategic planning & response



What are Decision Support Teams?

Transdisciplinary nested teams that will help deliver climate informed products and advice, specifically tailored to decision maker needs.




- 1) Understand current capacity for inclusion of climate informed advice
- 2) Identify near-term needs for CI-advice
- 3) Identify long-term needs for development
- 4) Link CEFI, IEA, Stock Assessment, and Other NOAA products to meet needs


Support and
co-leverage

not repeat or
re-invent


Decision Support Delivery Steps



MOM6




- Is there an existing tool/product that meets the need?
- Do the scales match advice?
- Does the output skillfully meet needs?




Types of Management Actions

Climate impacts ecosystems & food webs



Climate impacts on growth, survival & biomass



Changes to fish distributions (& fishing grounds)



Climate change (oceanography)



Climate Informed EBM advice




Draft Focal Areas for Alaska CEFI

FOCAL AREA 1: Web accessible and regionally tailored climate change products & trainings (cross cutting)


FOCAL AREA 2: Climate-integrated fisheries assessments and EBM

FOCAL AREA 3: Decision support for Climate aware regulatory frameworks

FOCAL AREA 4: Climate resilience community planning and response



Provide in addition to climate information on-ramps in ESPs, ESRs, Stock assessments, ACEPO, etc.



New this year: Prototype Climate information overview



Feedback welcome!

What is the ACT?

The ACT is the Alaska CEFI Regional Decision Support Team. The Alaska Climate CEFI Team (ACT), established in fall 2021, and expanding to include new members and partner institutions in 2024, guides regional development of publicly accessible CEFI tools and products to support climate-informed advice and adaptation planning.

Regional Decision Support Teams

Regional Decision Support Teams produce climate-related information and advice for effective management of fisheries, ecosystems, and protected species and industry, and community adaptation planning. They operate through NOAA's Regional Fisheries Science Centers to provide:

- Early warnings and projections of ecosystem conditions
- Risk assessments & scenario planning for fisheries and fishing communities
- Science support for climate-ready Ecosystem-Based Management

How will CEFI change advice?

Scientists from the Alaska Fisheries Science Center have been leading pilot projects such as the Alaska Climate Integrated Modeling project (ACIMI) and the Gulf of Alaska Integrated Modeling project (GOAIM). These projects serve as prototypes for the decision support tools and advice that CEFI may support. They demonstrate improved model performance with climate linkages and provide integrated climate advice that considers climate changes, biological and ecosystem responses, and alternative management and adaptation options to support climate-smart Ecosystem-Based Management and thriving climate-resilient communities in Alaska. Advice is designed to provide climate information via the existing Council process and advice pathways.

More Information

CEFI Fact Sheet
NOAA's Climate.gov
Alaska IEA Program
Alaska IEA Program

Questions? Email us!
Kirstin Holzman (ACT Lead), kirstin.holzman@noaa.gov

2024 Climate Science Update

ALASKA'S MARINE ECOSYSTEMS are undergoing climate-driven changes, including rising sea temperatures, shrinking sea ice, & shifts in ocean acidification & productivity, impacting species from snow crab to whales. Impacts on **FISHERIES & FISHING COMMUNITIES** have been widespread affecting economies, livelihoods, family structures, mental health, sharing networks, & food security. Future projections indicate further changes in species distributions, stock abundances, and ecosystem dynamics, requiring climate planning & adaptive, flexible, & ecosystem-based strategies.

CLIMATE IMPACTS: 2024 spotlight on the EBS

TEMPERATURE RISE: SSTs in the Eastern Bering Sea have risen by 1.0°C to 1.5°C, with a further increase of 1.5°C to 3.0°C expected by century's end. **SEA ICE DECLINE:** Arctic sea ice has declined precipitously over the observed record from 1978 to present. Eastern Bering Sea ice has remained stable, but experienced unprecedented lows in this region in 2018 and 2019 that have been attributed to human-caused climate change. **OCEAN ACIDIFICATION:** Global ocean surface pH has dropped by 0.1 units since 1750, with a further decline of 0.1-0.3 projected for the Bering Sea by the end of the century.

PACIFIC COD: Marine Heatwaves (MHW) were associated with a rapid redistribution of roughly half of Pacific cod biomass into the N. Bering Sea (NBS) in 2018-2019 as well as declines in biomass and recruitment. Future warming may push Pacific cod further north, expanding spawning habitat but potentially altering NBS carrying capacity. **SNOW CRAB** have also collapsed in response to marine heatwaves, & future warming is expected to further amplify impacts. Several WESTERN ALASKA **SALMON** stocks have declined in recent years, potentially linked to climate change, though impacts across across freshwater & marine life histories are complex. In contrast, Bristol Bay **SOCKEYE SALMON** & **SABLEFISH** have had increased abundance under warming conditions. The exact reasons for these divergent responses across species and stocks are still being evaluated.

CEFI: The Climate, Ecosystems, & Fisheries Initiative is a cross-NOAA effort to build the resiliency of Alaska's fisheries & provide science support needed to allow management & resource users to adapt to changing ocean conditions. CEFI aims to provide: (1) **Robust forecasts**, decadal predictions, and long-term projections of ocean conditions; (2) **Publicly available** climate-linked early warnings, climate-enhanced stock, ecosystem, & risk assessments, & evaluations of climate-resilient management response; (3) increased capacity to provide **climate-informed advice** to support long-term sustainability & resilience.

Alaska's Climate, Ecosystems, and Fisheries Initiative

Gene M. Watters
U.S. Secretary of Commerce

Richard W. Spring
Under Secretary of Commerce for Oceans and Atmosphere

Harold Cook
Assistant Administrator for Fisheries

National Marine Fisheries Service
Alaska Fisheries Science Center
7605 Sand Point Way NE, Seattle, WA 98148-3402
www.fisheries.noaa.gov

What we are planning & what we will do

Oceanographic

Region-specific oceanographic products will be developed for public use to support climate change adaptation. This includes the current Regional Ocean Modeling System (ROMS) and the advanced Modular Ocean Model 6 (MOM6). These high resolution, three dimensional ocean models effectively simulate past ocean conditions and lower trophic level dynamics, from phytoplankton to kill, and are instrumental in forecasting future changes.

Hindcasts: hindcasts are a powerful tool for reconstructing the climate and environmental conditions of the recent past. Through CEFI, hindcasts will be produced to recreate the oceanic and sea ice states over the past several decades. The ACT will evaluate the accuracy of these reconstructions by comparing them with historical ocean observations. Once validated, these model outputs will provide continuous data on ocean conditions from seabed to the surface. This information will enhance understanding of the ocean dynamics during different fishing and harvest seasons, and will help identify the drivers of species population changes, spatial shifts, and broader ecosystem responses to warming and marine heatwaves.

Forecasts: Forecasts from the same models will be updated each season to provide near-term projections of ocean conditions one to twelve months out.

Decadal predictions will provide data-driven outlook of potential ocean conditions up to 10 years into the future.

Long-term projections under high and low warming scenarios will deliver detailed information to support risk analyses and climate adaptation planning.

Kelly Kearney, kelly.kearney@noaa.gov
Wei Cheng, wei.cheng@noaa.gov

Biological

Develop dynamic climate-informed multispecies distribution models for Bering Sea groundfish, crab, and marine mammal species.

Assess changes in spatial overlap among species based on range projections under various climate scenarios.

Elizabeth McHuron, elizabeth.mchuron@noaa.gov
Maurice Goodman, goodman@uaf.edu

Link models and use ensemble approaches to quantify relationships between climate-ecosystem variables and population dynamics across fish species.

Identify best practices for selecting which climate-ecosystem variables can help predict stock dynamics.

Jennifer Bigman, jennifer.bigman@noaa.gov
Grant Adams, grant.adams@noaa.gov
André Punt, apunt@uaf.edu

Develop climate-linked harvest control rules to determine ADC buffers for the North Pacific Fisheries Management Council.

Develop software to produce climate-informed short-term growth and mortality forecasts for NPFMC harvest control rules.

Support climate-informed stock assessments and multispecies models for groundfish in the Gulf of Alaska and Bering Sea.

Grant Adams, grant.adams@noaa.gov
Kirstin Holzman, kirstin.holzman@noaa.gov
André Punt, apunt@uaf.edu

Social & Economic

Develop social hindcast methodology with pilot of community responses to climate-induced changes in Alaska fisheries (2016-2023) to better understand community adaptation strategies.

Identify possible social indicators using existing data sources to detect and monitor the effects of climate-induced changes associated with resilience and vulnerability of Alaska fishing communities.

Support development of publicly accessible, collaborative tools allowing communities to view CEFI information products on climate-induced shifts in species distributions and ecological and oceanographic conditions.

Explore pathways and examine barriers for utilizing social and economic information to inform Council decision-making on climate resilient fisheries management.

Angela Abolhasani, angela.abolhasani@noaa.gov
Sarah Wise, sarah.wise@noaa.gov

Collaboratively identify drivers of Yukon River Chinook salmon marine survival using ROMS ocean temperature outputs, wind hindcasts, and Traditional Knowledge in partnership with the Yukon River Drainage Fisheries Association.

Ellen Yasumishi, ellen.yasumishi@noaa.gov
Sarah Wise, sarah.wise@noaa.gov

Council Coordination

In coordination with the Council points of contact, develop a process for regular reporting of climate change impacts and responses to the Council through existing pathways (or "on-ramps"). Fisheries Initiative (CEFI) potential projects and outputs could also help support the Council's climate readiness planning. This information will highlight current and proposed synergies between CEFI products and Council planning. Coordination on the development of these products will involve collaboration among the Council, NOAA's Alaska CEFI team (ACT) and partners, and the Alaska Fisheries Information Network.

Facilitate Council planning for climate change impacts on fisheries and fishing communities in Alaska through iterative discussions, workshops, information reporting, and synthesis of medium to long-term impacts on Alaska marine ecosystems. Support management strategy evaluations (MSEs) to evaluate the performance of alternative management measures and assessments under future climate scenarios.

Annual climate-informed advice: Develop interactive and publicly accessible resources for decision making and exploring tradeoffs in annual fisheries management decisions. Support climate-informed biological reference points and harvest control rules.

Rapid response: Support the development and delivery of tools to aid in-season management and navigate emergent climate challenges.

Jodi Petrie, jodi.petrie@noaa.gov
Jason Gasper, jason.gasper@noaa.gov
Anne Marie Eich, annemarie.eich@noaa.gov

Katie Latanich, katie.latanich@noaa.gov
Diana Stram, diana.stram@noaa.gov

Kirstin Holzman, kirstin.holzman@noaa.gov
Angela Abolhasani, angela.abolhasani@noaa.gov


What is the ACT?

The ACT is the Alaska CEFI Regional Decision Support Team. The Alaska Climate CEFI Team (ACT), established in fall 2021, and expanding to include new members and partner liaisons in 2024, guides regional development of publicly accessible CEFI tools and products to support climate-informed advice and adaptation planning.

Regional Decision Support Teams

Regional Decision Support Teams produce climate-related information and advice for effective management of fisheries, ecosystems, and protected species and industry, and community adaptation planning. They operate through NOAA's Regional Fisheries Science Centers to provide:

- Early warnings and projections of ecosystem conditions
- Risk assessments & scenario planning for fisheries and fishing communities
- Science support for climate-ready Ecosystem-Based Management



How will CEFI change advice?

Scientists from the Alaska Fisheries Science Center have been leading pilot projects such as the Alaska Climate Integrated Modeling project (ACIM) and the Gulf of Alaska Integrated Modeling project (GOACLIM). These projects serve as prototypes for the decision support tools and advice that CEFI may support. They demonstrate improved model performance with climate linkages and provide integrated climate advice that considers climate changes, biological and ecosystem responses, and alternative management and adaptation options to support climate-smart Ecosystem-Based Management and thriving climate-resilient communities in Alaska. Advice is designed to provide climate information via the existing Council process and advice pathways.

More Information

CEFI Fact Sheet

NOAA's Climate.gov

Alaska IEA Program

Questions? Email us!

Kirstin Holsman (ACT Lead),
kirstin.holsman@noaa.gov

2024 Climate Science Update

ALASKA'S MARINE ECOSYSTEMS are undergoing climate-driven changes, including rising sea temperatures, shrinking sea ice, & shifts in ocean acidification & productivity, impacting species from snow crab to whales. Impacts on **FISHERIES & FISHING COMMUNITIES** have been widespread affecting economies, livelihoods, family structures, mental health, sharing networks, & food security. Future projections indicate further changes in species distributions, stock abundances, and ecosystem dynamics, requiring climate planning & adaptive, flexible, & ecosystem-based strategies.

CLIMATE IMPACTS: 2024 spotlight on the EBS

TEMPERATURE RISE: SSTs in the Eastern Bering Sea have risen by 1.0°C to 1.5°C, with a further increase of 1.5°C to 3.0°C expected by century's end. **SEA ICE DECLINE:** Arctic sea ice has declined precipitously over the observed record from 1978 to present. Eastern Bering Sea ice has remained stable, but experienced unprecedented lows in this region in 2018 and 2019 that have been attributed to human-caused climate change. **OCEAN ACIDIFICATION:** Global ocean surface pH has dropped by 0.1 units since 1750, with a further decline of 0.1 - 0.3 projected for the Bering Sea by the end of the century.

PACIFIC COD: Marine Heatwaves (MHW) were associated with a rapid redistribution of roughly half of Pacific cod biomass into the N. Bering Sea (NBS) in 2018 - 2019 as well as declines in biomass and recruitment. Future warming may push Pacific cod further north, expanding spawning habitat but potentially altering NBS carrying capacity. **SNOW CRAB** have also collapsed in response to marine heatwaves, & future warming is expected to further amplify impacts. Several **WESTERN ALASKA SALMON** stocks have declined in recent years, potentially linked to climate change, though impacts across across freshwater & marine life histories are complex. In contrast, Bristol Bay **SOCKEYE SALMON & SABLEFISH** have had increased abundance under warming conditions. The exact reasons for these divergent responses across species and stocks are still being evaluated.

CEFI: The Climate, Ecosystems, & Fisheries Initiative is a **cross-NOAA** effort to build the to build ocean modeling capacity & provide science support needed to allow management & resource users to adapt to changing ocean conditions. CEFI aims to provide: (1) **Robust forecasts**, decadal predictions, and long-term projections of ocean conditions; (2) **Publicly available** climate-linked early warnings, climate-enhanced stock, ecosystem, & risk assessments, & evaluations of climate-robust management response; (3) Increased capacity to provide **climate-informed advice** to support long-term sustainability & resilience.

Alaska's Climate, Ecosystems, and Fisheries Initiative



Gina M. Raimondo
U.S. Secretary of Commerce

Richard W. Spinrad
Under Secretary of Commerce
for Oceans and Atmosphere

Janet Coit
Assistant Administrator
for Fisheries

National Marine Fisheries Service
Alaska Fisheries Science Center
7600 Sand Point Way N.E., Seattle, WA 98115-6349
www.fisheries.noaa.gov



NOAA
FISHERIES

Alaska Fisheries
Science Center

Oceanographic

Region-specific oceanographic products will be developed for public use to support climate change adaptation. This includes the current [Bering10K](#) Regional Ocean Modeling System (ROMS) and the advanced [Modular Ocean Model 6 \(MOM6\)](#). These high resolution, three dimensional ocean models effectively simulate past ocean conditions and lower trophic level dynamics, from phytoplankton to krill, and are instrumental in forecasting future changes.



Hindcasts: hindcasts are a powerful tool for reconstructing the climate and environmental conditions of the recent past. Through CEFI, hindcasts will be produced to recreate the oceanic and sea ice states over the past several decades. The ACT will evaluate the accuracy of these reconstructions by comparing them with historical ocean observations. Once validated, these model outputs will provide continuous data on ocean conditions from seabed to the surface. This information will enhance understanding of the ocean dynamics during different fishing and harvest seasons, and will help identify the drivers of species population changes, spatial shifts, and broader ecosystem responses to warming and marine heatwaves.

Forecasts: Forecasts from the same models will be updated each season to provide near-term projections of ocean conditions one to twelve months out.

Decadal predictions will provide data-driven outlook of potential ocean conditions up to 10 years into the future.

Long-term projections under high and low warming scenarios will deliver detailed information to support risk analyses and climate adaptation planning.

Kelly Kearney, kelly.kearney@noaa.gov
Wei Cheng, wei.cheng@noaa.gov



Biological

Develop dynamic climate-informed **multispecies distribution models** for Bering Sea groundfish, crab, and marine mammal species.

Assess **changes in spatial overlap** among species based on range projections under various climate scenarios.

Elizabeth McHuron, liz.mchuron@noaa.gov
Maurice Goodman, goodmm2@uw.edu

Link models and use ensemble approaches to quantify relationships between climate-ecosystem variables and **population dynamics** across fish species.

Identify **best practices** for selecting which climate-ecosystem variables can help predict stock dynamics.


Jennifer Bigman, jennifer.bigman@noaa.gov
Grant Adams, grant.adams@noaa.gov
André Punt, aepunt@uw.edu

Develop **climate-linked harvest control rules** to determine ABC buffers for the North Pacific Fisheries Management Council.

Develop software to produce climate-informed short-term **growth and mortality forecasts** for NPFMC harvest control rules.

Support **climate-informed stock assessments and multispecies models** for groundfish in the Gulf of Alaska and Bering Sea.

Grant Adams, grant.adams@noaa.gov
Kirstin Holsman, kirstin.holsman@noaa.gov
André Punt, aepunt@uw.edu



Social & Economic

Develop **social hindcast methodology** with pilot of community responses to climate-induced changes in Alaska fisheries (2018-2023) to better understand community adaptation strategies.

Identify possible **social indicators** using existing data sources to detect and monitor the effects of climate-induced changes associated with resilience and vulnerability of Alaska fishing communities.


Support development of **publicly accessible, collaborative tools** allowing communities to view CEFI information products on climate-induced shifts in species distributions and ecological and oceanographic conditions.

Explore **pathways and examine barriers** for utilizing social and economic information to inform Council decision-making on climate resilient fisheries management.

Angela Abolhassani, angela.abolhassani@noaa.gov
Sarah Wise, sarah.wise@noaa.gov

Collaboratively identify drivers of Yukon River Chinook salmon marine survival using ROMS ocean temperature outputs, wind hindcasts, and Traditional Knowledge in partnership with the Yukon River Drainage Fisheries Association.

Ellen Yasumiishi, ellen.yasumiishi@noaa.gov
Sarah Wise, sarah.wise@noaa.gov



Council Coordination

In coordination with the Council points of contact, develop a process for **regular reporting of climate change impacts and responses to the Council** through existing pathways (or "on-ramps"). Fisheries Initiative (CEFI) potential projects and outputs could also help support the Council's climate readiness planning. This information will highlight current and proposed synergies between CEFI products and Council planning. Coordination on the development of these products will involve collaboration among the Council, NOAA's Alaska CEFI team (ACT) and partners, and the Alaska Fisheries Information Network.

Facilitate Council planning for climate change impacts on fisheries and fishing communities in Alaska through iterative discussions, workshops, information reporting, and synthesis of medium to long-term impacts on Alaska marine ecosystems. Support management strategy evaluations (MSEs) to evaluate the performance of alternative management measures and assessments under future climate scenarios.

Annual climate-informed advice: Develop interactive and publicly accessible resources for decision making and exploring tradeoffs in annual fisheries management decisions. Support climate-informed biological reference points and harvest control rules.

Rapid response: Support the development and delivery of tools to aid in-season management and navigate emergent climate challenges.

Jodi Pirtle, jodi.pirtle@noaa.gov
Jason Gasper, jason.gasper@noaa.gov
Anne Marie Eich, annemarie.eich@noaa.gov

Katie Latanich, katie.latanich@noaa.gov
Diana Stram, diana.stram@noaa.gov

Kirstin Holsman, kirstin.holsman@noaa.gov
Angela Abolhassani, angela.abolhassani@noaa.gov



Climate change is challenging but solvable

Overall, the latest studies on the net economic implications of decarbonisation – which also account for avoided climate damages – **point to overall benefit from the transition.**

-Prof Valentina Bosetti

If people are provided with opportunities to make choices supported by policies, infrastructure and technologies, there is an untapped mitigation potential to **bring down global emissions by between 40 and 70% by 2050** compared to a baseline scenario.

-Prof Joyashree Roy


The evidence is clear: there are now mitigation options available in all sectors that could together **halve global greenhouse gas emissions by 2030.** *-Dr Céline Guivarch*



Questions?

EXTRA SLIDES

National CEFI Component Workflow



ACLIM Publications (direct & indirect) 1 of 2


- (2024) McHuron et al. Current and future habitat suitability of northern fur seals and overlap with the commercial walleye pollock fishery in the eastern Bering Sea Movement Ecology
- (2024) Goodman et al. Climate covariate choice and uncertainty in projecting species range shifts: a case study in the Eastern Bering Sea. Fish and Fisheries (in press)
- (2024) Hollowed et al. Selecting climate linked decision relevant and adaptation informing community level scenarios for ecosystems through constituent engagement : A case study for the eastern Bering Sea. ICES JMS
- (2023) Punt et al. Capturing Uncertainty when Modelling Environmental Drivers of Fish Populations, with an Illustrative Application to Pacific Cod in the Eastern Bering Sea. Fisheries Research
- (2023) Bigman, JvS, B J Laurel, K Kearney, A J Hermann, W Cheng, K K Holsman, L A Rogers. Predicting Pacific cod thermal spawning habitat in a changing climate. ICES Journal of Marine Science, fsad096, <https://doi.org/10.1093/icesjms/fsad096>
- (2023) Hermann et al. Applications of biophysical modeling to Pacific high-latitude ecosystems. Oceanography
- (2023) Szuwalski, C.S. et al. The collapse of eastern Bering Sea snow crab. Science.
- (2023) Olmos, M., et al., Punt, A.E., Szuwalski, C.S. A step towards the integration of spatial dynamics in population dynamics models: Eastern Bering sea snow crab as a case study. Ecological Modelling 485: 110484.
- (2023) Szuwalski et al., Unintended consequences of climate-adaptive fisheries management targets. Fish and Fisheries. <https://doi.org/10.1111/faf.12737>
- (2022) Barnes, C.; Essington, T. E.; Pirtle, J; Rooper, C; Laman, E.; Holsman, K.; Aydin, K.; Thorson, J.. Climate-informed models benefit hindcasting but may present challenges when forecasting species-habitat associations. Ecography 2022: e06189 doi:10.1111/ecog.06189
- (2022) Pilcher, D.J., J.N. Cross, A. Hermann, K. Kearney, W. Cheng, J.T. Mathis. Dynamically downscaled projections of ocean acidification for the Bering Sea, *Deep-Sea Research II: Topical Studies in Oceanography* 198, 105055
- (2022) Hollowed, A. B., A. C. Haynie, A. J. Hermann, K. K. Holsman, A. E. Punt, C. S. Szuwalski. Implications of climate change on the Bering Sea and other cold water systems. Introduction to the special issue of Deep-Sea Research Part II: Topical Studies in Oceanography.
- (2022) Punt, A.E., et al., Szuwalski, C.S. 2022. A framework for assessing harvest strategy choice when considering multiple interacting fisheries and a changing environment: The example of eastern Bering Sea crab stocks. Fisheries Research. 252: 106338.
- (2022) Szuwalski, C.S.. Estimating time-variation in confounded processes in population dynamics modeling: a case study for snow crab in the eastern Bering Sea. Fisheries Research. 251: 106298.
- (2021) Hermann, A. J., Kearney, K., Cheng, W., Pilcher, D., Aydin, K., Holsman, K. K., & Hollowed, A. B.. Coupled modes of projected regional change in the Bering Sea from a dynamically downscaling model under CMIP6 forcing. Deep-Sea Research Part II: Topical Studies in Oceanography, 194 (Dec), 104974. <https://doi.org/10.1016/j.dsr2.2021.104974>

ACLIM Publications (direct & indirect) 1 of 2

- (2021) Cheng, W., Hermann, A. J., Hollowed, A. B., Holsman, K. K., Kearney, K. A., Pilcher, D. J., Stock, C. A., & Aydin, K. Y. Eastern Bering Sea shelf environmental and lower trophic level responses to climate forcing: Results of dynamical downscaling from CMIP6. *Deep-Sea Research Part II: Topical Studies in Oceanography*, 193, 104975. <https://doi.org/10.1016/j.dsr2.2021.104975>
- (2021) Punt, A., M G Dalton, W Cheng, A Hermann, K Holsman, T Hurst, J Ianelli, K Kearney, C McGilliard, D Pilcher, M Véron. Evaluating the impact of climate and demographic variation on future prospects for fish stocks: An application for northern rock sole in Alaska. *Deep Sea Research Part II: Topical Studies in Oceanography* 189–190:104951.
- (2021) Whitehouse, G. A., K. Y. Aydin, A. B. Hollowed, K. K. Holsman, W Cheng, A. Faig, A. C. Haynie, A. J. Hermann, K. A. Kearney, A. E. Punt, and T. E. Essington. Bottom-up impacts of forecasted climate change on the eastern Bering Sea food web. *Front. Mar. Sci.*, 03 February 2021 | <https://doi.org/10.3389/fmars.2021.624301>
- (2020) Holsman, K.K., A. Haynie, A. Hollowed, J. Reum, K. Aydin, A. Hermann, W. Cheng, A. Faig, J. Ianelli, K. Kearney, A. Punt. (2020) Ecosystem-based fisheries management forestalls climate-driven collapse. *Nature Communications*. DOI:10.1038/s41467-020-18300-3
- (2021) Thorson, J., M. Arimitsu, L. Barnett, W. Cheng, L. Eisner, A. Haynie, A. Hermann, K. Holsman, D. Kimmel, M. Lomas, J. Richar, E. Siddon. Forecasting community reassembly using climate-linked spatio-temporal ecosystem models. *Ecosphere* 44: 1–14, doi: 10.1111/ecog.05471
- (2020) Szuwalski, W. Cheng, R. Foy, A. Hermann, A. Hollowed, K. Holsman, J. Lee, W. Stockhausen, J. Zheng. Climate change and the future productivity and distribution of crab in the Bering Sea. *ICES J. Mar. Sci fsaa140*, <https://doi.org/10.1093/icesjms/fsaa140>
- (2020) Reum, J. C. P., J. L. Blanchard, K. K. Holsman, K. Aydin, A. B. Hollowed, A. J. Hermann, W. Cheng, A. Faig, A. C. Haynie, and A. E. Punt. 2020. Ensemble Projections of Future Climate Change Impacts on the Eastern Bering Sea Food Web Using a Multispecies Size Spectrum Model. *Frontiers in Marine Science* 7:1–17.
- (2020) Hollowed, A. B., K. K. Holsman, A. C. Haynie, A. J. Hermann, A. E. Punt, K. Aydin, J. N. Ianelli, S. Kasperski, W. Cheng, A. Faig, K. A. Kearney, J. C. P. Reum, P. Spencer, I. Spies, W. Stockhausen, C. S. Szuwalski, G. A. Whitehouse, and T. K. Wilderbuer. 2020. Integrated Modeling to Evaluate Climate Change Impacts on Coupled Social-Ecological Systems in Alaska. *Frontiers in Marine Science* 6. <https://doi.org/10.3389/fmars.2019.00775>
- (2019) Holsman, KK, EL Hazen, A Haynie, S Gourguet, A Hollowed, S Bograd, JF Samhour, K Aydin, Toward climate-resiliency in fisheries management. *ICES Journal of Marine Science*. 10.1093/icesjms/fsz031
- (2019) Hermann, A. J., G.A. Gibson, W. Cheng, I. Ortiz1, K. Aydin, M. Wang, A. B. Hollowed, and K. K. Holsman. Projected biophysical conditions of the Bering Sea to 2100 under multiple emission scenarios. *ICES Journal of Marine Science*, fsz043, <https://doi.org/10.1093/icesjms/fsz043>
- (2019) Reum, J., JL Blanchard, KK Holsman, K Aydin, AE Punt. Species-specific ontogenetic diet shifts attenuate trophic cascades and lengthen food chains in exploited ecosystems. *Okios* DOI: 10.1111/oik.05630
- (2019) Reum, J., K. Holsman, KK, Aydin, J. Blanchard, S. Jennings. Energetically relevant predator to prey body mass ratios and their relationship with predator body size. *Ecology and Evolution* (9):201–211 DOI: 10.1002/ece3.4715

Sea Ice Loss

Arctic Antarctic Start month: January Colors: Viridis Invert colors Vertical gridlines




ChArctic-v3.5.0

National Snow and Ice Data Center, Boulder, CO



National Snow and Ice Data Center

a part of CIRES at the University of Colorado Boulder



Sea Ice Loss

Arctic Antarctic

Start month: January


Colors: Viridis

Invert colors

Vertical gridlines

Arctic Sea Ice Extent

(Area of ocean with at least 15% sea ice)



ChArctic v3.5.0

National Snow and Ice Data Center, Boulder, CO



National Snow and Ice Data Center

a part of CIRES at the University of Colorado Boulder

