

# Ecosystem Status Report Aleutian Islands 2024 – SSC Dec

*Thank you!*

Ivonne Ortiz & Stephani Zador



**With contributions from:**

Mayumi Arimitsu, Kerim Aydin, Lewis Barnett, Sonia Batten, Jessica Beck, Shawn Bell, Thaddaeus Buser, Mathew W. Callahan, Wei Cheng, Christina Conrath, Alexandra Dowlin, Thomas Farrugia, Sarah Friedman, Sarah Gaichas, Tom Gelatt, Rebecca Howard, Robb Kaler, Mandy Keogh, Ned Laman, Geoffrey M. Lang, Emily Lemagie, Jackie Lindsey, Jackie McConnell, Susanne McDermott, Ivonne Ortiz, Clare Ostle, James Overland, Bianca Prohaska, Sean Rohan, Nora Rojek, Natalie Rouse, Greg Ruggerone, Margaret Siple, Phyllis Stabeno, Katie Sweeney, Cathy Tide, Rick Thoman, Muyin Wang, George A. Whitehouse, Bruce Wright, Stephani Zador, Adam Zaleski and Mark Zimmermann

photos: [photolib.noaa.gov](https://photolib.noaa.gov)



## Response to SSC comments (December 2023)

“...more focus on multi-year patterns...”

*We continue to address biennial patterns, and shifts in ecosystem structure potentially related to either the step increase of Eastern Kamchatka pink salmon as well as a shift in pelagic foragers now dominated by rockfish, which might be related but not driven by climate. This year, we continue looking at diet shifts and fish condition.*

## Response to SSC comments (December 2023)

“...more focus on multi-year patterns...”

*We continue to address biennial patterns, and shifts in ecosystem structure potentially related to either the step increase of Eastern Kamchatka pink salmon as well as a shift in pelagic foragers now dominated by rockfish, which might be related but not driven by climate. This year, we continue looking at diet shifts and fish condition.*

“...encourages ESR authors to collaborate with other social scientists about other potential indicators...”

*We continue to be a part of broader discussions on the identification of effective social and economic indicators and following the recommendation by the GPT this year we will be meeting with staff from the Department of Labor Research and Analysis.*

## Response to SSC comments (December 2023)

“...more focus on multi-year patterns...”

*We continue to address biennial patterns, and shifts in ecosystem structure potentially related to either the step increase of Eastern Kamchatka pink salmon as well as a shift in pelagic foragers now dominated by rockfish, which might be related but not driven by climate. This year, we continue looking at diet shifts and fish condition.*

“...encourages ESR authors to collaborate with other social scientists about other potential indicators...”

*We continue to be a part of broader discussions on the identification of effective social and economic indicators and following the recommendation by the GPT this year we will be meeting with staff from the Department of Labor Research and Analysis.*

“ ...SSC supports dedicated ecosystem studies in the AI... to see if predator/prey relationships have changed over time...determine if observed changes reflect a broader transition in the ecosystem...”

*We will continue to work towards a contribution on groundfish diets*

## Response to SSC comments (December 2023)

“...more focus on multi-year patterns...”

*We continue to address biennial patterns, and shifts in ecosystem structure potentially related to either the step increase of Eastern Kamchatka pink salmon as well as a shift in pelagic foragers now dominated by rockfish, which might be related but not driven by climate. This year, we continue looking at diet shifts and fish condition.*

“...encourages ESR authors to collaborate with other social scientists about other potential indicators...”

*We continue to be a part of broader discussions on the identification of effective social and economic indicators and following the recommendation by the GPT this year we will be meeting with staff from the Department of Labor Research and Analysis.*

“ ...SSC supports dedicated ecosystem studies in the AI... to see if predator/prey relationships have changed over time...determine if observed changes reflect a broader transition in the ecosystem...”

*We will continue to work towards a contribution on groundfish diets*

“... survey the AI during odd years (when Kamchatka Pink salmon abundances peak biennially) would be valuable...”

*We are hopeful of renewed efforts in collaboration with the Pacific marine Environmental Laboratory....and continue to explore other data sources available annually, such as seabird and fisheries-dependent data.*

## Response to SSC comments (December 2023)

“...more focus on multi-year patterns...”

*We continue to address biennial patterns, and shifts in ecosystem structure potentially related to either the step increase of Eastern Kamchatka pink salmon as well as a shift in pelagic foragers now dominated by rockfish, which might be related but not driven by climate. This year, we continue looking at diet shifts and fish condition.*

“...encourages ESR authors to collaborate with other social scientists about other potential indicators...”

*We continue to be a part of broader discussions on the identification of effective social and economic indicators and following the recommendation by the GPT this year we will be meeting with staff from the Department of Labor Research and Analysis.*

“ ...SSC supports dedicated ecosystem studies in the AI... to see if predator/prey relationships have changed over time...determine if observed changes reflect a broader transition in the ecosystem...”

*We will continue to work towards a contribution on groundfish diets*

“... survey the AI during odd years (when Kamchatka Pink salmon abundances peak biennially) would be valuable...”

*We are hopeful of renewed efforts in collaboration with the Pacific marine Environmental Laboratory....and continue to explore other data sources available annually, such as seabird and fisheries-dependent data.*

“...the authors consider moving East Kamchatka pink salmon indicators and POP indicators into the report cards. For pink salmon, it was discussed that additional information on biomass may be valuable...”

*We will be evaluating how to best update the report cards with new information, not only with POP/rockfish and pink salmon biomass, but temperature and seabird information as well. Biomass plots are now included in EK pink salmon contribution. This year, we placed that biomass in context with historical levels of EK pink salmon and biomass of AI groundfish stocks.*

# 2024 Ecosystem Status Report – Aleutian Islands

## Risk Table

### Environmental/Ecosystem Considerations

#### Sustained Level 2

Multiple indicators showing consistent adverse signals a) across the same trophic level as the stock, and/or b) up or down trophic levels (i.e., predators and prey of the stock)

- AI Pacific cod
- Atka mackerel

#### Sustained Level 1

No apparent ecosystem concerns related to biological status (e.g., environment, prey), or minor concerns with uncertain impacts on the stock.

- Pacific ocean perch
  - AI pollock
- Bogoslof pollock
  - R&B rockfish
  - Shortraker
  - Other rockfish

## Assessment 2024

1. Relaxation of prolonged warming

*Finally. But is it cool enough for the ecosystem to respond?*

2. Gradient of mostly poor productivity in west to good productivity in east

*Why different responses to the improved conditions?*

3. Few pink salmon, but this “low” year is now as high as previous high years

*All those EK Pinks are exporting energy, as low abundance = 60,000 mt  
current AI P. cod biomass = 70-80,000 mt*

4. Groundfish condition mostly poor throughout, even though environmental conditions would predict otherwise

*Is a rebalancing of the ecosystem keeping groundfish from finding enough food*



Sustained Level 2



AI Pacific cod



- Warm winter conditions
- Lower amount of fish in diet since ~2010
- Lower prey quality resulting in reduced fish condition.
- Decreased consumption of Atka as prey due to lower availability of Atka.

Sustained Level 2

Atka mackerel



- Warm winter conditions
- Smaller species in copepod communities
- Deeper mixed layer: potential impact on availability of prey in water column
- Lower than average fish condition across the entire chain
- Increased competition for prey (high rockfish abundance)






- School at Adak closed
- Decreasing trend even in Unalaska
- Akutan – only one increasing


# Noteworthy

# Changes in Pacific cod diets II



## Western Aleutians NMFS 543


- Less fish in diets
- Consumption of prey needs to be at least 1% of predator weight
- Correspondence between Atka mackerel biomass and total prey consumption
- Correspondence between total prey consumed and fish condition
- Atka mackerel biomass drives fish consumption in Western, Central Aleutians • higher temperature coincide with lower condition




# Noteworthy

# Changes in Pacific cod diets II


Pcod diet WAI




Prey biomass vs consumed prey



Fish condition




Bottom T °C




## Eastern Aleutians NMFS 541

- Less fish in diets
- Consumption of prey needs to be at least 1% of predator weight but of adequate caloric value?
- Less correspondence between Atka mackerel biomass and total prey consumption
- Less correspondence between total prey consumed and fish condition – quality of prey may partly explain this?
- Atka mackerel biomass drives fish consumption in Western, Central Aleutians • higher temperature coincide with lower condition



# Noteworthy

# Changes in Pacific cod diets II





NMFS 543

NMFS 542


NMFS 541

NMFS 518, 519




Aleutian Low –dominant atmospheric pressure system in winter Jan-Feb

- Defined by extent of pressure system & location of center on Aleutian Low; 2024: near average
- Strong winds and stormy conditions (mean NPI), deeper mixed layer, potentially changing availability of prey




data through 11-12-2023



NOAA Coral Reef Watch data, courtesy NOAA Pacific Islands Ocean Observing System (Updated: 11-10-2024)  
 Data are modeled satellite products and periodic discrepancies or gaps may exist across sensors and products.  
 Contact: matt.callahan@noaa.gov, Alaska Fisheries Science Center

Satellite SST


- Warm winter throughout, among 10 warmest winters
- Cooler spring & summer but still at or above 1985-2015 mean
- WAI – temperatures increased again in late summer and fall




• SST • BT

## Eddy Kinetic Energy


- Some increased flows in WAI beginning of year
- Currently below average in all three regions
- Lower transport of heat, salt and nutrients through passes
- EAI influenced by GOA weakening of coastal circulation and movement of eddies offshore related to the marine heatwave (warm events in general with increased upwelling) (Rallu De Malibrán et al. 2024 )






Continuous Plankton Recorder

- Large diatom biomass in 2023: above mean for third year biennial pattern – gone after 2012?




- Smaller size of copepod community **potentially due to warmer temperatures**
- below average since 2024



- First year with above average meso-zooplankton biomass






- Condition decreased across entire chain
- Atka mackerel condition decreased even in western and central Aleutians
- Pacific cod, northern rockfish and Pacific ocean perch below average condition since 2012
- Exceptions:
  - small pollock and southern rock sole

Werb and Rudnick, 2023. Remarkable changes in the dominant modes of North Pacific sea surface temperature. Modified from Figures 1b and 3b.


# Ortiz

# Biomass of Pelagic Foragers and Apex Predators





**Pelagic Foragers: Western AI**




**Central AI**




**Eastern AI**







**Apex Predators: Western AI**



**Central AI**



**Eastern AI**




- Stable or increasing biomass of pelagic foragers across the chain
- Slow decrease of apex predators, lowest and decreasing biomass in Western Aleutians


## Other related trends:

- Eelpouts, shrimp (common prey of apex predators) decreasing across the chain
- Structural epifauna is either decreasing or stable in the western and central Aleutians, sea pens increasing in the east
- Low abundance year for Eastern Kamchatka pink salmon, continuing cascading effects but more moderate?

Western Aleutians



Eastern Aleutians



2020-2024 Mean

- 1 s.d. above mean
- 1 s.d. below mean
- within 1 s.d. of mean
- fewer than 2 data points

2020-2024 Trend

- increase by 1 s.d. over time window
- decrease by 1 s.d. over time window
- change <1 s.d. over window
- fewer than 3 data points

### Seabird Reproductive Success

- Average or above for EAI seabirds: good foraging conditions in summer for plankton and fish foragers; poor in WAI

Site	Species													EAI	WAI	
	Primarily fish eaters						Primarily zooplankton eaters									
	glaucous winged gull	common murre	thick billed murre	horned puffin	tufted puffin	red-legged kittiwake	black-legged kittiwake	fork-tailed storm-petrel	Leach's storm-petrel	ancient murrelet	parakeet auklet	least auklet	whiskered auklet	crested auklet		
Aiktak						-	-				-	-	-	-		
Buldir		-								-						

Average & above

Average & below



Above average




Average




Below average

### Seabird Diets


Mostly capelin in the EAI, Atka mackerel in CAI, squid in WAI:





**Steller sea lions non-pups and pups:**







- western Distinct Population Segments non-pup and pup counts increased between 2008-2023
- Regionally:  
Western Aleutians non-pups and pups declined,  
Central Aleutians non-pups stable (but RCA 2,3 declined)  
pups declined, RCA 5 stable  
Eastern Aleutians non-pups and pups increased



# Rockfish as main pelagic foragers


# Multi-year patterns

- 63% vs 34% of pelagic forager biomass with lifespan of 88-104 in 1991 and 2022 respectively



- Last year prediction: sea ice should extend south of 60°N perhaps all the way to M2, largely borne out
- **60% chance of La Niña** during January – March 2025,
- Climate prediction center, NOAA [https://www.cpc.ncep.noaa.gov/products/analysis\\_monitoring/enso\\_advisory/ensodisc.shtml](https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/enso_advisory/ensodisc.shtml)
- **Warm** conditions for western Aleutians (NMME, Bond) for January - March
- Ensemble indicates conditions should not be extreme relative to the past 20-30 years


## Sea surface anomalies from National Multi-Model Ensemble



# Summary and implications



Jan – Mar among 10 warmest winter on record SST, cooler spring & summer except western Aleutian Islands; average bottom temperature  
*Potential concern for winter survival and winter spawners; lower bioenergetic costs but cooler temperature are just average*



Small copepod size, larger meso-zooplankton biomass (2023)

*More zooplankton but lower availability of large copepods as prey*



Pacific cod diets

*consumption of prey >1% of predator weight coherent with above average condition, Atka mackerel main fish prey in WAI, CAI*



Seabird reproductive success above average in EAI for plankton and fish-eating seabirds; poor for WAI

*Indicates potential availability of prey and good foraging conditions for both plankton and fish eating groundfish in EAI, but poor availability in prey in WAI*



Rockfish dominance of pelagic forage fish biomass

*Potential for increased competition for zooplankton; Rockfish are now 64% of the guild biomass with lifespans between 88-104 which years less fish prey for apex predators, slow response to environment which increases ecosystem stability*



Increasing Eastern Kamchatka pink salmon during both low abundance and high abundance years

*In a low abundance year, EK pink salmon is similar to current P. cod stock, peak abundance in 2019 reached 350,000 mt*

*Slow trends in the western and central Aleutians: decreasing structural epifauna, decreasing miscellaneous fish*

*potentially less alternative fish prey in the system and decreasing critical habitat*

***Transition of the ecosystem to a state where rockfish are the main pathway of zooplankton into food web, pink salmon export energy from the ecosystem.***

## Additional Information Available

---

Ecosystem Status Reports through 2024 are available [here](#):



ESR Reports (1999-2024)