
Climate work plan overview and Alternative Harvest Control Rule discussion paper

Diana Stram, NPFMC
Kirstin Holsman, AFSC

Ecosystem Committee: May 8, 2026



Outline of presentation

Climate work plan

- Overview of 4 elements
- Spreadsheet tracking tool
- Coordination with CEFI
- Current Council focus on Element 4

Harvest Control Rules (HCR) discussion paper

- Overview of HCRs (general)
- NPFMC Groundfish Assessment Tier system and OFL/ABC HCRs
- NPFMC BSAI Crab Assessment Tier system and OFL/ABCs
- How risk and/or uncertainty considered in NPFMC:
 - Stock Assessments
 - ABC-setting
 - TAC-setting
- Alternative HCRs under development
- Decisions points and scope for Ecosystem Committee recommendations



Climate work plan

Climate work plan adopted by the Council December 2024

- Spreadsheet tracking tool developed by staff
- Tracks combined efforts of CEFI, AFSC, AKRO and Council
- Initial summary focused on 2025-2026
- Items planned but not yet started ('potential') not included until they move into the 'new/in development' phase
- Current Council focus on Element 4 (HCR work)

Four initial Elements:

- A. Incorporate climate-forecast linked management advice
- B. Incorporate climate-driven interactions and cascading impacts through use of ecosystem indicators and models
- C. Consider and incorporate dynamic management tools to increase in-season adaptation capacity
- D. Review tier systems, consider climate-informed biomass targets and limits and climate-robust or forecast-informed harvest control rules



Work plan tracking tool

Key Element	Council Product *	Product Detail*	New, existing, modified	Relevant FMPs	Lead responsibility: Council staff/ RO/AFSC	
Element B: Incorporate climate-driven interactions and cascading impacts through use of ecosystem indicators and models						
B.6	Link to groundfish, crab, and marine mammal SDMs to MOM6 hindcasts and forecasts for EBS	EBS ESR	Develop climate informed SDMs and management trajectories based on MOM6 linked assessment models for groundfish and crab - provide contribution to the ESR	Modification to existing	BSAI Crab, BSAI Groundfish	Lead: AFSC, Partners: AKRO, NPFMC
B.7	Link to ice seal SDMs to MOM6 hindcasts and forecasts for EBS	EBS ESR (noteworthy or index)	Develop climate informed SDMs and management trajectories based on MOM6 linked assessment models for marine mammals	New/ in development	BSAI Crab, BSAI Groundfish	Lead: AFSC, Partners: AKRO, NPFMC
B.8	Coordination through SDM 6 month meetings and workshops	SDM and EFH updates	Coordination through SDM 6 mo meetings and workshops to facilitate cross pollination & tie into deliverables, reports and the dashboard	Existing/ operational	GOA Groundfish, GOA Crab, BSAI Crab, BSAI Groundfish	Lead: AFSC, Partners: AKRO, NPFMC
Element C: Consider and incorporate dynamic management tools to increase in-season adaptation capacity						
C.1	ACL in-season management authority	Regulatory impact review (RIR)	Analysis of modifying in-season management authority	New/ in development	BSAI Groundfish, GOA Groundfish	Co-leads NPFMC, AKRO
C.2	Stock Assessment Prioritization	ongoing discussions with NPFMC (PT, SSC)	Within typical stock assessment review process, across all stocks as needed	Modification to existing	BSAI Groundfish, GOA Groundfish	Lead: AFSC, Partners: NPFMC
Element D: Review tier systems, consider climate-informed biomass targets and limits and climate-robust or forecast-informed harvest control rules						
D.1		Presentation on Tier systems and flexibility within FMPs cor Groundfish and Crab FMPs	Presentation at SSC workshop	Completed	BSAI Groundfish, BSAI Crab, GOA Groundfish	Lead: NPFMC, Partners AFSC
D.2	Alternative Harvest Control Rule (HCRs) considerations	SSC Workshop on HCRs	SSC Workshop June 2025: https://meetings.npfmc.org/Meeting/Details/3093	Completed	BSAI Groundfish, BSAI Crab, GOA Groundfish, GOA Crab	Lead: NPFMC

Key Element	Council Product *	Product Detail*	New, existing, modified	Relevant FMPs	Lead responsibility: Council staff/ RO/AFSC	
Element A: Incorporate climate forecast linked management advice						
A.1	Create Science Brief; CEFI 1 pager	Climate Science Brief	Summarize new CEFI products, highlight new climate relevant information for the assessment season	Modification to existing	BSAI Groundfish	Lead: AFSC, Partners: NPFMC
A.4	Generate MOM6 indices (hindcast + forecast) for annual assessment cycle products	ESR EBS, Risk tables in stock assessments	Indices of cold pool, BT, SST, etc.	Modification to existing	BSAI Groundfish, BSAI Crab	Lead: AFSC, Partners: NPFMC
A.5		ESR AI, Risk tables in stock assessments	Indices of cold pool, BT, SST, etc.	New/ in development	BSAI Groundfish, BSAI Crab	Lead: AFSC, Partners: NPFMC
A.6		ESR GOA, Risk tables in stock assessments	Indices of cold pool, BT, SST, etc.	New/ in development	GOA Groundfish	Lead: AFSC, Partners: NPFMC
A.7		ESPs for BSAI stock assessments	Indices of cold pool, BT, SST, etc.	New/ in development	BSAI Groundfish, BSAI Crab	Lead: AFSC, Partners: NPFMC

3.0 Details from each 2025-2026 listed Element

Element A

A.1) Climate Science Brief (plus CEFI 1 pager)

A summary of new Changing Ecosystem and Fisheries Initiative (CEFI) products that highlights new climate-relevant information for the assessment season. The Climate Report is delivered annually and informs the groundfish and crab assessment cycles. A [pilot version of the report](#) was presented through the Fall 2024 assessment cycle (no report was generated in 2025 due to the government shutdown).

A.4, A.5, A.6, A.7) MOM6-derived indices for ESRs and ESPs:


Alaska regionally-tailored indices of cold pool, bottom temperature, sea surface temperature, mixed layer depth, Ocean Acidification (OA) and Aragonite Saturation rate, and additional variables are being generated from MOM6 hindcasts and forecasts and being provided publicly for use in various decision support tools including the EBS, AI, and GOA ESRs and to the ESPs in EBS and GOA SAFE chapters to support the annual assessment cycle.



Work Plan lists actions by AFSC/CEFI, AKRO and Council actions

As products are available they will be incorporated into the Council process (via assessments, ESRs, analyses, etc)

Council and RO staff also included in CEFI overall as well as Alaska Climate Team (ACT) to coordinate and communicate activities





Harvest Control Rules (HCR): treatment of risk and uncertainty




Considerations of Risk and Uncertainty

Council consideration of treatment of risk and uncertainty in harvest specifications for BSAI/GOA Groundfish and BSAI crab stocks for potential improvements.



Overfishing Limit (OFL)
Acceptable Biological Catch (ABC)



Total Allowable Catch (TAC)



Council (groundfish)
State of Alaska (crab)

Uncertainty

Relative lack of information or confidence in the reliability of that information

Risk

Considering the likelihood of an adverse event and the severity of the consequences



General sloping HCR (Fig 1)

Concepts


HCR: guideline for how much fishing can take place based upon an estimate of stock status

Spawning stock biomass (SSB): Best estimate of sexually mature biomass capable of reproducing

SSB_{LIMIT} SSB_{TARGET}

Fishing mortality rate (F): rate applied to SSB to determine catch. F_{LIMIT} F_{TARGET}

HCR






Groundfish Tier System and current OFL and ABC control rules (Section 2.3)



Groundfish Stock Assessment cycle (Fig 2)

Annual Groundfish Stock Assessment Cycle




Annual assessment review and recommendations process for BSAI and GOA groundfish



Groundfish Tier System (Fig 3)

Groundfish tier system



BSAI and GOA stocks by Tiers (Tables 1 and 2)


Stock	BSAI	Assessment Region	Tier
Pollock		BS	3
		AI	3
		Bogoslof	5
Pacific cod		BS	3
		AI	3
Sablefish		BSAI/GOA	3
Yellowfin sole		BSAI	1
Greenland turbot		BSAI	3
Arrowtooth flounder		BSAI	3
Kamchatka flounder		BSAI	3
Northern rock sole		BSAI	1
Flathead sole		BSAI	3
Alaska plaice		BSAI	3
Other flatfish		BSAI	5
Pacific Ocean perch		BSAI	3
Northern rockfish		BSAI	3
Blackspotted/Rougheye rockfish		BSAI	3
Shortraker rockfish		BSAI	5
Other rockfish		BSAI	5
Atka mackerel		BSAI	3
Skates		BSAI	3,5 ¹
Sharks		BSAI	6
Octopuses		BSAI	6

Stock	GOA	Assessment Region	Tier
Pollock		W/C/WYAK	3
		SEO	5
Pacific cod		GOA	3
Sablefish		BSAI/GOA	3
Shallow-Water flatfish		GOA	3,5 ¹
Deep-Water flatfish		GOA	3,6 ²
Rex sole		GOA	3
Arrowtooth flounder		GOA	3
Flathead sole		GOA	3
Pacific Ocean perch		GOA	3
Northern rockfish		GOA	3
Shortraker rockfish		GOA	5
Dusky rockfish		GOA	3
Rougheye and Blackspotted rockfish		GOA	3
Demersal Shelf rockfish		W/C/WYAK	6
		SEO	5,6 ³
Thornyhead rockfish		GOA	5
Other rockfish		GOA	4,5,6 ⁴
Atka mackerel		GOA	6
Big Skate		GOA	5
Longnose Skate		GOA	5
Other Skates		GOA	5
Sharks		GOA	5,6 ⁵
Octopuses		GOA	6



Groundfish OFL and ABC HCRs (Fig 4)

Groundfish Control Rule






BSAI Crab Tier System and current OFL and ABC control rules
(Section 2.4)




Crab Assessment and Review/Recommendations Cycle (Fig 6)

Annual Crab Stock Assessment Cycle




Crab Tier System (Fig 5)

Crab tier system



Crab OFL HCR

Crab control rule






Risk and Uncertainty:
Assessment, ABC-setting, TAC-setting, Alternative
HCRs (Section 3)




Science and policy in addressing risk and uncertainty

Scientific uncertainty: uncertainty in OFL


SSC Role




Scientific uncertainty & risk policy



Council Role



Management uncertainty



Management uncertainty: uncertainty in ability to constrain catch to ACL (= ABC)



Assessment changes that can impact ABC (Section 3.1)

- Model structure modifications:
 - Approaches reviewed at least Sept/Oct to Nov/Dec before adoption
- Biological reference points/new or revised data:
 - New data, revised time-series, selectivity changes
 - No change to model structure but can impact biological reference points
 - Not always reviewed in Sep/Oct prior to incorporation
- Tier level changes
 - Proposed changes reviewed at least Sept/Oct to Nov/Dec before adoption

Table 3 page 16

Stock	Proposed change/ rationale	Change in resulting maxABC	Time frame from proposal to adoption for use in specifications
EBS Pollock	Change Tier 1 to Tier 3, lack of data at low stock sizes results in uncertain Stock-Recruit relationship (SRR)	-1,666,000 t	Discussion paper reviewed in September/October 2024; Tier level change by SSC in December 2024
BSAI Yellowfin sole	Change Tier 1 to Tier 3, lack of data at low stock sizes results in uncertain SRR	Not yet available	Discussion paper September/October 2025; change pending 2026 assessment cycle
AI Pacific cod	Change from Tier 5 to Tier 3; approved age-structured model	-3,134	Multiple years in development and 2024 approval of age-structured model
BSAI Atka mackerel	Estimate of F_{OFL} and F_{ABC} 25% higher relative to previous year due to changes in the fishery selectivity	+10,290 t	Assessments in 2021 relative to 2020
Various	Revised Tier 3 proxies, age-(or sex) specific schedules updated. E.g., selectivity, weight-age, and maturity	Varies	Generally proposed in September, if approved by SSC, will be included for December meeting



ABC setting [Groundfish] Section 3.2

Tier specific HCR = maximum permissible ABC (maxABC)

maxABC is the upper bound on ABC but the SSC may recommend an ABC below the maxABC

No flexibility to recommend ABC >maxABC

Process in general:

Assessment author develops a recommendation for $ABC < \text{maxABC}$


Review by Plan Team and subsequent recommendation

Final recommendation by SSC

Reduction per FMP and NS1 Guidelines is based upon “whether conditions exist that warrant setting ABC at a value lower than the maximum permissible value” and accounts for “any other scientific uncertainty”

Reductions from maxABC have been documented recently through the use of risk tables

Groundfish Control Rule



Risk tables: Scientific uncertainty not directly captured in the stock assessments

Categories of concern

1. Assessment-related considerations
 2. Population dynamics considerations
 3. Ecosystem considerations
 4. Fishery-related stock considerations
- Risk table categories describe scientific uncertainty not directly captured in the stock assessments
 - Ranked 1-3 based on relative level of concern
 - Normal (1)
 - Increased concern (2)
 - Extreme concern (3)
 - Can be used to document recommendations of ABC < maxABC




Table 6, pg 22

Table 6: ABC < maxABC 2018-2024

Note categories and scoring levels have changed between 2018-2022

Stock	Assessment Year(s)	Risk Table Scores > 1	SSC reduction from maxABC
EBS Pollock	2018	P-2, E-2	30%
	2019	P-2, E-2, F-2	43%
	2020	E-2, F-2	30%
	2021	A-2, E-2, F-2	11%
	2022	A-2	43%
	2023	E-2	18%
EBS Pacific cod	2019	E-2	3%
AI Pacific cod	2023	P-2, E-2	10%
AK Sablefish	2018	A-2, P-4, E-2	45%
	2019	A-2, P-3, E-2, F-3	50%
	2020	A-3, P-3, E-2, F-3	44%
BSAI Yellowfin sole	2021	P-2, E-2	24%
BSAI Greenland turbot	2024	A-3, P-2, E-2	25%
BSAI Northern rock sole	2022	A-3	23%
	2023	rollover	23%
BSAI Blackspotted / <u>Rougheve</u> rockfish	2022	A-3, P-2, F-2	11%
	2023	rollover	11%
BSAI Sharks	2022	A-3, P-2	13%
	2023-2024	rollover	13%
GOA Pollock	2023	A-2	18%
GOA Pacific cod	2018	NA*	14%
	2019	A-2, P-2, E-2	NA**
GOA Dusky Rockfish	2020	A-2	24%
	2021	Rollover of stair-step	24%
GOA <u>Rougheve</u> /Blackspotted rockfish	2023	A-2, P-2	20%
	2024	NA***	10%
GOA Demersal shelf rockfish	2021	A-2, P-2, F-2	22%
	2022	A-2, P-2, F-2	15%

ABC setting Crab (Fig 10, Table 4)



Crab Stock	Tier	ABC Buffer by year				
		2021/22	2022/23	2023/24	2024/25	2025/26
EBS Snow Crab	3	25%	25%	50%	65%	40%
Bristol Bay Red King Crab	3	20%	20%	20%	20%	20%
EBS Tanner Crab	3	20%	20%	20%	20%	20%
Pribilof Islands red king crab	4	25%	25%	25%	25%	25%
Pribilof Islands blue king crab	4	25%	25%	25%	25%	25%
St. Matthew blue king crab	4	25%	25%	25%	25%	25%
Norton Sound red king crab	4	40%	40%	30%	30%	30%
Aleutian Islands golden king crab	3	30%	25%	25%	25%	25%
Pribilof Islands golden king crab	5	25%	25%	25%	25%	25%
Western Aleutian Islands red king crab	5	75%	75%	75%	75%	75%



Risk Table development for crab (Table 7)

Crab Stock	Tier	Risk Table/Year	Risk Table Score			
			Assessment	Population dynamics	Ecosystem	Fishery
EBS Snow Crab	3	Y 2025	2	2	1	1
Bristol Bay Red King Crab	3	Y 2025	1	2	2	1
EBS Tanner Crab	3	Y 2025	2	1	1	1
Pribilof Islands red king crab	4	Y 2025	2	2	1	1
Pribilof Islands blue king crab	4	Y 2025	1	1	1	1
St. Matthew blue king crab	4	N				
Norton Sound red king crab	4	N				
Aleutian Islands golden king crab	3	Y 2025	1	2	1	2
Pribilof Islands golden king crab	5	N				
Western Aleutian Islands red king crab	5	N				

TAC setting

Groundfish

- Annually recommends final harvest specifications to the Secretary of Commerce
- SUM of the recommended TACs must fall within the Optimum Yield (OY) range by FMP (specified in FMPs and implementing regs):
 - BSAI = 1.4 – 2.0 million mt
 - GOA 116,000 – 800,000 mt
- TAC must be set = or < ABC.
- The Council retains broad latitude to recommend TAC below ABC on an annual basis, consistent with the Groundfish FMPs and implementing regulations
 - drawing on socioeconomic information and concerns not addressed in the ABC-setting process provided the sum of the TACs is within the ranges specified in the respective FMPs

BSAI Crab

- State of Alaska as a Category 2 measure under the Crab FMP (framework management measures)



Benefits and Drawback across 3 approaches (Table 8)

Approach	Benefits	Drawbacks	Timing
Assessment changes	Highly targeted; can address stock-specific dynamics; scientific peer review provides quality control	Less transparent to the public; extended development time; changes within scientific purview, not direct Council action	Depends upon scope and scientific concurrence on modifications; can be done within two meetings of Plan Teams and SSC; sometimes may take multiple iterations of changes to eventually get approved (e.g., AI Pacific cod model and resulting Tier level change)
ABC changes from maxABC	Can be done within an assessment cycle drawing on information within the assessment (e.g., where recruitment estimates are modified) or with the use of risk table scoring and justifications for out of assessment model uncertainties	Qualitative; scoring and resulting reductions may be inconsistent across assessments; it is not possible to recommend an ABC above the maxABC	Modified by the SSC during their annual stock assessment review and ABC recommendations
TAC adjustments	Flexible and responsive on annual timescales; can draw on a broad range of information; no FMP amendment needed but sum of TACs must fall within the OY range by FMP.	Does not create lasting structural change; ad hoc; may lack analytical basis; subject to annual political dynamics; and structure (e.g., use of HCRs and risk tables)	Modified within a Council meeting cycle without additional analysis

Considerations for revising treatment of risk and uncertainty

Status quo

- Framework in place for many years
- BRP and ABC can change within individual assessments in response to:
 - New information
 - Shifts in information availability and Tiers
 - Changes in spatial distributions that may influence availability and/or catchability
 - Changes in selectivity and catchability (critical to assessment):
 - Gear changes
 - Time-area-gear regs influencing selectivity
 - Other factors impacting harvest practices (PSC caps, Discard mortality regs etc)



Concerns regarding population impacts to fish and crab under future environmental conditions lead to consideration of more adaptive HCRs

- Selectivity: probability fish is captured by gear (based on size, age, species)
- Catchability: efficiency of gear (how many fish are caught relative to population size)




Stock specific issues (Table 10)

Key Issues	Stocks
Recruitment variability, especially irregular large recruitment events	Sablefish, rockfish, pollock,
Increased natural mortality: elevated or event driven	GOA cod, AI cod, EBS snow crab
Prolonged recruitment failure	Greenland turbot
Environmental sensitivity of year-class strength	BSAI/GOA Pollock, Bristol Bay red king crab
Long-lived, slow recovery, declining productivity despite large biomass levels	BSAI/GOA Pacific ocean perch



Modified HCRS (Table 9, Fig 11)

HCR	GOAL
Status Quo (HCR #1)	Automatic rebuilding below target stock size (Status Quo baseline sloping CR used for BSAI and GOA Groundfish)
Lower fishing mortality at high stock size (HCR #5)	Maximize ecosystem biomass and SSB (intent to increase reserves, buffer against environmental shocks, and enhance long-term sustainability with lower fishing mortality)
Lower fishing mortality at high stock size (HCR #10)	Similar objectives as with HCR 5 (intent to provide increased buffering against environmental shocks through proportional reductions in fishing mortality)
Dynamic HCRs that respond to time-varying stock productivity (HCR #7)	Transition from qualitative risk tables to an explicit analytical approach in which recommended harvest rates respond to estimated temporal changes in stock productivity (obtained from population models with time-varying parameters, linkages with environmental covariates or other methods)



Stock specific issues (Table 10)

Key Issues	Stocks	Potential HCR approaches
Recruitment variability, especially irregular large recruitment events	Sablefish, rockfish, pollock,	HCR5 or 10, sablefish HCRs under MSE
Increased natural mortality: elevated or event driven	GOA cod, AI cod, EBS snow crab	HCR with environmental covariates (HCR7), HCR5 or 10?
Prolonged recruitment failure	Greenland turbot	HCR7
Environmental sensitivity of year-class strength	BSAI/GOA Pollock, Bristol Bay red king crab	HCR with environmental covariates (HCR7), HCR5 or 10?
Long-lived, slow recovery, declining productivity despite large biomass levels	BSAI/GOA Pacific ocean perch	Dynamic HCRs that adjusts reference points to reflect trends in stock productivity (variant of HCR 7)



Goals and Objectives (pg 28-29)

Overarching goal (draft)

- Increased resilience for long-term sustainability of fish and crab stocks

Objectives (draft)

- Sustainable biomass for groundfish and crab stocks
- Prevent overfishing
- Transition from use of risk tables to analytical approach (species whose productivity known to vary with environmental conditions)
- Increased buffering against environmental shocks
- Socio-economic sustainability
- Community-level objectives (to be defined)
- Transparency in the ABC –setting process 

Once objectives defined – performance metrics developed (e.g. appx 3)

Key decision points

- Priority management objectives identified
- Objectives stated in clear and specific manner
- Time frame over which performance metrics are calculated


Table A3.1
page 46

Category	Objective	Quantity of interest	Case study metrics
Status	Maintain biomass above biomass limit/ Minimize the chance of being overfished	Biomass (B) and B_{limit}	$P(B > 0.4B_{MSY}) \geq 0.95^a$ $P(B > 0.25B_0)^b$ $P(RSB < 20\%) < 0.05^c$ $P(SSB < 7.7\%SSB_0) < 0.10^d$
	Maintain spawning stock size above target reference points	Biomass and B_{target}	$P(B_{2052} > B_{target}) = 0.5^a$ $P(B > 0.4B_0)^b$ $P(RSB < 36\%) < 0.50$
	Maintain stock within the green zone of Kobe plot	Biomass and B_{limit} and Fishing mortality (F) and F_{limit}	$P(SSB \geq SSB_{MSY} \& F < F_{MSY}) \geq 0.6^d$
Yield	Maintain average/ median/minimum catch	Catch	$P(\text{Catch} > \min(\text{Catch})) > 0.5^b$ Average catch last five projection years ^b Average catch by fishery for various time frames ^d
	Maximize catch	Catch	Total catch $P(\text{Catch} > \min \text{ viable catch})^a$
	Minimize catch variability over time	Variation in catch	Median across replicates of average absolute annual change in landed catch ^a $P(AAV \leq 0.15)^b$
		Variation in output or input control (e.g., ABC or FABC)	$P(\text{Change in TCEY} > 15\%)$
Stability	Maintain age structure or size structure	Age Size (length or weight)	Change in mean age or size
Societal	Maximize long term profitability	Size (length or weight) Revenue	Mean size of directed fishery ^b Mean revenue (by fishery or region) ^e
	Maintain employment	Regional employment (number of jobs)	Mean percent change in the number of jobs ^e
	Maintain or minimize change in nutritional value	Size at age and nutrition metrics	Change in nutritional value of overall fishery ^e

Example performance metrics from sablefish MSE (appx 4; 4.7.1 pgs 52-59)

5 HCRs, 6 performance metrics shown in radar plot (Figure A4.4)

Harvest Control Rule
■ F40 ■ F50 ■ F40 5% Stability Constraint ■ F40 15k Harvest Cap ■ F40 Hybrid



Decision points/Considerations for the Committee

Defining goals and objectives

- Ideally defined overarching goal and associated objective to help Council draft a Purpose and Need statement for this analysis moving forward
- Objectives might be competing, analysis of performance against these objectives for different HCRs will indicate the degree to which different HCRs address objectives

Engagement and Communication

- Feedback on a communication and engagement strategy moving forward
- Complex concepts and analyses may benefit from a targeted communication strategy. This may include:
 - Additional primer materials/presentations on status quo considerations of risk and uncertainty in harvest specifications
 - Infographics on process and simulation testing/MSE of HCRs under development
 - Development of performance metrics and social indicators





Thank you!

Council HCR webpage: <https://www.npfmc.org/hcr-project-overview/>

