


An Overview of Stage 1 (2025-2026) Alternative HCR Evaluations Through ACLIM and GOACLIM



Prepared by:

K Holsman, M. Bryan, C. McGilliard, A. Rovellini, A. Hollowed and D. Stram

Contents

Overview.....	2
Introduction.....	3
Background on NPFMC HCRs.....	3
Considerations for HCRs under changing conditions.....	3
Stage 1 alternative HCRs.....	5
Table 1: Overview of ACLIM and GOACLIM 2025 HCR options.....	6
Figure 2. Subset of stage 1 HCRs for evaluation by ACLIM3 and GOACLIM2 during 2025-2026. Slide courtesy of K. Holsman.....	7
Description of stage 1 HCRs (1, 5, 7 and 10).....	7
Figure 1. Harvest Control Rule (HCR) Explorer available at https://kholsman.shinyapps.io/HCRshiny/	8
HCR 1: Status Quo.....	8
Simulation goal.....	8
Overview HCR 1: Status quo.....	8
Details HCR 1: Status quo.....	9
Figure 3. HCR 1: Status quo. This is the Tier 3 Harvest Control Rule, including the cutoff for certain species.....	9
HCR 5 & 10: Cap-like benefits / Maximize productivity/ increased reserve (buffer shocks)....	9
Simulation goal.....	9
General overview.....	10
Overview: HCR 5: Maximize productivity/ increased reserve (buffer shocks).....	10
Figure 4. Effective F given cap effects on pollock, Pacific cod, and arrowtooth flounder in the EBS. Figure from Holsman et al. 2020.....	11

Figure 5. Cap-like HCRs 10 (top) and 5 (bottom). Below or at Btarget this HCR is the same as the Status Quo (HCR1) Tier 3 Harvest Control Rule, including the cutoff for certain species at low SSB. Above Btarget catch is stabilized (F is reduced) with (HCR10) or without (HCR5) biomass offsets.....	12
Details: HCR 5.....	12
Figure 6: HCR 5: climate sensitivity reserve (buffer shocks). Cap-like HCRs 10 (top) and 5 (bottom): Below or at Btarget this HCR is the same as the Status Quo (HCR1) Tier 3 Harvest Control Rule, including the cutoff for certain species at low SSB. Above Btarget catch is stabilized (F is reduced) with optional biomass offsets.....	13
Details: HCR 10.....	13
Figure 7: HCR 10: gamma offset on inverse SSB decay rate.....	14
HCR 7: Risk Table Bridging: R/S variability covariate adjusted HCR.....	14
Simulation goal.....	14
Details HCR 7.....	15
Figure 8. HCR 7:Recruit per spawner biomass variability adjusted HCR.....	16
Alternative CAP Evaluations.....	16
Performance metrics.....	16
Figure 9: Potential performance metrics for HCR simulation evaluation.....	17
Appendix A: ACLIM and GOACLIM model overview.....	17
Table 2: Overview of CLIM models presently available for HCR and CAP evaluations in Stage 1 evaluations.....	18
References.....	19

Overview

This document serves as an white paper outlining the current set of HCRs under consideration for simulation testing by the ACLIM and GOACLIM teams. This document is informational only and HCRs outlined here are not "recommended HCRs", rather alternative formulations being evaluated for increased performance through coordinated modeling in 2025-2026. Four HCRs were noted in June 2025 by the SSC and Council as recommendations for future discussion but this does not preclude development of additional HCRs and/or any of the HCRs previously developed from consideration in the future. Rather this document focuses on contrasting the objectives and formulation of these HCRs to help the Council in development of overarching goals and objectives of moving to alternative HCRs in the future. Any evaluation of HCRs to be implemented for management purposes would be done by the NPFMC through the established Council process. The HCRs below build on previous modeling efforts. During ACLIM phase 2 (2019-2022), modelers evaluated a suite of Harvest Control Scenarios (1-5; Hollowed et al. 2022), in 2025 during phase 3 of the ACLIM project and in collaboration with GOACLIM phase 2 CLIM modelers added additional HCRs for evaluation. Below is more information on a subset of those standardized harvest control rules and the equations used to derive the curves. This subset of HCRs considered in this document will be referred to as "Stage 1 HCR" throughout

(Fig. 1). An interactive version of the full set of HCRs under consideration is available on line at <https://kholsman.shinyapps.io/HCRshiny>.

Introduction

Background on NPFMC HCRs

The North Pacific Fishery Management Council (NPFMC) has adopted a complex suite of harvest controls that, in aggregate, strive to achieve the goals and objectives of an Ecosystem Approach to Fisheries Management (EAFM). With the adoption of the Aleutian Islands Fisheries Ecosystem Plan ([AIFEP](#)) and the Bering Sea Fisheries Ecosystem Plan ([BSFEP](#)), the NPFMC is striving to adopt an Ecosystem Based Fisheries Management (EBFM) approach. These management systems include forage fish protections, prohibited species mortality caps, discard prohibitions, habitat protections, marine mammal and seabird provisions, rationalized fisheries, community development quotas, optimal yield system-level caps, and sustainable harvest control rules (HCRs) that encourage rebuilding when the reproductive potential of stocks (e.g., spawning biomass or proxies thereof) falls below target levels. The system limits annual catch levels to protect non-target groundfish and crab stocks incidentally captured in target fisheries. Collectively these provisions are designed to address National Standard 1 of the Magnuson Stevens Fishery Conservation and Management Act (MSA). Many other regulations address the additional National Standards for successful operation of a rationalized fishing system. These regional management controls are outlined in the Fishery Management Plan for Groundfish of the Bering Sea-Aleutian Islands ([BSAIGF FMP](#)), the Fishery Management Plan for the Groundfish Fisheries of the Gulf of Alaska ([GOAGF FMP](#)) and the Fishery Management Plan for Bering Sea-Aleutian Islands King and Tanner Crabs ([BSAI Crab FMP](#)).

While many of these management provisions have been amended several times, the regulations governing the setting of HCRs outlined in the FMPs have been in place for many years. In 1999, the NPFMC revised their frameworks for setting HCRs for groundfish in the Bering Sea-Aleutian Islands ([BSAI FMP Amendment 56](#)) and Gulf of Alaska ([GOA FMP Amendment 56](#)). These groundfish HCRs were reviewed by an independent panel in 2002 (Goodman et al., 2002). The NPFMC revised their HCRs for managing Bering Sea- Aleutian Islands crab ([BSAI Crab FMP Amendment 24](#)) in 2008. In 2011, the crab HCRs were revised to comply with revisions to the Magnuson Stevens Fishery Conservation and Management Act (in short, the Magnuson Stevens Act, MSA) that required setting Annual Catch Limits (ACLs, [BSAI Crab FMP Amendment 38](#)). The HCR frameworks comply with the MSA, National Guidelines for implementation of the MSA (National Standards) and guidelines for use of the Best Scientific Information Available ([BSIA](#)).

Considerations for HCRs under changing conditions

At its core, the current HCRs acknowledge that when viewed over long time-frames, most animal populations are governed by logistic population dynamics with associated concepts of maximum sustainable yield (MSY). Based on these concepts, scientists can identify biomass

targets, limits, fishing mortality, and biomass reference points for sustainable management. These reference points are set by the NPFMC Scientific and Statistical Committee (SSC) based on the [BSIA](#) regarding the reproductive potential of the stock and the associated fishing mortality target and limit reference points to achieve these biomass reference points. Due to knowledge gaps and data availability, uncertainty exists for many stocks regarding the relationship between spawning stock size and recruitment. To address this issue, the groundfish and crab HCR frameworks established management tiers based on the [BSIA](#) for a given stock. When information is insufficient to reliably define spawner recruit relationships, methods for calculating proxies for biological reference points are provided in the FMP. More recently, analysts have developed methods for estimating biological reference points under multiple axes of uncertainty (Szuwalski and Punt, 2025).

While the frameworks governing the HCRs for groundfish and crab have been static for many years, the calculation of the biological reference points has changed frequently in response to:

- new information on the key vital rates governing population dynamics of the stock
- shifts in BSIA and management tiers
- shifts in spatial distribution of the stock that influence availability and/or catchability
- changes in the selectivity and catchability of the stocks to fishing resulting from gear changes, time-area-gear regulations that influence selectivity, and other factors governing harvest practices of the fleet (including changes in prohibited species mortality caps and discard mortality restrictions).

The NPFMC's HCR framework proved flexible enough to accommodate these changes.

The NPFMC's current system of management has sustained fishing opportunities for many stocks, however, in recent decades shifts in ocean conditions have had profound impacts on some groundfish and crab with associated impacts on the fisheries and communities that depend on them (Bigman et al., 2023; Suryan et al., 2021). These impacts are harbingers of future changes in fish and crab abundance under projected states of nature under future environmental scenarios.

To address observed environmental changes in the Bering Sea-Aleutian Islands and Gulf of Alaska, the Plan Teams and SSC currently review the status of the ecosystem (Ecosystem Status Reports, ESRs) and current knowledge of the responses of stocks to ecosystem change (Ecosystem Socio-economic Profiles, ESPs) to make qualitative adjustments to annual harvest recommendations (ABCs) based on professional judgement using Risk Tables (Dorn and Zador, 2020). While the Risk Table approach is useful for some stocks, the qualitative nature of the application of the risk table approach underutilizes the mechanistic understanding currently available to analysts for key species (e.g., sablefish, pollock, cod, yellowfin sole, northern rock sole, arrowtooth flounder, red king crab, snow crab and Tanner crab).

For key species, the long legacy of monitoring and mechanistic research has enabled scientists to better understand the complex factors governing the productivity and distribution of groundfish and crab and the responses of fisheries to changes in the distribution and abundance of these species (Rogers et al., 2020; Bigman et al., 2023; Thalmann et al. 2024;

West et al., 2025). In addition, advancements in understanding the physical and bio-geo-chemical processes governing ocean conditions in the Gulf of Alaska and the Bering Sea allow hindcasts, forecasts and projections of ocean properties. These coupled high-resolution regional ocean models replicate observed ecosystem phenomena in the region with considerable skill (Hermann et al., 2019, 2020; Pilcher et al., 2022; Cheng et al., 2021; Kearney et al., 2020; Sullaway et al., 2024). When viewed together, these advances in scientific understanding and modeling enable scientists to construct mechanistically informed social and ecological models informed by output from next generation high-resolution regional ocean models to forecast and project future population trends, including fishing effects (Holsman et al., 2020; Punt et al. 2021; Hollowed et al. 2024; Wildermuth et al. 2024; Blamey et al. 2025; Champagnat et al. In Review). These environmentally-coupled socio-ecological models provide powerful platforms for exploring the performance of current and alternative harvest strategies relative to suites of desired outcomes (for example performance measures see Appendix A).

Stage 1 alternative HCRs

The SSC convened a [public workshop in June 2025](#) to review the frameworks for setting HCRs in the Fishery Management Plans given recent observations of environmental change on managed species and the availability of environmentally-coupled socio-ecological models for key species. The overarching goal of the workshop was to develop a scope of work and Terms of Reference for a workgroup (or several coordinated teams) to prepare a discussion paper of proposed HCR adjustments for consideration/recommendations for the Plan Teams/SSC/AP/Council. The workshop discussed and solicited input on adjustments to current Groundfish and Crab fishery management plan (FMP) harvest specification Tier-levels and resulting harvest control rules (HCRs) for determination of the overfishing level (OFL) and acceptable biological catch (ABC).

The June 2025 workshop was designed to address the Council's requests from October and December 2024. The Council's [October 2024 motion](#) encouraged researchers to 'Consider to what extent, and whether, to revise groundfish and crab harvest control rules (HCRs) to be more climate-resilient.' In [December 2024](#), the Council identified consideration of Tier-systems, climate-informed biomass targets and climate-robust or forecast-informed harvest control rules as part of their Climate Workplan. The SSC workshop was a starting point to frame how to approach and prioritize HCR adjustment opportunities.

The materials presented during the June 2025 workshop are available on the [HCR Workshop](#) website. Briefly, panelists and speakers provided an overview of current harvest control rules (HCRs) for crab and groundfish and introduced 10 alternative approaches (for details see [HCR Shiny APP](#)) as well as a suite of potential performance measures that could be considered by the Council. Upon review, CLIM modelers, the SSC and Council found the most promising candidates for future exploration were HCRs 1, 5, 7, and 10. The Council requested that the work-team should present a summary of the objectives for each HCR and draft projections under current and alternative environmental scenarios for suites of groundfish and crab species to the Plan Teams during the fall specifications cycle for comment and recommendations. Plan

Team and additional SSC recommendations will be reviewed by the Council. This document is intended to satisfy this request.

Table 1: Overview of ACLIM and GOACLIM 2025 HCR options. “Stage 2” denotes the HCRs that are not being evaluated as part of the Council’s request following the June SSC workshop, but have been or are being evaluated as part of the ACLIM and GOACLIM work.

HCR	Name	Detail	HCR Stage
HCR 1	Status quo	This HCR is the baseline sloping harvest control rule used for groundfish in Alaska	1
HCR 5	Maximize productivity/increased reserve (buffer shocks)	HCR 5 is designed to maximize ecosystem and spawning biomass productivity by increasing reserves, creating a buffer against environmental shocks and enhancing long-term sustainability.	1
HCR 7	Risk Table Bridging, R/S variability covariate adjusted HCR	This HCR provides a way to transition from qualitative risk tables to a more explicit, analytical approach for species whose productivity is known to vary with environmental conditions.	1
HCR 10	Maximize productivity/increased reserve, linear version (1/B_target) with offset	This HCR builds on HCR 5 by applying a proportional reduction in fishing mortality based on biomass levels, further enhancing stock and environmental productivity through strengthening the buffer against environmental shocks.	1
(Stage 2 HCRs Below)			
HCR 2	Lagged recovery to estimate emergency relief financing needs	Simulations with this HCR will mimic economic-driven fishery closures and delayed recovery in order to estimate emergency relief needs.	2
HCR 3	Long-term resilience (stronger reserve) B_target	This HCR aims to enhance long-term stock resilience by adjusting B_target (as a proportion of unfished biomass)	2
HCR 4	Environmental index informed sloping rate, e.g., MHW category alpha	Simulations with this HCR will assess whether adjusting harvest intensity based on poor forecasted conditions—such as marine heatwaves—can accelerate stock recovery following climate or environmental disturbances.	2
HCR 6	Combination of MHW (HCR4) + Maximize productivity (HCR5)	This HCR combines the approaches of HCR 4 and HCR 5 to address both immediate and long-term environmental impacts.	2
HCR 8	Adjust effective spawning biomass (simulate adjusted	This HCR adjusts the effective spawning biomass instead of the target biomass threshold, serving as a	2

	B_target)	sensitivity approach to explore variability in spawning stock biomass (SSB) estimates within a given assessment year or to evaluate alternative B_target values.	
HCR 9	Forecast informed version of HCR 5	This HCR builds on HCR 5 by using environmental forecasts to dynamically adjust reserves, enhancing ecosystem productivity and resilience to environmental shocks.	2



Figure 1. Subset of stage 1 HCRs for evaluation by ACLIM3 and GOACLIM2 during 2025-2026. Slide courtesy of K. Holsman.

Description of stage 1 HCRs (1, 5, 7 and 10)

Below we describe a general overview of the subset of environmentally linked HCRs that are identified for the first stage of ACLIM and GOACLIM coordinated evaluations in 2025/2026. A more detailed description (including the analytical formulation) of each HCR can be found on the [HCR Shiny app](#) (Figure 1) under the ‘detailed information’ tab (see options on landing page in upper right corner).




Figure 2. Harvest Control Rule (HCR) Explorer available at <https://kholsman.shinyapps.io/HCRshiny/>

HCR 1: Status Quo

Simulation goal

This HCR is the baseline (status quo) sloping harvest control rule used for groundfish in Alaska.

Overview HCR 1: Status quo

HCR 1 (Fig. 2 upper left; Fig. 3) is the baseline sloping harvest control rule evaluated in ACLIM phases 1 and 2 (Holsman et al. 2020, Reum et al. 2020, Whitehouse et al. 2020, Punt et al. 2024) and continues to be evaluated during ACLIM3 and GOACLIM2 simulations. It is the Tier 3 HCR currently used by the NPFMC to manage Alaska groundfish stocks, where F_{ABC} is reduced when current biomass falls below target biomass ($B_{40\%}$). There is a cutoff at $B_{20\%}$ for Alaska pollock, Atka mackerel, and Pacific cod where fishing falls to zero to prevent overfishing of these important Steller sea lion prey species. Including the baseline HCR allows us to determine its performance under future EBM, environmental, and climate scenarios.

Details HCR 1: Status quo

This is the basic sloping harvest control rule for groundfish in the EBS (Eq. 1). There is a $B_{20\%}$ cut-off for SSL (Atka, pollock, P. cod). $F_{ABC_{mxc}}$ is the HCR adjusted F rate that corresponds to ABC. The Tier 3 approach is to set the slope of the sloping HCR to $\alpha = 0.05$ and $B_{lim} = 0$ and $B_{target} = B_{40\%}$ or $B_{target} = 0.4$ (i.e., 40% of unfished biomass B_0 , as an MSY proxy) for most species except $B_{lim} = 0.2 B_0 (B_{20\%})$ for pollock, Pacific cod and Atka mackerel.

Equation 1. HCR 1: Status quo

$$F_{ABC_{mxc}} = \begin{cases} F_{ABC} & \frac{B_y}{B_{target}} > 1 \\ F_{ABC}((\frac{B_y}{B_{target}} - \alpha)/(1 - \alpha)) & \frac{B_{lim}}{B_{target}} \leq \frac{B_y}{B_{target}} < 1 \\ 0 & \frac{B_y}{B_{target}} < \frac{B_{lim}}{B_{target}} \end{cases}$$




Figure 3. HCR 1: Status quo. This is the Tier 3 Harvest Control Rule, including the cutoff for certain species

HCR 5 & 10: Cap-like benefits / Maximize productivity / increased biomass reserve (buffer environmental shocks)

Simulation goal

HCR 5 is designed to maximize ecosystem and spawning biomass productivity by increasing biomass reserves, creating a buffer against environmental shocks and enhancing long-term sustainability. These HCRs aim to enhance stock robustness to uncertainty in environmental

conditions and provide increased potential for higher recruitment levels and growth of fish to larger sizes through preserving higher levels of SSB when the population is abundant.

HCR 10 builds on HCR 5 by applying a proportional reduction in fishing mortality based on biomass levels. It also includes an optional SSB offset, ie. $B_{target+offset}$ (e.g., start slope at $B_{50\%}$).

General overview

HCR 5 and HCR 10 (Fig. 1 bottom left and bottom right) have the same shape as HCR 1 when $SSB \leq B_{target}$, but allow for a reduction in F_{target} when biomass is above B_{target} . These HCRs are inspired by the effective shape of the realized BSAI pollock harvest rate (F) under current management with the 2 million Ton annual cap on Bering sea groundfish harvest (Holsman et al. 2020, Ianelli et al. 2011). Despite strong environmental effects on stock productivity, BSAI pollock catch was relatively stable through recent marine heatwaves, perhaps due to the combined effect of the HCR and cap on catch above $B_{40\%}$ (i.e., population age structure reserve that helps stabilize ecosystem and stock productivity during unfavorable climate conditions). To test this hypothesis, CLIM modelers aim to test whether using an HCR that follows the shape of the effective pollock F could help impart a similar benefit to other stocks. Again, modelers are testing the hypothesis that by stabilizing catch above some point like B_{target} (left bottom, ski slope shape) or B_{target} with a biomass offset (right; HCR 10; plateau + slope-like shape) the HCR might:


- a) increase the age class diversity
- b) promote a higher productivity state in the fished biomass
- c) promote more larger fish
- d) increase ecosystem productivity and therefore fishery productivity

Performance of these HCRs relative to HCR 1 will be evaluated on fishery and management performance metrics described below. Initial results are promising with some indication that HCR 5 and HCR 10 result in an increase of spawning biomass, increased stability in annual catch, more diversity in the spawning biomass age structure, and prevention or forestalling of stock declines/crashes under variable environmental conditions. These are preliminary findings that may change as we expand the suite of models and species being evaluated with HCR 5 and 10. The simulation goals for HCRs 5 and 10 are to better understand the combined influence of the sloping HCR and the ecosystem cap in total catches, as well as evaluating whether buffering fishing mortality by maintaining larger stock sizes has additional benefits to the population against environmental shocks.

Overview: HCR 5: Maximize productivity / increased biomass reserve (buffer environmental shocks)

HCR 5 recreates the realized pollock cap effect on F_{target} when a stock is over B_{target} , which may maximize long-term ecosystem and stock productivity. The steepness of the cap effect

could be varied based on vulnerability analyses (e.g., Spencer (2019); Hare (2016)) (or approximated via MSE), with species that are known to be more sensitive to environmental variability, such as Pacific cod (Barbeaux et al. 2022), needing more reserve in the “bank”. Pollock is an example of HCR 5 in practice (via effects of the 2MT cap + sloping HCR; Holsman (2020)).



Supplementary Figure 3. Effective harvest rate F_y under the no cap and 2 MT cap scenarios. Ratio of effective F_y to F_{target} as a function of the ratio of biomass to unfished biomass reference points (B_y/B_0) for each pollock (left), Pacific cod (middle), and arrowtooth flounder (right). Scenarios without the cap (orange; follow the ABC harvest control rule exactly) and scenarios with the 2 MT cap (scatter, teal).

Figure 4. Effective F given cap effects on pollock, Pacific cod, and arrowtooth flounder in the EBS. Figure from Holsman et al. 2020.




Figure 5. Cap-like HCRs 10 (top) and 5 (bottom). Below or at B_{target} this HCR is the same as the Status Quo (HCR1) Tier 3 Harvest Control Rule, including the cutoff for certain species at low SSB. Above B_{target} catch is stabilized (F is reduced) with (HCR10) or without (HCR5) biomass offsets.

Details: HCR 5

As in HCR 1 with the 20% cutoff for SSL (Eq. 1), set the target to 40% of B_0 ($\alpha = 0.05$, $B_{lim} = B_{20\%}$, $B_{target} = B_{40\%}$). After $B_{40\%}$ have a slowly sloping F proportional to climate sensitivity to mimic realized F rates of pollock under the 2 MT cap. This approach is designed to maximize ecosystem productivity and build a reserve biomass for environmental or climate shocks (sensu Holsman (2020)). In this (Eq. 2) the climate sensitivity buffer γ is a value > 0 that controls the F decay rate (increase in reserve biomass) above B_{target} , i.e., $B_{40\%}$:

Equation 2: HCR 5

$$F_{ABC_{max}} = \begin{cases} F_{ABC} e^{(-\gamma(\frac{B_y}{B_{target}} - 1))} & \frac{B_y}{B_{target}} > 1 \\ F_{ABC}((\frac{B_y}{B_{target}} - \alpha)/(1 - \alpha)) & \frac{B_{lim}}{B_{target}} \leq \frac{B_y}{B_{target}} < 1 \\ 0 & \frac{B_y}{B_{target}} < \frac{B_{lim}}{B_{target}} \end{cases}$$

where, $0 \leq \gamma \leq 1$.

Example: Set the biological reference points as in HCR 1, i.e., 40% of B_0 , ($\alpha = 0.05$, $B_{lim} = B_{20\%}$, $B_{target} = B_{40\%}$). Above B_{target} apply the gamma / 2MT cap effects. Shown in Fig. 6, for low sensitivity stocks set $\gamma = 0.1$, for highly sensitive stocks set $\gamma = 0.2$. We're testing whether this would result in more stable biomass levels and catches through environmental shocks.




Figure 6: HCR 5: climate sensitivity reserve (buffer shocks). Below or at B_{target} this HCR is the same as the Status Quo (HCR1) Tier 3 Harvest Control Rule, including the cutoff for certain species at low SSB. Above B_{target} catch is stabilized (F is reduced).

Details: HCR 10

As in HCR 5 (Eq.2), HCR 10 aims to recreate the benefits of the cap effect on pollock via implementing and HCR that follows the effective F rates above B_{target} . The steepness of that cap effect here is inverse to SSB, with an optional B_{target} offset (γ , i.e. the point at which F reduction starts), more sensitive species might need a lower gamma (γ) to drive a larger age class reserve resulting in increased abundance of large spawners and benefits to recruitment and stock productivity sooner. BSAI Pollock are an example of the HCR 10 in practice (via effects of the 2MT cap + sloping HCR).

Equation 3: HCR 10

$$F_{ABC_{max}} = \begin{cases} F_{ABC}/\left(\frac{B_y}{B_{target}} \frac{1}{(1+\gamma)}\right) & \frac{B_y}{B_{target}} > (1+\gamma) \\ F_{ABC} & 1 < \frac{B_y}{B_{target}} < (1+\gamma) \\ F_{ABC}((\frac{B_y}{B_{target}} - \alpha)/(1-\alpha)) & \frac{B_{lim}}{B_{target}} \leq \frac{B_y}{B_{target}} < 1 \\ 0 & \frac{B_y}{B_{target}} < \frac{B_{lim}}{B_{target}} \end{cases}$$

where, $0 \leq \gamma \leq 1$.

Example: As in HCR1 (Eq. 1), set the target to 40% of B_0 , ($\alpha = 0.05$, $B_{lim} = B_{20\%}$ for amendment 80 species, $B_{target} = B_{40\%}$). Shown in Fig. 7, “small offset” where $\gamma = 0.5$, “medium offset” $\gamma = 1.5$, and “large offset” where $\gamma = 3$. We’re testing whether this would result in more stable biomass levels and catches through upcoming environmental shocks.




Figure 7: HCR 10: gamma offset on inverse SSB decay rate.

HCR 7: Risk Table Bridging: R/S variability covariate adjusted HCR

Simulation goal

This HCR provides a way to transition from qualitative risk tables to a more explicit, analytical approach for species whose productivity is known to vary with environmental conditions.

HCR 7 is a dynamic HCR that allows for adjustments in the F_{target} and B_{target} with changes in population productivity and is based on recent work of Dr. Paul Spencer on Tier 1 HCRs where he found that the realized F varied with sea surface temperature (SST) in the EBS. We

borrowed from that to adjust either F or B_{target} or a combination of both as a function of an index of annual bottom temperature. The goal of this HCR is to evaluate a HCR that allows for a transition from qualitative risk tables to a quantitative approach for species with known productivity variation due to environmental conditions. Currently, HCR 7 is parameterized such that productivity varies with an environmental covariate (Eq. 4). Future work will modify HCR 7 to be informed by retrospective analyses of stock-recruitment relationships.

HCR 7 is of particular interest to CLIM modelers. Preliminary evaluations reveal good performance; catch looks similar to Status quo (HCR 1) but SSB is higher with fewer catch or SSB crashes under high environmental variability (Holsman et al. in prep).

Details HCR 7

The HCR reference points for HCR 7 are adjusted based on variability in spawner-recruitment relationships using a covariate X_y (e.g., SST) such that:

Equation 4: HCR 7

$$F_{ABC_{max}} = \begin{cases} F_{ABC} e^{(\omega_1 * x_y)} & \frac{B_y}{\hat{B}_{target}} > 1 \\ F_{ABC} \left(\left(\frac{B_y}{\hat{B}_{target}} - \alpha \right) / (1 - \alpha) \right) e^{(\omega_1 * x_y)} & \frac{B_y}{\hat{B}_{lim}} \leq \frac{B_y}{\hat{B}_{target}} < 1 \\ 0 & \frac{B_y}{\hat{B}_{target}} < \frac{B_y}{\hat{B}_{lim}} \end{cases}$$

such that F_{ABC} is adjusted by covariate x_y according the parameter ω_1 and B_{target} and B_{lim} are adjusted based on the parameters ω_2 and ω_3 such that:

$$\hat{B}_{target} = B_{target} e^{(-\omega_2 * x_y)}$$

and

$$\hat{B}_{lim} = B_{lim} e^{(-\omega_3 * x_y)}$$

and ω_1, ω_2 and $\omega_3 \geq 0$.

For the ACLIM HCR 7 simulations ω_1 and ω_2 will eventually be fit using retrospective analyses of spawner-recruitment relationships across scaled (z-scored) SST (x_y) on EBS pollock by Spencer et al. in prep.

Example: Set the target to 40% of B_0 ($\alpha = 0.05, B_{lim} = B_{20\%}, B_{target} = B_{40\%}$). In the example below (x_y) = 2.4 and for case 7a all ω values were set to 0.0, in case 7b $\omega_1 = \omega_2 = \omega_3 = 0.1$, and in case 7c $\omega_1 = -0.1, \omega_2 = -0.1, \omega_3 = -0.1$.




Figure 8. HCR 7:Recruit per spawner biomass variability adjusted HCR.

Alternative ecosystem cap evaluations

As part of GOACLIM we investigated the effectiveness of the 800,000 mt Optimum Yield (OY) cap on annual groundfish catches in the Gulf of Alaska (GOA). Using the Atlantis end-to-end ecosystem model that simulates species interactions and ecosystem processes, we found that projected total GOA groundfish catches does not approach the current OY cap across a range of fishing scenarios and projected environmental conditions (Rovellini et al. 2025). Notably, underexploitation of arrowtooth flounder led to increased predation pressure on pollock and other groundfish, resulting in foregone catches and demonstrating how species interactions can influence total yield even when caps are not binding. These findings suggest that the current 800,000 mt cap is unlikely to constrain future GOA groundfish catches under predator-prey dynamics and projected environmental conditions. Building on this work, we are now developing new functionalities within Atlantis to simulate the implementation of alternative OY cap values, including a mechanistic method to preferentially rescale fishing mortality across stocks under constraining caps. We are currently conducting simulations that use these new functionalities, together with the status-quo HCR, to apply a range of OY caps to the GOA groundfish fisheries under various simulated environmental conditions. The purpose of this work is to evaluate ecosystem and fishery responses to alternative OY cap values, and to alternative fishing rescaling schemes when the cap is constraining. In parallel, OY estimation and simulation tools are being developed in the Rpath/Ecopath with Ecosim (EwE) food web modeling framework for both the GOA and the Bering Sea.

Performance metrics

The ACLIM and GOACLIM teams are in the process of developing a full set of performance metrics for simulation evaluations. We welcome feedback on additional performance metrics that may be of interest.

Performance criteria

Hollowed et al.
Alaska Integrated Climate Modeling

TABLE 6 | Suite of candidate performance indicators for ACLIM.

Core species abundance	Mean and variance for time block	Sustainable fishing index
Core species recruitment	Mean and variance for time block	Sustainable fishing index
Core species average size and age at maturity	Mean and variance for time block	Sustainable fishing index
Core species exploitation	Annual time trend F/F_{MSY}	Sustainable fishing index
Core species crab status	Annual trend reproductive potential vs. target reproductive potential	Sustainable fishing index
Core species crab catch	Mean and variance for time block	Sustainable fishing index
Centroid of distribution for core species	Annual time trend	Index distribution
Euphausiid biomass	Mean and variance for time block	Ecosystem stability index
Motile epifauna biomass	Mean and variance for time block	Trophic structure index
Benthic forager biomass	Mean and variance for time block	Trophic structure index
Pelagic forager biomass	Mean and variance for time block	Trophic structure index
Apex predator biomass	Mean and variance for time block	Trophic structure index
Species diversity index	Alpha and beta diversity indices	Ecosystem stability index
Mean trophic level of the catch	Mean and variance for time block	Ecosystem Based Fishery Management Index
Number of fishery closures by core species	Average for time block	Fishery efficiency index
Core species and fleet CPUE	Annual time trend of CPUE by species and fleet	Fishery catchability index
Fishing effort by fleet	Annual time trend of fishing effort	Fisheries participation and employment
Core species first-wholesale revenue index	Annual time trend	Economic index
Core species percent TAC utilization	Percentage of total allowable catch landed	Management index
Fleet species diversity index	Annual measure of diversity of target species revenues	Measure of fishery portfolio by sector
Fleet revenue variability	Coefficient of variations of fisheries revenue by sector	Financial risk index



Hollowed et al. 2020

Figure 9: Potential performance metrics for HCR simulation evaluation.

Appendix A: ACLIM and GOACLIM model overview

More information can be found about ACLIM and GOACLIM in the following places:

- ACLIM website:
<https://www.fisheries.noaa.gov/alaska/ecosystems/alaska-climate-integrated-modeling-project>
- GOACLIM
website:<https://www.fisheries.noaa.gov/alaska/ecosystems/gulf-alaska-climate-integrated-modeling-project>
- ACLIM overview (Hollowed et al. 2020):
<https://www.fisheries.noaa.gov/alaska/ecosystems/gulf-alaska-climate-integrated-modeling-project>
- GOACLIM overview for the SSC
(2025):<https://meetings.npfmc.org/CommentReview/DownloadFile?p=9db617a9-0b9a-4ab9-af00-de13f2fd91ce.pdf&fileName=GOA%20Climate%20Integrated%20Modeling.pdf>

These resources provide an overview of the multimodel interdisciplinary approach to the Alaska and Gulf of Alaska Integrated Modeling projects as well as more detailed information about each of the modeling platforms available for HCR and CAP evaluations. The table below further summarizes the core models that can be utilized for HCR evaluations (although this table is not

exhaustive, as multiple additional models are being developed and used within the CLIMs for additional purposes).

Table 2: Overview of CLIM models presently available for HCR and CAP evaluations in Stage 1 evaluations.

Model Name	EBS	GOA	Bottom-up dynamics included	Top- down dynamics included	Social/ Econ Included	Model type	Examples
ROMS-NPZ	X	X	X			Oceanographic and lower trophic (nutrient to zooplankton)	Hermann et al. 2016, 2019, 2021; Cheng et al. 2021; Kearney et al. 2020, Pilcher et al. 2019;
CE-Single spp models	X	X				Climate enhanced single species stock assessment models	Ianelli et al. 20XX; Spencer et al. 2019;
CEATTLE	X	X	[optional in rec; growth]	X	X (with ATTACH)	Multispecies stock assessment model	Holsman et al. 2016, 2020; Adams et al. 2022
Atlantis		X	X	X		End-to-end ecosystem model	Rovellini et al. (2024, 2025)
Rpath/EwE	X	X	X	X		Production to Biomass based end-to-end food-web model	Whitehouse et al. 2020;
MIZER	X (in progress)		X	X		Size-spectrum based food-web model	Reum et al. 2019b,c, 2020
SDMs	X	X (in progress)	X			Species distribution models	Goodman et al. 2024; Bigman et al. 2023; Kristiansen et al. in prep
ATTACH	X			X		Social-economic model of TAC and Catch	Faig et al. 2019
Conceptual models /	X	X	X	X	X	Semi-quantitative or qualitative	Reum et al. 2019a; Wise et al.

QNMs						network models	in prep
Food and nutrition security	X (in progress)					Social	Wise et al. in prep

References

Adams, G.D., K.K. Holsman, S.J. Barbeaux, M.W. Dorn, J.N. Ianelli, I. Spies, I. J. Stewart, A. E. Punt (2022). An ensemble approach to understand predation mortality for groundfish in the Gulf of Alaska. *Fisheries Research* vol 251, 106303, <https://doi.org/10.1016/j.fishres.2022.106303>

Barbeaux, S. J., Holsman, K., & Zador, S. (2020). Marine Heatwave Stress Test of Ecosystem-Based Fisheries Management in the Gulf of Alaska Pacific Cod Fishery. *Frontiers in Marine Science*, 7(703).

Bigman, J S, B J Laurel, K Kearney, A J Hermann, W Cheng, K K Holsman, and L A Rogers. "Predicting Pacific Cod Thermal Spawning Habitat in a Changing Climate." *ICES Journal of Marine Science* (2023). <https://doi.org/10.1093/icesjms/fsad096>. <https://doi.org/10.1093/icesjms/fsad096>.

Blamey, L.K., E.E. Plagány, T. Moore. 2025. iExploring dynamic reference points for a short-lived, highly variable species under future climate. *Can. J. Fish. Aquat. Sci.* 82: 1–16 (2025) | [dx.doi.org/10.1139/cjfas-2025-0070](https://doi.org/10.1139/cjfas-2025-0070)

Cheng, Wei, Albert J. Hermann, Anne B. Hollowed, Kirstin K. Holsman, Kelly A. Kearney, Darren J. Pilcher, Charles A. Stock, and Kerim Y. Aydin (2021). "Eastern Bering Sea Shelf Environmental and Lower Trophic Level Responses to Climate Forcing: Results of Dynamical Downscaling from CMIP6." *Deep Sea Research Part II: Topical Studies in Oceanography* 193 (2021). <https://doi.org/10.1016/j.dsr2.2021.104975>.

Champagnat, J., C.C. Monnahan, J.Y. Sullivan, J.T. Thorson, S.K. Shotwell, L. Rodgers, A.E. Punt. In Review. Causal models as a scientific framework for next-generation ecosystem and climate-linked stock assessments. [EcoVio-Rxiv](https://doi.org/10.1101/2023.05.10.535412)

Dorn, Martin W., and Stephani G. Zador. "A Risk Table to Address Concerns External to Stock Assessments When Developing Fisheries Harvest Recommendations." *Ecosystem Health and Sustainability* 6, no. 1 (2020/12/16 2020): 1813634. <https://doi.org/10.1080/20964129.2020.1813634>.

Faig, A., and Haynie, A. C. (2019). *catchfunction: BSAI Catch Function. R Package Version 1.5.3*.

Goodman, M. C., J. C. P. Reum, C. Barnes, A. E. Punt, J. N. Ianelli, E. A. McHuron, Giulio A. De Leo, K. K. Holsman (2024). Climate Covariate Choice and Uncertainty in Projecting Species Range Shifts: A Case Study in the Eastern Bering Sea. *Fish and Fisheries*, 2024; 0:1–21.
<https://doi.org/10.1111/faf.12875>

Hermann, Albert J., Georgina A. Gibson, Wei Cheng, Ivonne Ortiz, Kerim Aydin, Muyin Wang, Anne B. Hollowed, Kirstin K. Holsman, and Shubha Sathyendranath. "Projected Biophysical Conditions of the Bering Sea to 2100 under Multiple Emission Scenarios." *ICES Journal of Marine Science* (2019). <https://doi.org/10.1093/icesjms/fsz043>.

Hermann, Albert J., Kelly Kearney, Wei Cheng, Darren Pilcher, Kerim Aydin, Kirstin K. Holsman, and Anne B. Hollowed (2021). "Coupled Modes of Projected Regional Change in the Bering Sea from a Dynamically Downscaling Model under Cmip6 Forcing." *Deep Sea Research Part II: Topical Studies in Oceanography* 194 (2021/12/01/ 2021): 104974.

<https://doi.org/https://doi.org/10.1016/j.dsr2.2021.104974>.

<https://www.sciencedirect.com/science/article/pii/S0967064521000503>.

Hare, W. E. A. N., Jonathan A. AND Morrison. 2016. [A vulnerability assessment of fish and invertebrates to climate change on the northeast u.s. Continental shelf](#). PLOS ONE 11(2):1–30.

Hollowed, A. B., K. K. Holsman, A. Haynie, A. Hermann, A. Punt, K. Aydin, J. Ianelli, S. Kasperski, W. Cheng, A. Faig, K. Kearney, J. Reum, P. Spencer, I. Spies, W. Stockhausen, C. Szwalski, G. A. Whitehouse, T. Wilderbuer. 2020. Integrated modelling to evaluate climate change impacts on coupled social-ecological systems in Alaska. *Special Issue on Global Change and the Future Ocean, Frontiers in Marine Science*. 6:775.

<https://doi.org/10.3389/fmars.2019.00775>

Hollowed, A.B., K.K. Holsman, S.P. Wise. A.C. Haynie, W. Cheng, D.C.K. Evans, A.J. Herman, J.N. Ianelli, K.A. Kearney, A.E. Punt, J.C.P. Reum, D.L. Stram, C.S. Szwalski. 2024. Development of climate informed management scenarios for fisheries in the eastern Bering Sea. *ICES Journal of Marine Science*, 2024, Vol. 0, Issue 0, 1–14

<https://doi.org/10.1093/icesjms/fsa034>

Holsman, K.K., A.C. Haynie, A.B. Hollowed, J.C.P. Reum, K. Aydin, A.J. Hermann, W. Cheng, A. Faig, J.N. Ianelli, K.A. Kearney and A.E. Punt. 2020. [Ecosystem-based fisheries management forestalls climate-driven collapse](#). *Nature Communications* 11(1).

Kearney, Kelly, Albert Hermann, Wei Cheng, Ivonne Ortiz, and Kerim Aydin. "A Coupled Pelagic-Benthic-Sympagic Biogeochemical Model for Thebering Sea: Documentation and Validation of the Bestnpz Model (V2019.08.23) within a High-Resolution Regional Ocean Model." *Geoscientific Model Development* 13, no. 2 (2020): 597-650.

Pilcher, D. J., J. N. Cross, A. J. Hermann, K. A. Kearney, W. Cheng, and J. T. Mathis. "Dynamically Downscaled Projections of Ocean Acidification for the Bering Sea." *Deep Sea Research Part II: Topical Studies in Oceanography* 198 (2022). <https://doi.org/10.1016/j.dsrr.2022.105055>.

Punt, A. E., Dalton, M. G., Adams, G. D., Barbeaux, S. J., Cheng, W., Hermann, A. J., Holsman, K. K., Hulson, P.-J. F., Hurst, T. P., & Rovellini, A. (2024). Capturing uncertainty when modelling environmental drivers of fish populations, with an illustrative application to Pacific Cod in the eastern Bering Sea. *Fisheries Research*, 272, 106951.

Punt, A.E., M.G. Dalton, W. Cheng, A.J. Hermann, K.K. Holsman, T.P. Hurst, J.N. Ianelli, K.A. Kearney, C.R. McGilliard, D.J. Pilcher, M. Veron. 2021. Evaluating the impact of climate and demographic variation on future prospects for fish stocks: An application for northern rock sole in Alaska. *Deep-Sea Research II* 189-190 (2021) 104951

Reum, J. C. P., P. S. McDonald, W. C. Long, K. K. Holsman, L. Divine, D. Armstrong, and J. Armstrong. 2019. Rapid assessment of management options for promoting stock rebuilding in data-poor species under climate change. *Conservation Biology* 0:1–11.

Reum, J., JL Blanchard, KK Holsman, K Aydin, AE Punt. (2019) Species-specific ontogenetic diet shifts attenuate trophic cascades and lengthen food chains in exploited ecosystems. *Okios* DOI: 10.1111/oik.05630

Reum, J., K. Holsman, KK, Aydin, J. Blanchard, S. Jennings. Energetically relevant predator to prey body mass ratios and their relationship with predator body size. (2019) *Ecology and Evolution* (9):201–211 DOI: 10.1002/ece3.4715

Reum, J. C. P., J. L. Blanchard, K. K. Holsman, K. Aydin, A. B. Hollowed, A. J. Hermann, W. Cheng, A. Faig, A. C. Haynie, and A. E. Punt. (2020). Ensemble Projections of Future Climate Change Impacts on the Eastern Bering Sea Food Web Using a Multispecies Size Spectrum Model. *Frontiers in Marine Science* 7:1–17.

Rogers, Lauren A., Matthew T. Wilson, Janet T. Duffy-Anderson, David G. Kimmel, and Jesse F. Lamb. "Pollock and "the Blob": Impacts of a Marine Heatwave on Walleye Pollock Early Life Stages." *Fisheries oceanography* 30, no. 2 (2021): 142-58. <https://doi.org/10.1111/fog.12508>.

Rovellini, A, AE Punt, MD Bryan, Isaac Kaplan, M Dorn, K Aydin, EA Fulton, B Alglave, MR Baker, G Carroll, B Ferriss, MA Haltuch, A Hayes, A Hermann, P-Y Hernvann, K Holsman, OR Liu, E McHuron, HN Morzaria-Luna, J Moss, S Surma, MT Weise (2024) Linking climate stressors and ecological processes in ecosystem models, with a case study from the Gulf of Alaska. *ICES Journal of Marine Science*, fsae002

Rovellini, A., Punt, A. E., Dorn, M. W., Kaplan, I. C., Bryan, M. D., Adams, G., Aydin, K., Baker, M. R., Barnes, C. L., Ferriss, B. E., Fulton, E. A., Haltuch, M. A., Hermann, A. J., Holsman, K.

K., McGilliard, C. R., McHuron, E. A., Morzaria-Luna, H. N., & Surma, S. (2025). Evaluating ecosystem caps on fishery yield in the context of climate stress and predation. *Ecological Applications*, 35(3), e70036. <https://doi.org/10.1002/eap.70036>

Spencer, A. B. A. S., Paul D. AND Hollowed. 2019. [Trait-based climate vulnerability assessments in data-rich systems: An application to eastern bering sea fish and invertebrate stocks](#). *Global Change Biology* 25(11):3954–3971.

Sullaway, Genoa, Curry J. Cunningham, David Kimmel, Darren J. Pilcher, and James T. Thorson. "Evaluating the Performance of a System Model in Predicting Zooplankton Dynamics: Insights from the Bering Sea Ecosystem." *Fisheries Oceanography* 34, no. 1 (2024): e12691. <https://doi.org/https://doi.org/10.1111/fog.12691>. <https://onlinelibrary.wiley.com/doi/abs/10.1111/fog.12691>.

Suryan, Robert M., Mayumi L. Arimitsu, Heather A. Coletti, Russell R. Hopcroft, Mandy R. Lindeberg, Steven J. Barbeaux, Sonia D. Batten, *et al.* "Ecosystem Response Persists after a Prolonged Marine Heatwave." *Scientific Reports* 11, no. 1 (2021/03/18 2021): 6235. <https://doi.org/10.1038/s41598-021-83818-5>.

Szuwalski, Cody, and André E. Punt. "Spawning-Biomass-Per-Recruit Proxies for Fisheries Reference Points under Multiple Axes of Uncertainty." *Fisheries Research* 289 (2025/09/01/2025): 107494. <https://doi.org/https://doi.org/10.1016/j.fishres.2025.107494>. <https://www.sciencedirect.com/science/article/pii/S0165783625002310>.

Thalmann, Hillary L., Benjamin J. Laurel, L. Zoe Almeida, Kaitlyn E. Osborne, Kaylee Marshall, and Jessica A. Miller. "Marine Heatwaves Alter the Nursery Function of Coastal Habitats for Juvenile Gulf of Alaska Pacific Cod." *Scientific reports* 14, no. 1 (2024): 14018-17. <https://doi.org/10.1038/s41598-024-63897-w>.

West, Catherine F., Loren McClenachan, Steven J. Barbeaux, Ingrid Spies, Jason A. Addison, Bruce T. Anderson, Courtney A. Hofman, *et al.* "Integrating Marine Historical Ecology into Management of Alaska's Pacific Cod Fishery for Climate Readiness." *ICES journal of marine science* 82, no. 4 (2025). <https://doi.org/10.1093/icesjms/fsaf056>.

Wildermuth. R.P., D.T. Tommasi, P. Kuriyama, J. Smith, I. Kaplan. 2024. Evaluating robustness of harvest control rules to climate-driven variability in sardine recruitment. *Can. J. Fish. Aquat. Sci.* 81: 1029–1051 (2024) | [dx.doi.org/10.1139/cjfas-2023-0169](https://doi.org/10.1139/cjfas-2023-0169)

Whitehouse, G. A., K. Y. Aydin, A. B. Hollowed, K. K. Holsman, W Cheng, A. Faig, A. C. Haynie, A. J. Hermann, K. A. Kearney, A. E. Punt, and T. E. Essington (2021). Bottom-up impacts of forecasted climate change on the eastern Bering Sea food web. *Front. Mar. Sci.*, 03 February 2021 | <https://doi.org/10.3389/fmars.2021.624301>

Appendix A. Suite of candidate performance indicators for ACLIM (from Hollowed et al. 2022).

Name	Derivation	Purpose
Core species abundance	Mean and variance for time block	Sustainable fishing index
Core species recruitment	Mean and variance for time block	Sustainable fishing index
Core species average size and age at maturity	Mean and variance for time block	Sustainable fishing index
Core species exploitation	Annual time trend F/FMSY	Sustainable fishing index
Core species stock status	Annual time trend reproductive potential vs. target reproductive potential	Sustainable fishing index
Core species catch	Mean and variance for time block	Sustainable fishing index
Centroid of distribution for core species	Annual time trend	Index distribution
Euphausiid biomass	Mean and variance for time block	Ecosystem stability index
Motile epifauna biomass	Mean and variance for time block	Trophic structure index
Benthic forager biomass	Mean and variance for time block	Trophic structure index
Pelagic forager biomass	Mean and variance for time block	Trophic structure index
Apex predator biomass	Mean and variance for time block	Trophic structure index
Species diversity index	Alpha and beta diversity indices	Ecosystem stability index
Mean trophic level of the catch	Mean and variance for time block	Ecosystem Based Fishery Management index
Number of fishery closures by core species	Average for time block	Fishery efficiency index
Core species and fleet CPUE	Annual time trend of CPUE by species and fleet	Fishery catchability index
Fishing effort by fleet	Annual time trend of fishing effort	Fisheries participation and employment
Core species first-wholesale revenue index	Annual time trend	Economic index
Core species percent TAC utilization	Percentage of total allowable catch landed	Management index
Fleet species diversity index	Annual measure of diversity of target species revenues	Measure of fishery portfolio by sector
Fleet revenue variability	Coefficient of variations of fisheries revenue by sector	Financial risk index