

Norton Sound Red King Crab Stock Assessment for the fishing year 2024

Toshihide Hamazaki¹
Alaska Department of Fish and Game Commercial Fisheries Division
¹333 Raspberry Rd., Anchorage, AK 99518-1565
Phone: 907-267-2158
Email: Toshihide.Hamazaki@alaska.gov

Executive Summary

1. Stock. Red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska.
2. Catches. This stock supports three important fisheries: summer commercial, winter commercial, and winter subsistence. Of those, the summer commercial fishery accounts for 85% of total harvest. The summer commercial fishery started in 1977. Catch peaked in the late 1970s with retained catch of over 2.9 million pounds. Since 1994, the Norton Sound crab fishery has operated as superexclusive. During the 2023 fishery season commercial fisheries harvested, **3,580 crab (10,013 lb)** in winter and **146,087 (413,327 lb)** in summer. The winter subsistence fishery caught a total of **to be reported male crab (to be reported lb)** and retained **to be reported (to be reported lb)**. In total, **> 149,667 crab (>423,340 lb)** were harvested during the 2023 season. The estimated discard mortality from the model 21.0 was **21,350 lb (Assuming that Winter subsistence catch was the same as 2022)**, and the total fishing mortality was **0.446** million lb. This did not exceed ABC of 0.450 million lb. Thus, overfishing did not occur during the 2023 season.
3. Stock Biomass. Norton Sound red king crab is monitored not in biomass but in abundance. For the assessment model, biomass is calculated by multiplying the average weight of each length class. Abundance of the Norton Sound red king crab stock has been monitored by trawl surveys since 1976 by NMFS (1976-1991), NOAA NBS (2010-2022), and ADF&G (1996-2021). Historical survey abundance of Norton Sound red king crab of carapace length greater than 63 mm (CL > 63 mm) ranged from 1.41 million to 5.90 million crab. In 2023 abundance of crab estimated from the ADF&G trawl survey was 3.44 million crab with CV 0.325, and that from NOAA NBS survey was **to be reported** million crab with CV **to be reported** (Table 3).
4. Recruitment. Recruitment is not monitored directly. It is inferred by the assessment model. Model-estimated recruitment since the 1980s has averaged ~0.70 million, ranging from 0.20 to 1.60 million.

5. Management performance.

Status and catch specifications (million lb)

Year	MSST	Biomass (MMB)	GHL	Retained Catch	Total Catch	OFL	ABC
2019	2.24	3.12	0.15	0.08	0.08	0.24	0.19
2020	2.28	3.67	0.17	Conf.	Conf.	0.29	0.21
2021	2.25	5.05	0.31	0.007	0.007	0.59	0.35
2022	2.08	5.33	0.34	0.34	0.36	0.67	0.40
2023	2.65	5.29	0.392	0.423+	0.446	0.643	0.450
2024			TBD	TBD	TBD		

Note

MSST was calculated as $B_{MSY}/2$

OFL-ABC 2018-2020 are retained only

2019, 2020: Total catch equals retained catch. Discarded catch was estimated only for the summer commercial fishery, but the summer commercial fishery did not occur.

OFL-ABC 2023 -retained only

Status and catch specifications (k t)

Year	MSST	Biomass (MMB)	GHL	Retained Catch	Total Catch	OFL	ABC
2019	1.03	1.41	0.07	0.04	0.04	0.11	0.09
2020	1.04	1.66	0.08	Conf.	Conf.	0.13	0.09
2021	1.02	2.29	0.14	0.003	0.003	0.20	0.16
2022	0.95	2.42	0.15	0.15	0.16	0.30	0.18
2023	1.20	2.40	0.178	0.192+	0.202	0.292	0.204
2024			TBD	TBD	TBD		

Conversion to Metric ton: 1 Metric ton (t) = 2.2046×1000 lb

Biomass (million lb)

Year	Tier	B_{MSY}	Current MMB	B/B_{MSY} (MMB)	F _{OFL}	Years to define B_{MSY}	M	ABC Buffer	ABC
2019	4b	4.57	3.12	0.7	0.12	1980-2019	0.18	0.2	0.19
2020	4b	4.56	3.66	0.8	0.14	1980-2020	0.18	0.3	0.21
2021	4a	4.53	5.05	1.1	0.18	1980-2021	0.18	0.4	0.35
2022	4a	4.17	5.33	1.3	0.18	1980-2022	0.18	0.4	0.40
2023	4a	4.37	5.29	1.2	0.18	1980-2023	0.18	0.3	0.450
2024	4a					1980-2024	0.18	0.3	

Biomass in kt

Year	Tier	B _{MSY}	Current MMB	B/B _{MSY} (MMB)	F _{OFL}	Years to define B _{MSY}	M	ABC Buffer	ABC
2019	4b	2.06	1.41	0.7	0.12	1980-2019	0.18	0.2	0.09
2020	4b	2.07	1.66	0.8	0.14	1980-2020	0.18	0.3	0.09
2021	4a	2.05	2.29	1.1	0.18	1980-2021	0.18	0.4	0.16
2022	4a	1.90	2.42	1.3	0.18	1980-2022	0.18	0.4	0.18
2023	4a	1.98	2.40	1.2	0.18	1980-2023	0.18	0.3	0.204
2024	4a					1980-2024	0.18	0.3	

6. Probability Density Function of the OFL and mcmc estimates

To be reported for the final model in Jan 2024.

7. The basis for the ABC recommendation.

For Tier 4 stocks, the default maximum ABC is based on P*=49%: essentially identical to the OFL. The annual ABC buffer is determined by accounting for uncertainties in assessment and model results. **Criteria for determining the level of ABC buffer are qualitative.** The buffer was 10% from 2011 to 2014 that was increased to 20% in 2015, to 30% in 2020, and to 40% in 2021. In 2023 CPT recommended to reduce buffer to 30%

Year	ABC Buffer
2011-2014	10%
2015-2019	20%
2020	30%
2021-2022	40%
2023	30%

8. Summary of the results of any rebuilding analysis

NA: NSRKC is not overfished.

A. Summary of Major Changes in 2024 assessment model

1. Changes to the management of the fishery.

None.

2. Changes to the input data.

Input data update through 2023:

Winter subsistence, winter and summer commercial crab fishery harvest.

Trawl surveys: abundance, length-shell compositions: ADF&G summer trawl and NOAA NBS (**To be included**) in Aug 2023.

3. Changes to the assessment methodology.

a. **Model 21.0: Baseline model adopted in Sept 2021 for the final assessment for Jan 2022.**

b. **Model 23.0: Single length-independent M: Estimated**

4. Changes to the assessment results.

B. Response to SSC and CPT Comments

Following are SSC, CPT-SSC's requests/review (received in Jan-Feb 2023) and authors' responses, arranged by topic. Requests are italicized.

I. NSRKC Biology-Ecology

Size at maturity

SSC (Feb 2023): Test the sensitivity of the assessment model to a match lower size at maturity.

Authors reply:


The identical request was raised and answered in 2019 and 2022 (see: section E) (SAFE 2019, 2022). The assessment model is insensitive to the size at maturity in setting of F_{OFL} because Tier 4 F_{OFL} determination rule is based on the MMB/BMSY ratio.

Natural Mortality

SSC (Feb 2023): a variant of model 21.0 for next year's assessment with one estimated value of natural mortality for all size.

Authors reply:

The identical requests have been made and evaluated in 2013, 2015, 2016, and 2017 (see section E1) (NPFMC 2013, 2016, 2017). The model estimated M is > 2 times higher than 0.18 but the model fit was poorer than the default model.



Profile analyses from SAFE 2015 (left) and 2023 (right)

Author provided and evaluated model 23.0 with updated data.

II. NSRKC Assessment Surveys and Data

Discards

SSC (Feb 2023) A small-scale observer program should be considered for the NSRKC fishery.

Authors reply:

All the previous observer programs were small-scale. A full-scale observer program has never been operated in the NSRKC fisheries. Fisheries biologists and managers of both the ADF&G and the NSEDC indicated that estimates of discards mortality from the previous **small-**

scale programs were inaccurate and untrustworthy, which is due to great sampling bias in selection of vessels, fishing location, timing, and fishing behavior (NPFMC 2020). The NSRKC program lacks expertise in designing and operating an observer program that provide accurate estimates of discards mortality. We welcome experts, such as the North Pacific Observer Program, to design and operate a small-scale observer program.

VAST

SSC (Feb 2023) Continue to develop VAST or other model-based survey estimates of abundance.

Authors reply:

The authors look forward to Dr. Jon Richar (NOAA)'s progress on VAST.

GMACS

SSC (Feb 2023) Prioritize transitioning the model to GMACS.

Authors reply:


The authors look forward to Drs. Andre Punt and Matthieu Veron's (the University of Washington) progress.

Standardized CPUE

SSC (Jan 2023) Consider an update to the standardized commercial fishery CPUE model.

Authors reply:

Updating the model is a worthy endeavor, and the authors welcome any technical support for the model improvements. However, it is unlikely that this will result in a better assessment model estimate. The dynamics of the standardized CPUE is very similar to arithmetic CPUE. The model estimated additional variance is large to influence the model dynamics and fit (Figure 7). In fact, whether to include the standardized CPUE data has little impact on the model likelihood and projections (SAFE 2018). The standardized CPUE was originally included as supplemental to the triennial trawl survey, especially during the periods of no trawl survey. Given that the NOAA NBS trawl survey is conducted annually since 2021, importance of the data has been diminishing.



Standardized CPUE vs. arithmetic CPUE.

III. NSRKC Assessment model

NA

IV. NSRKC Management

LTK

SSC (Feb 2023)

Consider using NSRKC as a case study for the incorporation of local knowledge, traditional knowledge, and subsistence information for Council decision-informing analyses as previously suggested (see February 2018, 2019, 2020, and 2021 SSC Reports in BSAI Crab, BS FEP Climate Change Task Force, and BS FEP- LKTKS Taskforce sections). It is assumed that this work would be led by AFSC social science personnel (or other similarly qualified researchers) and would involve inputs from the Local Knowledge, Traditional Knowledge, and Subsistence (LKTKS) Task Force and the Climate Change Task Force (CCTF). The SSC would welcome a presentation on recent work done by the AFSC in this area.

Authors reply:

The authors look forward to the findings.

C. Introduction

1. Species:

Red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska.

2. General Distribution:

Norton Sound red king crab (NSRKC) is one of the northernmost red king crab populations that can support a commercial fishery (Powell et al. 1983). It is distributed throughout Norton Sound with a westward limit of 167-168° W. longitude, depths less than 30 m, and summer bottom temperatures above 4° C. The Norton Sound red king crab management area consists of two units: Norton Sound Section (Q3) and Kotzebue Section (Q4) (Menard et al. 2011). The Norton Sound Section (Q3) consists of all waters in Registration Area Q north of the latitude of Cape Romanzof, east of the International Dateline, and south of 66°N latitude (Figure 1). The Kotzebue Section (Q4) lies immediately north of the Norton Sound Section and includes Kotzebue Sound. Commercial fisheries have not occurred regularly in the Kotzebue Section. This report deals with the Norton Sound Section of the NSRKC management area.

3. Evidence of stock structure:

Based on variability at 15 SNP loci and in mtDNA sequences (COI, 665 bp), the NSRK stock belongs to the Okhotsk Sea–Norton Sound–Aleutian Islands evolutionary lineage (SNPs, FCT = 0.054; mtDNA FCT = 0.222) (Grant and Chen 2012). However, this does not indicate that NSRKC is a single stock. The study indicates it was incapable of detecting possible evolutionary stock differences within the NSRKC stock. No studies have investigated possible stock separation within the Norton Sound management area (Figure 1).

4. Life history characteristics relevant to management:

One of the unique life-history traits of NSRKC is that they spend their entire lives in shallow water since Norton Sound is generally less than 40 m in depth. Based on the 1976-2021 trawl surveys, NSRKC is found in areas with a mean depth range of 19 ± 6 (SD) m and bottom temperatures of $7.4^{\circ} \pm 2.5$ (SD) C during summer. NSRKC is consistently abundant offshore of Nome.

NSRKC migrates between deep offshore and shallow inshore waters within Norton Sound. The timing of the inshore mating migration is unknown but is assumed to be during late fall to winter (Powell et al. 1983). Offshore migration occurs in late May - July (Bell et al. 2016). The results from a study funded by North Pacific Research Board (NPRB) during 2012-2014 suggest that older/larger crab (> 104 mm CL) may stay offshore in the winter, based on findings that larger crabs are not found nearshore during spring offshore migration periods (Jenefer Bell, ADF&G, *personal comm*). Molting occurs in fall to winter. Double shelled crabs were often observed in late August commercial catch (Joyce Song ADF&G *personal comm*). Laboratory observation showed that male crab molted in August – November and female crab molted in Jan-March (Leah Zacher and Jennifer Gardner NOAA-AFSC *personal comm*). Trawl surveys show that crab distributions are patchy and dynamic. Functional maturity of NSRKC male crab is as small

as 79.4 mm CL (Leah Zacher NOAA Kodiak *personal comm*). Those small males could also fertilize eggs of ~ 4 females, which was comparable to the number of females larger (94-116 mm CL) crabs could fertilize. More interestingly, the largest crab (> 123 mm) was able to fertilize eggs of ~ 2 females.

5. Brief management history:

NSRKC fisheries consist of commercial and subsistence fisheries. The commercial red king crab fisheries occur in summer (June – August) and winter (December – May), and subsistence is open year-round. The majority of NSRKC is harvested during the offshore summer commercial fishery, whereas the winter commercial and subsistence fisheries occur nearshore through ice and take a much smaller harvest.

The distinguishing characteristic of the NSRKC fisheries is that all fisheries, surveys, researches, and management are conducted by local residents of Norton Sound. Commercial fisheries are designated as super-exclusive: a vessel registered for the Norton Sound crab fishery may not be used to take king crabs in any other registration areas. The ADF&G and the NSRKC crab research and management biologists are members of Nome community and are acquainted with many local fishermen (commercial and subsistence) and staff of community organizations such as Norton Sound Economic Development Corporation (NSEDC) and Kawerak Inc, exchanging information and research ideas about crab biology and fisheries management.

Summer Commercial Fishery

A large-vessel summer commercial crab fishery started in 1977 in the Norton Sound Section (Table 1) and continued from 1977 through 1990. No summer commercial fishery occurred in 1991 because there were no staff to manage the fishery. In March 1993, the Alaska Board of Fisheries (BOF) limited participation in the fishery to small boats. Then on June 27, 1994, a super-exclusive designation went into effect for the fishery. This designation states that a vessel registered for the Norton Sound crab fishery may not be used to take king crabs in any other registration areas during that registration year. A vessel moratorium was put into place before the 1996 season. This was intended to precede a license limitation program. In 1998, Community Development Quota (CDQ) groups were allocated a portion of the summer harvest; however, no CDQ harvest occurred until the 2000 season. On January 1, 2000, the North Pacific License Limitation Program (LLP) went into effect for the Norton Sound crab fishery. The program dictates that a vessel which exceeds 32 feet in length overall must hold a valid crab license issued under the LLP by the National Marine Fisheries Service. Changes in regulations and the location of buyers resulted in eastward movement of the harvest distribution in Norton Sound in mid-1990s. In Norton Sound, a legal crab is defined as $\geq 4 \frac{3}{4}$ -inch carapace width (CW, Menard et al. 2011), which is approximately equivalent to ≥ 104 mm carapace length (CL). In 2005 and 2006, commercial buyers, specifically Norton Sound Economic Development Corporation (NSEDC), accepted only legal crab of ≥ 5 inch CW. This preference became permanent in 2008.

Portions of the Norton Sound area are closed to commercial fishing for red king crab. Since the beginning of the commercial fisheries in 1977, waters approximately 5-10 miles offshore of southern Seward Peninsula from Port Clarence to St. Michael have been closed to protect nearshore subsistence fisheries and to act as a refuge for crab during the summer commercial crab fishery (Figure 2). The spatial extent of closed waters has varied historically, with the closure line being moved in to provide additional area to achieve harvest goals. In 2020 the BOF closed Norton Sound area east of 167 degrees W. longitude for the commercial summer crab fishery. In 2020 and 2021 the NSEDC did not purchase NSRKC resulting in small or no harvest. In 2022, the NSEDC resumed purchasing summer commercial catch.

CDQ Fishery

The Norton Sound and Lower Yukon CDQ groups divide the NSRKC CDQ allocation. Only fishers designated by the Norton Sound and Lower Yukon CDQ groups are allowed to participate in this portion of the king crab fishery. Fishers are required to have a CDQ fishing permit from the Commercial Fisheries Entry Commission (CFEC) and register their vessel with the Alaska Department of Fish and Game (ADF&G) before beginning fishing. Fishers operate under the authority of each CDQ group. CDQ harvest share is 7.5% of the guideline harvest level (GHL), and can be prosecuted in both summer and winter seasons.

Winter Commercial Fishery

The winter commercial crab fishery uses hand lines and pots through the nearshore ice. On average 10 permit holders harvested 2,500 crab during 1978-2009. From 2007 to 2015 the winter commercial catch increased from 3,000 crab to over 40,000 (Table 2). In 2015 the winter commercial catch reached 20% of total crab catch. The BOF responded in May 2015 by amending regulations to allocate 8% of the total commercial GHL to the winter commercial fishery, which has been in effect since the 2017 season. The timing of the winter red king crab commercial fishing season has changed over time to address ice stability. It was originally from January 1 to April 30, amended in 1985 to from November 15 to May 15. In 2015 the period was changed to January 15 to April 30 after fisheries opened in Nov 15 in 2014, so that January 15 starting date was into effect in 2016. In 2021 it was further amended to February 1 to April 30. The NSEDC terminated purchasing crab in 2019. Since 2019 all the winter commercial catches are by catcher-seller permit holders.

Year	Opening period
1977-1984	Jan 01 – Apr 30
1985- 2014	Nov 15 – May 15
2015	Nov 15 – Apr 30
2016-2020	Jan 15 – Apr 30
2021 -	Feb 01 – Apr 30

Subsistence Fishery

The winter subsistence fishery has a long history; however, harvest information is available only since the 1977/78 season. The majority of subsistence crab harvest occurs in winter using hand lines and pots through nearshore ice. The average annual winter subsistence harvest is 5,281 crabs (1977-2021). Subsistence harvesters need to obtain a permit before fishing and record daily effort and catch. There are no size or sex-specific harvest limits; however, the majority of retained catch is males of near legal size.

Summer subsistence crab fishery harvest has been monitored since 2004 with an average harvest of 1,145 crabs (2004-2020). The summer subsistence fishery was not included in the assessment model.

Harvest of both winter commercial and subsistence fisheries is influenced by availability of stable ice conditions. Small harvests can occur due to poor ice condition, regardless of crab abundance.

1. Brief description of the annual ADF&G harvest strategy

Since 1997 NSRKC has been managed based on a GHL. From 1999 to 2011 the GHL for the summer commercial fishery was determined using model estimated predicted biomass: (1) 0% harvest rate of legal crab when estimated legal biomass < 1.5 million lb; (2) $\leq 5\%$ of legal male biomass when the estimated legal biomass falls within the range 1.5-2.5 million lb; and (3) $\leq 10\%$ of legal male biomass when estimated legal biomass >2.5 million lb. In 2012 the summer commercial fishery GHL was revised to (1) 0% harvest rate of legal crab when estimated legal biomass < 1.25 million lb; (2) $\leq 7\%$ of legal male biomass when the estimated legal biomass falls within the range 1.25-2.0 million lb; (3) $\leq 13\%$ of legal male biomass when the estimated legal biomass falls within the range 2.0-3.0 million lb; and (3) $\leq 15\%$ of legal male biomass when estimated legal biomass >3.0 million lb.

In 2015 the BOF passed the following regulations regarding the winter commercial fisheries:

- 1) Revise GHL to include summer and winter commercial fisheries.
- 2) Set GHL for the winter commercial fishery (GHL_w) at 8% of the total GHL

NSRKC GHL is determined in early February after the final determination of ABC. GHL is determined not to exceed (e.g., 5-10% less) the retained portion of the total catch ABC.

Table: Brief NSRK fishery management history

Year	Notable historical management changes
1976	The abundance survey started
1977	Large vessel commercial fisheries began. Legal size was set to ≥ 5 inch CW
1978	Legal size was changed to ≥ 4.75 inch CW
1991	Fishery closed due to staff constraints
1993	Fishery is restricted to small boat. The end of large vessel commercial fishery operation.
1994	Super exclusive designation went into effect.
1998	Community Development Quota (CDQ) allocation went into effect
1999	Guideline Harvest Level (GHL) went into effect
2000	North Pacific License Limitation Program (LLP) went into effect.
2002	Change in closed water boundaries (Figure 2)
2006	The Statistical area Q3 section expanded (Figure 1)
2008	Start date of the open access fishery changed from July 1 to after June 15 by emergency order. Pot configuration requirement: at least 4 escape rings (> 4.5 inch diameter) per pot located within one mesh of the bottom of the pot, or at least $\frac{1}{2}$ of the vertical surface of a square pot or sloping side-wall surface of a conical or pyramid pot with mesh size > 6.5 inches.
2008	Market preferred size of ≥ 5 inch CW became a standard commercial retained size.
2012	The BOF adopted a revised GHL for summer fishery.
2016	Winter GHL for commercial fisheries was established and modified winter fishing season dates were implemented.
2019	The NSEDC stopped purchasing of the winter commercial crab.
2020	The BOF closed summer commercial fishery East of 167 longitude. Summer commercial
2021	fisheries opened but the NSEDC did not purchase the summer commercial crab.
2021	Summer commercial fishery Change winter fishery open date to February 1

2. Summary of the history of the B_{MSY} .

NSRKC is a Tier 4 crab stock. Direct estimation of the B_{MSY} is not possible. The B_{MSY} proxy is calculated as the mean model estimated mature male biomass (MMB) from 1980 to the present. The choice of this period was based on a hypothesized shift in stock productivity due to a climatic regime shift indexed by the Pacific Decadal Oscillation (PDO) in 1976-77.

D. Data

1. Summary of new information:

Winter commercial (Reported) and subsistence fisheries (to be reported in Jan 2024):

The winter commercial fishery retained catch in 2023 was 3,509 crab (10,013 lb). Winter subsistence total male crab catch was **to be reported** and retained male crab catch was **to be reported**, and total female catch was **to be reported** and retained female was **to be reported**. In total, **to be reported** crab were caught and **to be reported** crab were retained (Table 2).

Summer commercial fishery:

The summer commercial fishery opened on 6/15/2023 and closed on 7/24/2023. A total of 146,087 crab (413,327 lb) were harvested (Table 1).

Standardized CPUE (Appendix B).

Standardized CPUE for the years of 1991, 2020, 2021 were not calculated because a commercial fishery was closed (1991) or no crab was harvested during the commercial fishery (2020, 2021).

The standardized CPUE: 3 periods:

1977-1992: Large Scale commercial fishery (CL > 4.75 inches)

1993-2007: Small boat commercial fishery (CL > 4.75 inches)

2008-2023: Small boat commercial fishery with high grading (CL > 5.0 inches)

Available survey, catch, and tagging data

Available NSRKC data consist of the following: trawl survey that informs abundance and size composition, catch that informs size composition, and standardized CPUE that informs an index of abundance, and tag recovery that informs growth-transition.

Trawl survey

Trawl survey consists of 3 surveys: NMFS triennial survey: 1976-1992, ADF&G survey: 1996-2023, and NOAA NBS survey: 2010, 2017-2023.

NMFS triennial survey:

A Norton Sound trawl survey was initiated by NMFS in 1976 to assess stock status of crab and ground fish in Norton Sound and Kotzebue Sound. The survey established 10 nautical mile (nm) grid survey stations throughout the entire Norton Sound and 15 nm grids outside the Norton Sound area. The initial Norton Sound survey became the standard stations moving forward. The survey was conducted from mid-late August to September-October, except for 1979, which was in late July/early August. The survey used 83-112 Eastern Otter trawl gear, with tow distance of 1.3 – 1.7 nm (30 minutes tow). The survey was terminated in 1992.

ADF&G triennial -annual survey:

After the termination of the NMFS trawl survey, ADF&G began trawl surveys in 1996 using the same survey stations, but smaller boat and survey coverage. The survey started as

triennially but became an annual survey in 2017. The survey usually occurs in late July – mid August, using 400 Eastern Otter trawl gear with tow distance of 1.0 nm. The survey used to have a re-tow protocol: when the first tow caught more than 5 legal red king crab, the station was re-towed. This protocol was dropped in 2012 in favor of more coverage.

NOAA biennial-annual NBS survey:

NOAA NBS trawl survey started in 2010, and biennially since 2017. The survey occurs in late July-mid August, similar to the ADF&G survey. The survey has 20 nm grid using 83-112 Eastern Otter trawl gear, with tow distance of 1.3 – 2.5 nm (30 min tow).

Abundance estimation method

Methods of estimating red king crab abundance differed among the three surveys and throughout time periods. Abundance estimates have been revised many times.

Abundance and CV of the NMFS 1976-1991 surveys were provided by NOAA (Jon Richer NOAA *personal communication*). The abundance was estimated by averaging catch CPUE (#/nm²) of all stations (including survey stations out of Norton Sound) that was multiplied by standard Norton Sound Area (7600 nm²) (i.e., N = 7600*mean CPUE). The ADF&G survey abundance is calculated at each station (i.e., n= CPUE*100 nm²) and summed across all surveyed stations (i.e., N = sum of 100*CPUEs) (Bell and Hamazaki 2019). Extent of the ADF&G survey coverage differed among years due to survey conditions, and survey abundance has not been standardized. NOAA NBS survey abundance is estimated by the author in similar manner as ADF&G survey with the data limited to the Norton Sound survey area that overlaps the ADF&G survey area (5841 nm²) (Figure A).




Figure A. ADF&G trawl survey coverage (yellow shade) and NOAA NBS trawl survey coverage where abundance estimates were made (red hashed line),

Survey catchability appears to differ among ADF&G, NMFS, and NOAA NBS trawl surveys. ADF&G trawl survey abundance tends to be higher than NMFS and NOAA NBS trawl survey even though NMFS and NOAA NBS survey coverages are greater than ADF&G. The assessment model assumes (recommendation by CPT-SSC) that survey q of ADF&G trawl survey be 1.0, which resulted in $q = 0.7-0.8$ for NMFS and $0.7-0.96$ for NOAA NBS survey.

Trawl survey catches are highly patchy. The majority of catches occurred at 1 to 4 stations that caught 20% to 80% of crabs caught during the entire survey (Figure B). The most consistently abundant survey stations are near Nome (blue dots) outside of the summer commercial fishery area (red rectangular). Some offshore stations had high catches for a few years (orange dots: 1990s, yellow dots: 2020s) but they did not persist.




Figure B. Trawl survey stations where majority of catch occurred. Red rectangular indicates where the majority of summer commercial fishery occurs. Blue dots indicate the stations that had high catch consistently since 1976. Orange and yellow dots indicate high crab catch stations that occurred during the 1990s (orange) and 2020s (yellow).

Standardized CPUE

Standardized summer commercial fishery CPUE is included in the NSRKC assessment model as an index of NSRKC abundance that could supplement triennial trawl survey. In 2013, the CPUE standardization model was developed by Gretchen Bishop (ADF&G) (NPFMC 2013). Since then, the same model has been applied with updated data (Appendix B).


Tagging-recovery data

Norton Sound red king crab tagging was initially conducted in 1980 as a part of mark-recapture abundance survey (Brannian 1987). The study was conducted in 1980-1982 and 1985. From 1986 to 2012 crabs were tagged during the winter pot survey. The winter pot surveys tagged more smaller (sublegal) crabs; however, very few were recovered. Tagging resumed from 2012-2015 for a spring migration movement survey. In all the above studies, most of the tagged crabs were recovered by commercial fishermen, but subsistence fishermen also recovered a small number of tags.

Length-Shell proportion data

Length-shell data have been collected in every research and harvest monitoring survey. Of those, summer commercial harvest sampling, winter pot survey (terminated in 2012), and the trawl survey have been consistent.

Time series of the data used for the NSRKC assessment model are summarized in the following figure and table.



Time series of available data

Table A: List of survey data

	Years	Data Types	Tables
Summer trawl survey	76,79,82,85,88,91,96, 99, 02,06,08,10,11,14,17-23	Abundance Length-shell comp	3 6
Winter pot survey	81-87, 89-91,93,95-00,02-12	Length-shell comp	7
Summer commercial fishery	77-90,92-23	Retained catch Standardized CPUE, Length-shell comp	1 1, Appendix B 4
Summer Com total catch	12-23	Length-shell comp	9
Summer Com Discards	87-90,92,94	Length-shell comp	8
Winter subsistence fishery	76-23	Total & Retained catch	2
Winter commercial fishery	78-23	Retained catch	2
	15-18	Retained Length-Shell	5
Tag recovery	80-19	Recovered tagged crab	10

Table B: List of data available but not used for assessment

Data	Years	Data Types	Reason for not used
Summer pot survey	80-82,85	Abundance Length proportion	Uncertainties on how estimates were made.
Summer preseason survey	95	Length proportion	Just one year of data
Summer subsistence fishery	2005-2019	retained catch	Too few catches, ignored.
Winter Pot survey	87, 89-91,93,95- 00,02-12	CPUE	CPUE data unreliable.
Preseason Spring pot survey	2011-15	CPUE, Length proportion	Years of data too short
Postseason Fall pot survey	2013-15	CPUE, Length proportion	Years of data too short

Catches in other fisheries

In Norton Sound, the directed Pacific cod pot fishery was issued in 2018 under the CDQ permit. In 2018 and 2019 fishery seasons, a total of 8 and 13 kg (mortality applied) of NSRKC were taken in the groundfish fisheries (CPT 2020). However, all bycatch occurred to the west of 168.0 longitude where NSRKC survey has not been conducted. Norton Sound Fishery Management Area (Q3) extends to St. Lawrence Island and US-Russia border (Figure 1). In the absence of survey abundance extended to those area, it is questionable whether those bycatch mortalities should be included in the NSRKC population.

	Fishery	Data availability
Other crab fisheries	Does not exist	NA
Groundfish pot	Pacific cod	Y
Groundfish trawl	Does not exist	NA
Scallop fishery	Does not exist	NA

Other miscellaneous data:

- Tagging-recovery locations (2012-2019)
- Satellite tag migration tracking (NOAA 2016, ADF&G 2020-21)
- Spring offshore migration distance and direction (2012-2015)
- Monthly blood hormone level (indication of molting timing) (2014-2015)
- Functional maturity and mating success of captured crab (2021-22)

Data aggregated:

- Length data were aggregated by 10 mm range, starting from 64-73 mm. Crab length greater than 133 mm were aggregated in >133 mm class.
- Shell condition data were recorded in very new, new, old, very old, and very very old. Those were combined to newshell (very new, new) and oldshell (old, very old, very very old).
- Tag-recovery data were aggregated based on years in liberty, regardless of years of tagging and recovery.

Data estimated outside the model:

- Summer commercial catch standardized CPUE (Table 1, Appendix B)
- Average weight of crab by length class (Table 13)

E. Analytic Approach

1. History of the modeling approach and issues:

The Norton Sound red king crab stock was assessed using a length-based synthesis model (Zheng et al. 1998). Since adoption of the model, the model had the following model mismatches:

1. Model projects higher abundance-proportions of large size class (> 123mm CL) of crab than observed.
2. Poor model fit to trawl survey abundance.
3. Some model parameters hit boundaries

The issues of 1 and 2 are attributed to natural mortality specification. Natural mortality M specification was originally **specified** to be 0.2 for BSAI red king crab stocks including NSRKC (NPFMC 1998) and was **specified** to 0.18 with Amendment 24 (NPFMC 2011). In crab stock assessment modeling, natural

mortality is assumed to be the same across all individual lengths (i.e., length-independent M).

1. Model projects higher abundance-proportions of large size class ($> 123\text{mm CL}$) of crab.

This issue has been solved by assuming (3-4 times) higher M for the large crab (i.e., $M = 0.18$ for length classes ≤ 123 mm, and higher M for > 123 mm) (NPFMC 2012, 2013, 2014, 2015, 2016, 2017, 2018). However, because this solution is biologically suspect, several alternative assumptions have been considered in the past.

- a. Large crabs move out of the survey and fishing area

In modeling, this was dealt with by setting dome-shaped survey and commercial catch selectivity (i.e., lower catchability for large crabs). This modeling configuration resulted in estimating MMB two times higher than the default model, which indicates that true NSRKC abundance is twice larger than current trawl survey and commercial crab fishery indicate (NPFMC 2017). The NOAA NBS surveys (2010, 2017, 2019, 2021) did not find high numbers of red king crabs outside Norton Sound. The large crab could also be nearshore where the commercial fishery is closed, and a trawl survey is not conducted due to rocky bottom. However, spring tagging studies showed that most crabs migrated from near shore to offshore (fishing) area (Jenefer Bell, ADF&G personal comm.). There was little evidence that large crabs stay in nearshore waters during summer.

- b. Molting and growth of NSRKC are slower. (i.e., model overestimating molting and growth probability: transition matrix)

The model originally estimated the transition outside of the model. In 2014 the model was configured to estimate the transition matrix inside of the assessment model (NPFMC 2014). The transition matrix estimated inside of the model was similar to that estimated outside of the model. When length-specific molting probability was estimated individually, the shape of the probability curve was also similar to the default inverse logistic molting function (NPFMC 2016). A time varying molting function (random walk) process did not improve model fit. Laboratory studies showed that observed growth after molting was comparable to those from tag-recovery data, though sample size was limited and comparable tank-natural condition factors such as water temperature and food availability, were questionable (Leah Zacher of NOAA AFSC Kodiak personal comm),

- c. Higher natural mortality (M) than assumed $M = 0.18$

Profile analyses and estimating M across all length classes resulted in higher M (0.3-0.45) than default $M=0.18$ (NPFMC 2013, 2016, 2017). However, the model fit is slightly worse than the default model.

Formatted: Normal, Left, Right: 0.03", Line spacing: Multiple 1.03 li

- d. Higher natural mortality ($M > 0.18$) for small crab and large crab having higher mortality than small crab.

This model configuration had the best fit to data (NPFMC 2016, 2017).

- e. Gradual size dependent natural mortality.

The default assessment model assumes abrupt M increase at size CL 124mm or greater. An alternative model suggested that M gradually increasing from size as low as 94 mm CL; however, the overall model fit did not greatly improve from the default model (NPFMC 2017). In 2022, CPT requested estimating M for each length class, which also suggested length-dependent natural mortality.

- 2. Poor model fit to trawl survey abundance, especially NMFS survey (1976-1992) data

The NSRKC assessment model suggest higher crab abundance during the 1976-1990s period. The model deals with this issue by including survey q ($q < 1$), or the model assumes the NMFS trawl surveys **underestimated** NSRKC abundance. However, this assumption is arbitrary, which is also affected by other model configurations. For instance, when $M = 0.18$ is assumed for all length classes, the model suggests that survey q for NMFS is greater than 1.0, or that the NMFS trawl surveys **overestimated** NSRKC abundance (Model 21.1). This indicates the trawl surveys overestimated NSRKC abundance (NPFMC 2022). Alternatively, assuming the NMFS survey q to be 1.0 resulted in ADF&G trawl survey q greater than 1.0 (i.e., trawl survey overestimates abundance), even though ADF&G trawl survey area is generally smaller than NMFS and NOAA NBS survey areas.

This model fitting issue was also influenced by input sample sizes for size-shell compositions. Increasing the input sample size resulted in the model estimating lower abundance. Reducing the input sample sizes improved model fit to the trawl survey data but caused lower fit to size-shell composition data (NPFMC 2012, 2013, 2015). Alternative model weighting methods (e.g., Francis 2012) have been tried, but those did not improve model fit.

- 3. Some model parameters hit boundaries.

There are two model parameters that hits boundaries: Trawl survey selectivity ($\log_{\phi_{tl}}$), and the proportion of recruits ($r1, r2$).

1. Trawl Survey selectivity parameter

Trawl survey selectivity model is a one parameter logistic curve that reaches 1.0 at L_{max} (143.5 mm)

$$S_l = \frac{I}{I + e^{(\alpha(L_{max} - L) + \ln(1/0.999 - 1))}}$$

where $\alpha = \exp(\log_{\phi_{stl}})$, $L_{max} = 143.5$ mm L (63.5-143.5 mm)

Model estimated trawl survey selectivity is 1.0 across all size classes. This means that $e^{(\alpha(L_{max} - L) + \ln(1/0.999 - 1))} \approx 0$, $\alpha(L_{max} - L) + \ln(1/0.999 - 1) = -\infty$, $\alpha \approx 0$, and $\log_{\phi_{stl}} = -\infty$. Hence, the parameter will hit the boundary.

Alternative option is assuming $S_l = 1.0$ for all length classes; however, this also removes the model's ability to estimate S_l when all length classes are NOT 1.0.

2. The proportion of recruits

The proportion of recruits is a multinomial formula of $n=3$

$$p_l = \frac{\exp(r_l)}{1 + \sum_{i=1}^{n-1} \exp(r_i)} \text{ for } l = 1, \dots, n-1$$

$$p_n = 1 - \frac{\sum_{i=1}^{n-1} \exp(r_i)}{1 + \sum_{i=1}^{n-1} \exp(r_i)}$$

Model estimated recruit proportions for length classes 1, 2, 3 (P_1, P_2, P_3) are 0.592, 0.403, and 0.003. $P_3 \approx 0$ makes it extremely difficult for the model to estimate P_1 and P_2 , and thus model parameters r_1 and r_2 (Tables 11, 12). **Increasing the upper bound of the r parameters would still make r_1 hit the boundary and make estimates of P_1, P_2, P_3 to be closer to $P_1 = 0.60, P_2 = 0.40, \text{ and } P_3 = 0$.** Alternative option is assuming $P_3 = 0$; however, this also removes the model's ability to estimate P_3 when P_3 is far greater than 0.

Historical Model configuration progression:

2011 (NPFMC 2011)

- 1). $M = 0.18$.
- 2). M of the last length class = 0.288.
- 3). Include summer commercial discards mortality = 0.2.
- 4). Weight of fishing effort = 20.
- 5). The maximum effective sample size for commercial catch and winter surveys = 100.

2012 (NPFMC 2012)

- 1) M of the last length class = $3.6 \times M$.
- 2) The maximum effective sample size for commercial catch and winter surveys = 50.
- 3) Weight of fishing effort = 50.

2013 (NPFMC 2013)

- 3) Standardize commercial catch CPUE and replace the likelihood of commercial catch efforts with standardized commercial catch CPUE with weight = 1.0.
- 4) Eliminate summer pot survey data from likelihood.
- 5) Estimate survey q of 1976-1991 NMFS survey with a maximum of 1.0.
- 6) The maximum effective sample size for commercial catch and winter surveys = 20.

2014 (NPFMC 2014)

- 1) Modify the functional form of selectivity and molting probability to improve parameter estimates (2 parameter logistic to 1 parameter logistic).
- 2) Include additional variance for the standardized CPUE.
- 3) Include winter pot survey CPUE (was removed from the final model due to lack of fit).
- 4) Estimate growth transition matrix from tag-recovery data.

2015 (NPFMC 2015)

- 1) Winter pot survey selectivity is an inverse logistic, estimating selectivity of the smallest length group independently.
- 2) Reduce weight of tag-recovery: $W = 0.5$.
- 3) Model parsimony: one trawl survey selectivity and one commercial pot selectivity.

2016 (NPFMC 2016)

- 1) Length range extended from 74 mm – 124 mm above to 64 mm – 134 mm above.
- 2) Estimate multiplier for the largest (> 123 mm) length classes.

2017 (NPFMC 2017)

- 1) Change molting probability function from 1 to 2 parameter logistic. Assume molting probability not reaching 1 for the smallest length class.

2018 (NPFMC 2017) CPT-SSC suggested no model alternatives

2019 (NPFMC 2019)

- 1) Fit total catch length composition and estimate retention probability for summer and winter commercial fishery.
- 2) Include winter commercial retained length data.

2020 (NPFMC 2020) The CPT and SSC suggested no model alternatives

2021 (NPFMC 2021) Include discards data at the request of CPT and SSC

- 1) Models that bridge from the Model 19.0e to 21.0

- 2) Model 21.0 with natural mortality estimated by model.
- 3) Estimate size specific natural mortality.

2022 (NPFMC 2022)

- 1) Examine shell-based retention probability.
- 2) Estimate individual length class M .

2023 (NPFMC 2023)

- 1) Model with single M estimated.

2. Model Description

- a. Description of overall modeling approach:

The model is a male-only size structured model based on **abundance** that combines multiple sources of surveys, fishery catches and discards, and mark-recovery data using a maximum likelihood modeling framework to estimate population dynamics under fisheries. The model is an extension of the length-based model developed by Zheng et al. (1998) for NSRKC. A detailed description of the model is in Appendix A.

The model estimates abundances of male crab with $CL \geq 64$ mm and with 10 mm length intervals (8 length classes, ≥ 134 mm) because few crab measuring less than 64 mm CL were caught during surveys or fisheries.

The model assumes newshell crab as molted and oldshell crab as unmolted.

One critical characteristic of the model is that it does not estimate fishing mortality (F). Observed harvests were considered accurate and thus directly subtracted from the model estimated abundance.

The modeling scheme and data is described in the following figure.




Figure C. Norton Sound red king crab model and data scheme. Bold type indicate data that were fitted to the model. Boxes in dotted line indicate model estimated parameters and quantities. Natural mortality, M was set to 0.18 except for CL greater than 123mm that was estimated in the model.

Timeline of calendar events and crab modeling events:

- **Model year starts February 1st to January 31st of the following year.**
- **Initial Population Date: February 1st, 1976, consisting of only newshell crab.**
- **Instantaneous fishing mortality: winter (February 1st) and summer (July 1st) fisheries**
- **Instantaneous molting and recruitment occur on July 1st**

Critical model assumptions

NSRKC Crab Biology

1. Instantaneous annual natural mortality (M) is 0.18 and increases at the size greater than 123 mm CL. M is constant over time.

See History of the modeling approach and issues section for detailed discussion regarding this assumption

2. Male crab size at maturity is 94mm CL.

Size at maturity of NSRKC is highly uncertain (NPFMC 2018, 2019, 2020, 2021). First, maturity has two categories (biological and functional). Biological maturity indicates that male red king crab can produce viable sperm, whereas functional maturity indicates that male red king crab are large enough to mate. The former can be determined using the presence/absence of spermatophores in the vas deferens, whereas the latter can be inferred by measuring mating pairs *in situ* or in lab experiments. The current NSRKC functional maturity size (>94 mm) was inferred from Bristol Bay red king crab by incorporating the fact that Norton Sound red king crab are smaller.

SSC suggested investigating size at functional maturity of other stocks, such as of Barents Sea red king crab (NPFMC 2021). However, it is unlikely that those metadata analyses would provide insights about size at functional maturity of Norton Sound red king crab because NSRKC is the smallest among red king crab stocks. Author was not able to find any other red king crab stocks that are comparable to the size of NSRKC.

Recent laboratory studies reported that NSRKC male crab as small as 79.4 mm CL can fertilize females. Those small males could also fertilize eggs of ~ 4 females, which was comparable to the number of females larger (94-116 mm CL) crabs could fertilize. More interestingly, the largest crab (> 123 mm) was able to fertilize eggs of ~ 2 females (Leah Zacher NOAA Kodiak *personal comm*). However, reproductive success of those small crabs in the field is unknown.

Although determining size at functional maturity is important biologically, there is limited utility of this information for Tier 4 crab stock assessment. In Tier 4 stock assessment, size at maturity is used only for calculation of mature male biomass (MMB) and B_{MSY} (average MMB). Harvest control (F_{OFL}) is based on the ratio of projected MMB and B_{MSY} (projected MMB/ B_{MSY}).

Level	Criteria	F_{OFL}
a	$B / B_{MSY proxy} > 1$	$F_{OFL} = \gamma M$
b	$\beta < B / B_{MSY proxy} \leq 1$	$F_{OFL} = \gamma M (B / B_{MSY proxy} - \alpha) / (1 - \alpha)$
c	$B / B_{MSY proxy} \leq \beta$	$F_{OFL} = \text{bycatch mortality \& directed fishery } F = 0$

where B is a mature male biomass (MMB), B_{MSY} proxy is average mature male biomass over a specified time period, $M = 0.18$, $\gamma = 1$, $\alpha = 0.1$, and $\beta = 0.25$.

The MMB/B_{MSY} ratio is affected very little by changes of maturity size unless the ratio is very close to 1.0 (Tier 4a vs Tier 4b borderline). To illustrate this, we present 2022 assessment model results with various minimum size at maturity cutoffs, as follows.

Maturity size	94mm (default)	74mm	84mm	104mm	114mm	124mm	>134mm
B _{MSY} mil. lb	4.88	5.21	4.92	3.76	2.71	1.33	0.39
MMB(2022) mil. lb	5.21	5.91	5.61	4.42	2.86	1.03	0.27
MMB/B _{MSY}	1.16	1.13	1.14	1.18	1.06	0.77	0.70
Tier 4 level	a	a	a	a	a	b	b
F _{OFL}	0.18	0.18	0.18	0.18	0.18	0.13	0.12

As illustrated in the above table, changing minimum maturity size has little effect on MMB/B_{MSY} ratio and Tier 4 level designation. OFL and ABC are based on retained and unretained catch by size applied by F_{OFL}.

3. Molting occurs right after the summer fishery.

Molt timing of NSRKC was verified by field and laboratory survey. Double shelled crabs are often observed in September (Joyce Soong ADFG personal comm.), and crabs sent to Kodiak Lab molted in September-October (Leah Zacher NOAA personal comm).

4. Recruitment occurs in fall at the same time as molting.

In NSRKC assessment modeling, recruitment is not a function of mature males, but estimated model parameters entering to the immature length classes 64 mm - 93 mm. In modeling, this adjustment is done at the same time as molting-growth.

5. Molting probability is a descending logistic function of crab size. Molted crab become newshell and unmolted crab become oldshell crab.

Tag recovery data during the 2012-2014 study suggest lower molting probability for larger crabs. The table below shows the number of newshell crab tagged, released, and recaptured at 1 year of liberty. Crabs recaptured newshell is considered as molted and oldshell is considered as unmolted.

Length Class	Newshell	Oldshell	% molted
1 (64-73mm)	3	0	100
2 (74-83mm)	30	0	100
3 (84-93mm)	64	5	93
4 (94-103mm)	113	9	93
5 (104-113mm)	44	36	56
6 (114-123mm)	22	21	51
7 (124-133mm)	5	10	33
8 (>133mm)	0	4	0

This assumes that shell condition observations are correct, which is difficult to verify objectively. For instance, in tag-recovery data (2012-2016) below, 125 crabs had no growth (+/- 3 mm) in one year of liberty. Of those, 100 crabs were released as newshell and 25 crabs were released as oldshell. If no growth is considered unmolted, all those crabs should be recaptured as oldshell. However, 29% of crabs released as newshell were recaptured as newshell crab and 48% of crabs released as oldshell were recaptured as newshell.

	Recovered	
Released	Newshell	Oldshell
Newshell	28	70
Oldshell	12	13

This could be caused by (1) inaccurate length measurement, (2) inaccurate shell condition assessment, or (3) no growth after molting.

- Growth increment is a function of length, constant over time. Molted crab does not shrink.

Tag recovery data showed that growth increment of large crab tend to be smaller than that of small crab (Figure D). The data also showed negative growth increment, at the largest length class.




Figure D. Growth increment by tagged length class of molted (newshell recovered) crab with 1 year at liberty.

NSRKC Surveys

1. ADF&G trawl survey (1996-2021) abundance has the same scale as the population (i.e., catchability $q = 1.0$). Abundances by historical NMFS (1976-1991) and NOAA NBS (2010-present) survey are biased low (i.e., $q < 1.0$).

Survey $q = 1.0$ for ADF&G trawl survey and lower survey q for NOAA survey was adopted in 2013 assessment (NPFMC 2013). However, it is possible that ADF&G surveys are overestimates of abundance. Model estimated survey q for ADF&G trawl survey was greater than 1.0 (NPFMC 2013, 2019).

2. Size selectivity is an asymptotic one parameter logistic function of 1.0 at the length class 134 mm CL and the same across years and survey agencies.

$$S_l = \frac{1}{1 + e^{(\alpha(L_{\max} - L) + \ln(1/0.999 - 1))}}$$

This logistic function form was adopted during the crab workshop in 2005 to reduce model parameters and increase parameter estimation stability.

Although the surveys differ among NOAA (1976-1991), ADF&G (1996-2021), and NOAA NBS (2010-present) in terms of survey vessel and trawl net structure, selectivity of all surveys were assumed to be identical. Model fits separating and combining the surveys were examined in 2015; however, selectivity was essentially identical (1.0 across all size classes) (NPFMC 2015). For model parsimony, the SSC recommended using only one selectivity.

3. Winter pot survey selectivity is a dome shaped function: a combination of a reverse logistic function starting from length class 84 mm CL and model estimate for CL < 84 mm length classes. The selectivity is constant over time.

$$S_{w,l} = \frac{I}{I + e^{\alpha(L-\beta)}}$$

This assumption is based on the low proportion of large crab that are caught in the nearshore area where winter surveys occur. This does not necessarily imply that the crab pots are less selective to large crabs. Alternatively, this may imply that fewer large crab migrate into nearshore waters in winter.

NSRKC Fisheries

1. Fisheries occur twice on July 01 and Feb 01 and are instantaneous.
2. Summer commercial fishery size selectivity is an asymptotic one parameter logistic function of length, with the selectivity in length class 134 mm CL set to 1. Selectivity is constant over time.

$$S_l = \frac{I}{I + e^{(\alpha(L_{max}-L)+\ln(1/0.999-1))}}$$

This logistic function form was adopted during the crab workshop in 2005 to reduce model parameters and parameter estimation stability. Although summer commercial fishery changed greatly between the periods (1977-1992, 1993-present) in terms of fishing vessel composition, and pot configuration, the selectivity of each period is assumed to be identical. Model fits of separating and combining the two periods were examined in 2015 and showed no difference between the two (NPFMC 2015). For model parsimony, the SSC recommended using only one selectivity.

3. Not all legal sized crabs are retained. Retention probability is an asymptotic logistic function.

Legal size of NSRKC is defined as carapace width (CW) greater than 4.75 inches that was conventionally equated as greater than 104 mm CL. Since 1996 ADF&G has started noting legal size crab based on carapace width in trawl, commercial fishery observer, and other miscellaneous surveys to complement the carapace length measurement. Originally, the proportion was based solely from the trawl survey. As more data are collected from commercial observer surveys, recent proportions are based on more observer data.

Proportion of legal (CW>4.75 inch) crab in Trawl survey

size class	64	74	84	94	104	114	124	134
1996	0.00	0.00	0.00	0.18	0.93	1.00	1.00	1.00
1999	0.00	0.00	0.00	0.40	0.98	0.98	1.00	1.00
2002	0.00	0.00	0.00	0.28	0.97	1.00	1.00	1.00
2006	0.00	0.00	0.00	0.18	1.00	1.00	1.00	1.00
2008	0.00	0.00	0.00	0.19	0.96	1.00	1.00	1.00
2011	0.00	0.00	0.00	0.24	0.99	1.00	1.00	1.00
2014	0.00	0.00	0.00	0.21	0.98	1.00	1.00	1.00
2017	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00
2018	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00
2019	0.00	0.00	0.00	0.33	1.00	1.00	1.00	1.00
2020	0.00	0.00	0.00	0.22	1.00	1.00	1.00	1.00
Average	0.00	0.00	0.00	0.25	0.98	1.00	1.00	1.00

Proportion of legal (CW>4.75 inch) crab in Observer survey

size class	64	74	84	94	104	114	124	134
2012	0.00	0.01	0.02	0.22	0.90	1.00	1.00	1.00
2013	0.00	0.00	0.00	0.44	0.98	1.00	1.00	1.00
2014	0.00	0.00	0.00	0.22	0.91	1.00	1.00	1.00
2015	0.00	0.00	0.00	0.38	0.98	1.00	1.00	1.00
2016	0.00	0.00	0.00	0.46	1.00	1.00	1.00	1.00
2017	0.00	0.00	0.00	0.13	0.91	1.00	1.00	1.00
2018	0.00	0.00	0.00	0.16	0.95	0.99	1.00	1.00
2019	0.00	0.00	0.00	0.18	0.93	1.00	1.00	1.00
Average	0.00	0.00	0.00	0.30	0.95	1.00	1.00	1.00

The proportion of legal crab used in the assessment model is an average proportion based on observer survey data. In the assessment model, this proportion is used to estimate the number of retained crab in winter and summer commercial fisheries prior to 2008. It is assumed prior to 2008, all legal sized crab were retained.

Since 2008 commercially retained crab size is CW> 5.0 inches and retention probability is estimated from the observer survey.

The table below shows the proportion of legal vs. retained crab during the 2012-2019 observer survey, in response to request from the public.

Year		64	74	84	94	104	114	124	134
2012	Legal	0	0.01	0.02	0.22	0.9	1	1	1
	Retained	0	0	0	0.05	0.46	0.63	0.64	0.85
2013	Legal	0	0	0	0.44	0.98	1	1	1
	Retained	0	0	0	0.14	0.86	0.99	1	1
2014	Legal	0	0	0	0.22	0.91	1	1	1
	Retained	0	0	0	0.04	0.74	0.97	0.99	1
2015	Legal	0	0	0	0.38	0.98	1	1	1
	Retained	0	0	0	0.11	0.74	0.91	0.94	0.89
2016	Legal	0	0	0	0.46	1	1	1	1
	Retained	0	0	0	0.13	0.89	0.99	1	1
2017	Legal	0	0	0	0.12	0.91	1	1	1
	Retained	0	0	0	0.02	0.75	0.99	1	1
2018	Legal	0	0	0	0.16	0.95	0.99	1	1
	Retained	0	0	0	0.14	0.92	0.99	1	0.99
2019	Legal	0	0	0	0.18	0.93	1	1	1
	Retained	0	0	0	0.15	0.93	1	1	1

The proportion of legal sized crab retained from observer survey 2012-2019

Year	64	74	84	94	104	114	124	134
2012	0	0	0	0.23	0.51	0.63	0.64	0.85
2013	0	0	0	0.31	0.88	0.99	1	1
2014	0	0	0	0.19	0.82	0.97	0.99	1
2015	0	0	0	0.28	0.76	0.91	0.94	0.89
2016	0	0	0	0.28	0.89	0.99	1	1
2017	0	0	0	0.14	0.82	0.99	1	1
2018	0	0	0	0.87	0.98	1	1	0.99
2019	0	0	0	0.86	1	1	1	1

The above data justifies using logistic function as selection criteria.

Fishery	Model retention	Data
Summer: 1977-2007	Logistic retention prob	Discard, retained size prop
Summer: 2008-2022	Logistic retention prob	Total, retained size prop
Winter: 1977-2007	Mean legal crab proportion	No data
Winter: 2008-2022	Logistic retention prob	Retained size prop
Winter: Subsistence	All crab > 94mm retained	No data (No legal size limit)

3. Winter commercial pot selectivity is the same as the selectivity of the winter pot survey.

This assumption is based on the survey pot being similar to the one used for subsistence, and that many commercial fishermen are also subsistence harvesters. However, by regulation winter commercial king crab pots can be any dimension (5AAC 34.925(d)) and recent popularity of winter commercial fishery may have deviated this assumption.

4. Winter subsistence fishery retains crab size greater than 94 mm CL.

This was based on the assumption that subsistence fishermen would keep crab smaller than legal crab size. By regulation, subsistence fishery had no size limit for retention. Size of crab caught by subsistence fishery has never been monitored.

5. Discards handling mortality rate for all fisheries is 20%.

Discards mortality rate was specified by CPT. No empirical estimates are available.

Model data weighting

Survey data	Input sample size
Summer commercial, winter pot, and summer observer	minimum of $0.1 \times$ actual sample size or 10
Summer trawl and pot survey	minimum of $0.5 \times$ actual sample size or 20
Tag recovery	$0.5 \times$ actual sample size

Recruitment SD: 0.5.

Discards CV: 0.3

“Implied” effective sample sizes were calculated as

$$n = \sum_l \hat{P}_{y,l} (1 - \hat{P}_{y,l}) / \sum_l (P_{y,l} - \hat{P}_{y,l})^2$$

Where $P_{y,l}$ and $\hat{P}_{y,l}$ are observed and estimated length compositions in year y and length group l , respectively. Estimated implied effective sample sizes vary greatly over time.

Data-weighting for NSRKC model is aimed at achieving a balance between various data sets. The current model data weighting schemes, although arbitrary, were deemed appropriate by the CPT-SSC (NPFMC 2011, 2012, See Section E. 1. *Historical Model configuration progression* section). As illustrated in the figure below, increasing weight of size composition data (input sample size: from minimum) would lower model fit to the trawl survey abundance data.




Figure E. Model 21.0 default input sample size (20: trawl, 10: others) (black) vs. increased input sample size (200, 100) (blue dash line), and reduce input size (10, 5) (red dash line).

Thus far, there is no objective criteria for determining the balance (i.e., how much a model should fit observed trawl abundance data vs. size composition data). The author has tried alternative weighting schemes (NPFMC 2019, 2020, 2021) and found current ones are most appropriate.

Changes of assumptions since last assessment:

None

3. Model Selection and Evaluation

- a. Description of alternative model configurations.

For the 2024 draft assessment, the following alternative models are presented.

Model 21.0: Adopted by CPT-SSC for the 2023 final assessment.

Model 23.0: M estimated for all length classes (SSC request)

Model	M
21.0	0.18+est (L)
23.0	Est

4. Results

Same to the results from the 2016 (NPFMC 2016), model 23.0 estimated higher $M = 0.4116$ that was slightly lower than $M = 0.42$ estimated in 2015. Similar also to the 2015, the biggest difference between the two models (21.0 vs. 23.0) were higher likelihoods for trawl and commercial catch size-shell composition and tag-recovery size distribution. The single M model had lower fit than the default (21.0) model.

For selection of an assessment model for the final draft, a selection criterion is whether to choose better model fit (21.0) or biological authenticity (23.0). The author recommends model 21.0 over 23.0 for better model fit, which was the same recommendation that the CPT and SSC made in 2015.

Evaluation of negative log-likelihood values.

Model	21.0	23.0
Additional Parameters		0
AIC change		
Total	362.3	375.9
TSA	11.0	12.1
DIS	3.4	3.2
St.CPUE	-14.8	-14.8
TLP	134.0	142.2
WLP	39.6	40.1
CLP	49.5	54.4
OBS	24.3	28.0
WCLP	2.8	2.2
REC	19.4	20.7
TAG	85.0	87.8
M	0.18 0.615	0.4116
RMSE Trawl	0.34	0.33
RMSE CPUE	0.44	0.44
Total OFL	0.72	1.21

TSA: Trawl Survey Abundance

St. CPUE: Summer commercial catch standardized CPUE

TLP: Trawl survey length composition:

WLP: Winter pot survey length composition

CLP: Summer commercial retention catch length composition

REC: Recruitment deviation

OBS: Summer commercial catch observer discards and total catch length composition

TAG: Tagging recovery data composition

WCLP: Winter commercial length-shell composition

DIS: Summer commercial discards abundance

F. Calculation of the OFL

1. Specification of the Tier level and stock status.

NSRKC stock is placed in Tier 4. It is not possible to estimate the spawner-recruit relationship, but some abundance and harvest estimates are available to build a computer simulation model that captures the essential population dynamics. Tier 4 stocks are assumed to have reliable estimates of current survey biomass and instantaneous M ; however, the estimates of M for NSRKC stock are uncertain.

At the Tier 4 level the OFL is determined by the F_{MSY} proxy, B_{MSY} proxy, and estimated legal male abundance and biomass:

Level	Criteria	F_{OFL}
a	$B / B_{MSY proxy} > 1$	$F_{OFL} = \gamma M$
b	$\beta < B / B_{MSY proxy} \leq 1$	$F_{OFL} = \gamma M (B / B_{MSY proxy} - \alpha) / (1 - \alpha)$
c	$B / B_{MSY proxy} \leq \beta$	$F_{OFL} = \text{bycatch mortality \& directed fishery } F = 0$

where B is a mature male biomass (MMB), B_{MSY} proxy is average mature male biomass over a specified time period, $M = 0.18$, $\gamma = 1$, $\alpha = 0.1$, and $\beta = 0.25$.

For NSRKC, MMB is defined as the biomass of males > 94 mm CL on February 01 (Appendix A). B_{MSY} proxy is

B_{MSY} proxy = average model estimated MMB from 1980-2024.

Estimated B_{MSY} proxy: To be reported in Jan 2024

Predicted mature male biomass in 2024 on February 01

Mature male biomass: To be reported in Jan 2024

Since the projected MMB is above B_{MSY} proxy,

The NSRKC status is Tire 4a

And F_{OFL} for calculation of the OFL is

$F_{OFL} = \gamma \cdot M$ for M is length invariant of 0.18 and $F_{OFL,l} = \gamma \cdot M_l$ for length-dependent M

2. Calculation formula of NSRKC OFL.

OFL of NSRKC is total OFL (OFL_T) that is a sum of the retained and unretained OFL (OFL_r , OFL_{nr}).

$$OFL_T = OFL_r + OFL_{ur}$$

where

$$OFL_r = \text{retained_}B \cdot F_{OFL} \text{ and } OFL_{ur} = \text{unretained_}B \cdot F_{OFL} \cdot hm$$

retained_B is a biomass of crab subject to fisheries that is a sum of the products of crab abundance ($N_{w,l} + O_{w,l}$), fishery selectivity ($S_{s,l}$), retention probability ($S_{r,l}$), and average weight lb (wm_l) by length class (l).

$$\text{retained_}B = \sum_l (N_{w,l} + O_{w,l}) S_{s,l} S_{r,l} wm_l$$

unretained_B is a biomass of crab subject to fisheries and is a sum of the products of crab abundance ($N_{w,l} + O_{w,l}$), fishery selectivity ($S_{s,l}$), 1 minus retention probability ($S_{r,l}$), and average weight lb (wm_l) by length class (l).

$$\text{unretained_}B = \sum_l (N_{w,l} + O_{w,l}) S_{s,l} (1 - S_{r,l}) wm_l$$

hm is handling mortality, default 0.2

The NSRKC fishery consists of two distinct fisheries: winter and summer. The two fisheries are discontinuous with 5 months (0.42 year) between the two fisheries during which natural mortality occurs. To estimate the OFL for the two fisheries, the CPT in 2016 recommended the following formula that the sum of winter and summer catch (H_w, H_s) equals total OFL ($OFL = H_w + H_s$) and that winter catch is a fraction (p) of total OFL: $H_w = p \cdot OFL$, where ***p* is predetermined fraction of** the winter fishery to total fishery. In NSRKC fishery $p = 0.16$ is used.

$$H_w = B_w (1 - \exp(-x \cdot F_{OFL})),$$

$$H_s = B_s (1 - \exp((1-x) \cdot F_{OFL})), \text{ and}$$

$$B_s = (B_w - H_w) e^{-0.42 \cdot M}$$

where

B_w is the winter NSRKC biomass, B_s is the summer NSRKC biomass, and x is a fraction parameter,

Solving x of the above (see Appendix A for derivation), retained and unretained OFL is calculated as:

(1)

and

$$OFL_{nr} = \text{unretained_}B_w \cdot FOFL_a \cdot hm$$

$$\text{where } FOFL_a = \left(1 - e^{-(F_{OFL} + 0.42M)} - (1 - e^{-0.42M}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL} + 0.42M)})}{1 - p \cdot (1 - e^{-0.42M})} \right) \right) \quad (2)$$

Because M of NSRKC is length-dependent, the proper calculation of NSRKC OFL should account for length-dependent M as:

$$OFL_r = \sum_l [\text{retained_}B_{w,l} \cdot FOFL_{a,l}]$$

$$\text{where } FOFL_{a,l} = \left(1 - e^{-(F_{OFL,l} + 0.42M_l)} - (1 - e^{-0.42M_l}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL,l} + 0.42M_l)})}{1 - p \cdot (1 - e^{-0.42M_l})} \right) \right) \quad (3)$$

and

$$OFL_{ur} = \sum_l [\text{unretained_}B_{w,l} \cdot FOFL_{a,l}] \cdot hm$$

$$\text{where } FOFL_{a,l} = \left(1 - e^{-(F_{OFL,l} + 0.42M_l)} - (1 - e^{-0.42M_l}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL,l} + 0.42M_l)})}{1 - p \cdot (1 - e^{-0.42M_l})} \right) \right) \quad (4)$$

where M_l is a size specific natural mortality,

Determination of Total catch OFL

Total catch OFL is calculable by adding retained and unretained portion of the OFL (i.e., Total OFL = $OFL_r + OFL_{nr}$). Standard calculation of OFL for Tier 4 crab is $F_{OFL} = \gamma M = \gamma 0.18$.

3. Determination of NSRKC OFL for the 2024 fishery season.

Projected NSRKC biomass catchable to fishery in 2024

Length independent F_{OFL} .

Retained catch OFL (OFL_r) = To be reported in Jan 2024
Unretained catch OFL (OFL_{nr}) = To be reported in Jan 2024
Total catch OFL (OFL_T) = To be reported in Jan 2024

G. Calculation of the ABC

1. Specification of the probability distribution of the OFL.

ABC is calculated as $(1-ABC \text{ buffer}) \cdot OFL$

For 2024 fishery season, CPT recommended 30 % buffer.
ABC for the 2024 fishery is

ABC = To be reported in Jan 2024.

H. Rebuilding Analyses

Not applicable

I. Data Gaps and Research Priorities

The major data gap of NSRKC is an incomplete understanding of NSRKC biology, including natural mortality, and fate of oldshell and large crabs. Additionally, research should focus on female abundance and fecundity as well as their reproductive potential. As for management, the number and length-shell composition of unretained crab in fisheries are needed for calculation of total catch. Viability of unretained crab need to be researched to examine current default 20% mortality. Incorporation of local and traditional knowledge (LK/TK) and socio-economic impacts of NSRKC fisheries on the region, could bring further insights about NSRKC biology and management.

Acknowledgments

We thank many ADF&G, CPT, SSC for review of the assessment model and suggestions for improvements and diagnoses.

J. References

Bell, J., J. M. Leon, T. Hamazaki, S. Kent, and W. W. Jones. 2016. Red king crab movement, growth, and size composition within eastern Norton Sound, Alaska, 2012-2014. Alaska

Department of Fish and Game, Fishery Data Series No. 16-37, Anchorage.

- Bell, J. and T. Hamazaki 2019. Summary of 2017 and 2018 Norton Sound red king crab bottom trawl survey. Fishery Data Series No. 19-33. Alaska Department of Fish and Game Anchorage
- Fournier, D., and C.P. Archibald. 1982. A general theory for analyzing catch at age data. *Can. J. Fish. Aquat. Sci.* 39:1195-1207.
- Fournier, D.A., H.J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J. Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim. Methods Softw.* 27:233-249.
- Grant, S.W. and W. Chen. 2012. Incorporating deep and shallow components of genetic structure into the management of Alaskan red king crab. *Evolutionary Applications.* 5:820-837.
- Menard, J., J. Soong, and S. Kent 2011. 2009 Annual Management Report Norton Sound, Port Clarence, and Kotzebue. Fishery Management Report No. 11-46. Alaska Department of Fish and Game Anchorage
- Methot, R.D. 1989. Synthetic estimates of historical abundance and mortality for northern anchovy. *Amer. Fish. Soc. Sym.* 6:66-82.
- Mohn, R. 1999. The retrospective problem in sequential population analysis: An investigation using cod fishery and simulated data. *ICES Journal of Marine Science*, 56:473-488.
- NPFMC 1998. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 1998 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2010. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2010 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2011. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2011 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2012. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2012 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2013. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2013 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2014. Stock assessment and fishery evaluation report for the King and Tanner crab

- fisheries of the Bering Sea and Aleutian Islands regions. 2014 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2015. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2015 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2016. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2016 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2017. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2017 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2018. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2018 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2019. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2019 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- NPFMC 2020. Stock assessment and fishery evaluation report for the King and Tanner crab fisheries of the Bering Sea and Aleutian Islands regions. 2020 Crab SAFE. North Pacific Fishery Management Council, Anchorage, AK, USA.
- Powell, G.C., R. Peterson, and L. Schwarz. 1983. The red king crab, *Paralithodes camtschatica* (Tilesius), in Norton Sound, Alaska: History of biological research and resource utilization through 1982. Alaska Dept. Fish and Game, Inf. Leaflet. 222. 103 pp.
- Zheng, J., G.H. Kruse, and L. Fair. 1998. Use of multiple data sets to assess red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska: A length-based stock synthesis approach. Pages 591-612 *In* Fishery Stock Assessment Models, edited by F. Funk, T.J. Quinn II, J. Heifetz, J.N. Ianelli, J.E. Powers, J.F. Schweigert, P.J. Sullivan, and C.-I. Zhang, Alaska Sea Grant College Program Report No. AK-SG-98-01, University of Alaska Fairbanks.

Table 2. Historical winter commercial and subsistence red king crab fisheries, Norton Sound Section, eastern Bering Sea. Bold typed data are used for the assessment model.

Model Year	Year ^a	Commercial		Winter ^b	Subsistence			Total Crab	
		# of Fishers	# of Crab Harvested		Permits			Caught ^c	Retained ^d
					Issued	Returned	Fished		
1978	1978	37	9,625	1977/78	290	206	149	NA	12,506
1979	1979	1 ^f	221^f	1978/79	48	43	38	NA	224
1980	1980	1 ^f	22^f	1979/80	22	14	9	NA	213
1981	1981	0	0	1980/81	51	39	23	NA	360
1982	1982	1 ^f	17^f	1981/82	101	76	54	NA	1,288
1983	1983	5	549	1982/83	172	106	85	NA	10,432
1984	1984	8	856	1983/84	222	183	143	15,923	11,220
1985	1985	9	1,168	1984/85	203	166	132	10,757	8,377
1986	1985/86	5	2,168	1985/86	136	133	107	10,751	7,052
1987	1986/87	7	1,040	1986/87	138	134	98	7,406	5,772
1988	1987/88	10	425	1987/88	71	58	40	3,573	2,724
1989	1988/89	5	403	1988/89	139	115	94	7,945	6,126
1990	1989/90	13	3,626	1989/90	136	118	107	16,635	12,152
1991	1990/91	11	3,800	1990/91	119	104	79	9,295	7,366
1992	1991/92	13	7,478	1991/92	158	105	105	15,051	11,736
1993	1992/93	8	1,788	1992/93	88	79	37	1,193	1,097
1994	1993/94	25	5,753	1993/94	118	95	71	4,894	4,113
1995	1994/95	42	7,538	1994/95	166	131	97	7,777	5,426
1996	1995/96	9	1,778	1995/96	84	44	35	2,936	1,679
1997	1996/97	2 ^f	83^f	1996/97	38	22	13	1,617	745
1998	1997/98	5	984	1997/98	94	73	64	20,327	8,622
1999	1998/99	5	2,714	1998/99	95	80	71	10,651	7,533
2000	1999/00	10	3,045	1999/00	98	64	52	9,816	5,723
2001	2000/01	3	1,098	2000/01	50	27	12	366	256
2002	2001/02	11	2,591	2001/02	114	61	45	5,119	2,177
2003	2002/03	13	6,853	2002/03	107	70	61	9,052	4,140
2004	2003/04	2 ^f	522^f	2003/04 ^h	96	77	41	1,775	1,181
2005	2004/05	4	2,091	2004/05	170	98	58	6,484	3,973
2006	2005/06	1 ^f	75^f	2005/06	98	97	67	2,083	1,239
2007	2006/07	8	3,313	2006/07	129	127	116	21,444	10,690
2008	2007/08	9	5,796	2007/08	139	137	108	18,621	9,485
2009	2008/09	7	4,951	2008/09	105	105	70	6,971	4,752
2010	2009/10	10	4,834	2009/10	125	123	85	9,004	7,044
2011	2010/11	5	3,365	2010/11	148	148	95	9,183	6,640
2012	2011/12	35	9,157	2011/12	204	204	138	11,341	7,311
2013	2012/13	26	22,639	2012/13	149	148	104	21,524	7,622
2014	2013/14	21	14,986	2013/14	103	103	75	5,421	3,252
2015	2014/15	44	41,062	2014/15	155	153	107	9,840	7,651
2016	2015/16	25	29,792	2015/16	139	97	64	6,468	5,340
2017	2017	43	26,008	2017	163	163	109	7,185	6,039
2018	2018	28	9,180	2018	123	120	82	5,767	4,424
2019	2019	6	1,050	2019	101	101	60	2,080	1,545
2020	2020	1	conf	2020	79	79	50	813	548
2021	2021	5	320	2021	103	103	76	4,655	2,892
2022	2022	8	2,424	2022	113	63	42	10,686	7,630
2023	2023	2	conf	2023					

a Prior to 1985 the winter commercial fishery occurred from January 1 - April 30. As of March 1985, fishing may occur from November 15 - May 15.

b The winter subsistence fishery occurs during months of two calendar years (as early as December, through May).

c The number of crab actually caught, **including females**

d The number of crab retained is the number of crab caught and kept, **including females**

f Confidentiality was waived by the fishers.9963+

h Prior to 2005, permits were only given out of the Nome ADF&G office. Starting with the 2004-5 season, permits were given out in Elim, Golovin, Shaktoolik, and White Mountain.

Table 3. Summary of Norton Sound red king crab trawl survey abundance estimates (x 1000) (CL ≥ 64mm). NMFS and ADF&G trawl survey abundance estimate is based on 10×10 nm² grids, and NBS trawl survey is based on 20×20 nm² grids. Bold typed data are used for the assessment model.

Year	Dates	Survey Agency	Survey method	Abundance ≥64 mm		Female			% clutch full 95% CI
				CV	N	% barren	% clutch full		
1976	9/02 - 9/25	NMFS	Trawl	4301.8	0.31	181	2.6	66.7	62.4-71.0
1979	7/26 - 8/05	NMFS	Trawl	1457.4	0.22	42	25.0	79.9	64.8-94.8
1980	7/04 - 7/14	ADF&G	Pots	2092.3	N/A				
1981	6/28 - 7/14	ADF&G	Pots	2153.4	N/A				
1982	7/06 - 7/20	ADF&G	Pots	1140.5	N/A				
1982	9/05 - 9/11	NMFS	Trawl	3548.9	0.25	269	0	84.3	81.5-87.2
1985	7/01 - 7/14	ADF&G	Pots	2320.4	0.08				
1985	9/16 -10/01	NMFS	Trawl	2424.9	0.26	151	0	87.5	NA
1988	8/16 - 8/30	NMFS	Trawl	2702.3	0.29	219	1.0	80.7	77.3-84.2
1991	8/22 - 8/30	NMFS	Trawl	3132.5	0.43	105	0	69.3	57.7-80.8
1996	8/07 - 8/18	ADF&G	Trawl	1283.0	0.25	168	30.8	71.9	65.9-77.9
1999	7/28 - 8/07	ADF&G	Trawl	2608.0	0.24	81	4.7	80.4	76.0-84.7
2002	7/27 - 8/06	ADF&G	Trawl	2056.0	0.36	168	4.7	76.8	73.4-80.2
2006	7/25 - 8/08	ADF&G	Trawl	3336.0	0.39	194	3.6	67.3	63.2-71.5
2008	7/24 - 8/11	ADF&G	Trawl	2894.2	0.31	116	3.3	56.1	48.5-61.7
2010	7/27 - 8/09	NBS	Trawl	1980.1	0.44	28	0	70.2	63.8-78.5
2011	7/18 - 8/15	ADF&G	Trawl	3209.3	0.29	135	9.8	67.2	61.7-72.6
2014	7/18 - 7/30	ADF&G	Trawl	5934.6	0.47	60	0	60.4	54.3-66.6
2017	7/28 - 8/08	ADF&G	Trawl	1762.1	0.22	43	21.4	71.6	60.0-82.7
2017	8/18 - 8/29	NBS	Trawl	1035.8	0.40	58	0	80.0	72.5-87.5
2018	7/22 - 7/29	ADF&G	Trawl	1108.9	0.25	424	15.8	76.3	59.7-83.5
2019	7/17-7/29	ADF&G	Trawl	4660.8	0.60	386	47.8	50.6	43.1-56.4
2019	8/04-8/07	NBS	Trawl	2532.4	0.26	94	17.6	47.9	36.8-58.9
2020	7/31-8/14	ADF&G	Trawl	1716.5	0.27	186	4.5	66.2	61.6-70.8
2021	7/19-8/03	ADF&G	Trawl	2400.0	0.60	90	3.4	59.8	54.9-64.6
2021	7/29-8/07	NBS	Trawl	2370.0	0.43	139	2.6	61.1	58.8-63.4
2022	8/03-8/12	NBS	Trawl	2103.0	0.37	387	3.5	66.5	64.2-68.7
2023	7/21-7/30	ADF&G	Trawl	3441.7	0.33	47	0	80.0	74.1-85.8

Abundance of NMFS survey was estimated by NMFS, by multiplying the mean CPUE (# NRKC/nm²) across all hauls (including re-tows) to a standard survey area (7600nm²). Abundance of ADF&G and NBS survey was estimated by ADF&G by multiplying CPUE (# NRKC/nm²) of each station to the grid represented by the station and summing across all surveyed station (ADF&G: 4700 – 5200nm². NBS 5841 nm²).

Norton Sound Red King Crab Stock Assessment Sept 2023

%barren is calculated by dividing the number of mature females with no eggs by total number of mature females.

Mean and 95% CI of % clutch full is calculated among non-barren mature females. Clutch fullness of each non-barren female was assigned by fullness index that was converted to percentage in the table below.

Clutch fullness index of both NMFS-NBS and ADF&G were converted as follows

NMFS and NBS Code	NMFS and NBS Fullness	Assigned %	ADF&G code	ADF&G Fullness	Assigned %
1	barren	0	1	barren	
2	0-1/8	6.25	2	0 (post-release)	
3	1/8-1/4	18.75	3	1-29%	15
4	1/4 – 1/2	27.5	4	30-59%	45
5	1/2 – 3/4	62.5	5	60-89%	75
6	3/4 – 1	87.5	6	90-100%	95
7	>1	100			

Table 4. Summer commercial retained catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1977	1549	0	0	0	0.00	0.42	0.34	0.08	0.05	0	0	0	0.00	0.06	0.04	0.01	0.00
1978	389	0	0	0	0.01	0.19	0.47	0.26	0.04	0	0	0	0.00	0.01	0.01	0.01	0.00
1979	1660	0	0	0	0.03	0.23	0.38	0.26	0.07	0	0	0	0.00	0.03	0.00	0.00	0.01
1980	1068	0	0	0	0.00	0.10	0.31	0.37	0.18	0	0	0	0.00	0.00	0.01	0.02	0.01
1981	1784	0	0	0	0.00	0.07	0.15	0.28	0.23	0	0	0	0.00	0.00	0.05	0.12	0.09
1982	1093	0	0	0	0.04	0.19	0.16	0.22	0.29	0	0	0	0.00	0.01	0.02	0.03	0.03
1983	802	0	0	0	0.04	0.41	0.36	0.06	0.03	0	0	0	0.00	0.04	0.01	0.02	0.02
1984	963	0	0	0	0.10	0.42	0.28	0.06	0.01	0	0	0	0.01	0.07	0.05	0.01	0.00
1985	2691	0	0	0.00	0.06	0.31	0.37	0.15	0.02	0	0	0	0.00	0.03	0.03	0.01	0.00
1986	1138	0	0	0	0.03	0.36	0.39	0.12	0.02	0	0	0	0.00	0.02	0.04	0.02	0.00
1987	1985	0	0	0	0.02	0.18	0.29	0.27	0.11	0	0	0	0.00	0.03	0.06	0.03	0.01
1988	1522	0	0.00	0	0.02	0.20	0.30	0.18	0.04	0	0	0	0.01	0.06	0.10	0.07	0.02
1989	2595	0	0	0	0.01	0.16	0.32	0.17	0.05	0	0	0	0.00	0.06	0.12	0.09	0.02
1990	1289	0	0	0	0.01	0.14	0.35	0.26	0.07	0	0	0	0.00	0.04	0.07	0.05	0.01
1991																	
1992	2566	0	0	0	0.02	0.20	0.27	0.14	0.09	0	0	0	0.00	0.08	0.13	0.06	0.02
1993	17804	0	0	0	0.01	0.23	0.39	0.23	0.03	0	0	0	0.00	0.02	0.04	0.03	0.01
1994	404	0	0	0	0.02	0.09	0.08	0.07	0.02	0	0	0	0.02	0.19	0.25	0.20	0.05
1995	1167	0	0	0	0.04	0.26	0.29	0.15	0.05	0	0	0	0.01	0.05	0.07	0.06	0.01
1996	787	0	0	0	0.03	0.22	0.24	0.09	0.05	0	0	0	0.01	0.12	0.14	0.08	0.02
1997	1198	0	0	0	0.03	0.37	0.34	0.10	0.03	0	0	0	0.00	0.06	0.04	0.03	0.01
1998	1055	0	0	0	0.03	0.23	0.24	0.08	0.03	0	0	0	0.02	0.11	0.14	0.08	0.03
1999	562	0	0	0	0.06	0.29	0.24	0.18	0.09	0	0	0	0.00	0.02	0.05	0.04	0.00
2000	17213	0	0	0	0.02	0.30	0.39	0.11	0.02	0	0	0	0.00	0.05	0.07	0.04	0.01
2001	20030	0	0	0	0.02	0.22	0.37	0.21	0.07	0	0	0	0.00	0.02	0.05	0.02	0.01
2002	5219	0	0	0	0.04	0.23	0.28	0.25	0.07	0	0	0	0.00	0.03	0.04	0.03	0.01
2003	5226	0	0	0	0.02	0.37	0.32	0.12	0.03	0	0	0	0.00	0.02	0.05	0.05	0.01
2004	9606	0	0	0	0.01	0.38	0.39	0.11	0.03	0	0	0	0.00	0.03	0.03	0.01	0.01
2005	5360	0	0	0	0.00	0.25	0.47	0.16	0.02	0	0	0	0.00	0.02	0.05	0.02	0.01
2006	6707	0	0	0	0.00	0.18	0.35	0.17	0.02	0	0	0	0.00	0.05	0.14	0.07	0.01
2007	6125	0	0	0	0.01	0.36	0.34	0.14	0.03	0	0	0	0.00	0.02	0.06	0.03	0.01
2008	5766	0	0	0	0.00	0.35	0.35	0.06	0.01	0	0	0	0.00	0.09	0.09	0.04	0.01
2009	6026	0	0	0	0.01	0.34	0.33	0.11	0.02	0	0	0	0.00	0.08	0.08	0.02	0.01
2010	5902	0	0	0	0.01	0.39	0.36	0.10	0.01	0	0	0	0.00	0.05	0.05	0.02	0.00
2011	2552	0	0	0	0.00	0.32	0.40	0.12	0.02	0	0	0	0.00	0.06	0.06	0.02	0.00
2012	5056	0	0	0	0.00	0.24	0.46	0.18	0.02	0	0	0	0.00	0.03	0.04	0.02	0.00
2013	6072	0	0	0	0.00	0.24	0.37	0.24	0.06	0	0	0	0.00	0.01	0.04	0.02	0.00
2014	4682	0	0	0	0.01	0.28	0.24	0.18	0.07	0	0	0	0.00	0.04	0.09	0.07	0.02
2015	4173	0	0	0	0.01	0.48	0.28	0.10	0.03	0	0	0	0.00	0.02	0.03	0.03	0.01
2016	1543	0	0	0	0.00	0.25	0.47	0.16	0.03	0	0	0	0.00	0.02	0.02	0.03	0.01
2017	3412	0	0	0	0.00	0.18	0.39	0.21	0.03	0	0	0	0.01	0.03	0.12	0.05	0.01
2018	2609	0	0	0	0.00	0.11	0.32	0.32	0.08	0	0	0	0	0.01	0.08	0.08	0.02
2019	1136	0	0	0	0.01	0.32	0.23	0.13	0.03	0	0	0	0	0.02	0.10	0.14	0.03
2020																	
2021																	
2022	2981	0	0	0	0.02	0.46	0.30	0.03	0.00	0	0	0	0.00	0.12	0.05	0.01	0.00
2023	2458	0	0	0	0.00	0.26	0.42	0.13	0.01	0	0	0	0.00	0.07	0.09	0.01	0.02

Table 5. Winter commercial catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
2015	576	0	0	0	0.07	0.50	0.24	0.06	0.01	0	0	0	0.01	0.04	0.03	0.03	0.01
2016	1016	0	0	0	0.03	0.45	0.31	0.03	0.00	0	0	0	0.01	0.09	0.04	0.02	0.01
2017	540	0	0	0	0.00	0.20	0.30	0.13	0.02	0	0	0	0.00	0.08	0.19	0.06	0.02
2018	401	0	0	0	0.00	0.11	0.25	0.27	0.05	0	0	0	0.04	0.16	0.10	0.02	

Table 6. Summer Trawl Survey length-shell compositions.

Year	Survey	Sample	New Shell										Old Shell							
			64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+		
1976	NMFS	1326	0.01	0.02	0.10	0.19	0.34	0.18	0.02	0.00	0.00	0.00	0.01	0.02	0.03	0.04	0.01	0.01		
1979	NMFS	220	0.01	0.01	0.00	0.02	0.05	0.05	0.03	0.01	0.01	0.00	0.01	0.04	0.14	0.40	0.19	0.03		
1982	NMFS	327	0.22	0.07	0.16	0.23	0.17	0.03	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.02	0.02	0.03		
1985	NMFS	350	0.11	0.11	0.19	0.17	0.16	0.06	0.01	0.00	0.00	0.00	0.00	0.02	0.05	0.08	0.05	0.01		
1988	NMFS	366	0.16	0.19	0.12	0.13	0.11	0.06	0.03	0.00	0.00	0.00	0.01	0.01	0.03	0.07	0.05	0.03		
1991	NMFS	340	0.18	0.08	0.02	0.03	0.06	0.03	0.01	0.01	0.03	0.06	0.02	0.08	0.16	0.14	0.09	0.02		
1996	ADF&G	269	0.29	0.21	0.13	0.09	0.05	0.00	0.00	0.01	0.00	0.00	0.03	0.03	0.04	0.04	0.04	0.03		
1999	ADF&G	283	0.03	0.01	0.10	0.29	0.26	0.13	0.03	0.01	0.00	0.00	0.00	0.03	0.05	0.04	0.02	0.00		
2002	ADF&G	244	0.09	0.12	0.14	0.11	0.02	0.03	0.02	0.01	0.01	0.03	0.07	0.10	0.09	0.09	0.05	0.02		
2006	ADF&G	373	0.18	0.26	0.21	0.11	0.06	0.04	0.02	0.00	0.00	0.00	0.00	0.02	0.04	0.04	0.01	0.00		
2008	ADF&G	275	0.12	0.15	0.21	0.11	0.10	0.03	0.02	0.01	0.00	0.01	0.04	0.06	0.08	0.01	0.04	0.00		
2010	NOAA	69	0.01	0.04	0.06	0.17	0.06	0.03	0.00	0.00	0.00	0.03	0.09	0.20	0.19	0.07	0.03	0.01		
2011	ADF&G	315	0.13	0.11	0.09	0.11	0.18	0.14	0.03	0.01	0.00	0.00	0.01	0.02	0.09	0.04	0.03	0.00		
2014	ADF&G	387	0.08	0.15	0.24	0.18	0.09	0.02	0.01	0.01	0.00	0.00	0.03	0.10	0.05	0.04	0.01	0.00		
2017	ADF&G	116	0.14	0.12	0.05	0.09	0.10	0.04	0.00	0.00	0.01	0.02	0.02	0.02	0.07	0.18	0.04	0.00		
2017	NOAA	58	0.09	0.10	0.14	0.05	0.05	0.05	0.05	0.03	0.03	0.00	0.03	0.05	0.03	0.19	0.05	0.03		
2018	ADF&G	73	0.37	0.10	0.11	0.03	0.01	0.03	0.04	0.01	0	0.07	0.01	0.04	0.03	0.03	0.10	0.03		
2019	ADF&G	307	0.55	0.30	0.03	0	0.00	0.00	0.00	0	0.00	0.00	0.01	0.02	0.01	0.02	0.03	0.01		
2019	NOAA	135	0.36	0.30	0.08	0.04	0.01	0	0.01	0.01	0.04	0.01	0.04	0.02	0.01	0.01	0.04	0.01		
2020	ADF&G	111	0.13	0.22	0.30	0.06	0.05	0.01	0	0	0.03	0.08	0.05	0.02	0.02	0.02	0	0.01		
2021	ADF&G	158	0.06	0.17	0.22	0.22	0.22	0.04	0.01	0.01	0	0.01	0	0.02	0.01	0.01	0.01			
2021	NOAA	82	0.05	0.16	0.21	0.16	0.10	0.02	0	0	0.01	0.05	0.11	0.06	0.06	0.01	0	0		
2022	NOAA	378	0.16	0.17	0.11	0.10	0.07	0.03	0.01	0.01	0.02	0.02	0.07	0.08	0.087	0.05	0.02	0.01		
2023	ADF&G	240	0.00	0.03	0.09	0.20	0.21	0.07	0.00	0	0	0	0.01	0.03	0.17	0.16	0.03	0		

Table 7. Winter pot survey length-shell compositions.

Year	CPUE	Sample	New Shell									Old Shell								
			64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+		
1981/82	NA	719	0.00	0.10	0.23	0.21	0.07	0.02	0.02	0.00	0.00	0.05	0.11	0.11	0.04	0.02	0.02	0.00		
1982/83	24.2	2583	0.03	0.08	0.28	0.28	0.21	0.07	0.01	0.00	0.00	0.00	0.00	0.00	0.02	0.01	0.01	0.01		
1983/84	24.0	1677	0.01	0.16	0.26	0.23	0.15	0.06	0.01	0.00	0.00	0.00	0.00	0.02	0.06	0.03	0.01	0.01		
1984/85	24.5	789	0.02	0.09	0.25	0.35	0.16	0.06	0.01	0.00	0.00	0.00	0.00	0.01	0.03	0.02	0.00	0.00		
1985/86	19.2	594	0.04	0.12	0.17	0.24	0.19	0.08	0.01	0.00	0.00	0.00	0.00	0.01	0.06	0.04	0.01	0.00		
1986/87	5.8	144	0.00	0.06	0.15	0.19	0.07	0.04	0.00	0.00	0.00	0.00	0.01	0.04	0.30	0.11	0.03	0.00		
1987/88																				
1988/89	13.0	500	0.02	0.13	0.15	0.13	0.19	0.17	0.03	0.00	0.00	0.00	0.00	0.05	0.08	0.03	0.00	0.00		
1989/90	21.0	2076	0.00	0.05	0.21	0.26	0.18	0.12	0.06	0.01	0.00	0.00	0.00	0.00	0.03	0.06	0.02	0.00		
1990/91	22.9	1283	0.00	0.01	0.09	0.29	0.27	0.10	0.01	0.00	0.00	0.00	0.00	0.00	0.03	0.12	0.07	0.02		
1992/93	5.5	181	0.00	0.01	0.03	0.06	0.13	0.12	0.03	0.00	0.00	0.00	0.00	0.02	0.19	0.27	0.10	0.05		
1993/94																				
1994/95	6.2	858	0.01	0.06	0.08	0.10	0.26	0.23	0.07	0.01	0.00	0.00	0.00	0.00	0.03	0.07	0.06	0.02		
1995/96	9.9	1580	0.06	0.14	0.20	0.19	0.11	0.07	0.03	0.00	0.00	0.00	0.00	0.01	0.06	0.07	0.03	0.01		
1996/97	2.9	398	0.07	0.21	0.22	0.11	0.15	0.11	0.05	0.01	0.00	0.00	0.00	0.00	0.02	0.03	0.01	0.01		
1997/98	10.9	881	0.00	0.14	0.41	0.27	0.05	0.02	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.02	0.02	0.01		
1998/99	10.7	1307	0.00	0.02	0.12	0.36	0.36	0.08	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.01	0.01	0.00		
1999/00	6.2	575	0.02	0.09	0.10	0.16	0.33	0.18	0.03	0.00	0.00	0.00	0.00	0.00	0.05	0.02	0.01	0.00		
2000/01	3.1	44																		
2001/02	13.0	828	0.05	0.29	0.26	0.17	0.06	0.06	0.04	0.01	0.01	0.00	0.01	0.01	0.00	0.01	0.00	0.00		
2002/03	9.6	824	0.02	0.10	0.22	0.28	0.18	0.06	0.02	0.00	0.00	0.01	0.01	0.02	0.02	0.03	0.02	0.01		
2003/04	3.7	296	0.00	0.02	0.16	0.26	0.32	0.14	0.01	0.00	0.00	0.00	0.01	0.02	0.02	0.01	0.02	0.01		
2004/05	4.4	405	0.00	0.07	0.14	0.18	0.22	0.19	0.07	0.00	0.00	0.00	0.00	0.00	0.04	0.06	0.01	0.00		
2005/06	6.0	512	0.00	0.14	0.23	0.21	0.16	0.05	0.02	0.00	0.00	0.01	0.01	0.02	0.04	0.07	0.03	0.01		
2006/07	7.3	159	0.07	0.14	0.19	0.35	0.13	0.04	0.00	0.00	0.00	0.00	0.01	0.01	0.02	0.04	0.00	0.00		
2007/08	25.0	3552	0.01	0.14	0.25	0.17	0.14	0.07	0.01	0.00	0.01	0.04	0.07	0.03	0.03	0.01	0.01	0.00		
2008/09	21.9	525	0.00	0.07	0.13	0.35	0.20	0.08	0.01	0.00	0.00	0.00	0.00	0.00	0.04	0.10	0.00	0.00		
2009/10	25.3	578	0.01	0.05	0.13	0.21	0.24	0.11	0.02	0.00	0.00	0.00	0.01	0.06	0.10	0.05	0.01	0.00		
2010/11	22.1	596	0.02	0.08	0.13	0.20	0.17	0.13	0.05	0.00	0.00	0.00	0.01	0.03	0.11	0.05	0.01	0.00		
2011/12	29.4	675	0.03	0.11	0.23	0.19	0.12	0.13	0.04	0.00	0.00	0.00	0.00	0.01	0.05	0.05	0.03	0.00		

Table 8. Summer commercial 1987-1994 observer discards length-shell compositions.

Year Sample	New Shell								Old Shell							
	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1987	1146	0.06	0.19	0.32	0.33	0.03	0.00	0.00	0.00	0.00	0.00	0.02	0.04	0.00	0.00	0.00
1988	722	0.01	0.04	0.15	0.48	0.14	0.00	0.00	0.00	0.00	0.01	0.03	0.10	0.04	0.00	0.00
1989	1000	0.07	0.19	0.24	0.22	0.03	0.00	0.00	0.00	0.02	0.03	0.07	0.11	0.03	0.00	0.00
1990	507	0.08	0.23	0.27	0.27	0.04	0.00	0.00	0.00	0.02	0.02	0.02	0.05	0.01	0.00	0.00
1992	580	0.11	0.17	0.30	0.29	0.03	0.00	0.00	0.00	0.01	0.02	0.02	0.04	0.01	0.00	0.00
1994	850	0.07	0.06	0.11	0.15	0.02	0.00	0.00	0.00	0.07	0.07	0.15	0.24	0.05	0.00	0.00

Table 9. Summer commercial observer total catch length-shell compositions.

Year Sample	New Shell								Old Shell							
	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
2012	3055	0.10	0.05	0.08	0.15	0.15	0.17	0.06	0.01	0.00	0.00	0.00	0.03	0.08	0.09	0.03
2013	4762	0.19	0.16	0.09	0.10	0.16	0.16	0.09	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.01
2014	3506	0.02	0.05	0.13	0.22	0.22	0.12	0.08	0.03	0.00	0.00	0.00	0.02	0.03	0.03	0.02
2015	1671	0.01	0.04	0.09	0.23	0.37	0.14	0.05	0.01	0.00	0.00	0.00	0.01	0.02	0.02	0.01
2016	2114	0.01	0.01	0.03	0.12	0.29	0.36	0.08	0.02	0.00	0.00	0.00	0.01	0.03	0.03	0.02
2017	2748	0.02	0.03	0.03	0.06	0.19	0.33	0.18	0.02	0.00	0.00	0.00	0.00	0.02	0.07	0.03
2018	1628	0.03	0.06	0.12	0.11	0.09	0.17	0.18	0.04	0.00	0.00	0.01	0.01	0.15	0.07	0.08
2019	236	0.13	0.06	0.06	0.13	0.08	0.05	0.01	0.01	0	0	0.00	0.04	0.11	0.14	0.14

Table 10. The observed proportion of tagged crab by each size class released and recovered after 1 -3 year of liberty 1980-2019 periods.

Year at liberty 1

	64-73	74-83	84-93	94-103	104-113	114-123	124-33	> 134	n
64-73	0	0.2	0.8	0	0	0	0	0	5
74-83		0	0.44	0.47	0.09	0	0	0	47
84-93			0	0.32	0.62	0.05	0.01	0	146
94-103				0.03	0.62	0.34	0.01	0.00	317
104-113					0.31	0.59	0.09	0	241
114-123						0.42	0.47	0.11	210
124-133							0.69	0.31	81
>134								1	26

Year at liberty 2

	64-73	74-83	84-93	94-103	104-113	114-123	124-33	> 134	n
64-73	0	0	0.09	0.55	0.36	0	0	0	11
74-83		0	0	0.11	0.85	0.04	0	0	113
84-93			0	0.04	0.32	0.61	0.03	0	114
94-103				0.02	0.36	0.41	0.20	0	94
104-113					0.06	0.71	0.22	0	108
114-123						0.17	0.72	0.11	65
124-133							0.36	0.64	25
>134								1	8

Year at liberty 3

	64-73	74-83	84-93	94-103	104-113	114-123	124-33	> 134	n
64-73	0	0	0	0	0.5	0.5	0	0	22
74-83	0	0	0	0	0.26	0.66	0.082	0	73
84-93	0	0	0	0.04	0.26	0.53	0.17	0	53
94-103	0	0	0	0	0.06	0.67	0.27	0	52
104-113	0	0	0	0	0	0.26	0.62	0.12	34
114-123	0	0	0	0	0	0	0.79	0.21	14
124-133	0	0	0	0	0	0	0.1	0.9	10
>134	0	0	0	0	0	0	0	1	1

Table 11. Summary of bounds and model estimated parameters for a length-based population model of Norton Sound red king crab. Parameters with “log_” indicate log scaled parameters

Parameter	Parameter description	Lower	Upper
log q ₁	Commercial fishery catchability (1977-93)	-20.5	20
log q ₂	Commercial fishery catchability (1994-2007)	-20.5	20
log q ₃	Commercial fishery catchability (2008-2019)	-20.5	20
log N ₇₆	Initial abundance	2.0	15.0
R ₀	Mean Recruit	2.0	12.0
log σ _R ²	Recruit standard deviation	-40.0	40.0
a ₁₋₇	Intimal length proportion	0	10.0
r _{1,2}	Proportion of length class 1 for recruit	0	5.0
log α	Inverse logistic molting parameter	-5.0	-1.0
log β	Inverse logistic molting parameter	1.0	5.5
log φ _{st1}	Logistic trawl selectivity parameter	-5.0	1.0
log φ _{wra}	Inverse logistic winter pot selectivity parameter	-5.0	1.0
log φ _{wrb}	Inverse logistic winter pot selectivity parameter	0.0	6.0
Sw _{1,2}	Winter pot selectivity of length class 1,2	0.1	1.0
log φ _l	Logistic commercial catch selectivity parameter	-5.0	1.0
log φ _{ra}	Logistic summer commercial retention selectivity Newshell (1976-2007, 2008-2022)	-5.0	1.0
log φ _{rb}	Logistic summer commercial retention selectivity Newshell (1976-2007, 2008-2022)	0.0	6.0
log φ _{wra}	Logistic winter commercial retention selectivity p	-5.0	1.0
log φ _{wrb}	Logistic winter commercial retention selectivity	0.0	6.0
w _t ²	Additional variance for standard CPUE	0.0	6.0
m1-8	Natural mortality multipliers	0	5.0
q.1	Survey q for NMFS trawl 1976-91	0.1	1.0
q.2	Survey q for NMFS NBS trawl 2010,17,19	0.1	1.0
σ	Growth transition sigma	0.0	30.0
β ₁	Growth transition mean	0.0	20.0
β ₂	Growth transition increment	0.0	20.0

*: Parameter was unestimable because model estimated trawl survey selectivity was 1.0 across all size classes.

Norton Sound Red King Crab Stock Assessment Sept 2023

Name	21.0		23.0	
	Estimate	std.dev	Estimate	std.dev
log q_1	-7.310	0.194	-7.174	0.191
log q_2	-6.721	0.165	-6.575	0.169
log q_3	-6.850	0.151	-6.740	0.157
log N_{76}	9.128	0.137	9.461	0.158
R_0	6.448	0.080	7.091	0.150
a_1	1.002	4.445	2.605	4.511
a_2	1.734	4.174	2.903	4.316
a_3	3.477	3.914	4.370	4.077
a_4	3.966	3.891	4.674	4.056
a_5	4.232	3.883	4.820	4.047
a_6	3.493	3.911	3.905	4.079
a_7	2.054	4.183	2.037	4.368
r1	5.000	0.002	5.000	0.003
r2	4.635	0.162	4.508	0.166
log a	-2.746	0.088	-2.765	0.094
log b	4.833	0.015	4.816	0.016
log ϕ_{11}	-5.000	0.035	-2.392	0.080
log ϕ_{wa}	-2.400	0.424	-1.858	0.425
log ϕ_{wb}	4.772	0.069	4.860	0.028
Sw1	0.061	0.034	0.045	0.022
Sw2	0.424	0.147	0.373	0.088
Sw3	0.732	0.237	0.733	0.141
log ϕ_l	-2.056	0.043	-1.942	0.041
log ϕ_{ra1}	-0.854	0.143	-0.886	0.142
log ϕ_{rb1}	-0.496	0.270	-0.484	0.266
log ϕ_{ra2}	4.641	0.008	4.648	0.009
log ϕ_{rb2}	4.654	0.013	4.655	0.013
log ϕ_{wra}	-0.951	0.559	-0.922	0.589
log ϕ_{wrb}	4.654	0.038	4.651	0.039
w^2_t	0.142	0.039	0.144	0.040
q.1	0.719	0.128	0.712	0.124
q.2	0.812	0.157	0.796	0.154
σ	3.801	0.207	3.792	0.202
β_1	11.874	0.692	12.807	0.721
β_2	7.795	0.169	7.557	0.176
M			0.412	0.027
$m1$				
$m2$				
$m3$				
$m4$				
$m5$				
$m6$				
$m7$				
$m8$	3.416	0.262		

Table 12. Annual abundance estimates of legal crab (million crab) and mature male biomass (Feb 01) (MMB, million lb) for Norton Sound red king crab estimated by a length-based analysis.

MMB

Year	Model 21.0	Model 23.0
1976	17.33	17.45
1977	19.15	19.24
1978	16.71	16.76
1979	11.86	11.90
1980	6.42	6.44
1981	4.36	4.38
1982	3.40	3.42
1983	4.30	4.33
1984	5.04	5.07
1985	5.70	5.73
1986	6.05	6.09
1987	5.96	5.99
1988	5.88	5.91
1989	5.62	5.65
1990	5.31	5.33
1991	5.00	5.02
1992	4.76	4.78
1993	4.45	4.47
1994	3.85	3.87
1995	3.29	3.30
1996	2.90	2.91
1997	2.88	2.90
1998	3.49	3.52
1999	4.38	4.42
2000	4.62	4.66
2001	4.18	4.21
2002	3.98	4.01
2003	4.07	4.10
2004	4.08	4.11
2005	3.81	3.83
2006	3.57	3.60
2007	3.74	3.78
2008	4.26	4.31
2009	4.72	4.78
2010	4.92	4.98
2011	4.60	4.65
2012	4.15	4.19
2013	3.96	3.99
2014	4.36	4.40
2015	4.91	4.96
2016	4.50	4.54
2017	3.67	3.70
2018	2.82	2.85
2019	2.39	2.42
2020	3.00	3.08
2021	4.45	4.64
2022	5.31	5.52
2023	5.29	5.39

Legal abundance (x million) (\geq 104 mm CL)

Year	Model 21.0 Final	Model 21.0	Model 22.0	Model 22.1	Model 22.2
1976	5.02	7.39	7.33	8.98	8.89
1977	6.15	7.14	7.08	8.32	8.25
1978	5.22	5.59	5.54	6.23	6.19
1979	3.51	3.77	3.74	4.09	4.07
1980	1.83	2.02	2.01	2.19	2.18
1981	1.15	1.46	1.45	1.59	1.58
1982	0.84	1.34	1.33	1.50	1.48
1983	1.14	1.76	1.74	1.97	1.94
1984	1.40	2.03	2.00	2.27	2.24
1985	1.60	2.25	2.21	2.51	2.47
1986	1.76	2.31	2.27	2.56	2.51
1987	1.71	2.22	2.18	2.43	2.39
1988	1.70	2.15	2.12	2.35	2.32
1989	1.62	2.01	1.99	2.18	2.16
1990	1.51	1.90	1.89	2.05	2.04
1991	1.43	1.78	1.77	1.92	1.91
1992	1.36	1.68	1.68	1.79	1.78
1993	1.27	1.57	1.57	1.65	1.65
1994	1.10	1.35	1.36	1.42	1.42
1995	0.93	1.16	1.17	1.21	1.22
1996	0.80	1.05	1.06	1.10	1.10
1997	0.78	1.09	1.10	1.13	1.13
1998	0.89	1.40	1.40	1.47	1.47
1999	1.24	1.74	1.75	1.83	1.83
2000	1.40	1.71	1.72	1.77	1.77
2001	1.22	1.49	1.50	1.52	1.52
2002	1.09	1.46	1.47	1.50	1.50
2003	1.11	1.54	1.56	1.61	1.61
2004	1.17	1.53	1.55	1.60	1.60
2005	1.11	1.40	1.42	1.45	1.46
2006	0.98	1.34	1.36	1.41	1.41
2007	0.99	1.48	1.50	1.58	1.57
2008	1.16	1.71	1.74	1.84	1.84
2009	1.32	1.88	1.92	2.04	2.03
2010	1.44	1.90	1.94	2.04	2.04
2011	1.37	1.69	1.73	1.80	1.80
2012	1.18	1.51	1.54	1.59	1.59
2013	1.07	1.51	1.52	1.60	1.59
2014	1.13	1.76	1.77	1.91	1.90
2015	1.41	1.95	1.97	2.11	2.11
2016	1.38	1.66	1.68	1.76	1.76
2017	1.11	1.27	1.29	1.31	1.32
2018	0.81	0.97	0.98	0.97	0.98
2019	0.65	0.86	0.86	0.85	0.85
2020	0.68	1.26	1.26	1.26	1.26
2021	1.19	1.92	1.93	1.92	1.92
2022	1.58	2.13	2.13	2.06	2.06
2023	1.54	1.98	1.98	1.88	1.88

Table 13. Summary of observed catch (million lb) for Norton Sound red king crab.

Year	Summer Com	Winter Com	Winter Sub	Discards Winter Sub	Total
1977	0.52	0.000	0.000	0	0.520
1978	2.09	0.024	0.025	0.008	2.147
1979	2.93	0.001	0.000	0	2.931
1980	1.19	0.000	0.000	0	1.190
1981	1.38	0.000	0.001	0	1.381
1982	0.23	0.000	0.003	0.001	0.234
1983	0.37	0.001	0.021	0.006	0.398
1984	0.39	0.002	0.022	0.005	0.419
1985	0.43	0.003	0.017	0.002	0.452
1986	0.48	0.005	0.014	0.004	0.503
1987	0.33	0.003	0.012	0.002	0.347
1988	0.24	0.001	0.005	0.001	0.247
1989	0.25	0.000	0.012	0.002	0.264
1990	0.19	0.010	0.024	0.004	0.228
1991	0	0.010	0.015	0.002	0.027
1992	0.07	0.021	0.023	0.003	0.117
1993	0.33	0.005	0.002	0	0.337
1994	0.32	0.017	0.008	0.001	0.346
1995	0.32	0.022	0.011	0.002	0.355
1996	0.22	0.005	0.003	0.001	0.229
1997	0.09	0.000	0.001	0.001	0.092
1998	0.03	0.002	0.017	0.012	0.061
1999	0.02	0.007	0.015	0.003	0.045
2000	0.3	0.008	0.011	0.004	0.323
2001	0.28	0.003	0.001	0	0.284
2002	0.25	0.007	0.004	0.003	0.264
2003	0.26	0.017	0.008	0.005	0.290
2004	0.34	0.001	0.002	0.001	0.344
2005	0.4	0.006	0.008	0.003	0.417
2006	0.45	0.000	0.002	0.001	0.453
2007	0.31	0.008	0.021	0.011	0.350
2008	0.39	0.015	0.019	0.009	0.433
2009	0.4	0.012	0.010	0.002	0.424
2010	0.42	0.012	0.014	0.002	0.448
2011	0.4	0.009	0.013	0.003	0.425
2012	0.47	0.025	0.015	0.004	0.514
2013	0.35	0.061	0.015	0.014	0.440
2014	0.39	0.035	0.007	0.002	0.434
2015	0.40	0.099	0.019	0.005	0.523
2016	0.42	0.080	0.011	0.001	0.512
2017	0.41	0.078	0.012	0.001	0.501
2018	0.30	0.029	0.008	0.001	0.338
2019	0.08	0.032	0.003	0.001	0.116
2020	0	Conf.	0.001	0.000	Conf
2021	0	0.0	0.004	0.002	0.006
2022	0.32	0.070	0.006	0.003	0.400
2023	0.41	0.01	TBD	TBD	TBD

Table 14: Jittering : **To be reported in Jan 2024**




Figure 1. King crab fishing districts and sections of Statistical Area Q.




Figure 2. Closed water regulations in effect for the Norton Sound commercial crab fishery. Line around the coastline delineates the 3-mil state waters zone.




Figure 3. Model estimated natural mortality, annual molting probability, selectivity for trawl survey, winter pot survey, and summer commercial fishery, and retention probability for winter commercial and summer commercial. X-axis is carapace length (mm).




Figure 4. Model estimated transition probability for each size classes.




Figure 5. Model estimated abundances of total, legal (CL>104 mm) and recruit (CL 64-94 mm) males during 1976-2022.




Figure 5. Estimated MMB during 1976-2023. Horizontal line Bmsy (Average MMB of 1980-2023).




Figure 6. Observed (open circle) (White: NMFS, Red: ADF&G) and model trawl survey male abundances with 95% lognormal Confidence Intervals (crab \geq 64 mm CL).




Figure 7. Observed (open circle) with 95% lognormal Confidence Intervals with additional variance (gray), and model estimated standardized CPUE.




Figure 8. Predicted (line) vs. observed (bar New Shell: blue, Old Shell: green) length class proportions for the summer commercial harvest 1977-2022.




Figure 9. Predicted (line) vs. observed (bar New Shell: blue, Old Shell: green) length class proportions for trawl survey 1976-2022.




Figure 10. Predicted (line) vs. observed (bar New Shell: blue, Old Shell: green) length class proportions for winter pot survey 1982-2012.




Figure 11. Predicted (line) vs. observed (bar New Shell: left blue, Old Shell: right green) length class proportions for summer commercial total and discards (1987-1994, 2012-2019) and winter commercial retained fishery 2015-2018




Figure 12. Predicted (line) vs. observed (bar) length class proportions for tag recovery data.





Figure 13. Input vs. model implied effective sample size. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the harmonic mean of implied sample size. Figures in the second column show input sample sizes (x-axis) vs. implied effective sample sizes (y-axis). Dashed line indicates the linear regression slope, and solid line is 1:1 line. Figures in the third column show years (x-axis) vs. implied effective sample sizes (y-axis). Horizontal solid line is the harmonic mean of implied sample size.



Norton Sound Red King Crab Stock Assessment Sept 2023






Figure 14. Bubble plots of predicted and observed length proportions. Black circles indicate model underestimates compared to observed, and white circles indicate model overestimates compared to observed. Size of circle indicates degree of deviance (larger circle = larger deviance). In ideal model fit case, distribution of sizes and colors of circles should be random (i.e., no systematic model misfits).



Norton Sound Red King Crab Stock Assessment Sept 2023



Norton Sound Red King Crab Stock Assessment Sept 2023






Figure 15. Standardized Pearson residual plots for trawl survey, summer commercial retained catch, winter pot survey, and observer for length size classes 1-8.





Norton Sound Red King Crab Stock Assessment Sept 2023




Figure 16. QQ Plot of Trawl survey and Commercial CPUE.




Figure 17. Retrospective Analyses of Norton Sound Red King Crab MMB from 2012 to 2024.
Solid black line: 2023 assessment model results. **(To be reported in Jan 2024)**

