

Norton Sound Red King Crab Stock Assessment for the fishing year 2020

Toshihide Hamazaki¹ and Jie Zheng²
Alaska Department of Fish and Game Commercial Fisheries Division
¹333 Raspberry Rd., Anchorage, AK 99518-1565
Phone: 907-267-2158
Email: Toshihide.Hamazaki@alaska.gov
²P.O. Box 115526, Juneau, AK 99811-5526
Phone : 907-465-6102
Email : Jie.Zheng@alaska.gov

Executive Summary

1. Stock. Red king crab, *Paralithodes camtschaticus*, in Norton Sound, Alaska.
2. Catches. This stock supports three important fisheries: summer commercial, winter commercial, and winter subsistence fisheries. Of those, the summer commercial fishery accounts for 85% of total harvest. The summer commercial fishery started in 1977. Catch peaked in the late 1970s with retained catch of over 2.9 million pounds. Since 1994, Norton Sound Crab fishery operated as super exclusive. For 2019 fishery season, Norton Sound Red King Crab harvest consisted of: 1,050 crabs (3,295 lb.) by winter commercial, 1,548 (~3,000 lb) by winter subsistence. Summer commercial fishery was closed on September 05 and harvest numbers has not been finalized but ~25,000 crab (~75,000 lb) by summer commercial, totaling ~42,000 crab (~82,000 lb.). Total harvests were below ABC of 0.19 million lb.
3. Stock Biomass. Norton Sound Red King Crab stock has been monitored by triennial survey since 1976 by NOAA (1976-1991) and ADF&G (1996-present), ranged from 1.41 million to 5.9 million crab. In 2019, abundance by trawl survey was 4.66 million crab with CV 0.60.
4. Recruitment. Model estimated recruitment was weak during the late 1970s and high during the early 1980s, with a slightly downward trend from 1983 to 1993. Estimated recruitment has been highly variable but on an increasing trend in recent years.
5. Management performance.

1
2 *Status and catch specifications (million lb.)*

Year	MSST	Biomass (MMB)	GHL	Retained Commercial Catch	Total Retained Catch	Retained OFL	Retained ABC
2016	2.26 ^A	5.87	0.52	0.51	0.52	0.71 ^A	0.57
2017	2.31 ^B	5.14	0.50	0.49	0.50	0.67 ^B	0.54
2018	2.41 ^C	4.08	0.30	0.31	0.34	0.43 ^C	0.35
2019	2.24 ^D	3.12	0.15	TBD	TBD	0.24 ^D	0.19
2020							

3
4
5
6 *Status and catch specifications (1000t)*

Year	MSST	Biomass (MMB)	GHL	Retained Commercial Catch	Total Retained Catch	Retained OFL	Retained ABC
2016	1.03 ^A	2.66	0.24	0.23	0.24	0.32 ^A	0.26
2017	1.05 ^B	2.33	0.23	0.22	0.24	0.30 ^B	0.24
2018	1.09 ^C	1.85	0.13	0.14	0.15	0.20 ^C	0.16
2019	1.03 ^D	1.41	0.09	TBD	TBD	0.11 ^D	0.09
2020							

8
9
10 Notes:
11 MSST was calculated as $B_{MSY}/2$
12 A-Calculated from the assessment reviewed by the Crab Plan Team in May 2016
13 B-Calculated from the assessment reviewed by the Crab Plan Team in May 2017
14 C-Calculated from the assessment reviewed by the Crab Plan Team in Jan 2018
15 D-Calculated from the assessment reviewed by the Crab Plan Team in Jan 2019
16 E-Calculated from the assessment reviewed by the Crab Plan Team in Jan 2020

17
18 Conversion to Metric ton: 1 Metric ton (t) = 2.2046×1000 lb

19
20
21 *Biomass in millions of pounds*

Year	Tier	B_{MSY}	Current MMB	B/ B_{MSY} (MMB)	F_{OFL}	Years to define B_{MSY}	M	1-Buffer	Retained ABC
2016	4a	4.53	5.87	1.3	0.18	1980-2016	0.18	0.8	0.57
2017	4a	4.62	5.14	1.1	0.18	1980-2017	0.18	0.8	0.54
2018	4b	4.82	4.08	0.9	0.15	1980-2018	0.18	0.8	0.35
2019	4b	4.57	3.12	0.7	0.12	1980-2019	0.18	0.8	0.19
2020	4b					1980-2020	0.18	0.8	

22
23 *Biomass in 1000t*

Year	Tier	B _{MSY}	Current MMB	B/B _{MSY} (MMB)	F _{OFL}	Years to define B _{MSY}	M	1-Buffer	Retained ABC
2016	4a	2.06	2.66	1.3	0.18	1980-2016	0.18	0.8	0.26
2017	4a	2.10	2.33	1.1	0.18	1980-2017	0.18	0.8	0.24
2018	4b	2.18	1.85	0.9	0.15	1980-2018	0.18	0.8	0.16
2019	4b	2.06	1.41	0.7	0.12	1980-2019	0.18	0.8	0.09
2020	4b					1980-2020	0.18		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

6. Probability Density Function of the OFL, OFL profile, and mcmc estimates.

To be completed for the Jan 2020 final assessment.

7. The basis for the ABC recommendation

For Tier 4 stocks, the default maximum ABC is based on P*=49% that is essentially identical to the OFL. Accounting for uncertainties in assessment and model results, the SSC chose to use 90% OFL (10% Buffer) for the Norton Sound red king crab stock from 2011 to 2014. In 2015, the buffer was increased to 20% (ABC = 80% OFL).

8. A summary of the results of any rebuilding analysis

N/A

A. Summary of Major Changes in 2019

1. Changes to the management of the fishery:

None

2. Changes to the input data

a. Data update: To be finalized in Jan 2020

i. 1977-2019 standardized commercial catch CPUE and CV. No changes in standardization methodology (NPFMC 2013). (to be finalized for Jan 2020 assessment)

ii. Winter and Summer commercial fishery harvest, discards, and length composition data. Retained size was not collected for 2019 winter commercial. Catch and length data (to be finalized for Jan 2020 assessment)

iii. Tag recovery data 2019 (to be included for Jan 2020 assessment)

iv. Trawl survey: abundance, length-shell composition

ADFG 2019 (included)

1 NOAA 2019 (to be included for Jan 2020 assessment)

2

3 3. Changes to the assessment methodology:

4 None

5 4. Changes to the assessment results.

6 None

7

8 **B. Response to SSC and CPT Comments**

9

10 Crab Plan Team – January 23-25, 2019

11

- 12 • Continue to evaluate methods to improved ADF&G bottom trawl survey biomass
13 estimation, including model based approaches such as VAST.

14 Authors' reply: VAST modeling approach has been considered. **Authors request experts'**
15 **instruction and assistance for implementation.**

16

17

- 18 • Conduct a sensitivity analysis to evaluate the effect of mark-recapture data by fitting the
19 model only marks that are liberty for one year.

20

21 Authors' reply:

22

23 Alternative model: 19.1

24

25

- 26 • Evaluate potential differences in survey Q between NOAA and ADFG bottom trawl
27 surveys.

28

29 Authors' reply: Alternative model 19.2 and 19.3

30


- 31 • Collect more chela-carapace data, especially at the small size ranges, to improve the size
32 at maturity estimate.

33

34 Author's reply

35

36 In 2019 97 male samples were collected during the annual bottom trawl survey. No distinctive
37 break point has been present.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

SSC – February 4-6 2019

- The model choice does not have much impact on the results, or on the Tier 4 reference points, hence the focus for the stock assessment should be on the input data.

Author Reply

We fully concur. We are collecting more data as budget allows.

- Bring forward total catch OFLs and ABCs or provide rationale why the retained catch OFL and ABC are still more appropriate at this time.

Author Reply

Estimating total catch OFL requires estimating the number of discards in summer commercial fisheries. Thus far, no formal estimates of discards has not been established for NSRKC. See Appendix C for 2002-2018 preliminary discards estimates.

- 1 • Include options with an estimated constant M across size classes (including the largest
2 class) and a dome-shaped selectivity for the summer commercial fishery and for the
3 summer survey.

4 Authors' reply: Alternative model 19.4 and 19.5
5

- 6 • Spatial distribution and modeling. a thorough examination of the spatial distribution of
7 red king crab, in particular spatial differences in size composition, across the northern
8 Bering Sea beyond Norton Sound would be helpful. Available data include the 2010 and
9 2017-2018 NMFS bottom trawl surveys.

10 Authors' reply: This task is more appropriate for NMFS.

- 11 • Spatial modeling: Compare the ADF&G and NMFS surveys using appropriate methods
12 for zero-inflated distributions, such as those offered in various R packages (e.g., pscl,
13 gamlss, INLA, VAST, glmmfields).

14 Author's reply:

15 We are not familiar with those packages and spatial modeling. **Authors request experts**
16 **instruction and assistance for implementation.**

17 It should also be noted that ADF&G and NMFS surveys are NOT "paired" (i.e., side-by-side
18 survey). ADF&G and NMFS surveys differ in **survey protocols (e.g., tow distance), trawl**
19 **gears, survey spatial extent and timing. Thus, it is expected that the two surveys would**
20 **differ in abundance and spatial distribution.** Thus, model expected survey abundance also
21 differ (based on survey timing, incorporating amount of harvest by the mid-point survey date) in
22 the assessment model.

- 23 • Survey time series: Explore using two catchability parameters for the differing time
24 blocks of the survey time series shown in Figure 7 which uses a different length range
25 after 1995 to compute the abundance index.

26 Author's reply:

27 The NMFS survey abundance prior to 1995 were provided by NMFS (NPFMC 2014) when
28 NSRKC model was based on 74mm and above. When this was changed to 64mm and above
29 survey abundance after 1995 were updated by the authors (NPFMC2016), but not for the pre-
30 1995 NMFS surveys. This was because the assessment model was already estimating q ($q \sim 0.7$)
31 for pre-1995 survey abundance. In this assessment, the pre-1995 survey abundance was updated
32 to 64mm and above. We also included differences in abundance estimation methodologies
33 between pre-1995 NMFS and post 1995 trawl surveys (Table 3). Combining with application of
34 VAST, we will further explore improvement of trawl survey abundance.

35

- 36 • Local and traditional knowledge: Encourage through collaborations at the local level to
37 consider these sources of knowledge

38 Author's reply:

1 **Authors request SSC and experts' instructions how to collaborate and incorporate local and**
2 **traditional knowledge into assessment.**
3

- 4 • Male maturity: new maturity studies are clearly needed to improve the assessment.
5 Explore Russian data on maturity if available. Also, the relationship between maturity
6 and temperature across stocks should be explored for potential predictive capability for
7 Norton Sound.

8 Authors' reply:

9 We are eager to incorporate SSC's suggestions on data weighting; however, we are not familiar
10 with the dataset mentioned. **Authors request experts' instruction and assistance for**
11 **implementation.**


12

13

- 14 • Consider estimating observer length composition weighted by catch/strata.

15 Author's reply:

16 While weighted length composition is considered more accurate than simple unweighted one,
17 there is little difference between the two.



18
19

- 20 • Consider data weighting based on iterative tuning, number of hauls, or other approaches.

21 Author's reply:

1 We are eager to incorporate SSC's suggestions on data weighting; however, we are not familiar
2 with those. **Authors request experts' instruction and assistance for implementation.**

- 3
- 4 • Include before/after variables in CPUE standardization to account for a change in
5 commercially acceptable size limit. Clarify if the time series of CPUE is showing
6 different measures of CPUE for the time periods prior to and after 1995.

7 Author's reply:

8 In the original CPUE standardization, the CPUE data were separated in two periods: 1976-1992
9 and 1993-present, and two regressions were run. In this revision, we included time stage
10 variables PD, 1976-1992, 1993-2014, 2015-present, and ran a single regression model. The PD
11 variable turned out to be insignificant and was removed from the final regression model.
12 Furthermore, this also increased model sd, so that model estimated additional variance (advar)
13 became 0.

- 14
- 15 • Use revised Mohn's rho.

16 Author's reply:

17 Will be implemented for final assessment. However, more fundamental note, CPT has not
18 established standardized criterion for Mohn's rho (e.g., min-max rho value) for selection of the
19 best alternative model, or an adjustment of predicted biomass or determining OFL/ABC buffer.
20 In that sense, Mohn's rho whether original or revised is just a relative index that can inform
21 relative retrospective performance among alternative candidate models. Since retrospective
22 analyses are done for identical years (e.g. past 10 years) for each candidate models, both sum and
23 mean deviation function the same way for evaluating relative performance. **Authors**
24 **appreciate SSC's directive for potential application of revised Mohn's rho for improvement**
25 **of the NSRKC assessment model.**

- 26 • Parameters r_1 and $\log\text{-}\phi_{st1}$ hitting bounds.

27 Author's reply:

28 r_1 is a parameter for normalization for estimating proportion, $p_i = \exp(r_i) / [1 + \sum(\exp(r))]$, (see
29 equation 2 of Appendix A), so that hitting bounds is acceptable. $\log\text{-}\phi_{st1}$ is the trawl survey
30 selectivity curve in log scale (see equation (16) Appendix A). Since trawl selectivity was
31 estimated to be 1.0 across all lengths, hitting bound does not affect results of the assessment
32 model. SSC (NPFMC 2017) suggested setting trawl survey selectivity to 1.0 for all length.

33

34 Crab Plan Team – April 29, 2019

- 35
- 36 • Draft assessment in GMACS will potentially be provided in September 2019.

37 Author Reply:

38

39 We are eager to incorporate SSC's suggestions on data weighting; however, we are not familiar
40 with GMACS. **Authors request experts' instruction and assistance for implementation.**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

C. Introduction

1. Species: red king crab (*Paralithodes camtschaticus*) in Norton Sound, Alaska.
2. General Distribution: Norton Sound red king crab is one of the northernmost red king crab populations that can support a commercial fishery (Powell et al. 1983). It is distributed throughout Norton Sound with a westward limit of 167-168° W. longitude, depths less than 30 m, and summer bottom temperatures above 4°C. The Norton Sound red king crab management area consists of two units: Norton Sound Section (Q3) and Kotzebue Section (Q4) (Menard et al. 2011). The Norton Sound Section (Q3) consists of all waters in Registration Area Q north of the latitude of Cape Romanzof, east of the International Dateline, and south of 66°N latitude (Figure 1). The Kotzebue Section (Q4) lies immediately north of the Norton Sound Section and includes Kotzebue Sound. Commercial fisheries have not occurred regularly in the Kotzebue Section. This report deals with the Norton Sound Section of the Norton Sound red king crab management area.
3. Evidence of stock structure: Thus far, no studies have investigated possible stock separation within the putative Norton Sound red king crab stock.
4. Life history characteristics relevant to management: One of the unique life-history traits of Norton Sound red king crab is that they spend their entire lives in shallow water since Norton Sound is generally less than 40 m in depth. Distribution and migration patterns of Norton Sound red king crab have not been well studied. Based on the 1976-2006 trawl surveys, red king crab in Norton Sound are found in areas with a mean depth range of 19 ± 6 (SD) m and bottom temperatures of 7.4 ± 2.5 (SD) °C during summer. Norton Sound red king crab are consistently abundant offshore of Nome.

Norton Sound red king crab migrate between deeper offshore and inshore shallow waters. Timing of the inshore mating migration is unknown, but is assumed to be during late fall to winter (Powell et al. 1983). Offshore migration occurs in late May - July (Jenefer Bell, ADF&G, personal communication). The results from a study funded by North Pacific Research Board (NPRB) during 2012-2014 suggest that older/large crab (> 104mm CL) stay offshore in winter, based on findings that large crab are not found nearshore during spring offshore migration periods (Jenefer Bell, ADF&G, personal communication). Molt timing is unknown but likely occurs in late August – September, based on increase catches of newly-molted crab late in the fishing season (August- September) (Joyce Soong, ADF&G personal communication) and evaluation of molting hormone profiles in the hemolymph (Jenefer Bell, ADF&G, personal communication). Recent observations also indicate that mating may be biennial (Robert Foy, NOAA, personal communication). Trawl surveys show that crab distribution is dynamic with recent surveys showing high abundance on the southeast side of Norton Sound, offshore of Stebbins and Saint Michael.

- 1 5. Brief management history: Norton Sound red king crab fisheries consist of commercial and
2 subsistence fisheries. The commercial red king crab fishery started in 1977 and occurs in
3 summer (June – August) and winter (December – May). The majority of red king crab harvest
4 occurs offshore during the summer commercial fishery, whereas the winter commercial and
5 subsistence fisheries occur nearshore through ice.

6
7 Summer Commercial Fishery

8 A large-vessel summer commercial crab fishery started in 1977 in the Norton Sound Section
9 (Table 1) and continued from 1977 through 1990. No summer commercial fishery occurred in
10 1991 because there were no staff to manage the fishery. In March 1993, the Alaska Board of
11 Fisheries (BOF) limited participation in the fishery to small boats. Then on June 27, 1994, a
12 super-exclusive designation went into effect for the fishery. This designation stated that a
13 vessel registered for the Norton Sound crab fishery may not be used to take king crabs in any
14 other registration areas during that registration year. A vessel moratorium was put into place
15 before the 1996 season. This was intended to precede a license limitation program. In 1998,
16 Community Development Quota (CDQ) groups were allocated a portion of the summer
17 harvest; however, no CDQ harvest occurred until the 2000 season. On January 1, 2000 the
18 North Pacific License Limitation Program (LLP) went into effect for the Norton Sound crab
19 fishery. The program dictates that a vessel which exceeds 32 feet in length overall must hold a
20 valid crab license issued under the LLP by the National Marine Fisheries Service. Changes in
21 regulations and the location of buyers resulted in eastward movement of the harvest
22 distribution in Norton Sound in the mid-1990s. In Norton Sound, a legal crab is defined as \geq
23 4-3/4 inch carapace width (CW, Menard et al. 2011), which is approximately equivalent to \geq
24 104 mm carapace length mm CL. Since 2005, commercial buyers (Norton Sound Economic
25 Development Corporation) started accepting only legal crab of \geq 5 inch CW. This may have
26 increased discards; however, because discards have not been monitored until 2012, impact of
27 this change on discards is unknown. This issue was also examined in assessment model
28 selection, which showed no difference in estimates of selectivity functions before and after
29 2005 (NPFMC 2016).

30 Portions of Norton Sound area are closed to commercial fishing for red king crab. Since the
31 beginning of the commercial fisheries in 1977, waters approximately 5-10 miles offshore of
32 southern Seward Peninsula from Port Clarence to St. Michael have been closed to protect crab
33 nursery grounds during the summer commercial crab fishery (Figure 2). The spatial extent of
34 closed waters has varied historically.

35
36 CDQ Fishery

37 The Norton Sound and Lower Yukon CDQ groups divide the CDQ allocation. Only fishers
38 designated by the Norton Sound and Lower Yukon CDQ groups are allowed to participate in
39 this portion of the king crab fishery. Fishers are required to have a CDQ fishing permit from
40 the Commercial Fisheries Entry Commission (CFEC) and register their vessel with the Alaska
41 Department of Fish and Game (ADF&G) before begin fishing. Fishers operate under the
42 authority of each CDQ group. The CDQ fishery may open at any time by emergency order.
43 CDQ harvest share is 7.5% of total projected harvest.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

Winter Commercial Fishery

The winter commercial crab fishery is a small fishery using hand lines and pots through the nearshore ice. On average 10 permit holders harvested 2,500 crabs during 1978-2009. From 2007 to 2015 the winter commercial catch increased from 3,000 crabs to over 40,000 (Table 2). In 2015 winter commercial catch reached 20% of total crab catch. The BOF responded in May 2015 by amending regulations to allocate 8% of the total commercial guideline harvest level (GHL) to the winter commercial fishery, which became in effect since 2017 season. The winter red king crab commercial fishing season was also set from January 15 to April 30, unless changed by emergency order. The new regulation became in effect since the 2016 season.

Subsistence Fishery

While the winter subsistence fishery has a long history, harvest information is available only since the 1977/78 season. The majority of the subsistence crab fishery harvest occurs using hand lines and pots through nearshore ice. Average annual winter subsistence harvest was 5,400 crab (1977-2010). Subsistence harvesters need to obtain a permit before fishing and record daily effort and catch. There are no size or sex specific harvest limits; however, the majority of retained catches are males of near legal size. The subsistence fishery catch is influenced not only by crab abundance, but also by changes in distribution, changes in gear (e.g., more use of pots instead of hand lines since 1980s), and ice conditions (e.g., reduced catch due to unstable ice conditions: 1987-88, 1988-89, 1992-93, 2000-01, 2003-04, 2004-05, and 2006-07).

The summer subsistence crab fishery harvest has been monitored since 2004 with an average harvest of 712 crab per year. Since this harvest is very small, the summer subsistence fishery was not included in the assessment model.

6. Brief description of the annual ADF&G harvest strategy

Since 1997 Norton Sound red king crab has been managed based on a guideline harvest level (GHL). From 1999 to 2011 the GHL for the summer commercial fishery was determined by a prediction model and the model estimated predicted biomass: (1) 0% harvest rate of legal crab when estimated legal biomass < 1.5 million lb; (2) $\leq 5\%$ of legal male abundance when the estimated legal biomass falls within the range 1.5-2.5 million lb; and (3) $\leq 10\%$ of legal male when estimated legal biomass >2.5 million lb.

In 2012 a revised GHL for the summer commercial fishery was implemented: (1) 0% harvest rate of legal crab when estimated legal biomass < 1.25 million lb; (2) $\leq 7\%$ of legal male abundance when the estimated legal biomass falls within the range 1.25-2.0 million lb; (3) $\leq 13\%$ of legal male abundance when the estimated legal biomass falls within the range 2.0-3.0 million lb; and (3) $\leq 15\%$ of legal male biomass when estimated legal biomass >3.0 million lb.

In 2015 the Alaska Board of Fisheries passed the following regulations regarding winter commercial fisheries:

1. Revised GHL to include summer and winter commercial fisheries.

- 1 2. Set guideline harvest level for winter commercial fishery (GHL_w) at 8% of the total
- 2 GHL
- 3 3. Dates of the winter red king crab commercial fishing season are from January 15 to
- 4 April 30.
- 5

Year	Notable historical management changes
1976	The abundance survey started
1977	Large vessel commercial fisheries began (Legal size \geq 5 inch CW)
1978	Legal size changes to \geq 4.75 inch CW
1991	Fishery closed due to staff constraints
1994	Super exclusive designation went into effect. The end of large vessel commercial fishery operation.
1998	Community Development Quota (CDQ) allocation went into effect
1999	Guideline Harvest Level (GHL) went into effect
2000	North Pacific License Limitation Program (LLP) went into effect.
2002	Change in closed water boundaries (Figure 2)
2005	Commercially accepted legal crab size changed from \geq 5 inch CW
2006	The Statistical area Q3 section expanded (Figure 1)
2008	Start date of the open access fishery changed from July 1 to after June 15 by emergency order. Pot configuration requirement: at least 4 escape rings (>4.5 inch diameter) per pot located within one mesh of the bottom of the pot, or at least $\frac{1}{2}$ of the vertical surface of a square pot or sloping side-wall surface of a conical or pyramid pot with mesh size > 6.5 inches.
2012	The Board of Fisheries adopted a revised GHL for summer fishery.
2016	Winter GHL for commercial fisheries was established and modified winter fishing season dates were implemented.

- 6
- 7 7. Summary of the history of the B_{MSY} .
- 8 NSRKC is a Tier 4 crab stock. Direct estimation of the B_{MSY} is not possible. The B_{MSY} proxy
- 9 is calculated as mean model estimated mature male biomass (MMB) from 1980 to present.
- 10 Choice of this period was based on a hypothesized shift in stock productivity a due to a climatic
- 11 regime shift indexed by the Pacific Decadal Oscillation (PDO) in 1976-77. Stock status of the
- 12 NSRKC was Tier 4a until 2013. In 2014 the stock fell to Tier 4b, but came back to Tier 4a for
- 13 the 2015-2017 seasons. In 2018 the stock again fell to Tier 4b.
- 14

15 **D. Data**

- 16 1. Summary of new information:
- 17

18 Winter commercial and subsistence fishery:

19

20 Winter commercial fishery catch in 2019 was 9,189 crab (20,118 lb.). Subsistence retained

21 crab catch was 4,424 and unretained was 1,343 or 23 % of total catch (Table 2).

22

23 Summer commercial fishery:

24

1 The summer commercial fishery opened on 6/25/2019 and closed on 9/03/2019. Total of
 2 75,023 crab (24,506 lb.) were harvested (Table 1). This is the lowest harvest since 2000.


3
 4 Total retained harvest for 2019 season was 88,646 crab (34,811 lb. or 0.035 million lb) and
 5 did not exceed the 2019 ABC of 0.19 million lb.

6
 7 Summer Trawl abundance survey ADFG (7/22-7/29) was estimated to be 4.67 million (CV
 8 60%). This was the second highest abundance, recorded.
 9 (Table 3).

10

11

12 2. Available survey, catch, and tagging data



13

	Years	Data Types	Tables
Summer trawl survey	76,79,82,85,88,91,96, 99,	Abundance	3
	02,06,08,10,11,14,17, 18,19	Length-shell comp	6
Winter pot survey	81-87, 89-91,93,95-00,02-12	Length-shell comp	7
Summer commercial fishery	77-90,92-19	Retained catch	1
		Standardized CPUE,	1
		Length-shell comp	4
Summer Com total catch	12-18	Length-shell comp	9
Summer Com Discards	87-90,92,94	Length-shell comp	8
Winter subsistence fishery	76-19	Total & Retained catch	2

Winter commercial fishery	78-19	Retained catch	2
	15-18	Retained Length-Shell	5
Tag recovery	80-18	Recovered tagged crab	10

1
2
3

Data available but not used for assessment

Data	Years	Data Types	Reason for not used
Summer pot survey	80-82,85	Abundance Length proportion	Uncertainties on how estimates were made.
Summer preseason survey	95	Length proportion	Just one year of data
Summer subsistence fishery	2005-2013	retained catch	Too few catches compared to commercial
Winter Pot survey	87, 89-91,93,95-00,02-12	CPUE	CPUE data Not reliable due to ice conditions
Preseason Spring pot survey	2011-15	CPUE, Length proportion	Years of data too short
Postseason Fall pot survey	2013-15	CPUE, Length proportion	Years of data too short

4
5
6
7

Catches in other fisheries

8 In Norton Sound, directed Pacific Cod pot fishery was issued in 2018 under the CDQ permit.
9 However, the fishery did not occur. This fishery may develop in the future.

	Fishery	Data availability
Other crab fisheries	Does not exist	NA
Groundfish pot	Pacific Cod (Planned, but not executed)	NA
Groundfish trawl	Does not exist	NA
Scallop fishery	Does not exist	NA

10
11
12
13
14
15
16
17
18
19

3. Other miscellaneous data:

Satellite tag migration tracking (NOAA 2016)

Spring offshore migration distance and direction (2012-2015)

Monthly blood hormone level (indication of molting timing) (2014-2015)

Data aggregated:

Proportion of legal size crab, estimated from trawl survey and observer data. (Table 13)

Data estimated outside the model:

Summer commercial catch standardized CPUE (Table 1, Appendix B)

1 ***E. Analytic Approach***

2

3 **1. History of the modeling approach.**

4 The Norton Sound red king crab stock was assessed using a length-based synthesis model
5 (Zheng et al. 1998). Since adoption of the model, the major challenge is a conflict between
6 model projection and data, specifically the model projects higher abundance-proportion of
7 large size class (> 123mm CL) of crab than observed. This problem was further exasperated
8 when natural mortality M was set to 0.18 from previous $M = 0.3$ in 2011 (NPFMC 2011).
9 This issue has been resolved by assuming (3-4 times) higher M for the length crabs (i.e., M
10 = 1.8 for length classes ≤ 123 mm, and higher M for > 123 mm) (NPFMC 2012, 2013, 2014,
11 2015, 2016, 2017, 2018). Alternative assumptions have been explored, such as changing
12 molting probability (i.e., crab matured quicker or delayed maturation), higher natural
13 mortality, and dorm shaped selectivity (i.e., large crab are not caught, or moved out of
14 fishery/survey grounds). However, those alternative assumptions did not produce better
15 model fits. Model estimated length specific molting probability was similar to inverse
16 logistic curve, and did not improve model fit (NPFMC 2016). Constant M across all length
17 classes resulted in higher M (0.3-0.45) (NPFMC 2013, 2017). Dome shaped selectivity
18 (i.e., assume large crabs were not caught/not surveyed/moved out of survey and fishing
19 area) increased MMB twice higher than other models. A model with gradual increase of
20 M across length classes resulted in M increase starting at size 94mm. However, this did not
21 improve overall model fit and was rejected for model consideration (NPFMC 2018). With
22 addition of total catch length data in summer and retention length data in winter
23 commercial fisheries, 2019 model specification examined estimation of retention curve for
24 both summer and winter fishery, and evaluation of OFL under Tier 3 formula.

25

26

27 Historical Model configuration progression:

28

29 2011 (NPFMC 2011)

30 1. $M = 0.18$

31 2. M of the last length class = 0.288

32 3. Include summer commercial discards mortality = 0.2

33 4. Weight of fishing effort = 20,

34 5. The maximum effective sample size for commercial catch and winter surveys = 100,

35

36 2012 (NPFMC 2012)

37 1. M of the last length class = $3.6 \times M$

38 2. The maximum effective sample size for commercial catch and winter surveys = 50,

39 3. Weight of fishing effort = 50.

40

41 2013 (NPFMC 2013)

42 1. Standardize commercial catch cpue and replace likelihood of commercial catch efforts
43 to standardized commercial catch cpue with weight = 1.0

44 2. Eliminate summer pot survey data from likelihood

3. Estimate survey q of 1976-1991 NMFS survey with maximum of 1.0
4. The maximum effective sample size for commercial catch and winter surveys = 20.

2014 (NPFMC 2014)

1. Modify functional form of selectivity and molting probability to improve parameter estimates (2 parameter logistic to 1 parameter logistic)
2. Include additional variance for the standardized cpue.
3. Include winter pot survey cpue (But was removed from the final model due to lack of fit)
4. Estimate growth transition matrix from tagged recovery data.

2015 (NPFMC 2015)

1. Winter pot survey selectivity is an inverse logistic, estimating selectivity of the smallest length group independently
2. Reduce Weight of tag-recovery: $W = 0.5$
3. Model parsimony: one trawl survey selectivity and one commercial pot selectivity

2016 (NPFMC 2016)

1. Length range extended from 74mm – 124mm above to 64mm – 134mm above.
2. Estimate multiplier for the largest ($> 123\text{mm}$) length classes.

2017 (NPFMC 2017)

1. Change molting probability function from 1 to 2 parameter logistic. Assume molting probability not reaching 1 for the smallest length class.

2019 (NPFMC 2019)

1. Fit total catch length composition and estimate retention probability for summer and winter commercial fishery
2. Include winter commercial retained length data

2. Model Description

- a. Description of overall modeling approach:

The model is a male-only size structured model that combines multiple sources of survey, catch, and mark-recovery data using a maximum likelihood approach to estimate abundance, recruitment, catchability of the commercial pot gear, and parameters for selectivity and molting probabilities (See Appendix A for full model description).

Unlike other crab assessment models, NSRK modeling year is starts from February 1st to January 31st of the following year. This schedule was selected because Norton Sound winter crab fisheries can start when Norton Sound ice become thick enough to operate fishery safely, which can be as earliest as mid-late January.

1 b-f. See Appendix A.

2
3 g. Critical assumptions of the model:

4
5 i. Male crab matures at CL length 94mm.

6 Size at maturity of NSRKC (CL 94 mm) was determined by adjusting that of BBRKC (CL
7 120mm) reflect the slower growth and smaller size of NSRKC.

8 ii. Molting occurs in the fall after the summer fishery

9 iii. Instantaneous natural mortality M is 0.18 for all length classes, except for the last
10 length group ($> 123\text{mm}$).

11 iv. Trawl survey selectivity is a logistic function with 1.0 for length classes 5-6. .
12 Selectivity is constant over time.

13
14 v. Winter pot survey selectivity is a dome shaped function: Reverse logistic function
15 of 1.0 for length class CL 84mm, and model estimate for CL $< 84\text{mm}$ length classes.
16 Selectivity is constant over time.

17 This assumption is based on the fact that a low proportion of large crab are caught
18 in the nearshore area where winter surveys occur. Causes of this pattern may be
19 that (1) large crab do not migrate into nearshore waters in winter or (2) large crab
20 are fished out by winter fisheries where the survey occurs (i.e., local depletion).
21 Recent studies suggest that the first explanation is more likely than second (Jenefer
22 Bell, ADFG, personal communication).

23
24
25 vi. Summer commercial fisheries selectivity is an asymptotic logistic function of 1.0
26 at the length class CL 134mm. While the fishery changed greatly between the
27 periods (1977-1992 and 1993-present) in terms of fishing vessel composition and
28 pot configuration, the selectivity of each period was assumed to be identical. Model
29 fits of separating and combining the two periods were examined in 2015, and
30 showed no difference between the two models (NPFMC 2015). For model
31 parsimony, the two were combined.

32
33 vii. Summer trawl survey selectivity is an asymptotic logistic function of 1.0 at the
34 length of CL 124mm. While the survey changed greatly between NOAA (1976-
35 1991) and ADF&G (1996-present) in terms of survey vessel and trawl net structure,
36 selectivity of both periods was assumed to be identical. Model fits separating and
37 combining the two surveys were examined in 2015. No differences between the
38 two models were observed (NPFMC 2015) and for model parsimony the two were
39 combined.

40 viii. Winter commercial and subsistence fishery selectivity and length-shell conditions
41 are the same as those of the winter pot survey. All winter commercial and
42 subsistence harvests occur February 1st.

1 Winter commercial king crab pots can be any dimension (5AAC 34.925(d)). No
2 length composition data exists for crab harvested in the winter commercial or
3 subsistence fisheries. However, because commercial fishers are also subsistence
4 fishers, it is reasonable to assume that the commercial fishers used crab pots that
5 they use for subsistence harvest, and hence both fisheries have the same
6 selectivity.
7

- 8 ix. Growth increments are a function of length, are constant over time, estimated
9 from tag recovery data.
10
11 x. Molting probability is an inverse logistic function of length for males.
12
13 xi. A summer fishing season for the directed fishery is short. All summer commercial
14 harvests occur July 1st.
15
16 xii. Discards handling mortality rate for all fisheries is 20%.
17 No empirical estimate is available.
18
19 xiii. Annual retained catch is measured without error.
20
21 xiv. All legal size crab ($\geq 4\text{-}3/4$ inch CW) are retained, and sublegal size crab or
22 commercially unacceptable size crab (< 5 inch CW, since 2005) are discarded
23
24 Since 2005, buyers announced that only legal crab with ≥ 5 inch CW are acceptable for
25 purchase. Since samples are taken at a commercial dock, it was anticipated that this
26 change would lower the proportion of legal crab. However, the model was not sensitive
27 to this change (NPFMC 2013, 2017).
28
29 xv. Length compositions have a multinomial error structure and abundance has a log-
30 normal error structure.
31

32 h. Changes of assumptions since last assessment:

33 None.
34

35 **3. Model Selection and Evaluation** 36

37 a. Description of alternative model configurations.
38

- 39 • For 2020 preliminary assessment, we explored all alternative modeling suggestions by CPT
40 and SSC (See Authors' responses). Baseline model (Model 19.0) is the Model 18.2b
41 adopted for the 2019 assessment. Model 19.1 explores the effects of tagging data on
42 molting and growth transition matrix. Model 19.2 and 19.3 reexamine validity of
43 assumptions about trawl survey q set in 2013 (NPFMC 2013). Finally, Model 19.4
44 reexamines the assumption of size dependent mortality (i.e., higher M for larger crab) by
45 estimating natural mortality and dome shape selectivity, which was examined in 2017

1 (NPFMC 2017). In 2017 model assessment, estimating size invariant M resulted higher
 2 M, and dome shaped selectivity resulted in assuming large number of crab never observed
 3 and caught by fishery. The Model 19.4-19.5 combines that two alternatives examined
 4 previously. Same as 2017 selectivity for each size class was estimated directly with
 5 selectivity of one size class assumed to be 1.0. Smoothing penalty was also included in
 6 likelihood.

7 We examined alternative models of
 8 Model 19.0: Baseline: Model 18.2b
 9 Model 19.1: Model 19.0 + Tag recovery data just for 1 year
 10 Model 19.2: Model 19.0 + NOAA trawl survey Q =1.0, Est: ADFG survey Q
 11 Model 19.3: Model 19.0 + Est survey Qs NOAA and ADFG
 12 Model 19.4: Model 19.0 + Est M equal for all lengths + Dome shape selectivity for trawl
 13 and summer commercial (max sel 94-103 for trawl, 104-113 for com)
 14 Model 19.5: Model 19.0 + Est M equal for all lengths + Dome shape selectivity for trawl
 15 and summer commercial (max sel 94-103 for trawl, 104-113 for com)
 16
 17

18 b. Evaluation of negative log-likelihood alternative models results:
 19

Model	Model 19.0	Model 19.1	Model 19.2	Model 19.3	Model 19.4	Model 19.5
Additional Parameters				+1	+14	+14
Total	306.1	254.4	306.2	305.8	296.5	288.6
TSA	9.8	9.6	9.9	9.7	8.8	9.4
St.CPUE	-24.1	-24.1	-24.1	-23.8	-23.2	-23.2
TLP	110.8	109.7	110.5	110.6	108.4	105.4
WLP	39.0	39.6	38.6	38.8	41.4	42.5
CLP	48.4	48.9	48.3	48.3	54.1	50.2
OBS	20.4	19.9	20.3	20.4	19.4	20.2
REC	2.6	2.7	2.4	2.5	1.8	1.9
WN	18.1	18.3	18.1	18.1	18.8	18.8
TAG	81.2	30.0	81.2	81.2	65.0	61.8
BMSY(mil.lb)	4.66	4.70	3.40	4.00	6.72	5.13
MMB(mil.lb)	3.98	3.87	2.86	3.35	5.45	4.66
Legal crab Catchable (mil.lb)	2.53	2.46	1.78	2.10	2.37	2.18
OFL(mil.lb)	0.31	0.29	0.22	0.26	0.46	0.60
NOAA q	0.70	0.68	1	0.81	0.66	0.71
ADFG q	1	1	1.40	1.20	1	1
M	018/0.58	018/0.64	018/0.52	018/0.55	0.31	0.43

20
 21 TSA: Trawl Survey Abundance
 22 St. CPUE: Summer commercial catch standardized CPUE
 23 TLP: Trawl survey length composition:
 24 WLP: Winter pot survey length composition

- 1 CLP: Summer commercial retention catch length composition
- 2 REC: Recruitment deviation
- 3 OBS: Summer commercial catch observer discards (Baseline) or total catch (Alternative models) length composition
- 4 TAG: Tagging recovery data composition
- 5 WN: Winter Commercial length-shell composition

6
7 See Appendix C1-C3 for standard output figures and estimated parameters.
8
9

10 Search for balance:

11 SSC noted in 2019 that model choice does not have much impact on the results, or on the Tier
12 4 reference points, which was also true for the 2020 assessment. The only meaningful change
13 occurs when we change assumptions about survey and fishery data selectivity and q , natural
14 mortality, and fate of large crabs, in other words, changing assumptions and understandings
15 about biology of the NSRKC that are significantly lacking.

16 Using only 1st year molting (Model 19.0 vs. 19.1) resulted in slight changes in transition matrix
17 (Table 14), and this did not improve model fit, MMB, and likelihood (Figure 4,8,9,11). Thus,
18 including more than 1 years of recovery data appeared to have little effects on estimation of
19 size transition matrix and the NSRKC assessment model. Estimating ADFG survey q was
20 greater than 1.0 (Model 19.2, 19.3), indicating that ADFG trawl survey overestimates NSRKC
21 abundance (Figure 7). This lowered MMB and OFL from the baseline (Figure 5). Assuming
22 dome shape selectivity and M (Model 19.4, 19.5) resulted in higher natural mortality and
23 higher MMB (Figure 6), indicating that NSRKC having greater natural mortality than assumed
24 0.18 and that larger crabs exists in Norton Sound that have never been observed or caught by
25 summer trawl survey or summer commercial fisheries. Under Tier 4 harvest control rule higher
26 natural mortality result in higher OFL (though they are lower than Tier 3 OFL: NPFMC 2019).
27 It is possible that the above observations could change with addition data that need to be
28 finalized and incorporated into the final assessment in January 2020. At this time, we
29 recommend either Model 19.0 or 19.1 for January 2020 final assessment.
30

31
32 **4. Results :**

33
34 1. List of effective sample sizes and weighting factors (Figure 4)

35 “Implied” effective sample sizes were calculated as

36
$$n = \sum_l \hat{P}_{y,l} (1 - \hat{P}_{y,l}) / \sum_l (P_{y,l} - \hat{P}_{y,l})^2$$

37 Where $P_{y,l}$ and $\hat{P}_{y,l}$ are observed and estimated length compositions in year y and length
38 group l , respectively. Estimated effective sample sizes vary greatly over time.
39

40 Maximum sample sizes for length proportions:
41

Survey data	Sample size
Summer commercial, winter pot, and summer observer	minimum of $0.1 \times$ actual sample size or 10
Summer trawl and pot survey	minimum of $0.5 \times$ actual sample size or 20
Tag recovery	$0.5 \times$ actual sample size

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

Weighting factor

Recruitment SD 0.5

2. Tables of estimates.

- a. Model parameter estimates (Tables 11, 12). (To be completed for final assessment Jan 2020)
- b. Abundance and biomass time series (Table 15) (To be completed for final assessment Jan 2020)
- c. Recruitment time series (Table 15). (To be completed for final assessment Jan 2020)
- d. Time series of catch/biomass (Tables 16) (To be completed for final assessment Jan 2020)

3. Graphs of estimates.

- a. Molting probability and trawl/pot selectivity
- b. Estimated male abundances (recruits, legal, and total)
- c. Estimated mature male biomass
- e. Time series of catch and estimated harvest rate

4. Evaluation of the fit to the data.

- a. Fits to observed and model predicted catches.
Not applicable. Catch is assumed to be measured without error.
- a. Model fits to survey numbers.
 - 1. Time series of trawl survey
 - 2. Time series of standardized cpue for the summer commercial fishery
- d. Model fits to catch and survey proportions by length
- e. Marginal distribution for the fits to the composition data

f. Plots of implied versus input effective sample sizes and time-series of implied effective sample size. Will be presented on Jan 2019 final assessment.

g. RMSEs of trawl survey and standardized CPUE. Will be presented on Jan 2019 final assessment.

QQ plots and histograms of residuals of trawl survey and standardized CPUE Will be presented on Jan 2019 final assessment.

h.

5. Retrospective analyses. Will be presented on Jan 2019 final assessment.

6. Uncertainty and sensitivity analyses.

A. Calculation of the OFL

1. Specification of the Tier level and stock status.

The Norton Sound red king crab stock is placed in Tier 4. It is not possible to estimate the spawner-recruit relationship, but some abundance and harvest estimates are available to build a computer simulation model that captures the essential population dynamics. Tier 4 stocks are assumed to have reliable estimates of current survey biomass and instantaneous M ; however, the estimates for the Norton Sound red king crab stock are uncertain.

Tire 4 level and the OFL are determined by the F_{MSY} proxy, B_{MSY} proxy, and estimated legal male abundance and biomass:

Level	Criteria	F_{OFL}
a	$B / B_{MSY\ proxy} > 1$	$F_{OFL} = \gamma M$
b	$\beta < B / B_{MSY\ proxy} \leq 1$	$F_{OFL} = \gamma M (B / B_{MSY\ proxy} - \alpha) / (1 - \alpha)$
c	$B / B_{MSY\ proxy} \leq \beta$	$F_{OFL} = \text{bycatch mortality \& directed fishery } F = 0$

1 where B is a mature male biomass (MMB), B_{MSY} proxy is average mature male biomass over a
 2 specified time period, $M = 0.18$, $\gamma = 1$, $\alpha = 0.1$, and $\beta = 0.25$

3

4 For Norton Sound red king crab, MMB is defined as the biomass of males > 94 mm CL on February
 5 01 (Appendix A). B_{MSY} proxy is

6

7 B_{MSY} proxy = average model estimated MMB from 1980-2020

8 **Estimated B_{MSY} proxy is: million lb.**

9

10 Predicted mature male biomass in 2020 on February 01

11

12 **Mature male biomass : million lb.**

13

14 Since projected MMB is less than B_{MSY} proxy,

15 **Norton Sound red king crab stock status is Tier 4b**

16 Where FOFL is calculated by

$$17 \quad F_{OFL} = \gamma M (B / B_{MSY\ proxy} - \alpha) / (1 - \alpha)$$

18 **FOFL of for all length classes**

19

20 2. Calculation of OFL.

21

22 OFL was calculated for retained (OFL_r), un-retained (OFL_{ur}), and total (OFL_T) for legal sized crab,
 23 $Legal_B$, by applying F_{OFL} .

24

25 $Legal_B$ is a biomass of legal crab subject to fisheries and is calculated as: Projected abundance by
 26 length crab \times fishing selectivity by length class \times Proportion of legal crab per length class \times Average
 27 lb per length class.

28 For the Norton Sound red king crab assessment, $Legal_B$ was defined as winter biomass catchable to
 29 summer commercial pot fishery gear $Legal_B_w$, as

$$30 \quad Legal_B_w = \sum_l (N_{w,l} + O_{w,l}) S_{s,l} P_{lg,l} w m_l$$

31 The Norton Sound red king crab fishery consists of two distinct fisheries: winter and summer. The
 32 two fisheries are discontinuous with 5 months between the two fisheries during which natural
 33 mortalities occur. To incorporate this fishery, the CPT in 2016 recommended the following formula:

$$34 \quad Legal_B_s = Legal_B_w (1 - \exp(-x \cdot F_{OFL})) e^{-0.42M}$$

1 $OFL_r = (1 - \exp(-(1 - x) \cdot F_{OFL}))Legal_B_s$

2 And $p = \frac{Legal_B_w(1 - \exp(-x \cdot F_{OFL}))}{OFL_r}$

3 Where p is a specific proportion of winter crab harvest to total (winter + summer) harvest.

4
5 Solving x of the above, a revised retained OFL is

6 $OFL = Legal_B_w \left(1 - e^{-(F_{OFL} + 0.42M)} - (1 - e^{-0.42M}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL} + 0.42M)})}{1 - p \cdot (1 - e^{-0.42M})} \right) \right)$

7
8 Accounting for difference in length specific natural mortality

9 $OFL_r = \sum_l \left[Legal_B_{w,l} \left(1 - e^{-(F_{OFL,l} + 0.42M_l)} - (1 - e^{-0.42M_l}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL,l} + 0.42M_l)})}{1 - p \cdot (1 - e^{-0.42M_l})} \right) \right) \right]$

10

11 Unretained OFL (OFL_{ur}) is a sub-legal crab biomass catchable to summer commercial pot fisheries
12 calculated as: Projected legal abundance (Feb 1st) \times Commercial pot selectivity \times Proportion of sub-
13 legal crab per length class \times Average lb per length class \times handling mortality ($hm = 0.2$)

14

15 $OFL_{ur} = \sum_l \left[Sub_legal_B_{w,l} \left(1 - e^{-(F_{OFL,l} + 0.42M_l)} - (1 - e^{-0.42M_l}) \left(\frac{1 - p \cdot (1 - e^{-(F_{OFL,l} + 0.42M_l)})}{1 - p \cdot (1 - e^{-0.42M_l})} \right) \right) \right] \cdot hm$

16

17 The total male OFL is

18 $OFL_T = OFL_r + OFL_{ur}$

19

20 For calculation of the OFL 2020, we specified $p = 0.16$.

21

22 Legal male biomass catchable to fishery (Feb 01): 2. million lb

23 **OFL_r = TBD**

24 **OFL_{nr} = TBD**

25 **OFL_T = TBD**

26

27 **B. Calculation of the ABC**

28

1 1. Specification of the probability distribution of the OFL.
2 Probability distribution of the OFL was determined based on the CPT recommendation in
3 January 2015 of 20% buffer:

4 Retained ABC for legal male crab is 80% of OFL

5
6 **ABC = TBD**

7 8 9 ***C. Rebuilding Analyses***

10 Not applicable

11

12 ***D. Data Gaps and Research Priorities***

13

14 The major data gap is the fate of crab greater than 123 mm.

15

16

17

Acknowledgments

18 We thank all CPT members for all review of the assessment model and suggestions for
19 improvements and diagnoses. We also thank Dr. Shareef Siddeek for critical review of draft.

20

21

22 **References**

23

24 Fournier, D., and C.P. Archibald. 1982. A general theory for analyzing catch at age data. Can. J. Fish.
25 Aquat. Sci. 39:1195-1207.

26 Fournier, D.A., H.J. Skaug, J. Ancheta, J. Ianelli, A. Magnusson, M.N. Maunder, A. Nielsen, and J.
27 Sibert. 2012. AD Model Builder: using automatic differentiation for statistical inference of
28 highly parameterized complex nonlinear models. Optim. Methods Softw. 27:233-249.

29 Menard, J., J. Soong, and S. Kent 2011. 2009 Annual Management Report Norton Sound, Port
30 Clarence, and Kotzebue. Fishery Management Report No. 11-46.

31 Methot, R.D. 1989. Synthetic estimates of historical abundance and mortality for northern anchovy.
32 Amer. Fish. Soc. Sym. 6:66-82.

33 Mohn, R. 1999. The retrospective problem in sequential population analysis: An investigation using
34 cod fishery and simulated data. ICES Journal of Marine Science, 56:473-488.

35 NPFMC 2011. Stock assessment and fishery evaluation report for the King and Tanner crab
36 fisheries of the Bering Sea and Aleutian Islands regions. 2011 Crab SAFE. North Pacific
37 Fishery Management Council, Anchorage, AK, USA

- 1
2 NPFMC 2012. Stock assessment and fishery evaluation report for the King and Tanner crab
3 fisheries of the Bering Sea and Aleutian Islands regions. 2012 Crab SAFE. North Pacific
4 Fishery Management Council, Anchorage, AK, USA
5
6 NPFMC 2013. Stock assessment and fishery evaluation report for the King and Tanner crab
7 fisheries of the Bering Sea and Aleutian Islands regions. 2013 Crab SAFE. North Pacific
8 Fishery Management Council, Anchorage, AK, USA
9
10 NPFMC 2014. Stock assessment and fishery evaluation report for the King and Tanner crab
11 fisheries of the Bering Sea and Aleutian Islands regions. 2014 Crab SAFE. North Pacific
12 Fishery Management Council, Anchorage, AK, USA
13
14 NPFMC 2015. Stock assessment and fishery evaluation report for the King and Tanner crab
15 fisheries of the Bering Sea and Aleutian Islands regions. 2015 Crab SAFE. North Pacific
16 Fishery Management Council, Anchorage, AK, USA
17
18 NPFMC 2016. Stock assessment and fishery evaluation report for the King and Tanner crab
19 fisheries of the Bering Sea and Aleutian Islands regions. 2016 Crab SAFE. North Pacific
20 Fishery Management Council, Anchorage, AK, USA
21
22 NPFMC 2017. Stock assessment and fishery evaluation report for the King and Tanner crab
23 fisheries of the Bering Sea and Aleutian Islands regions. 2017 Crab SAFE. North Pacific
24 Fishery Management Council, Anchorage, AK, USA
25
26 NPFMC 2018. Stock assessment and fishery evaluation report for the King and Tanner crab
27 fisheries of the Bering Sea and Aleutian Islands regions. 2018 Crab SAFE. North Pacific
28 Fishery Management Council, Anchorage, AK, USA
29
30 Powell, G.C., R. Peterson, and L. Schwarz. 1983. The red king crab, *Paralithodes camtschatica*
31 (Tilesius), in Norton Sound, Alaska: History of biological research and resource utilization
32 through 1982. Alaska Dept. Fish and Game, Inf. Leaflet 222. 103 pp.
- 33 Zheng, J., G.H. Kruse, and L. Fair. 1998. Use of multiple data sets to assess red king crab,
34 *Paralithodes camtschaticus*, in Norton Sound, Alaska: A length-based stock synthesis
35 approach. Pages 591-612 *In* Fishery Stock Assessment Models, edited by F. Funk, T.J. Quinn
36 II, J. Heifetz, J.N. Ianelli, J.E. Powers, J.F. Schweigert, P.J. Sullivan, and C.-I. Zhang, Alaska
37 Sea Grant College Program Report No. AK-SG-98-01, University of Alaska Fairbanks

Table 1. Historical summer commercial red king crab fishery economic performance, Norton Sound Section, eastern Bering Sea, 1977-2017. Bold type shows data that are used for the assessment model.

Year	Guideline Harvest Level (lb) ^b	Commercial Harvest (lb) ^{a,b}		Number Harvest	Total Number (Open Access)			Total Pots		ST CPUE		Season Length		Mid-day from July
		Open Access	CDQ		Vessels	Permits	Landings	Registered	Pulls	CPUE	SD	Days	Dates	
1977	^c	517.787		195,877	7	7	13		5,457	3.29	0.68	60	^c	0.049
1978	3,000.000	2,091.961		660,829	8	8	54		10,817	4.68	0.65	60	6/07-8/15	0.142
1979	3,000.000	2,931.672		970,962	34	34	76		34,773	2.87	0.64	16	7/15-7/31	0.088
1980	1,000.000	1,186.596		329,778	9	9	50		11,199	3.07	0.65	16	7/15-7/31	0.066
1981	2,500.000	1,379.014		376,313	36	36	108		33,745	0.86	0.64	38	7/15-8/22	0.096
1982	500.000	228.921		63,949	11	11	33		11,230	0.20	0.62	23	8/09-9/01	0.151
1983	300.000	368.032		132,205	23	23	26	3,583	11,195	0.90	0.65	3.8	8/01-8/05	0.096
1984	400.000	387.427		139,759	8	8	21	1,245	9,706	1.59	0.65	13.6	8/01-8/15	0.110
1985	450.000	427.011		146,669	6	6	72	1,116	13,209	0.50	0.66	21.7	8/01-8/23	0.118
1986	420.000	479.463		162,438	3	3		578	4,284	1.74	0.70	13	8/01-8/25	0.153
1987	400.000	327.121		103,338	9	9		1,430	10,258	0.61	0.64	11	8/01-8/12	0.107
1988	200.000	236.688		76,148	2	2		360	2,350	2.36	0.86	9.9	8/01-8/11	0.110
1989	200.000	246.487		79,116	10	10		2,555	5,149	1.21	0.61	3	8/01-8/04	0.096
1990	200.000	192.831		59,132	4	4		1,388	3,172	1.08	0.68	4	8/01-8/05	0.099
1991	340.000			0	No Summer Fishery									
1992	340.000	74.029		24,902	27	27		2,635	5,746	0.17	0.60	2	8/01-8/03	0.093
1993	340.000	335.790		115,913	14	20	208	560	7,063	0.90	0.35	52	7/01-8/28	0.093
1994	340.000	327.858		108,824	34	52	407	1,360	11,729	0.81	0.34	31	7/01-7/31	0.044
1995	340.000	322.676		105,967	48	81	665	1,900	18,782	0.42	0.34	67	7/01-9/05	0.093
1996	340.000	224.231		74,752	41	50	264	1,640	10,453	0.51	0.34	57	7/01-9/03	0.101
1997	80.000	92.988		32,606	13	15	100	520	2,982	0.84	0.35	44	7/01-8/13	0.074
1998	80.000	29.684	0.00	10,661	8	11	50	360	1,639	0.79	0.36	65	7/01-9/03	0.110
1999	80.000	23.553	0.00	8,734	10	9	53	360	1,630	0.92	0.36	66	7/01-9/04	0.104
2000	336.000	297.654	14.87	111,728	15	22	201	560	6,345	1.24	0.34	91	7/01-9/29	0.126
2001	303.000	288.199	0	98,321	30	37	319	1,200	11,918	0.64	0.34	97	7/01-9/09	0.104
2002	248.000	244.376	15.226	86,666	32	49	201	1,120	6,491	1.23	0.34	77	6/15-9/03	0.060
2003	253.000	253.284	13.923	93,638	25	43	236	960	8,494	0.85	0.34	68	6/15-8/24	0.058
2004	326.500	314.472	26.274	120,289	26	39	227	1,120	8,066	1.27	0.34	51	6/15-8/08	0.033
2005	370.000	370.744	30.06	138,926	31	42	255	1,320	8,867	1.19	0.34	73	6/15-8/27	0.058
2006	454.000	419.191	32.557	150,358	28	40	249	1,120	8,867	1.31	0.34	68	6/15-8/22	0.052
2007	315.000	289.264	23.611	110,344	38	30	251	1,200	9,118	1.02	0.34	52	6/15-8/17	0.036
2008	412.000	364.235	30.9	143,337	23	30	248	920	8,721	1.32	0.34	73	6/23-9/03	0.079
2009	375.000	369.462	28.125	143,485	22	27	359	920	11,934	0.84	0.34	98	6/15-9/20	0.090
2010	400.000	387.304	30	149,822	23	32	286	1,040	9,698	1.22	0.34	58	6/28-8/24	0.074
2011	358.000	373.990	26.851	141,626	24	25	173	1,040	6,808	1.58	0.34	33	6/28-7/30	0.038
2012	465.450	441.080	34.91	161,113	40	29	312	1,200	10,041	1.29	0.34	72	6/29-9/08	0.093
2013	495.600	373.278	18.585	130,603	37	33	460	1,420	15,058	0.67	0.33	74	7/3-9/14	0.110
2014	382.800	360.860	28.148	129,657	52	33	309	1,560	10,127	1.12	0.34	52	6/25-8/15	0.052
2015	394.600	371.520	29.595	144,255	42	36	251	1,480	8,356	1.45	0.34	26	6/29-7/24	0.033
2016	517.200	416.576	3,583	138,997	36	37	220	1,520	8,009	1.27	0.34	25	6/27-7/21	0.025
2017	496,800	411,736	0	135,322	36	36	270	1,640	9,401	1.10	0.34	30	6/26-7/25	0.027
2018	319,400	298,396	0	89,613	34	34	256	1,400	8,797	0.64	0.34	35	6/24-7/29	0.038
2019	150,600	73,784	1,239	24,506	24	26	146	1,096	5,438	0.26	0.34	62	6/25-9/03	

^a Deadloss included in total. ^b Millions of pounds. ^c Information not available.

Table 2. Historical winter commercial and subsistence red king crab fisheries, Norton Sound Section, eastern Bering Sea, 1977-2016. Bold typed data are used for the assessment model.

Model Year	Year ^a	Commercial			Subsistence			Total Crab	
		# of Fishers	# of Crab Harvested	Winter ^b	Issued	Permits Returned	Fished	Caught ^c	Retained ^d
1978	1978	37	9,625	1977/78	290	206	149	NA	12,506
1979	1979	1 ^f	221^f	1978/79	48	43	38	NA	224
1980	1980	1 ^f	22^f	1979/80	22	14	9	NA	213
1981	1981	0	0	1980/81	51	39	23	NA	360
1982	1982	1 ^f	17^f	1981/82	101	76	54	NA	1,288
1983	1983	5	549	1982/83	172	106	85	NA	10,432
1984	1984	8	856	1983/84	222	183	143	15,923	11,220
1985	1985	9	1,168	1984/85	203	166	132	10,757	8,377
1986	1985/86	5	2,168	1985/86	136	133	107	10,751	7,052
1987	1986/87	7	1,040	1986/87	138	134	98	7,406	5,772
1988	1987/88	10	425	1987/88	71	58	40	3,573	2,724
1989	1988/89	5	403	1988/89	139	115	94	7,945	6,126
1990	1989/90	13	3,626	1989/90	136	118	107	16,635	12,152
1991	1990/91	11	3,800	1990/91	119	104	79	9,295	7,366
1992	1991/92	13	7,478	1991/92	158	105	105	15,051	11,736
1993	1992/93	8	1,788	1992/93	88	79	37	1,193	1,097
1994	1993/94	25	5,753	1993/94	118	95	71	4,894	4,113
1995	1994/95	42	7,538	1994/95	166	131	97	7,777	5,426
1996	1995/96	9	1,778	1995/96	84	44	35	2,936	1,679
1997	1996/97	2 ^f	83^f	1996/97	38	22	13	1,617	745
1998	1997/98	5	984	1997/98	94	73	64	20,327	8,622
1999	1998/99	5	2,714	1998/99	95	80	71	10,651	7,533
2000	1999/00	10	3,045	1999/00	98	64	52	9,816	5,723
2001	2000/01	3	1,098	2000/01	50	27	12	366	256
2002	2001/02	11	2,591	2001/02	114	61	45	5,119	2,177
2003	2002/03	13	6,853	2002/03	107	70	61	9,052	4,140
2004	2003/04	2 ^f	522^f	2003/04 ^h	96	77	41	1,775	1,181
2005	2004/05	4	2,091	2004/05	170	98	58	6,484	3,973
2006	2005/06	1 ^f	75^f	2005/06	98	97	67	2,083	1,239
2007	2006/07	8	3,313	2006/07	129	127	116	21,444	10,690
2008	2007/08	9	5,796	2007/08	139	137	108	18,621	9,485
2009	2008/09	7	4,951	2008/09	105	105	70	6,971	4,752
2010	2009/10	10	4,834	2009/10	125	123	85	9,004	7,044
2011	2010/11	5	3,365	2010/11	148	148	95	9,183	6,640
2012	2011/12	35	9,157	2011/12	204	204	138	11,341	7,311
2013	2012/13	26	22,639	2012/13	149	148	104	21,524	7,622
2014	2013/14	21	14,986	2013/14	103	103	75	5,421	3,252
2015	2014/15	44	41,062	2014/15	155	153	107	9,840	7,651
2016	2015/16	25	29,792	2015/16	139	97	64	6,468	5,340
2017	2017	43	26,008	2017	163	163	109	7,185	6,039
2018	2018	28	9,180	2018	123	120	82	5,767	4,424
2019	2019	6	1,050	2019	101	101	60	2,080	1,545

a Prior to 1985 the winter commercial fishery occurred from January 1 - April 30. As of March 1985, fishing may occur from November 15 - May 15.

b The winter subsistence fishery occurs during months of two calendar years (as early as December, through May).

c The number of crab actually caught; some may have been returned.

d The number of crab retained is the number of crab caught and kept.

f Confidentiality was waived by the fishers.

h Prior to 2005, permits were only given out of the Nome ADF&G office. Starting with the 2004-5 season, permits were given out in Elim, Golovin, Shaktoolik, and White Mountain.

Table 3. Summary of triennial trawl survey Norton Sound male red king crab abundance estimates (CL \geq 64mm) . Trawl survey abundance estimate is based on 10 \times 10 nm² grid, except for 2010 and 2017 (20 \times 20 nm²). Bold typed data are used for the assessment model.

Year	Dates	Survey Agency	Survey method	Survey coverage			Abundance \geq 64 mm	
				Total surveyed hauls	Stations w/ NSRKC	n mile ² expanded		CV
1976	9/02 – 9/25	NMFS	Trawl	117	61	7600	4301.8	0.31
1979	7/26 - 8/05	NMFS	Trawl	115	33	7600	1457.4	0.22
1980	7/04 - 7/14	ADFG	Pots				2092.3	N/A
1981	6/28 - 7/14	ADFG	Pots				2153.4	N/A
1982	7/06 - 7/20	ADFG	Pots				1140.5	N/A
1982	9/05 - 9/11	NMFS	Trawl	57	46	7600	3548.9	0.25
1985	7/01 - 7/14	ADFG	Pots				2320.4	0.083
1985	9/16 -10/01	NMFS	Trawl	78	58	7600	2424.9	0.26
1988	8/16 - 8/30	NMFS	Trawl	82	45	7600	2702.3	0.29
1991	8/22 - 8/30	NMFS	Trawl	51	38	7600	4049.1	0.40
1996	8/07 - 8/18	ADFG	Trawl	50	30	4938	1283.0	0.25
1999	7/28 - 8/07	ADFG	Trawl	52	31	5221	2608.0	0.24
2002	7/27 - 8/06	ADFG	Trawl	57	37	5621	2056.0	0.36
2006	7/25 - 8/08	ADFG	Trawl	114	45	6000	3336.0	0.39
2008	7/24 - 8/11	ADFG	Trawl	86	44	7330	2894.2	0.31
2010 ^a	7/27 - 8/09	NMFS	Trawl	35	15	5841	1980.1	0.44
2011	7/18 - 8/15	ADFG	Trawl	65	34	6447	3209.3	0.29
2014	7/18 - 7/30	ADFG	Trawl	47	34	4700	5934.6	0.47
2017	7/28 - 8/08	ADFG	Trawl	60	41	6000	1762.1	0.22
2017	8/18 - 8/29	NMFS	Trawl	35	18	5841	1035.8	0.40
2018	7/22 - 7/29	ADFG	Trawl	60	34	6000	1108.9	0.25
2019	7/17-7/29	ADFG	Trawl	52	27	5221	4660.8	0.60

Abundance of NMFS survey (1976-1991) was estimated by NMFS, multiplying the mean CPUE (# NRKC/NM²) across all hauls (including re-tows) to a standard survey area (7600NM²).

In contrast, abundance of ADFG (1996-2019) and NMFS (2010,2017) survey were estimated by ADFG by multiplying CPUE (# NRKC/NM²) of each station to an area represented by the station (~100NM²) and summing across all surveyed station (ADFG: 4700 – 5200NM². NOAA 5841 NM²).

Table 4. Summer commercial retained catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1977	1549	0	0	0	0.00	0.42	0.34	0.08	0.05	0	0	0	0.00	0.06	0.04	0.01	0.00
1978	389	0	0	0	0.01	0.19	0.47	0.26	0.04	0	0	0	0.00	0.01	0.01	0.01	0.00
1979	1660	0	0	0	0.03	0.23	0.38	0.26	0.07	0	0	0	0.00	0.03	0.00	0.00	0.01
1980	1068	0	0	0	0.00	0.10	0.31	0.37	0.18	0	0	0	0.00	0.00	0.01	0.02	0.01
1981	1784	0	0	0	0.00	0.07	0.15	0.28	0.23	0	0	0	0.00	0.00	0.05	0.12	0.09
1982	1093	0	0	0	0.04	0.19	0.16	0.22	0.29	0	0	0	0.00	0.01	0.02	0.03	0.03
1983	802	0	0	0	0.04	0.41	0.36	0.06	0.03	0	0	0	0.00	0.04	0.01	0.02	0.02
1984	963	0	0	0	0.10	0.42	0.28	0.06	0.01	0	0	0	0.01	0.07	0.05	0.01	0.00
1985	2691	0	0	0.00	0.06	0.31	0.37	0.15	0.02	0	0	0	0.00	0.03	0.03	0.01	0.00
1986	1138	0	0	0	0.03	0.36	0.39	0.12	0.02	0	0	0	0.00	0.02	0.04	0.02	0.00
1987	1985	0	0	0	0.02	0.18	0.29	0.27	0.11	0	0	0	0.00	0.03	0.06	0.03	0.01
1988	1522	0	0.00	0	0.02	0.20	0.30	0.18	0.04	0	0	0	0.01	0.06	0.10	0.07	0.02
1989	2595	0	0	0	0.01	0.16	0.32	0.17	0.05	0	0	0	0.00	0.06	0.12	0.09	0.02
1990	1289	0	0	0	0.01	0.14	0.35	0.26	0.07	0	0	0	0.00	0.04	0.07	0.05	0.01
1991																	
1992	2566	0	0	0	0.02	0.20	0.27	0.14	0.09	0	0	0	0.00	0.08	0.13	0.06	0.02
1993	17804	0	0	0	0.01	0.23	0.39	0.23	0.03	0	0	0	0.00	0.02	0.04	0.03	0.01
1994	404	0	0	0	0.02	0.09	0.08	0.07	0.02	0	0	0	0.02	0.19	0.25	0.20	0.05
1995	1167	0	0	0	0.04	0.26	0.29	0.15	0.05	0	0	0	0.01	0.05	0.07	0.06	0.01
1996	787	0	0	0	0.03	0.22	0.24	0.09	0.05	0	0	0	0.01	0.12	0.14	0.08	0.02
1997	1198	0	0	0	0.03	0.37	0.34	0.10	0.03	0	0	0	0.00	0.06	0.04	0.03	0.01
1998	1055	0	0	0	0.03	0.23	0.24	0.08	0.03	0	0	0	0.02	0.11	0.14	0.08	0.03
1999	562	0	0	0	0.06	0.29	0.24	0.18	0.09	0	0	0	0.00	0.02	0.05	0.04	0.00
2000	17213	0	0	0	0.02	0.30	0.39	0.11	0.02	0	0	0	0.00	0.05	0.07	0.04	0.01
2001	20030	0	0	0	0.02	0.22	0.37	0.21	0.07	0	0	0	0.00	0.02	0.05	0.02	0.01
2002	5219	0	0	0	0.04	0.23	0.28	0.25	0.07	0	0	0	0.00	0.03	0.04	0.03	0.01
2003	5226	0	0	0	0.02	0.37	0.32	0.12	0.03	0	0	0	0.00	0.02	0.05	0.05	0.01
2004	9606	0	0	0	0.01	0.38	0.39	0.11	0.03	0	0	0	0.00	0.03	0.03	0.01	0.01
2005	5360	0	0	0	0.00	0.25	0.47	0.16	0.02	0	0	0	0.00	0.02	0.05	0.02	0.01
2006	6707	0	0	0	0.00	0.18	0.35	0.17	0.02	0	0	0	0.00	0.05	0.14	0.07	0.01
2007	6125	0	0	0	0.01	0.36	0.34	0.14	0.03	0	0	0	0.00	0.02	0.06	0.03	0.01
2008	5766	0	0	0	0.00	0.35	0.35	0.06	0.01	0	0	0	0.00	0.09	0.09	0.04	0.01
2009	6026	0	0	0	0.01	0.34	0.33	0.11	0.02	0	0	0	0.00	0.08	0.08	0.02	0.01
2010	5902	0	0	0	0.01	0.39	0.36	0.10	0.01	0	0	0	0.00	0.05	0.05	0.02	0.00
2011	2552	0	0	0	0.00	0.32	0.40	0.12	0.02	0	0	0	0.00	0.06	0.06	0.02	0.00
2012	5056	0	0	0	0.00	0.24	0.46	0.18	0.02	0	0	0	0.00	0.03	0.04	0.02	0.00
2013	6072	0	0	0	0.00	0.24	0.37	0.24	0.06	0	0	0	0.00	0.01	0.04	0.02	0.00
2014	4682	0	0	0	0.01	0.28	0.24	0.18	0.07	0	0	0	0.00	0.04	0.09	0.07	0.02
2015	4173	0	0	0	0.01	0.48	0.28	0.10	0.03	0	0	0	0.00	0.02	0.03	0.03	0.01
2016	1543	0	0	0	0.00	0.25	0.47	0.16	0.03	0	0	0	0.00	0.02	0.02	0.03	0.01
2017	3412	0	0	0	0.00	0.18	0.39	0.21	0.03	0	0	0	0.01	0.03	0.12	0.05	0.01
2018	2609	0	0	0	0.00	0.11	0.32	0.32	0.08	0	0	0	0	0.01	0.08	0.08	0.02
2019	1136	0	0	0	0.01	0.32	0.23	0.13	0.03	0	0	0	0	0.02	0.10	0.14	0.03

Table 5. Winter commercial catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
2015	576	0	0	0	0.07	0.50	0.24	0.06	0.01	0	0	0	0.01	0.04	0.03	0.03	0.01
2016	1016	0	0	0	0.03	0.45	0.31	0.03	0.00	0	0	0	0.01	0.09	0.04	0.02	0.01
2017	540	0	0	0	0.00	0.20	0.30	0.13	0.02	0	0	0	0.00	0.08	0.19	0.06	0.02
2018	401	0	0	0	0.00	0.11	0.25	0.27	0.05	0	0	0	0	0.04	0.16	0.10	0.02
2019	0																

Table 6. Summer Trawl Survey length-shell compositions.

Year	Survey	Sample	New Shell								Old Shell							
			64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1976	NMFS	1326	0.01	0.02	0.10	0.19	0.34	0.18	0.02	0.00	0.00	0.00	0.01	0.02	0.03	0.04	0.01	0.01
1979	NMFS	220	0.01	0.01	0.00	0.02	0.05	0.05	0.03	0.01	0.01	0.00	0.01	0.04	0.14	0.40	0.19	0.03
1982	NMFS	327	0.22	0.07	0.16	0.23	0.17	0.03	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.02	0.02	0.03
1985	NMFS	350	0.11	0.11	0.19	0.17	0.16	0.06	0.01	0.00	0.00	0.00	0.00	0.02	0.05	0.08	0.05	0.01
1988	NMFS	366	0.16	0.19	0.12	0.13	0.11	0.06	0.03	0.00	0.00	0.00	0.01	0.01	0.03	0.07	0.05	0.03
1991	NMFS	340	0.18	0.08	0.02	0.03	0.06	0.03	0.01	0.01	0.03	0.06	0.02	0.08	0.16	0.14	0.09	0.02
1996	ADFG	269	0.29	0.21	0.13	0.09	0.05	0.00	0.00	0.01	0.00	0.00	0.03	0.03	0.04	0.04	0.04	0.03
1999	ADFG	283	0.03	0.01	0.10	0.29	0.26	0.13	0.03	0.01	0.00	0.00	0.00	0.03	0.05	0.04	0.02	0.00
2002	ADFG	244	0.09	0.12	0.14	0.11	0.02	0.03	0.02	0.01	0.01	0.03	0.07	0.10	0.09	0.09	0.05	0.02
2006	ADFG	373	0.18	0.26	0.21	0.11	0.06	0.04	0.02	0.00	0.00	0.00	0.00	0.02	0.04	0.04	0.01	0.00
2008	ADFG	275	0.12	0.15	0.21	0.11	0.10	0.03	0.02	0.01	0.00	0.01	0.04	0.06	0.08	0.01	0.04	0.00
2010	NMFS	69	0.01	0.04	0.06	0.17	0.06	0.03	0.00	0.00	0.00	0.03	0.09	0.20	0.19	0.07	0.03	0.01
2011	ADFG	315	0.13	0.11	0.09	0.11	0.18	0.14	0.03	0.01	0.00	0.00	0.01	0.02	0.09	0.04	0.03	0.00
2014	ADFG	387	0.08	0.15	0.24	0.18	0.09	0.02	0.01	0.01	0.00	0.00	0.03	0.10	0.05	0.04	0.01	0.00
2017	ADFG	116	0.14	0.12	0.05	0.09	0.10	0.04	0.00	0.00	0.01	0.02	0.02	0.02	0.07	0.18	0.04	0.00
2017	NMFS	58	0.09	0.10	0.14	0.05	0.05	0.05	0.05	0.03	0.03	0.00	0.03	0.05	0.03	0.19	0.05	0.03
2018	ADFG	73	0.37	0.10	0.11	0.03	0.01	0.03	0.04	0.01	0	0.07	0.01	0.04	0.03	0.03	0.10	0.03
2019	ADFG	307	0.55	0.30	0.03	0	0.00	0.00	0.00	0	0.00	0.00	0.01	0.02	0.01	0.02	0.03	0.01
2019	NMFS																	

Table 7. Winter pot survey length-shell compositions.

Year	CPUE	Sample	New Shell								Old Shell							
			64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1981/82	NA	719	0.00	0.10	0.23	0.21	0.07	0.02	0.02	0.00	0.00	0.05	0.11	0.11	0.04	0.02	0.02	0.00
1982/83	24.2	2583	0.03	0.08	0.28	0.28	0.21	0.07	0.01	0.00	0.00	0.00	0.00	0.00	0.02	0.01	0.01	0.01
1983/84	24.0	1677	0.01	0.16	0.26	0.23	0.15	0.06	0.01	0.00	0.00	0.00	0.00	0.02	0.06	0.03	0.01	0.01
1984/85	24.5	789	0.02	0.09	0.25	0.35	0.16	0.06	0.01	0.00	0.00	0.00	0.00	0.01	0.03	0.02	0.00	0.00
1985/86	19.2	594	0.04	0.12	0.17	0.24	0.19	0.08	0.01	0.00	0.00	0.00	0.00	0.01	0.06	0.04	0.01	0.00
1986/87	5.8	144	0.00	0.06	0.15	0.19	0.07	0.04	0.00	0.00	0.00	0.00	0.00	0.01	0.04	0.30	0.11	0.03
1987/88																		
1988/89	13.0	500	0.02	0.13	0.15	0.13	0.19	0.17	0.03	0.00	0.00	0.00	0.00	0.05	0.08	0.03	0.00	0.00
1989/90	21.0	2076	0.00	0.05	0.21	0.26	0.18	0.12	0.06	0.01	0.00	0.00	0.00	0.00	0.03	0.06	0.02	0.00
1990/91	22.9	1283	0.00	0.01	0.09	0.29	0.27	0.10	0.01	0.00	0.00	0.00	0.00	0.00	0.03	0.12	0.07	0.02
1992/93	5.5	181	0.00	0.01	0.03	0.06	0.13	0.12	0.03	0.00	0.00	0.00	0.00	0.02	0.19	0.27	0.10	0.05
1993/94																		
1994/95	6.2	858	0.01	0.06	0.08	0.10	0.26	0.23	0.07	0.01	0.00	0.00	0.00	0.00	0.03	0.07	0.06	0.02
1995/96	9.9	1580	0.06	0.14	0.20	0.19	0.11	0.07	0.03	0.00	0.00	0.00	0.00	0.01	0.06	0.07	0.03	0.01
1996/97	2.9	398	0.07	0.21	0.22	0.11	0.15	0.11	0.05	0.01	0.00	0.00	0.00	0.00	0.02	0.03	0.01	0.01
1997/98	10.9	881	0.00	0.14	0.41	0.27	0.05	0.02	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.02	0.02	0.01
1998/99	10.7	1307	0.00	0.02	0.12	0.36	0.36	0.08	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.01	0.01	0.00
1999/00	6.2	575	0.02	0.09	0.10	0.16	0.33	0.18	0.03	0.00	0.00	0.00	0.00	0.00	0.05	0.02	0.01	0.00
2000/01	3.1	44																
2001/02	13.0	828	0.05	0.29	0.26	0.17	0.06	0.06	0.04	0.01	0.01	0.00	0.01	0.01	0.00	0.01	0.00	0.00
2002/03	9.6	824	0.02	0.10	0.22	0.28	0.18	0.06	0.02	0.00	0.00	0.01	0.01	0.02	0.02	0.03	0.02	0.01
2003/04	3.7	296	0.00	0.02	0.16	0.26	0.32	0.14	0.01	0.00	0.00	0.00	0.01	0.02	0.02	0.01	0.02	0.01
2004/05	4.4	405	0.00	0.07	0.14	0.18	0.22	0.19	0.07	0.00	0.00	0.00	0.00	0.00	0.04	0.06	0.01	0.00
2005/06	6.0	512	0.00	0.14	0.23	0.21	0.16	0.05	0.02	0.00	0.00	0.01	0.01	0.02	0.04	0.07	0.03	0.01
2006/07	7.3	159	0.07	0.14	0.19	0.35	0.13	0.04	0.00	0.00	0.00	0.00	0.01	0.01	0.02	0.04	0.00	0.00
2007/08	25.0	3552	0.01	0.14	0.25	0.17	0.14	0.07	0.01	0.00	0.01	0.04	0.07	0.03	0.03	0.01	0.01	0.00
2008/09	21.9	525	0.00	0.07	0.13	0.35	0.20	0.08	0.01	0.00	0.00	0.00	0.00	0.00	0.04	0.10	0.00	0.00
2009/10	25.3	578	0.01	0.05	0.13	0.21	0.24	0.11	0.02	0.00	0.00	0.00	0.01	0.06	0.10	0.05	0.01	0.00
2010/11	22.1	596	0.02	0.08	0.13	0.20	0.17	0.13	0.05	0.00	0.00	0.00	0.01	0.03	0.11	0.05	0.01	0.00
2011/12	29.4	675	0.03	0.11	0.23	0.19	0.12	0.13	0.04	0.00	0.00	0.00	0.00	0.01	0.05	0.05	0.03	0.00

Table 8. Summer commercial 1987-1994, 2012-2018 observer discards length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
1987	1146	0.06	0.19	0.32	0.33	0.03	0.00	0.00	0.00	0.00	0.00	0.02	0.04	0.00	0.00	0.00	0.00
1988	722	0.01	0.04	0.15	0.48	0.14	0.00	0.00	0.00	0.00	0.01	0.03	0.10	0.04	0.00	0.00	0.00
1989	1000	0.07	0.19	0.24	0.22	0.03	0.00	0.00	0.00	0.02	0.03	0.07	0.11	0.03	0.00	0.00	0.00
1990	507	0.08	0.23	0.27	0.27	0.04	0.00	0.00	0.00	0.02	0.02	0.02	0.05	0.01	0.00	0.00	0.00
1992	580	0.11	0.17	0.30	0.29	0.03	0.00	0.00	0.00	0.01	0.02	0.02	0.04	0.01	0.00	0.00	0.00
1994	850	0.07	0.06	0.11	0.15	0.02	0.00	0.00	0.00	0.07	0.07	0.15	0.24	0.05	0.00	0.00	0.00
2012	939	0.21	0.11	0.19	0.32	0.10	0.01	0.00	0.00	0.00	0.00	0.00	0.02	0.02	0.00	0.00	0.00
2013	2617	0.34	0.29	0.16	0.16	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2014	1755	0.05	0.10	0.26	0.41	0.12	0.01	0.00	0.00	0.00	0.00	0.01	0.03	0.01	0.00	0.00	0.00
2015	824	0.01	0.08	0.18	0.44	0.23	0.02	0.00	0.00	0.00	0.00	0.00	0.02	0.02	0.00	0.00	0.00
2016	426	0.04	0.05	0.17	0.50	0.17	0.02	0.00	0.00	0.00	0.00	0.00	0.03	0.01	0.00	0.00	0.00
2017	544	0.10	0.16	0.13	0.31	0.26	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00
2018	532	0.10	0.17	0.36	0.30	0.02	0.00	0.00	0.00	0.00	0.01	0.02	0.01	0.00	0.00	0.00	0.00
2019																	

Table 9. Summer commercial 2012-2018 observer total catch length-shell compositions.

Year	Sample	New Shell								Old Shell							
		64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
2012	3055	0.10	0.05	0.08	0.15	0.15	0.17	0.06	0.01	0.00	0.00	0.00	0.03	0.08	0.09	0.03	0.00
2013	4762	0.19	0.16	0.09	0.10	0.16	0.16	0.09	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.01	0.00
2014	3506	0.02	0.05	0.13	0.22	0.22	0.12	0.08	0.03	0.00	0.00	0.00	0.02	0.03	0.03	0.02	0.01
2015	1671	0.01	0.04	0.09	0.23	0.37	0.14	0.05	0.01	0.00	0.00	0.00	0.01	0.02	0.02	0.01	0.00
2016	2114	0.01	0.01	0.03	0.12	0.29	0.36	0.08	0.02	0.00	0.00	0.00	0.01	0.03	0.03	0.02	0.00
2017	2748	0.02	0.03	0.03	0.06	0.19	0.33	0.18	0.02	0.00	0.00	0.00	0.00	0.02	0.07	0.03	0.01
2018	1628	0.03	0.06	0.12	0.11	0.09	0.17	0.18	0.04	0.00	0.00	0.01	0.01	0.15	0.07	0.08	0.02
2019																	

Table 10. The number of tagged data released and recovered after 1 year (Y1) – 3 year (Y3) during 1980-1992 and 1993-2019 periods.

Release Length Class	Recap Length Class	1980-1992			1993-2019		
		Y1	Y2	Y3	Y1	Y2	Y3
64 – 73	64 – 73						
64 – 73	74 - 83	1					
64 – 73	84 - 93	1	1		3		
64 – 73	94 - 103		1			5	
64 – 73	104 – 113		1			4	11
64 – 73	114 – 123						11
64 – 73	124 – 133						
64 – 73	134+						
74 - 83	74 - 83						
74 - 83	84 - 93	3			21		
74 - 83	94 - 103	7			22	12	
74 - 83	104 – 113		13		4	94	19
74 - 83	114 – 123		1	2		5	46
74 - 83	124 – 133						6
74 - 83	134+						
84 - 93	84 - 93						
84 - 93	94 - 103	15	1		41	5	2
84 - 93	104 – 113	19	5	1	81	34	14
84 - 93	114 – 123		5	2	7	69	27
84 - 93	124 – 133				1	3	9
84 - 93	134+						6
94 - 103	94 - 103	4	1		7	2	
94 - 103	104 – 113	53	5	1	165	33	2
94 - 103	114 – 123	31	5	7	82	38	32
94 - 103	124 – 133	2	2	2		19	13
94 - 103	134+				1		
104 – 113	104 – 113	18			59	7	
104 – 113	114 – 123	38	15	3	109	64	9
104 – 113	124 – 133	7	8	4	15	18	18
104 – 113	134+						
114 – 123	114 – 123	17	2		72	9	
114 – 123	124 – 133	27	10	2	72	38	10
114 – 123	134+	5	1		19	6	3
124 – 133	124 – 133	15			41	9	1
124 – 133	134+	10	4	2	15	12	7
134+	134+	15	6	1	11	2	

Table 11. Summary of initial input parameter values and bounds for a length-based population model of Norton Sound red king crab. Parameters with “log_” indicate log scaled parameters.

Parameter	Parameter description	Lower	Upper
log $q_{1,2}$	Commercial fishery catchability (1977-92, 1993-2017)	-20.5	20
log N_{76}	Initial abundance	2.0	15.0
R_0	Mean Recruit	2.0	12.0
log σ_R^2	Recruit standard deviation	-40.0	40.0
a_{1-7}	Intimal length proportion	0	10.0
r_1	Proportion of length class 1 for recruit	0	10.0
log α	Inverse logistic molting parameter	-5.0	-1.0
log β	Inverse logistic molting parameter	1.0	5.5
log ϕ_{st1}	Logistic trawl selectivity parameter	-5.0	1.0
log ϕ_{wa}	Inverse logistic winter pot selectivity parameter	-5.0	1.0
log ϕ_{wb}	Inverse logistic winter pot selectivity parameter	0.0	6.0
$SW_{1,2}$	Winter pot selectivity of length class 1,2	0.1	1.0
log ϕ_1	Logistic commercial catch selectivity parameter	-5.0	1.0
log ϕ_2	Logistic commercial catch selectivity parameter	0.0	6.0
log acr	Logistic summer commercial retention selectivity parameter	-5.0	1.0
log bcr	Logistic summer commercial retention selectivity parameter	0.0	6.0
log awr	Logistic winter commercial retention selectivity parameter	-5.0	1.0
log bwr	Logistic winter commercial retention selectivity parameter	0.0	6.0
w^2_t	Additional variance for standard CPUE	0.0	6.0
ms	Natural mortality multipliers	0.5	5.0
q	Survey q for NMFS trawl 1976-91	0.1	1.0
σ	Growth transition sigma	0.0	30.0
β_1	Growth transition mean	0.0	20.0
β_2	Growth transition increment	0.0	20.0

Table 12. Estimated molting probability incorporated transition matrix (Model 19.0, 19.1).

Model 19.0								
Pre-molt Length Class	Post-molt Length Class							
	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
64 - 73	0.02	0.09	0.79	0.10	0.00	0.00	0.00	0.00
74 - 83		0.04	0.23	0.70	0.03	0.00	0.00	0.00
84 - 93			0.08	0.42	0.49	0.01	0.00	0.00
94 - 103				0.15	0.58	0.26	0.00	0.00
104 - 113					0.29	0.61	0.10	0.00
114 - 123						0.50	0.47	0.03
124 - 133							0.73	0.27
134+								1.00

Model 19.1

Pre-molt Length Class	Post-molt Length Class							
	64-73	74-83	84-93	94-103	104-113	114-123	124-133	134+
64 - 73	0.02	0.13	0.74	0.10	0.00	0.00	0.00	0.00
74 - 83		0.04	0.25	0.66	0.04	0.00	0.00	0.00
84 - 93			0.09	0.40	0.50	0.01	0.00	0.00
94 - 103				0.16	0.53	0.31	0.00	0.00
104 - 113					0.29	0.56	0.15	0.00
114 - 123						0.47	0.47	0.06
124 - 133							0.68	0.32
134+								1.00




Figure 1. King crab fishing districts and sections of Statistical Area Q.




Figure 2. Closed water regulations in effect for the Norton Sound commercial crab fishery. Line around the coastline delineates the 3-mile state waters zone.




Figure 3. Model estimated annual molting probability, and selectivity for trawl survey, winter pot survey, summer commercial fishery, and summer and winter commercial retention. X-axis is carapace length (mm). Line colors black, red, and blue correspond to model 19.0, model 19.4, and model 19.5, respectively.

MMB Feb 01




Figure 4. Estimated MMB during 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2020). Line colors black, red, and blue correspond to model 19.0, model 19.1 respectively.

MMB Feb 01




Figure 5. Estimated MMB during 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2020). Line colors black, red, and blue correspond to model 19.0, model 19.2, and model 19.3, respectively.

MMB Feb 01




Figure 6. Estimated MMB during 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019). Line colors black, red, and blue correspond to model 19.0, model 19.4, and model 19.5, respectively.




Figure 7. Observed (open circle) (White: NMFS, Red ADF&G) and model estimated (dots) trawl survey male abundances with 95% lognormal Confidence Intervals (crab \geq 64 mm CL) Line colors black, red, and blue correspond to model 19.0, model 19.2, and model 19.3, respectively.

Summer commercial standardized cpue




Figure 8. Observed (open circle bar) r and model estimated (lines) standardized CPUE. Line colors black and red correspond to model 19.0 and model 19.1, respectively.




Figure 9. Predicted (line) vs. observed (dots: black New Shell, red Old Shell) length class proportions across years. Line colors black and red correspond to model 19.0 and model 19.1, respectively.




Figure 10. Predicted (line) vs. observed (dots: black New Shell, red Old Shell) length class proportions across years. Line colors black, red, and blue correspond to model 19.0, model 19.4, and 19.5 respectively.





Figure 11. Predicted (line) vs. observed length class proportions tag recovery data. Line colors black and red correspond to model 19.0 and model 19.1, respectively.

Appendix A. Description of the Norton Sound Red King Crab Model

a. Model description.

The model is an extension of the length-based model developed by Zheng et al. (1998) for Norton Sound red king crab. The model has 8 male length classes with model parameters estimated by the maximum likelihood method. The model estimates abundances of crab with CL ≥ 64 mm and with 10-mm length intervals (8 length classes, ≥ 134 mm) because few crab measuring less than 64 mm CL were caught during surveys or fisheries and there were relatively small sample sizes for trawl and winter pot surveys. The model treats newshell and oldshell male crab separately but assumes they have the same molting probability and natural mortality.

Norton Sound Red King Crab Modeling Scheme



Timeline of calendar events and crab modeling events:

- **Model year starts February 1st to January 31st of the following year.**
- **All winter fishery harvest occurs on February 1st**
- **Molting and recruitment occur on July 1st**
- **Initial Population Date: February 1st 1976**

Initial pre-fishery summer crab abundance on February 1st 1976

Abundance of the initial pre-fishery population was assumed to consist of newshell crab to reduce the number of parameters, and estimated as

$$N_{l,1} = p_l e^{\log_{-} N_{76}} \quad (1)$$

where, length proportion of the first year (p_l) was calculated as

$$p_l = \frac{\exp(a_l)}{1 + \sum_{l=1}^{n-1} \exp(a_l)} \text{ for } l = 1, \dots, n-1$$

$$p_n = 1 - \frac{\sum_{l=1}^{n-1} \exp(a_l)}{1 + \sum_{l=1}^{n-1} \exp(a_l)} \quad (2)$$

for model estimated parameters a_l .

Crab abundance on July 1st

Summer (01 July) crab abundance of new and oldshells consists of survivors of winter commercial and subsistence crab fisheries and natural mortality from 01Feb to 01July:

$$N_{s,l,t} = (N_{w,l,t-1} - C_{w,t-1} P_{w,n,l,t-1} - C_{p,t} P_{p,n,l,t-1} - D_{w,n,l,t-1} - D_{p,n,l,t-1}) e^{-0.42M_l}$$

$$O_{s,l,t} = (O_{w,l,t-1} - C_{w,t-1} P_{w,o,l,t-1} - C_{p,t} P_{p,o,l,t-1} - D_{w,o,l,t-1} - D_{p,o,l,t-1}) e^{-0.42M_l} \quad (3)$$

where

$N_{s,l,t}$, $O_{s,l,t}$: summer abundances of newshell and oldshell crab in length class l in year t ,
 $N_{w,l,t-1}$, $O_{w,l,t-1}$: winter abundances of newshell and oldshell crab in length class l in year $t-1$,
 $C_{w,t-1}$, $C_{p,t-1}$: total winter commercial and subsistence catches in year $t-1$,
 $P_{w,n,l,t-1}$, $P_{w,o,l,t-1}$: Proportion of newshell and oldshell length class l crab in year $t-1$, harvested by winter commercial fishery,
 $P_{p,n,l,t-1}$, $P_{p,o,l,t-1}$: Proportion of newshell and oldshell length class l crab in year $t-1$, harvested by winter subsistence fishery,
 $D_{w,n,l,t-1}$, $D_{w,o,l,t-1}$: Discard mortality of newshell and oldshell length class l crab in winter commercial fishery in year $t-1$,

$D_{p,n,l,t-1}, D_{p,o,l,t-1}$: Discard mortality of newshell and oldshell length class l crab in winter subsistence fishery in year $t-1$,

M_l : instantaneous natural mortality in length class l ,

0.42 : proportion of the year from Feb 1 to July 1 is 5 months.

Length proportion compositions of winter commercial catch ($P_{w,n,l,t}, P_{w,o,l,t}$) in year t were estimated as:

$$\begin{aligned} P_{w,n,l,t} &= N_{w,l,t} S_{w,l} P_{lg,l} / \sum_{l=1} [(N_{w,l,t} + O_{w,l,t}) S_{w,l} P_{lg,l}] \\ P_{w,o,l,t} &= O_{w,l,t} S_{w,l} P_{lg,l} / \sum_{l=1} [(N_{w,l,t} + O_{w,l,t}) S_{w,l} P_{lg,l}] \end{aligned} \quad (4)$$

where

$P_{lg,l}$: the proportion of legal males in length class l ,

$S_{w,l}$: Selectivity of winter fishery pot.

Subsistence fishery does not have a size limit; however, crab of size smaller than length class 3 are generally not retained. Hence, we assumed proportion of length composition $l = 1$ and 2 as 0, and estimated length compositions ($l \geq 3$) as follows

$$\begin{aligned} P_{p,n,l,t} &= N_{w,l,t} S_{w,l} / \sum_{l=3} [(N_{w,l,t} + O_{w,l,t}) S_{w,l}] \\ P_{p,o,l,t} &= O_{w,l,t} S_{w,l} / \sum_{l=3} [(N_{w,l,t} + O_{w,l,t}) S_{w,l}] \end{aligned} \quad (5)$$

Crab abundance on Feb 1st

Newshell Crab: Abundance of newshell crab of year t and length-class l ($N_{w,l,t}$) year- t consist of:

(1) new and oldshell crab that survived the summer commercial fishery and molted, and (2) recruitment ($R_{l,t}$).

$$N_{w,l,t} = \sum_{l'=1}^{l'=l} G_{l',l} [(N_{s,l',t-1} + O_{s,l',t-1}) e^{-y_c M_l} - C_{s,t} (P_{s,n,l',t-1} + P_{s,o,l',t-1}) - D_{l',t-1}] m_r e^{-(0.58-y_c) M_l} + R_{l,t} \quad (6)$$

Oldshell Crab: Abundance of oldshell crabs of year t and length-class l ($O_{w,l,t}$) consists of the non-molting portion of survivors from the summer fishery:

$$O_{w,l,t} = [(N_{s,l,t-1} + O_{s,l,t-1}) e^{-y_c M_l} - C_{s,t} (P_{s,n,l,t-1} + P_{s,o,l,t-1}) - D_{l,t-1}] (I - m_l) e^{-(0.58-y_c) M_l} \quad (7)$$

where

$G_{l',l}$: a growth matrix representing the expected proportion of crabs growing from length class l' to length class l

$C_{s,t}$: total summer catch in year t

$P_{s,n,l,t}$, $P_{s,o,l,t}$: proportion of summer catch for newshell and oldshell crabs of length class l in year t ,

$D_{l,t}$: summer discard mortality of length class l in year t ,

m_l : molting probability of length class l ,

y_c : the time in year from July 1 to the mid-point of the summer fishery,

0.58: Proportion of the year from July 1st to Feb 1st is 7 months is 0.58 year,

$R_{l,t}$: recruitment into length class l in year t .

Discards

Discards are crabs that were caught by fisheries but were not retained, which consists of summer commercial, winter commercial and winter subsistence.

Summer and winter commercial discards

In summer ($D_{l,t}$) and winter ($D_{w,n,l,t}$, $D_{w,o,l,t}$) commercial fisheries, sublegal males (<4.75 inch CW and <5.0 inch CW since 2005) are discarded. Those discarded crabs are subject to handling mortality. The number of discards was not directly observed, and thus was estimated from the model as: Observed Catch x (estimated abundance of crab that are not caught by commercial pot)/(estimated abundance of crab that are caught by commercial pot)

Model discard mortality in length-class l in year t from the summer and winter commercial pot fisheries is given by

$$D_{l,t} = C_{s,t} \frac{(N_{s,l,t} + O_{s,l,t}) S_{s,l} (1 - P_{lg,l})}{\sum_l (N_{s,l,t} + O_{s,l,t}) S_{s,l} P_{lg,l}} hm_s \quad (\text{Baseline model}) \quad (8)$$

$$D_{l,t} = C_{s,t} \frac{(N_{s,l,t} + O_{s,l,t}) S_{s,l} (1 - S_{r,l})}{\sum_l (N_{s,l,t} + O_{s,l,t}) S_{s,l} S_{r,l}} hm_s \quad (\text{Alternative model})$$

$$D_{w,n,l,t} = C_{w,t} \frac{N_{w,l,t} S_{w,l} (1 - P_{lg,l})}{\sum_l (N_{w,l,t} + O_{w,l,t}) S_{w,l} P_{lg,l}} hm_w \quad (9)$$

$$D_{w,o,l,t} = C_{w,t} \frac{O_{w,l,t} S_{w,l} (1 - P_{lg,l})}{\sum_l (N_{w,l,t} + O_{w,l,t}) S_{w,l} P_{lg,l}} hm_w \quad (10)$$

where

hm_s : summer commercial handling mortality rate assumed to be 0.2,

hm_w : winter commercial handling mortality rate assumed to be 0.2,

$S_{s,l}$: Selectivity of the summer commercial fishery,

$S_{w,l}$: Selectivity of the winter commercial fishery,
 $S_{r,l}$: Retention selectivity of the summer commercial fishery,

Winter subsistence Discards

Discards (unretained) of winter subsistence fishery is reported in a permit survey ($C_{d,t}$), though its size composition is unknown. We assumed that subsistence fishers discarded all crabs of length classes 1 -2.

$$D_{p,n,l,t} = C_{d,t} \frac{N_{w,l,t} S_{w,l}}{\sum_{l=1}^2 (N_{w,l,t} + O_{w,l,t}) S_{w,l}} hm_w \quad (11)$$

$$D_{p,o,l,t} = C_{d,t} \frac{O_{w,l,t} S_{w,l}}{\sum_{l=1}^2 (N_{w,l,t} + O_{w,l,t}) S_{w,l}} hm_w \quad (12)$$

$C_{d,t}$: Winter subsistence discards catch,

Recruitment

Recruitment of year t , R_t , is a stochastic process around the geometric mean, R_0 :

$$R_t = R_0 e^{\tau_t}, \tau_t \sim N(0, \sigma_R^2) \quad (13)$$

R_t of the last year was assumed to be an average of previous 5 years: $R_t = (R_{t-1} + R_{t-2} + R_{t-3} + R_{t-4} + R_{t-5})/5$.

R_t was assumed to be newshell crab of immature (< 94mm) length classes 1 to r :

$$R_{r,t} = p_r R_t \quad (14)$$

where r takes multinomial distribution, same as the equation (2)

Molting Probability

Molting probability for length class l , m_l , was estimated as an inverse logistic function of length-class mid carapace length (L) and parameters (α, β) where β corresponds to L_{50} .

$$m_l = \frac{I}{1 + e^{\alpha(L-\beta)}} \quad (15)$$

Trawl net, summer commercial pot, retention selectivity

Trawl and summer commercial pot selectivity was assumed to be a logistic function of mid-length-class, constrained to be 0.999 at the largest length-class (L_{max}):

$$S_l = \frac{I}{1 + e^{(\alpha(L_{max}-L) + \ln(1/0.999-1))}} \quad (16)$$

Alternative Summer commercial pot, retention selectivity

Summer pot selectivity was assumed to be a logistic function of length-class mid carapace length (L) and parameters (α, β) where β corresponds to L_{50} .

$$S_{c,l} = \frac{I}{1 + e^{-\alpha(L-\beta)}} \quad (16')$$

Winter pot selectivity

Winter pot selectivity was assumed to be a dome-shaped with inverse logistic function of length-class mid carapace length (L) and parameters (α, β) where β corresponds to L_{50} .

$$S_{w,l} = \frac{I}{1 + e^{\alpha(L-\beta)}} \quad (17)$$

Selectivity of the length classes $S_{w,s}$ ($S = l_1, l_2$) were individually estimated.

Growth transition matrix

The growth matrix $G_{l',l}$ (the expected proportion of crab molting from length class l' to length class l) was assumed to be normally distributed:

$$G_{l',l} = \begin{cases} \frac{\int_{l_{m_l-h}}^{l_{m_l+h}} N(L | \mu_{l'}, \sigma^2) dL}{\sum_{l=1}^n \int_{l_{m_l-h}}^{l_{m_l+h}} N(L | \mu_{l'}, \sigma^2) dL} & \text{when } l \geq l' \\ 0 & \text{when } l < l' \end{cases} \quad (18)$$

Where

$$N(x | \mu_l, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(L - \mu_l)^2}{\sigma^2}\right)$$

$$lm_l = L_1 + st \cdot l$$

$$\mu_l = L_1 + \beta_0 + \beta_1 \cdot l$$

Observation model

Summer trawl survey abundance

Modeled trawl survey abundance of year t ($B_{st,t}$) is July 1st abundance subtracted by summer commercial fishery harvest occurring from July 1st to the mid-point of summer trawl survey, multiplied by natural mortality occurring between the mid-point of commercial fishery date and trawl survey date, and multiplied by trawl survey selectivity. For the first year (1976) trawl survey, the commercial fishery did not occur.

$$\hat{B}_{st,t} = \sum_l [(N_{s,l,t} + O_{s,l,t})e^{-y_c M_l} - C_{s,t} P_{c,t} (P_{s,n,l,t} + P_{s,o,l,t})] e^{-(y_{st} - y_c) M_l} S_{st,l} \quad (19)$$

where

y_{st} : the time in year from July 1 to the mid-point of the summer trawl survey,

y_c : the time in year from July 1 to the mid-point for the catch before the survey, ($y_{st} > y_c$: Trawl survey starts after opening of commercial fisheries),

$P_{c,t}$: the proportion of summer commercial crab harvested before the mid-point of trawl survey date.

$S_{st,l}$: Selectivity of the trawl survey.

Winter pot survey CPUE

Winter pot survey cpue (f_{wt}) was calculated with catchability coefficient q and exploitable abundance:

$$\hat{f}_{wt} = q_w \sum_l [(N_{w,l,t} + O_{w,l,t})S_{w,l}] \quad (20)$$

Summer commercial CPUE

Summer commercial fishing CPUE (f_t) was calculated as a product of catchability coefficient q and mean exploitable abundance minus one half of summer catch, A_t :

$$\hat{f}_t = q_i (A_t - 0.5C_t) \quad (21)$$

Because the fishing fleet and pot limit configuration changed in 1993, q_l is for fishing efforts before

1993, q_2 is from 1994 to present.

Baseline model

Where A_t is exploitable legal abundance in year t , estimated as

$$\begin{aligned} A_t &= \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{s,l} P_{lg,l}] \quad (\text{Baseline model}) \\ A_t &= \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{s,l} S_{r,l}] \quad (\text{Alternative model}) \end{aligned} \quad (22)$$

Summer pot survey abundance (Removed from likelihood components)
Abundance of t -th year pot survey was estimated as

$$\hat{B}_{p,t} = \sum_l [(N_{s,l,t} + O_{s,l,t}) e^{-y_p M_t}] S_{p,l} \quad (23)$$

Where

y_p : the time in year from July 1 to the mid-point of the summer pot survey.

Length composition

Summer commercial catch

Length compositions of the summer commercial catch for new and old shell crabs $P_{s,n,l,t}$ and $P_{s,o,l,t}$, were modeled based on the summer population, selectivity, and legal abundance:

$$\begin{aligned} \hat{P}_{s,n,l,t} &= N_{s,l,t} S_{s,l} P_{lg,l} / A_t \\ \hat{P}_{s,o,l,t} &= O_{s,l,t} S_{s,l} P_{lg,l} / A_t \\ \hat{P}_{s,n,l,t} &= N_{s,l,t} S_{s,l} S_{r,l} / A_t \\ \hat{P}_{s,o,l,t} &= O_{s,l,t} S_{s,l} S_{r,l} / A_t \end{aligned} \quad \begin{array}{l} (\text{Baseline model}) \\ (\text{Alternative model}) \end{array} \quad (24)$$

Summer commercial fishery discards (Base model)

Length/shell compositions of observer discards were modeled as

$$\begin{aligned} \hat{P}_{b,n,l,t} &= N_{s,l,t} S_{s,l} (I - P_{lg,l}) / \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{s,l} (I - P_{lg,l})] \\ \hat{P}_{b,o,l,t} &= O_{s,l,t} S_{s,l} (I - P_{lg,l}) / \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{s,l} (I - P_{lg,l})] \end{aligned} \quad (25)$$

Summer commercial fishery total catch (Alternative model)

Length/shell compositions of observer discards were modeled as

$$\begin{aligned}\hat{P}_{t,n,l,t} &= N_{s,l,t} S_{s,l} / \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{s,l}] \\ \hat{P}_{t,o,l,t} &= O_{s,l,t} S_{s,l} / \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{s,l}]\end{aligned}\quad (25')$$

Summer trawl survey

Proportions of newshell and oldshell crab, $P_{st,n,l,t}$ and $P_{st,o,l,t}$ were given by

$$\begin{aligned}\hat{P}_{st,n,l,t} &= \frac{[N_{s,l,t} e^{-y_c M_l} - C_{s,t} P_{c,t} \hat{P}_{s,n,l',t}] e^{-(y_{st} - y_c) M_l} S_{st,l}}{\sum_l [(N_{s,l,t} + O_{s,l,t}) e^{-y_c M_l} - C_{s,t} P_{c,t} (\hat{P}_{s,n,l',t} + \hat{P}_{s,o,l',t})] e^{-(y_{st} - y_c) M_l} S_{st,l}} \\ \hat{P}_{st,o,l,t} &= \frac{[O_{s,l,t} e^{-y_c M_l} - C_{s,t} \hat{P}_{s,o,l',t} P_{c,t}] e^{-(y_{st} - y_c) M_l} S_{st,l}}{\sum_l [(N_{s,l,t} + O_{s,l,t}) e^{-y_c M_l} - C_{s,t} P_{c,t} (\hat{P}_{s,n,l,t} + \hat{P}_{s,o,l,t})] e^{-(y_{st} - y_c) M_l} S_{st,l}}\end{aligned}\quad (26)$$

Winter pot survey

Winter pot survey length compositions for newshell and oldshell crab, $P_{sw,n,l,t}$ and $P_{sw,o,l,t}$ ($l \geq 1$) were calculated as

$$\begin{aligned}\hat{P}_{sw,n,l,t} &= N_{w,l,t} S_{w,l} / \sum_l [(N_{w,l,t} + O_{w,l,t}) S_{w,l}] \\ \hat{P}_{sw,o,l,t} &= O_{w,l,t} S_{w,l} / \sum_l [(N_{w,l,t} + O_{w,l,t}) S_{w,l}]\end{aligned}\quad (27)$$

Spring Pot survey 2012-2015

Winter pot survey length compositions for newshell and oldshell crab, $P_{sw,n,l,t}$ and $P_{sw,o,l,t}$ ($l \geq 1$) were assumed to be supper crab population caught by winter pot survey gears

$$\begin{aligned}\hat{P}_{sp,n,l,t} &= N_{s,l,t} S_{w,l} / \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{w,l}] \\ \hat{P}_{sp,o,l,t} &= O_{s,l,t} S_{w,l} / \sum_l [(N_{s,l,t} + O_{s,l,t}) S_{w,l}]\end{aligned}\quad (28)$$

Estimates of tag recovery

The proportion of released tagged length class l' crab recovered after t -th year with length class of l

by a fishery of s -th selectivity (S_l) was assumed to be proportional to the growth matrix, catch selectivity, and molting probability (m_l) as

$$\hat{P}_{l',l,s} = \frac{S_l \cdot [X']_{l',l}}{\sum_{l=1}^n S_l \cdot [X']_{l',l}} \quad (29)$$

where X is a molting probability adjusted growth matrix with each component consisting of

$$X_{l',l} = \begin{cases} m_{l'} \cdot G_{l',l} & \text{when } l' \neq l \\ m_l \cdot G_{l',l} + (1-m_l) & \text{when } l' = l \end{cases} \quad (30)$$

b. Software used: AD Model Builder (Fournier et al. 2012).

c. Likelihood components.

Under assumptions that measurement errors of annual total survey abundances and summer commercial fishing efforts follow lognormal distributions and each type of length composition has a multinomial error structure (Fournier and Archibald 1982; Methot 1989), the log-likelihood function is

$$\begin{aligned}
& \sum_{i=1}^{i=4} \sum_{t=1}^{t=n_i} K_{i,t} \left[\sum_{l=1}^{l=n} P_{i,l,t} \ln(\hat{P}_{i,l,t} + \kappa) - \sum_{l=1}^{l=n} P_{i,l,t} \ln(P_{i,l,t} + \kappa) \right] \\
& - \sum_{t=1}^{t=n_i} \frac{[\ln(q \cdot \hat{B}_{i,t} + \kappa) - \ln(B_{i,t} + \kappa)]^2}{2 \cdot \ln(CV_{i,t}^2 + 1)} \\
& - \sum_{t=1}^{t=n_i} \left[\frac{\ln[\ln(CV_t^2 + 1) + w_t]}{2} + \frac{[\ln(\hat{f}_t + \kappa) - \ln(f_t + \kappa)]^2}{2 \cdot [\ln(CV_t^2 + 1) + w_t]} \right] \\
& - \sum_{t=1}^{t=n_i} \frac{\tau_t^2}{2 \cdot SDR^2} \\
& + W \sum_{s=1}^{s=2} \sum_{t=1}^{t=3} \sum_{l'=1}^{l'=n} K_{l',t,s} \left[\sum_{l=1}^{l=n} P_{l',l,t} \ln(\hat{P}_{l',l,t,s} + \kappa) - \sum_{l=1}^{l=n} P_{l',l,t} \ln(P_{l',l,t,s} + \kappa) \right]
\end{aligned} \tag{32}$$

where

i : length/shell compositions of :

- 1 triennial summer trawl survey,
- 2 annual winter pot survey,
- 3 summer commercial fishery retained catch,
- 4 observer discards or total catch during the summer fishery
- 5 spring pot survey.

$K_{i,t}$: the effective sample size of length/shell compositions for data set i in year t ,

$P_{i,l,t}$: observed and estimated length compositions for data set i , length class l , and year t .

κ : a constant equal to 0.0001,

CV : coefficient of variation for the survey abundance,

$B_{i,k,t}$: observed and estimated annual total abundances for data set i and year t ,

f_t : observed and estimated summer fishing CPUE,

w_t^2 : extra variance factor,

SDR : Standard deviation of recruitment = 0.5,

$K_{l',t}$: sample size of length class l' released and recovered after t -th in year,

$P_{l',l,t,s}$: observed and estimated proportion of tagged crab released at length l' and recaptured at length l , after t -th year by commercial fishy pot selectivity s ,

W : weighting for the tagging survey likelihood

It is generally believed that total annual commercial crab catches in Alaska are fairly accurately reported. Thus, total annual catch was assumed known.

d. Parameter estimation framework:

i. Parameters Estimated Independently

The following parameters were estimated independently: natural mortality ($M = 0.18$), proportions of legal males by length group.

Natural mortality was based on an assumed maximum age, t_{max} , and the 1% rule (Zheng 2005):

$$M = -\ln(p)/t_{max},$$

where p is the proportion of animals that reach the maximum age and is assumed to be 0.01 for the 1% rule (Shepherd and Breen 1992, Clarke et al. 2003). The maximum age of 25, which was used to estimate M for U.S. federal overfishing limits for red king crab stocks results in an estimated M of 0.18. Among the 199 recovered crabs from the tagging returns during 1991-2007 in Norton Sound, the longest time at liberty was 6 years and 4 months from a crab tagged at 85 mm CL. The crab was below the mature size and was likely less than 6 years old when tagged. Therefore, the maximum age from tagging data is about 12, which does not support the maximum age of 25 chosen by the CPT.

Proportions of legal males ($CW > 4.75$ inches) by length group were estimated from the ADF&G trawl data 1996-2011 (Table 11).

ii. Parameters Estimated Conditionally

Estimated parameters are listed in Table 10. Selectivity and molting probabilities based on these estimated parameters are summarized in Tables 11.

A likelihood approach was used to estimate parameters

e. Definition of model outputs.

- i. Estimate of mature male biomass (MMB) is on **February 1st** and is consisting of the biomass of male crab in length classes 4 to 8

$$MMB = \sum_{l=4} (N_{w,l} + O_{w,l})wm_l$$

wm_l : mean weight of each length class (Table 11).

- ii. Projected legal male biomass for winter and summer fishery OFL was calculated as

$$Legal_B = \sum_l (N_{w,l} + O_{w,l})S_{s,l}P_{lg,l}wm_l \text{ Baseline model}$$

$$Legal_B = \sum_l (N_{w,l} + O_{w,l}) S_{s,l} S_{r,l} w m_l \text{ Alternative model}$$

iii. Recruitment: the number of males in length classes 1, 2, and 3.

iv.

f. OFL

The Norton Sound red king crab fishery consists of two distinct fisheries: winter and summer. The two fisheries are discontinuous with 5 months between the two fisheries during which natural mortalities occur. To incorporate this fishery, the CPT in 2016 recommended the following formula:

$$OFL_r = \text{Winter harvest (Hw)} + \text{Summer harvest (Hs)} \quad (1)$$

And

$$p = \frac{Hw}{OFL_r} \quad (2)$$

Where p is a specific proportion of winter crab harvest to total (winter + summer) harvest

At given fishery mortality (F_{OFL}), Winter harvest is a fishing mortality

$$Hw = (1 - e^{-x \cdot F}) B_w \quad (3)$$

$$Hs = (1 - e^{-(1-x) \cdot F}) B_s \quad (4)$$

where B_s is a summer crab biomass after winter fishery and x ($0 \leq x \leq 1$) is a fraction that satisfies equation (2)

Since B_s is a summer crab biomass after winter fishery and 5 months of natural mortality ($e^{-0.42M}$)

$$\begin{aligned} B_s &= (B_w - Hw) e^{-0.42M} \quad (5) \\ &= (B_w - (1 - e^{-x \cdot F}) B_w) e^{-0.42M} \\ &= B_w e^{-x \cdot F - 0.42M} \end{aligned}$$

Substituting $0.42M$ to m , summer harvest is

$$\begin{aligned} Hs &= (1 - e^{-(1-x) \cdot F}) B_s \quad (6) \\ &= (1 - e^{-(1-x) \cdot F}) B_w e^{-x \cdot F - m} = (e^{-(x \cdot F + m)} - e^{-(F + m)}) B_w \end{aligned}$$

Thus, OFL is

$$\begin{aligned}
OFL &= Hw + Hs = (1 - e^{-xF})B_w + (e^{-(x \cdot F + m)} - e^{-(F+m)})B_w \\
&= (1 - e^{-xF} + e^{-(x \cdot F + m)} - e^{-(F+m)})B_w \\
&= [1 - e^{-(F+m)} - (1 - e^{-m})e^{-xF}]B_w
\end{aligned} \tag{7}$$

Combining (2) and (7),

$$p = \frac{Hw}{OFL_r} = \frac{(1 - e^{-xF})B_w}{[1 - e^{-(F+m)} - (1 - e^{-m})e^{-xF}]B_w} \tag{8}$$

Solving (8) for x

$$\begin{aligned}
(1 - e^{-xF}) &= p[1 - e^{-(F+m)} - (1 - e^{-m})e^{-xF}] \\
e^{-xF} - p(1 - e^{-m})e^{-xF} &= 1 - p[1 - e^{-(F+m)}] \\
[1 - p(1 - e^{-m})]e^{-xF} &= 1 - p[1 - e^{-(F+m)}] \\
e^{-xF} &= \frac{1 - p[1 - e^{-(F+m)}]}{1 - p(1 - e^{-m})}
\end{aligned} \tag{9}$$

Combining (7) and (9), and substituting back,

revised retained OFL is

$$OFL = Legal - B_w \left(1 - e^{-(F_{OFL} + 0.42M)} - (1 - e^{-0.42M}) \left(\frac{1 - p(1 - e^{-(F_{OFL} + 0.42M)})}{1 - p(1 - e^{-0.42M})} \right) \right)$$

Further combining (3) and (9), Winter fishery harvest rate (Fw) i

$$\begin{aligned}
Fw &= (1 - e^{-xF}) = 1 - \frac{1 - p[1 - e^{-(F+m)}]}{1 - p(1 - e^{-m})} = \frac{1 - p(1 - e^{-m}) - 1 + p[1 - e^{-(F+m)}]}{1 - p(1 - e^{-m})} \\
&= \frac{p(e^{-m} - e^{-(F+m)})}{1 - p(1 - e^{-m})} = \frac{p(1 - e^{-F})e^{-0.42M}}{1 - p(1 - e^{-0.42M})}
\end{aligned} \tag{10}$$

Summer fishery harvest rate (Fs) is

$$\begin{aligned}
Fs &= (e^{-(x \cdot F+m)} - e^{-(F+m)}) = (e^{-x \cdot F} - e^{-F})e^{-m} & (11) \\
&= \left(\frac{1 - p[1 - e^{-(F+m)}]}{1 - p(1 - e^{-m})} - e^{-F} \right) e^{-m} \\
&= \left(\frac{1 - p[1 - e^{-(F+m)}] - e^{-F} + p(e^{-F} - e^{-(F+m)})}{1 - p(1 - e^{-m})} \right) e^{-m} \\
&= \left(\frac{1 - p + pe^{-(F+m)} - e^{-F} + pe^{-F} - pe^{-(F+m)}}{1 - p(1 - e^{-m})} \right) e^{-m} \\
&= \frac{(1 - p)(1 - e^{-F})e^{-m}}{1 - p(1 - e^{-m})} = \frac{(1 - p)(1 - e^{-F})e^{-0.24M}}{1 - p(1 - e^{-0.24M})}
\end{aligned}$$

Appendix B

Norton Sound Red King Crab CPUE Standardization

Note: This is an update of model by G. Bishop (SAFE 2013).

Methods

Data Source & Cleaning

Commercial fishery harvest data were obtained from a fish ticket database, which included: Landing Date, Fish Ticket Number, Vessel Number, Permit Fishery ID, Statistical Area(s) fished, Effort, and Number and Pounds of Crab harvested (Table A2-1,2,3, Figure A2-1). Fish ticket database may have multiple entries of identical Fish Ticket Number, Vessel Number, Permit Fishery ID, and Statistical Area. In those cases, at least one Effort data are missing or zero with the Number and Pounds of Crab harvested. These entries indicate that crabs were either retained from commercial fishery (i.e., not sold), or dead loss.

Following data cleaning and combining methods were conducted.

1. Sum crab number and efforts by Fish Ticket Number, Vessel Number, Permit Fishery ID, Statistical Area
2. Remove data of missing or zero Efforts, Number of Crab, Pounds of Crab (Those are considered as true missing data)
3. Calculate CPUE as Number of Crab/Effort

Data Censoring

During 1977-92 period, vessels of 1 year of operation and/or 1 delivery per year harvested 20-90% of crabs (Table A2-5, Figure A2-2). For instance, all vessels did only 1 delivery in 1989, and in 1988 64% of crabs were harvested by 1 vessel that did only 1 delivery. On the other hand, during the 1993-2017

Norton Sound red king crab CPUE standardization

period of post super-exclusive fishery status, the majority of commercial crab fishery and harvest was done by vessels with more than 5 years of operations and more than 5 deliveries per year. For 1977 – 1992, censoring was made for vessels of more than 2 years of operations. Increasing deliveries to more than one would result in no estimates for some years. For 1993 – 2018, censoring was made for vessels of more than 5 years of operations and 5 deliveries per year.

Analyses

A GLM was constructed as

$$\ln(CPUE) = YR + PD + VSL + MSA + WOY + PF + PD$$

Where YR: Year, PD: Fishery periods (1977-1992, 1993-2004,2005-2018), VSL: Vessel, MSA: Statistical Area, WOY: Week of Year, PF: Permit vs open fishery (Table 1). All variables were treated as categorical. Inclusion of interaction terms were not considered because they were absent (SAFE 2013).

For selection of the best model, forward and backward stepwise selection was conducted. (R step function)

```
fit <- glm(L.CPUE.NO ~ factor(YR) + factor(VSL) + factor(WOY) +  
factor(MSA) + factor(PF) + factor(PD),,data=NSdata.C)  
step <- step(fit, direction='both', trace = 10)  
best.glm<-glm(formula(step), data=NSdata.C)
```

Norton Sound red king crab CPUE standardization

Table B-1. List of variables in the fish ticket database. Variables in bold face were used for generalized linear modeling.

Variable	Description
YR	Year of commercial fishery
VSL	Unique vessel identification number
Fish Ticket Number	Unique delivery to a processor by a vessel.
PF	Unique Permit Fishery categories
PD	Fishery period: 1977-1992, 1993-2004,2005-2018
Statistical Area	Unique fishery area.
MOA	Modified statistical area, combining each statistical area into 4 larger areas: Inner, Mid, Outer, Outer North
Fishing beginning date	Date of pots set
Landing date	Date of crab landed to processor
WOY	Week of Landing Date (calculated)
Effort	The number of pot lift
Crab Numbers	Total number of crabs harvested from pots
Crab Pounds	Total pounds of crab harvested from pots
ln(CPUE)	ln(Crab Numbers/Effort) (calculated)

Table B-2. Permit fisheries, descriptions, and years with deliveries for Norton Sound summer commercial red king crab harvest data.

Permit fishery	Type	Description	Years
K09Q	Open access	KING CRAB , POT GEAR VESSEL UNDER 60', BERING SEA	1994–2002
K09Z	Open access	KING CRAB , POT GEAR VESSEL UNDER 60', NORTON SOUND	1992–2017
K09ZE	CDQ	KING CRAB , POT GEAR VESSEL UNDER 60', NORTON SOUND CDQ, NSEDC	2000–2017
K09ZF	CDQ	KING CRAB , POT GEAR VESSEL UNDER 60', NORTON SOUND CDQ, YDFDA	2002–2004
K91Q	Open access	KING CRAB , POT GEAR VESSEL 60' OR OVER, BERING SEA	1978–1989
K91Z	Open access	KING CRAB , POT GEAR VESSEL 60' OR OVER, NORTON SOUND	1982–1994

Table B-3. Modified statistical area definitions used for analysis of Norton Sound summer commercial red king crab harvest data.

Modified statistical area	Statistical areas included
Inner	616331, 616401, 626331, 626401, 626402
Mid	636330, 636401, 636402, 646301, 646330, 646401, 646402
Outer	656300, 656330, 656401, 656402, 666230, 666300, 666330, 666401
Outer North	666402, 666431, 676300, 676330 ,676400, 676430, 676501, 686330

Norton Sound red king crab CPUE standardization

Table B-4. Final generalized linear model formulae and AIC selected for Norton Sound summer commercial red king crab fishery. The dependent variable is ln(CPUE) in numbers.

Var	Df	Deviance	Resid DF	Resid Dev	AIC
YR	41	1312.43	6274	5082.7	
VSL	90	574.57	6143	3770.3	
WOY	15	82.89	6129	3195.7	
MSA	3	65.83	6125	3047.0	
PF	6	20.14	6119	3026.9	13547
+PD+MOY	3				13547.67

Norton Sound red king crab CPUE standardization

Table B-5. Standardized (Censored/full data), and scaled arithmetic observed CPUE indices

Year	Censored	
	CPUE	SE
1977	3.29	0.68
1978	4.68	0.65
1979	2.87	0.64
1980	3.07	0.65
1981	0.86	0.64
1982	0.20	0.62
1983	0.90	0.65
1984	1.59	0.65
1985	0.50	0.66
1986	1.74	0.70
1987	0.61	0.64
1988	2.36	0.86
1989	1.21	0.61
1990	1.08	0.68
1991		
1992	0.17	0.60
1993	0.90	0.35
1994	0.81	0.34
1995	0.42	0.34
1996	0.51	0.34
1997	0.84	0.35
1998	0.79	0.36
1999	0.92	0.36
2000	1.24	0.34
2001	0.64	0.34
2002	1.23	0.34
2003	0.85	0.34
2004	1.27	0.34
2005	1.19	0.34
2006	1.31	0.34
2007	1.02	0.34
2008	1.32	0.34
2009	0.84	0.34
2010	1.22	0.34
2011	1.58	0.34
2012	1.29	0.34
2013	0.67	0.33
2014	1.12	0.34
2015	1.45	0.34
2016	1.27	0.34
2017	1.10	0.34
2018	0.64	0.34

Norton Sound red king crab CPUE standardization




Figure A2-1. Closed area and statistical area boundaries used for reporting commercial harvest information for red king crab in Registration Area Q, Northern District, Norton Sound Section and boundaries of the new *Modified Statistical Areas* used in this analysis.

Appendix C

Norton Sound Red King Crab Summer Commercial fishery Discards Estimation

Formal methodologies have not been established for estimating Red King Crab discards by Norton Sounds Summer commercial fishery from observer data. Here, I describe a few methods and discuss pros and cons of each method.

Data source and description of survey protocols

Norton Sound Summer Commercial fishery observer survey started in 2009 as a potential feasibility project, and formal data collection started since 2012. The observer survey in Norton Sound is voluntary. Due to small boat size, the boat that can take an fishery observer is limited. Fishery observer often work as a crew member helping fishing.

Methods

Data Source & Cleaning

From 2012 to 2018, crab catches of 3-4 volunteer crab fishing vessels were observed. Annual observed pots ranged 69 to 199 and total observed crabs ranging from 2200 to 5300 (Table 1). All observed data were combined.

Estimation Methods

Two methods were considered: CPUE and Proportion method. CPUE method expands observed CPUE (Observed number of crab)/(observed pots) to all fisheries pot lifts, whereas proportional method

Norton Sound red king crab CPUE standardization

expands observed proportion of discards to retained: (observed number of discards)/(observed number of retained) to all fisheries retained catch.

CPUE has two methods: LNR and Subtraction. LNR simply expands CPUE of discards, whereas Subtraction expands CPUE of total catch and subtract total retained catch.

LNR method

LNR method simply expands CPUE of discards to total pot lifts

$$D_{LNR} = \frac{(N_{obs,sub} + N_{obs,ld})}{P_{obs}} P_{total}$$

Subtraction method

Subtraction method expands total catch CPUE and subtract total retained catch

$$D_{Sub} = \frac{(N_{obs,sub} + N_{obs,ld} + N_{obs,lr})}{P_{obs}} P_{total} - N_{total}$$

Proportion method that expands the proportion of discards to retained.

$$D_{prop} = \frac{(N_{obs,sub} + N_{obs,ld})}{N_{obs,lr}} N_{total,lr}$$

Results

While general annual discards trends were similar among the 3 methods, the number of discards differed (Table 2). Overall, the Subtraction method estimated the highest and the Proportional method estimated the lowest.

Discussion

The CPUE method assumes that observed CPUE would represent total CPUE or that **every fisherman catches crab in the same catch efficiency**, whereas the proportional method assumes that observed discards proportions would represent total proportion or that **every fisherman has similar crab composition**. Given than observed fishermen and vessels are limited to those who has larger boat and high catchers, one could assume that they would be more efficient in catching legal crabs with fewer discards than those with small boats. In fact, retained legal crab CPUE by observed fishermen was on average 25% (-11% - 46%) higher than those from the average retained CPUE from fish all fishermen in fish ticket database (Table 3). Furthermore, observed fishermen caught more crabs in stat areas where discards proportion was the lowest (646401) than those of all fishermen (Table 4). Those would explain higher discards estimate for Subtraction method and lower estimates by LNR and Proportion estimate. Higher legal retained crab CPUE inflated Total catch and thus discards by Subtraction method, while lower sublegal and discards proportion may have underestimated discards.

Table 1. Observed pot lifts, catch, and total pot lifts and catch from 2012 to 2018

Year	Observer Survey			Fish Tickets			
	Observed pots	Sublegal	Legal Retained	Legal Discarded	Female	Total pot lifts	Total Leal Retained
2012	78	898	1055	177	152	10041	161113
2013	199	2775	2166	258	123	15058	130603
2014	147	1504	1838	341	104	10127	129656
2015	69	969	1676	577	224	8356	144224
2016	67	264	1700	169	878	8,009	138997
2017	110	432	2174	122	373	9440	135322
2018	78	547	1096	10	574	8797	89613

Norton Sound red king crab CPUE standardization

Table 2. The number of discarded crab estimated by 3 methods.

Year	LNR	Sub	Prop
2012	138386	113084	164167
2013	229502	262797	182880
2014	127104	124070	130150
2015	187223	245965	133037
2016	51760	115976	35403
2017	47543	98790	34484
2018	62820	96816	45542

Table 3. Retained Crab CPUE between observed (CPUE.ob) and Fish ticket (CPUE.ft)

Year	CPUE.ob	CPUE.ft	Ratio
2012	13.53	16.05	0.84
2013	10.88	8.67	1.25
2014	12.50	12.80	0.98
2015	24.29	17.26	1.41
2016	25.37	17.36	1.46
2017	19.76	14.33	1.38
2018	14.05	10.19	1.38

Table 4. Average legal crab proportion caught by 2012-2018 trawl survey and Summer commercial harvest proportion in major fishing stat area

STAT Area	Trawl % Legal	Trawl Catch Prop	Total Legal Catch Prop	Ob. Catch %Legal	Ob. Catch Prop	CPUE
666401	62%	11%	15%	72%	7%	12.79
656401	21%	19%	21%	62%	18%	12.03
646401	12%	24%	19%	58%	46%	14.70
636401	31%	18%	33%	68%	19%	15.78
626401	53%	28%	15%	80%	2%	11.78

Norton Sound red king crab CPUE standardization




Figure 1. The number of discarded crab estimated by 3 methods.

Model 19.0




Figure D1-1. QQ Plot of Trawl survey and Commercial CPUE.




Figure D1-2: Implied effective samples. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the mean implied effective sample size. The second column show input sample size (x-axis) vs. implied effective sample size (y-axis). Dashed line indicates linear regression slope, and solid line is 1:1 line. The third column show year (x-axis) vs. implied effective sample size (y-axis).




Figure D1-3. Molting probability and trawl/pot selectivity. X-axis is carapace length.

Trawl survey crab abundance




Figure D1-4. Estimated trawl survey male abundance (crab \geq 64 mm CL). Observed: White: NOAA Trawl Survey, Red: ADG&G Trawl Survey

Modeled crab abundance Feb 01




Figure D1-5. Estimated abundance of legal males from 1976-2015.

MMB Feb 01




Figure D1-6. Estimated abundance of Mature Male Biomass from 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019).

Summer commercial standardized cpue




Figure D1-7. Summer commercial standardized cpue 1977-2018.




Figure D1-8. Total catch and estimated harvest rate 1976-2018.




Figure D1-9. Predicted (dashed line) vs. observed (dots) length class proportions for commercial catch. Black: New Shell, Red: Old Shell




Figure D1-10. Predicted (dashed line) vs. observed (black dots) length class proportions for the winter and spring pot survey.




Figure D1-11. Predicted (dashed) vs. observed (dots) length class proportions for Trawl survey.




Figure D1-13. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D1-12. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D1-13. Predicted vs. observed length class proportions for tag recovery data.




Figure D1-13. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).




Figure D1-14. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).

Table D1. Summary of parameter estimates for a length-based stock synthesis population model of Norton Sound red king crab.

name	Estimate	std.dev
log_q1	-6.783	0.111
log_q2		
log_N76	9.122	0.109
R0	6.478	0.083
a1	1.752	4.587
a2	2.769	4.260
a3	3.934	4.107
a4	4.072	4.094
a5	4.300	4.085
a6	3.537	4.114
a7	2.101	4.383
r1	10.000	0.283
r2	9.655	0.332
log_a	-2.682	0.090
log_b	4.835	0.015
log_φ _{st1}	-5.000	0.051
log_φ _{wa}	-2.206	0.301
log_φ _{wb}	4.796	0.032
Sw1	0.072	0.035
Sw2	0.499	0.126
log_φ _l	-2.086	0.057
log_φ _{ra}	-0.787	0.129
log_φ _{rb}	4.646	0.008
log_φ _{wra}	-0.965	0.553
log_φ _{wrb}	4.654	0.038
w ² _t	0.000	0.000
q	0.700	0.113
σ	3.886	0.208
β ₁	12.393	0.700
β ₂	7.661	0.171
ms78	3.248	0.255

Model 19.1




Figure D2-1. QQ Plot of Trawl survey and Commercial CPUE.




Figure D2-2: Implied effective samples. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the mean implied effective sample size. The second column show input sample size (x-axis) vs. implied effective sample size (y-axis). Dashed line indicates linear regression slope, and solid line is 1:1 line. The third column show year (x-axis) vs. implied effective sample size (y-axis).




Figure D2-3. Molting probability and trawl/pot selectivity. X-axis is carapace length.

Trawl survey crab abundance




Figure D2-4. Estimated trawl survey male abundance (crab \geq 64 mm CL). Observed: White: NOAA Trawl Survey, Red: ADG&G Trawl Survey

Modeled crab abundance Feb 01




Figure D2-5. Estimated abundance of legal males from 1976-2015.

MMB Feb 01




Figure D2-6. Estimated abundance of Mature Male Biomass from 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019).

Summer commercial standardized cpue




Figure D2-7. Summer commercial standardized cpue 1977-2018.




Figure D2-8. Total catch and estimated harvest rate 1976-2018.




Figure D2-9. Predicted (dashed line) vs. observed (dots) length class proportions for commercial catch. Black: New Shell, Red: Old Shell




Figure D2-10. Predicted (dashed line) vs. observed (black dots) length class proportions for the winter and spring pot survey.




Figure D2-11. Predicted (dashed) vs. observed (dots) length class proportions for Trawl survey.




Figure D2-13. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D2-12. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D2-13. Predicted vs. observed length class proportions for tag recovery data.




Figure D2-13. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).




Figure D2-14. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).

Table D2. Summary of parameter estimates for a length-based stock synthesis population model of Norton Sound red king crab.

name	Estimate	std.dev
log_q1	-6.775	0.112
log_q2		
log_N76	9.171	0.112
R0	6.526	0.084
a1	2.214	5.073
a2	3.308	4.774
a3	4.334	4.654
a4	4.373	4.646
a5	4.566	4.637
a6	3.777	4.663
a7	2.265	4.871
r1	10.000	0.312
r2	9.616	0.362
log_a	-2.733	0.099
log_b	4.837	0.016
log_φ _{st1}	-5.000	0.080
log_φ _{wa}	-2.130	0.297
log_φ _{wb}	4.808	0.030
Sw1	0.071	0.034
Sw2	0.490	0.120
log_φ _l	-2.093	0.055
log_φ _{ra}	-0.798	0.128
log_φ _{rb}	4.648	0.008
log_φ _{wra}	-0.953	0.561
log_φ _{wrb}	4.653	0.038
w ² _t	0.000	0.000
q	0.677	0.109
σ	4.232	0.255
β ₁	11.829	0.926
β ₂	7.919	0.221
ms78	3.554	0.280

Model 19.2




Figure D3-1. QQ Plot of Trawl survey and Commercial CPUE.




Figure D3-2: Implied effective samples. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the mean implied effective sample size. The second column show input sample size (x-axis) vs. implied effective sample size (y-axis). Dashed line indicates linear regression slope, and solid line is 1:1 line. The third column show year (x-axis) vs. implied effective sample size (y-axis).




Figure D3-3. Molting probability and trawl/pot selectivity. X-axis is carapace length.

Trawl survey crab abundance




Figure D3-4. Estimated trawl survey male abundance (crab \geq 64 mm CL). Observed: White: NOAA Trawl Survey, Red: ADG&G Trawl Survey

Modeled crab abundance Feb 01




Figure D3-5. Estimated abundance of legal males from 1976-2015.

MMB Feb 01




Figure D3-6. Estimated abundance of Mature Male Biomass from 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019).

Summer commercial standardized cpue




Figure D3-7. Summer commercial standardized cpue 1977-2018.




Figure D3-8. Total catch and estimated harvest rate 1976-2018.




Figure D3-9. Predicted (dashed line) vs. observed (dots) length class proportions for commercial catch. Black: New Shell, Red: Old Shell




Figure D3-10. Predicted (dashed line) vs. observed (black dots) length class proportions for the winter and spring pot survey.




Figure D3-11. Predicted (dashed) vs. observed (dots) length class proportions for Trawl survey.





Figure D3-13. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.



Proportion

CL mm

Figure D3-12. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D3-13. Predicted vs. observed length class proportions for tag recovery data.




Figure D3-13. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).




Figure D3-14. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).

Table D3. Summary of parameter estimates for a length-based stock synthesis population model of Norton Sound red king crab.

name	Estimate	std.dev
log_q1	-6.471	0.123
log_q2		
log_N76	8.895	0.091
R0	6.206	0.095
a1	2.091	4.628
a2	3.055	4.325
a3	4.093	4.166
a4	4.189	4.152
a5	4.400	4.142
a6	3.609	4.172
a7	2.110	4.440
r1	10.000	0.335
r2	9.671	0.376
log_a	-2.665	0.089
log_b	4.829	0.015
log_φ _{st1}	-5.000	0.113
log_φ _{wa}	-2.198	0.316
log_φ _{wb}	4.805	0.032
Sw1	0.072	0.035
Sw2	0.497	0.124
log_φ _l	-2.082	0.056
log_φ _{ra}	-0.796	0.128
log_φ _{rb}	4.647	0.008
log_φ _{wra}	-0.988	0.536
log_φ _{wrb}	4.656	0.037
w ² _t	0.004	0.019
q ADFG	1.400	0.217
σ	3.870	0.209
β ₁	12.524	0.705
β ₂	7.636	0.173
ms78	2.883	0.259

Model 19.3




Figure D4-1. QQ Plot of Trawl survey and Commercial CPUE.




Figure D4-2: Implied effective samples. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the mean implied effective sample size. The second column show input sample size (x-axis) vs. implied effective sample size (y-axis). Dashed line indicates linear regression slope, and solid line is 1:1 line. The third column show year (x-axis) vs. implied effective sample size (y-axis).




Figure D4-3. Molting probability and trawl/pot selectivity. X-axis is carapace length.

Trawl survey crab abundance




Figure D4-4. Estimated trawl survey male abundance (crab ≥ 64 mm CL). Observed: White: NOAA Trawl Survey, Red: ADG&G Trawl Survey

Modeled crab abundance Feb 01




Figure D4-5. Estimated abundance of legal males from 1976-2015.

MMB Feb 01




Figure D4-6. Estimated abundance of Mature Male Biomass from 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019).

Summer commercial standardized cpue




Figure D4-7. Summer commercial standardized cpue 1977-2018.




Figure D4-8. Total catch and estimated harvest rate 1976-2018.




Figure D4-9. Predicted (dashed line) vs. observed (dots) length class proportions for commercial catch. Black: New Shell, Red: Old Shell




Figure D4-10. Predicted (dashed line) vs. observed (black dots) length class proportions for the winter and spring pot survey.




Figure D4-11. Predicted (dashed) vs. observed (dots) length class proportions for Trawl survey.




Figure D4-13. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D4-12. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D4-13. Predicted vs. observed length class proportions for tag recovery data.




Figure D4-13. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).




Figure D4-14. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).

Table D4. Summary of parameter estimates for a length-based stock synthesis population model of Norton Sound red king crab.

name	Estimate	std.dev
log_q1	-6.627	0.227
log_q2		
log_N76	9.008	0.174
R0	6.341	0.191
a1	1.968	4.606
a2	2.959	4.289
a3	4.020	4.140
a4	4.124	4.127
a5	4.344	4.117
a6	3.570	4.146
a7	2.106	4.414
r1	10.000	0.305
r2	9.663	0.351
log_a	-2.674	0.090
log_b	4.832	0.016
log_φ _{st1}	-5.000	0.067
log_φ _{wa}	-2.203	0.307
log_φ _{wb}	4.800	0.032
Sw1	0.072	0.035
Sw2	0.498	0.125
log_φ _l	-2.085	0.056
log_φ _{ra}	-0.791	0.129
log_φ _{rb}	4.647	0.008
log_φ _{wra}	-0.977	0.543
log_φ _{wrb}	4.655	0.037
w ² _t	0.000	0.000
q NOAA	0.811	0.197
q ADFG	1.200	0.290
σ	3.878	0.209
β ₁	12.453	0.707
β ₂	7.649	0.173
ms78	3.083	0.342

Model 19.4




Figure D5-1. QQ Plot of Trawl survey and Commercial CPUE.




Figure D5-2: Implied effective samples. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the mean implied effective sample size. The second column show input sample size (x-axis) vs. implied effective sample size (y-axis). Dashed line indicates linear regression slope, and solid line is 1:1 line. The third column show year (x-axis) vs. implied effective sample size (y-axis).




Figure D5-3. Molting probability and trawl/pot selectivity. X-axis is carapace length.

Trawl survey crab abundance




Figure D5-4. Estimated trawl survey male abundance (crab \geq 64 mm CL). Observed: White: NOAA Trawl Survey, Red: ADG&G Trawl Survey

Modeled crab abundance Feb 01




Figure D5-5. Estimated abundance of legal males from 1976-2015.

MMB Feb 01




Figure D5-6. Estimated abundance of Mature Male Biomass from 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019).

Summer commercial standardized cpue




Figure D5-7. Summer commercial standardized cpue 1977-2018.




Figure D5-8. Total catch and estimated harvest rate 1976-2018.




Figure D5-9. Predicted (dashed line) vs. observed (dots) length class proportions for commercial catch. Black: New Shell, Red: Old Shell




Figure D5-10. Predicted (dashed line) vs. observed (black dots) length class proportions for the winter and spring pot survey.




Figure D5-11. Predicted (dashed) vs. observed (dots) length class proportions for Trawl survey.




Figure D5-13. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D5-12. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D5-13. Predicted vs. observed length class proportions for tag recovery data.




Figure D5-13. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).




Figure D5-14. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).

Model 19.5




Figure D6-1. QQ Plot of Trawl survey and Commercial CPUE.




Figure D6-2: Implied effective samples. Figures in the first column show implied effective sample size (x-axis) vs. frequency (y-axis). Vertical solid line is the mean implied effective sample size. The second column show input sample size (x-axis) vs. implied effective sample size (y-axis). Dashed line indicates linear regression slope, and solid line is 1:1 line. The third column show year (x-axis) vs. implied effective sample size (y-axis).




Figure D6-3. Molting probability and trawl/pot selectivity. X-axis is carapace length.

Trawl survey crab abundance




Figure D6-4. Estimated trawl survey male abundance (crab \geq 64 mm CL). Observed: White: NOAA Trawl Survey, Red: ADG&G Trawl Survey

Modeled crab abundance Feb 01




Figure D6-5. Estimated abundance of legal males from 1976-2015.

MMB Feb 01




Figure D6-6. Estimated abundance of Mature Male Biomass from 1976-2019. Dash line shows Bmsy (Average MMB of 1980-2019).

Summer commercial standardized cpue




Figure D6-7. Summer commercial standardized cpue 1977-2018.




Figure D6-8. Total catch and estimated harvest rate 1976-2018.




Figure D6-9. Predicted (dashed line) vs. observed (dots) length class proportions for commercial catch. Black: New Shell, Red: Old Shell




Figure D6-10. Predicted (dashed line) vs. observed (black dots) length class proportions for the winter and spring pot survey.




Figure D6-11. Predicted (dashed) vs. observed (dots) length class proportions for Trawl survey.




Figure D6-13. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D6-12. Predicted (dashed) vs. observed (dots) length class proportions for the observer survey.




Figure D6-13. Predicted vs. observed length class proportions for tag recovery data.




Figure D6-13. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).




Figure D6-14. Bubble plots of predicted and observed length proportions. Black circle indicates model estimates lower than observed, white circle indicates model estimates higher than observed. Size of circle indicates degree of deviance (larger circle = larger deviance).

