

# Including ecosystem information in assessments and management advice

---

André E. Punt  
School of Aquatic and Fishery Sciences  
University of Washington



AQUATIC AND FISHERY SCIENCES  
UNIVERSITY *of* WASHINGTON

3 things we assume to be true



I. All reference points are wrong  
some reference points are useful.

*“All models are wrong, but some are useful”.*

George E.P. Box

## II. Stationarity is a key premise of the underpinning of passive adaptive management.

**Definition:** standing still; not moving. having a fixed position; not movable. established in one place; not itinerant or migratory.

**More formally:** in a stationary system, if we observe the system at two times, we cannot, based on the observation, know with any certainty which of the two times came earlier.


**Not correct.** Stationery is a mass noun referring to commercially manufactured writing materials, ...

II. Stationarity is a key premise of the underpinning of passive adaptive management.


---

In a stationary system, the longer we observe the system the more we learn (i.e., we get better precision).

This is not the case (in general) for non-stationary systems.



III. We know that the estimates of parameters vary over time




# Environmental-driven changes in biological parameters are common


Drivers	Dome-shaped		Monotonic			Regimes		
	Env	SpBio	Env	SpBio	SpBio and/or Env	Stocks with/shifts in average	Mean length	SD
Totals	31	7	105	29	52	160	10.3	8.1
Clupeiformes	0	3	14	0	13	25	11.0	7.8
Pleuronectiformes	8	2	23	4	6	34	9.5	7.4
Scorpaeniformes	4	0	20	5	2	21	11.0	7.1
Gadiformes	5	1	25	10	15	39	10.4	8.4
Perciformes	10	1	19	5	14	32	10.7	9.3
Decapoda	1	0	0	4	1	2	7.1	6.1
Ophidiiformes	3	0	3	1	0	5	9.8	7.5
Osmeriformes	0	0	1	0	0	1	9.3	5.1
Tetraodontiformes	0	0	0	0	1	1	7.7	3.8

# What do we know about time-variation in parameters and can we link it to climate?

Most links between climate change (rather than environmental variation) and biological parameters are (currently) inferences based on models. Empirical evidence is depressingly (but perhaps pleasingly) limited.



Perry et al. Science (2005)




Denechaud et al. Glob Chang Biol (2020)




It is time to recognize that the assumption of stationarity is **flawed** and it is necessary to adjust assessments and HCRs to account for environmental variation (process error) and the impact of systematic changes in parameters due to climate change.

**But** do we know how? And what are the consequences of the “wrong call”?



# All environmental effects are not created equal

We need deal with the following types  
of impacts




- Random (perhaps with autocorrelation)
- Regime-shift-like
- Trend (permanent change in the mean)

# Harvest Control Rules

The default approach is to “ignore” changes in productivity except to project weight-at-age and selectivity-at-age when computing OFLs and ABCs, etc.

But this ignores environmentally/  
climate-driven changes.



MSY is “Maximum sustainable yield (MSY) is the largest long-term average catch or yield that can be taken from a stock or stock complex under prevailing ecological and environmental conditions”.

# Short-term and long-term considerations

## Shorter-term considerations:

- Values for biological parameters and MCRs.
- How to assess stock and recruitment.
- How to deal with issues unrelated to fishing.

## Longer-term considerations:

- Changes in strategic targets and limits.
- **New** trade-offs among (new) conflicting objectives.
- Changes in distribution (and hence allocation issues).

The future may be here already!



# Short-term considerations


The key steps we need to address in the short-term (aka now) are:

- When to decide that the (whole) past is inadequate to characterize the future.
- Which parameters to allow to be time-varying
  - Growth – certainly
  - Selectivity – definitely
  - Recruitment - probably
  - Catchability – probably
  - Natural mortality – perhaps
- What does this imply for existing HCRs and reference points?



# Current ways to account for time-varying parameters when defining reference points and applying HCRs (short-term)


- Ignore time-varying parameters
- Use methods to select **“under prevailing ecological and environmental conditions”**
  1. Calculate reference points based on a moving average
  2. Explicitly allow for time-varying parameters when conducting assessments (e.g. multispecies models)
  3. Dynamic  $B_0$  approach



# Reference points

Control rules can be modified by basing reference points on “recent conditions”. What could we mean by this?


- Base  $F_{MSY}$  (or  $F_{SPR}$ ) on current values for growth, natural mortality?, selectivity etc.



# Reference points

Control rules can be modified by basing reference points on “recent conditions”. What could we mean by this?



- Base  $F_{MSY}$  (or  $F_{SPR}$ ) on current values for growth, natural mortality?, selectivity etc.
- Define biomass reference points on time-varying stock-recruitment relationships or dynamic  $B_0$ .
- Change unfished biomass based on selecting a current regime.





# Use methods to select “under prevailing ecological and environmental conditions”

Zheng and Siddeek (2019) used  $AIC_c$  to select when the stock-recruitment relationship for Bristol Bay red king crab changed. The underlying model was a change-point regression of the form  $\log(R_y/SSB_y)$  vs  $SSB_y$




# This can be controversial!

Jackass morwong has a “regime shift” in recruitment in 1988 supported by independent evidence for changes and in the residuals about the stock-recruitment relationship.

Allowing for the regime shift meant that a stock that was overfished is now in the “precautionary zone”.


Application of AIC to select stock-recruitment regimes for Eastern Bering Sea Tanner crab led to a “rebuilt” designation.




# Explicitly allow for time-varying parameters when applying HCRs:

This is perhaps the most complicated option and the one that has been adopted least often in practice. Examples could be:

- Survival for larval and juvenile crab depends on pH such that  $F_{MSY}$  is a declining function of time.
- Pacific sardine where  $F_{MSY}$  is a function of temperature (selected using MSE)




PFMC, 2013



Hurtado and Punt, mimeo

Adopt a control rule that depends on an environmental covariate (e.g., Pacific sardine)



# Long-term considerations

This will involve a rethinking (re-envisaging) of:

- Data collection schemes
- Stock assessments
  - More integration of environmental variables
- Harvest control rules
  - Selected based on single-species MSEs (extended single-species models)
  - Selected based on multi-species / ecosystem MSEs?
- Capacity and expertise.



# Harvest Control Rules (longer-term)

Control rules can be modified to include environmental variables

$$F_{\text{targ}} = \begin{cases} 0 & \text{if } B_y \leq B_{\text{lim}} \\ F_{\text{MSY},y} \frac{B_y - B_{\text{lim}}}{B_{\text{targ}} - B_{\text{lim}}} & \text{if } B_{\text{lim}} < B_y \leq B_{\text{targ}} \\ F_{\text{MSY},y} & \text{if } B_y > B_{\text{targ}} \end{cases}$$

$$F_{\text{MSY},y} = F_{\text{MSY}} \left( 1 - \lambda + 2\lambda \frac{I_y - I_{\text{min}}}{I_{\text{max}} - I_{\text{min}}} \right)$$



$I_{\text{min}}$  and  $I_{\text{max}}$  are the minimum and maximum values of an indicator,  $\lambda$  is the “environmental range”

c.f. Bentley et al. Front Mar Sci 2021




# Explicitly allow for time-varying parameters when applying HCRs

Accounting for multispecies interactions increases the complexity related to defining reference points – within the US reference points are not (directly) based on multi-species models.




Explicit multispecies assessments linked to control rules is the (currently unattainable?) gold standard.


But using ecosystem model-based indicators to change single-species target F (within a range that is compatible with management goals) may be a pragmatic way forward.





# How do we evaluate candidate management options?





The operating model component of an MSE can be set up to include environmental drivers of population dynamic processes (usually the parameters of the stock-recruitment relationship, but in principle natural mortality, growth, etc.)

The management strategy could also make use of environmental data (e.g., the MSE for Pacific sardine).

$$F = G \frac{m_1 m_2}{d^2}$$

$$-E + V = Z$$

$$i\hbar \frac{\partial}{\partial t} \psi = \hat{H} \psi$$

$$\phi(x) = \frac{1}{\sqrt{2\pi\sigma}}$$

$$E = mc^2$$


$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

$$\frac{df}{dt}$$

Extended single-species models:


- Identify parameters that may be time-varying (in principle)
- Identify hypotheses that have been postulated / speculated / tested linking environmental variables to parameters
- Triage the environmental variables to those that can be hind and forecasted
- Fit the assessment model to quantify the relationship between the environmental variables and the parameters
- Select a set of harvest control rules to permit projections

# Bottom Temperature (°C)




- GCM\_scen
- cesm\_ssp126
  - cesm\_ssp585
  - gfdl\_ssp126
  - gfdl\_ssp585
  - miroc\_ssp126
  - miroc\_ssp585

**Goal: To address climate information needs with best available science & tools**



# An ACLIM vision



**Goal: To address climate information needs with best available science & tools**



Climate impacts on growth & condition

Climate impacts on survival & abundance

Climate impacts on food-web dynamics

Climate driven changes to fish distributions (& fishing grounds)

Changes in biomass & TAC


Changes in allocation

Changes in value

Changes in catch, revenue, & profit


Changes to the ecosystem

Changes to wellbeing, & communities of place or practice



**Downscaled hindcast/projections:**  
 CORE-CFSR Hindcast (1960-2017)  
 ECHO-G (AR4 A1B)  
 MIROC3.2 med res. (AR4 A1B)  
 CGCM3-t47 (AR4 A1B)  
 CCSM4-NCAR- PO (AR5 RCP 4.5 & 8.5)  
 CCSM4-NCAR- PON (AR5 RCP 8.5)  
 MIROCESM-C- PO (AR5 RCP 4.5 & 8.5)  
 GFDL-ESM2M\*- PO (AR5 RCP 4.5 & 8.5)  
 GFDL-ESM2M\*- PON (AR5 RCP 8.5)

**Bering Sea Models**




Characterize confidence around findings through multi-model approach.

What do we know about MS  
performance


# What have we learned / what do we know?

- Including environmental covariates when defining HCRs is not guaranteed to lead to improved management performance.
  - Pacific sardine yes; other cases marginally.
- Results will be sensitive to how (and to what extent) biological parameters change over time.
  - Amar et al. (2009) linked recruitment to environmental parameters and found sensitivity to the choice of IPCC model.
  - Multi model inferences!!



## What have learned / what do know?

- Including environmental covariates when defining HCRs is not guaranteed to lead to improved management performance.
  - Pacific sardine yes; other cases marginally.
- Results will be sensitive to how (and to what extent) biological parameters change over time.
  - Amar et al. (2009) linked recruitment to environmental parameters and found sensitivity to the choice of IPCC model.
- Results will be sensitive to how the model structure.
  - Multi-model MSE has been conducted for Pacific sardine.





# What have we learned / what do we know?

- Including environmental covariates when defining HCRs is not guaranteed to lead to improved management performance.
  - Pacific sardine yes; other cases marginally.
- Results will be sensitive to how (and to what extent) biological parameters change over time.
  - Amar et al. (2009) linked recruitment to environmental parameters and found sensitivity to the choice of IPCC model.
- The environment can impact both the target fishing mortality (e.g. Pacific sardine) or the target biomass (e.g. crab and pollock)
  - Kell et al. (2005) found that management based on fishing mortality was more robust than management based on biomass when growth and recruitment were driven by climate.

Table 3. Summary of long-term fishing mortality reference points in 2001. For reference,  $F_{2000-2002} = 0.96$ ,  $F_{lim} = 0.86$ .



	2001	$\alpha$		$\beta$	
		HadCM3	Constant increase	HadCM3	Constant increase
$F_{c_{max}}$	0.98	0.91	0.84	0.98	0.98
$F_{MSY}$	0.51	0.47	0.43	0.51	0.51


Table 5. Summary of relative long-term biomass, given as a percentage of  $B_{pu}$  in 2001.

	2001	$\alpha$		$\beta$	
		HadCM3	Constant increase	HadCM3	Constant increase
$B_{F_{2000-2002}}$ (%)	32	6	0	24	20
$B_{F_{pu}}$ (%)	141	114	90	106	86
$B_{F=0.45}$ (%)	255	228	203	192	155

## What have learned / what do know?

- Including environmental covariates when defining HCRs is not guaranteed to lead to improved management performance.
  - Trends matter more than random variation and regimes.
- The risk of ignoring trends in environmental drivers is not symmetric
  - Ignoring time-trends in environmental drivers is more consequential than trying to account for environmental drivers when this is not needed.
  - It can be useful to only use an environmental driver when it explains a “reasonable” amount of variance (e.g. 50% De Oliveira and Butterworth, 2005).





There be  
dragons here

---

**Table 2**

Summary weight of evidence per example species.


Stock	Observed change in a productivity indicator	Understanding of assessment model input data	Understanding of assessment model structural assumptions	Explanatory hypothesis	Weight of evidence
Southern Gulf of St. Lawrence Atlantic cod	18 years of low biomass (3)	Stock biology well characterized. Catch and its composition is well known and the survey exhibits low CVs (3)	Modeled values of total mortality suggest that natural mortality has increased for a number of sub-stocks (2)	Increase in natural mortality is mostly attributed to increased predation by seals, but no direct measurement from field studies (2)	10
Gulf of Maine/Georges Bank Atlantic herring stock complex	16 years of relatively high natural mortality (2)	Some uncertainty in stock boundaries and catch levels that have not been characterised (1)	A modeled increase in natural mortality since 1995 eliminates a strong retrospective pattern and better matches consumption estimates (2)	A multispecies model that estimates herring consumption based on a long time series of stomach contents across a range of species was used to inform the <i>M</i> increase hypothesis (2)	7
SNE/Mid-Atlantic yellowtail flounder	21 years of low biomass (2)	Some uncertainty in stock boundaries and catch levels that have not been characterised (1)	Modeled annual recruitment residuals below average for past 22 years (1)	Effect of oceanographic processes on recruitment is poorly understood. Environmental correlate with poor recruitment sought but not found (0)	4
Jackass morwong	12 years of low biomass (2)	Stock boundaries well defined, total catch well estimated and uncertainty in CPUE accounted for. Percentage of catch covered by size/age sampling is relatively low (2)	Modeled annual recruitment residuals below average for past 19 years. Fit to the data is improved by productivity shift; lower steepness cannot account for decline (2)	Some knowledge of larval distribution and recent changes in ocean circulation conditions lead to a plausible mechanism. Corroboration from some other species (1)	7
Gemfish	22 years of low biomass (3)	Stock boundaries well defined, total catch well estimated and uncertainty in CPUE accounted for. Percentage of catch covered by size/age sampling is relatively low (2)	Modeled annual recruitment residuals generally below average for past 24 years (1)	Larval biology poorly understood. Mechanism for poor recruitment not understood (0)	6



Good news (increased average recruitment)  
may not be..

Bad news (lower average recruitment may not  
be either)

“contrary to the recognition of recruitment failure and to  
all indicators precipitously declining, this fishery remains  
categorized as ‘not overfished’ and is listed as  
‘sustainable’.” (Edgar et al., Aquatic Conservation)



What the data  
giveth the  
data may take away

I review the role of environmental variability in the survival of juvenile fish and shellfish by examining the success of previously published environment–recruitment correlations when tested with new data. The proportion of published correlations that have been verified upon retest is low. There is one generalization that stands out: correlations for populations at the limit of a species' geographical range have often remained statistically significant when re-examined. An examination of environment–recruitment correlations that were reviewed 13 years ago by Shepherd and co-workers shows that only 1 out of 47 reviewed studies is currently used in the estimation of recruitment in routine assessments. The results suggest that future progress will require testing general hypotheses using data from many populations.

*Reviews in Fish Biology and Fisheries* **8**, 285–305 (1998)

## **When do environment–recruitment correlations work?**

RANSOM A. MYERS

*Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada B3H 4J1. E-mail: Ransom.Myers@Dal.Ca*



We need to think about the consequences about the productivity paradox.

Coming up from Cody!

# Tentative Recommendations

- We need to recognize that short- and long-term environmental factors will influence population dynamics, including species distributions.
- The default for developing harvest control rules and management plans should be the assumption of non-stationarity dynamics:
  - Stationarity should only be assumed given a lack of data (is this cutting your throat with Occam's Razor)
  - We need operational ways to decide when to allow reference points to change over time.
- It is time **now** to construct frameworks to evaluate alternative management systems (think training, data needs, and stakeholder engagement)





A blue ballpoint pen with a silver tip is positioned diagonally across the top left of the frame. The background is a document with a grid and a bar chart with several blue bars of varying heights. The text is overlaid on the right side of the image.

# Data demands?

What are the data demands of supporting climate-based models and HCRs?

- Will existing surveys and other monitoring schemes be sufficient given changing distributions? Are we ready for the monitoring systems needed to track species as they move in response to climate-driven movement?
- How many regions have sufficient data on population dynamics (e.g. weight-at-age, diet?)

# Process demands?

---

- How do we conduct reviews for multispecies / environmental assessment frameworks and HCRs?
- What are the processes for obtaining stakeholder feedback on trade-offs among objectives

## A guinea pig's tale: learning to review end-to-end marine ecosystem models for management applications

Isaac C. Kaplan<sup>1\*</sup> and Kristin N. Marshall<sup>2</sup>

<sup>1</sup>Northwest Fisheries Science Center, National Marine Fisheries Service, NOAA, 2725 Montlake Boulevard E., Seattle, WA 98112, USA

<sup>2</sup>University of Washington School of Aquatic and Fisheries Sciences, PO Box 355020, Seattle, WA 98195, USA

\*Corresponding author: tel: +1-206-302-2446; fax: +1-206-860-3394; e-mail: [isaac.kaplan@noaa.gov](mailto:isaac.kaplan@noaa.gov)

Kaplan, I. C., and Marshall, K. N. A guinea pig's tale: learning to review end-to-end marine ecosystem models for management applications. – *ICES Journal of Marine Science*, 73: 1715–1724.


Received 10 June 2015; revised 3 March 2016; accepted 7 March 2016.

A shift towards ecosystem-based management in recent decades has led to new analytical tools such as end-to-end marine ecosystem models. End-to-end models are complex and typically simulate full ecosystems from oceanography to foodwebs and fisheries, operate on a spatial framework, and link to physical oceanographic models. Most end-to-end approaches allow multiple ways to implement human behaviours involving fishery catch, fleet movement, or other impacts such as nutrient loading or climate change effects. Though end-to-end ecosystem models were designed specifically for marine management, their novelty makes them unfamiliar to most decision makers. Before such models can be applied within the context of marine management decisions, additional levels of vetting will be required, and a dialogue with decision makers must be initiated. Here we summarize a review of an Atlantis end-to-end model, which involved a multi-day, expert review panel with local and international experts, convened to challenge models and data used in the management context. We propose nine credibility and quality control standards for end-to-end models intended to inform management, and suggest two best practice guidelines for any end-to-end modelling application. We offer our perspectives (as recent test subjects or “guinea pigs”) on how a review could be motivated and structured and on the evaluation criteria that should be used, in the most specific terms possible.

# Final thoughts

- To date, climate-enhanced assessment and MSE frameworks have not been adopted for tactical and strategic decision making.
- Approaches such as ACLIM require much expanded **collaboration** networks (oceanographers, biologists, modellers, economists, social scientists, and communicators) – **communicators** may be our most critical need.
- We need to start now to develop partnerships, identify knowledge gaps, and create frameworks.

## Ecosystems say good management pays off

Elizabeth A. Fulton<sup>1,2</sup>  | André E. Punt<sup>1,3</sup> | Catherine M. Dichmont<sup>4,5</sup> | Chris J. Harvey<sup>6</sup> | Rebecca Gorton<sup>1</sup>

<sup>1</sup>CSIRO, Hobart, Tasmania, Australia

<sup>2</sup>Centre for Marine Socioecology, University of Tasmania, Hobart, Tasmania, Australia

<sup>3</sup>School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington

<sup>4</sup>Cathy Dichmont Consulting, Banksia Beach, Queensland, Australia

<sup>5</sup>The College of Science and Engineering, James Cook University, Douglas, Queensland, Australia

<sup>6</sup>NOAA Fisheries, Northwest Fisheries Science Center, Seattle, Washington

### Correspondence

Elizabeth A. Fulton, CSIRO, Hobart, TAS, Australia.  
Email: beth.fulton@csiro.au

### Funding information

Fisheries Research and Development Corporation, Grant/Award Number: 2012/202

### Abstract

Understanding the strengths and weaknesses of alternative assessment methods, harvest strategies and management approaches are an important part of operationalizing single-species and ecosystem-based fisheries management. Simulations run using two variants of a whole-of-ecosystem model for the Southern and Eastern Scalefish and Shark Fishery (SESSF) area shows that (a) data-rich assessments outperform data-poor assessments for target species and that this performance is reflected in the values of many system-level ecosystem indicators; (b) ecosystem and multispecies management outperforms single-species management applied over the same domain; (c) investment in robust science-based fisheries management pays dividends even when there are multiple jurisdictions, some of which are not implementing effective management; and (d) that multispecies yield-oriented strategies can deliver higher total catches without a notable decline in overall system performance, although the resulting system structure is different to that obtained with other forms of ecosystem-based management.

### KEYWORDS

Atlantis, ecosystem-based management, fisheries, harvest strategies, risk equivalency





**QUESTIONS?**



# Use methods to select “under prevailing ecological and environmental conditions”

Many variants of the basic approach exist.

- Kapur et al. (2020) used the first derivative of the spatial/temporal smoothing term of a generalized additive model to identify spatial/temporal zones of variation in fish length-at-age.
- This is one way to select when a new regime has started.

