## **1** Executive Summary

2 Pacific salmon (Onchorhynchus spp.) are an essential part of Alaska's commercial, recreational, and 3 subsistence fisheries, providing economic opportunities to communities in Alaska as well as supplying important food and traditional and cultural practices for tens of thousands of Indigenous and rural 4 people and communities. Alaska maintains some of the best freshwater and marine habitats for salmon 5 health and resilience. Despite this, some Alaska salmon populations are facing sustained and dramatic 6 declines, with devastating impacts to food security and traditional ways of life for the people that 7 depend on them. In addition, some salmon stocks are experiencing wide fluctuations in returns that 8 lead to high inter-annual variation creating uncertain economic outcomes. Because of these declines, 9 Congress enacted the Alaska Salmon Research Task Force Act (AKSRTF; Appendix 1) to identify the 10 gaps in knowledge that are needed to understand the variability and declining trends. The purposes of 11 the act are to: 1) ensure that Pacific salmon productivity and abundance trends in Alaska are 12 characterized and that research needs are identified; 2) prioritize scientific research needs for Pacific 13 salmon in Alaska; 3) address the increased variability or decline in Pacific salmon returns in Alaska 14 by creating a coordinated salmon research strategy; and 4) support collaboration and coordination for 15 16 Pacific salmon conservation efforts in Alaska.

17 In this regard, the AKSRTF recommends the following:

# 18 Gravel to Gravel (G2G) life history research strategy

Salmon life begins and ends in the gravel and throughout their life history they depend on freshwater and marine habitats to grow and thrive. Through G2G, Tribes, State and Federal agencies and institutions and others work together to build a strong foundation for co-stewardship, where Traditional and Indigenous Knowledge along with western science play important parts in resilient habitats and communities within Alaska.

# Potential Drivers influencing Pacific salmon production in Alaska and recommended applied research needs

Based on a review of existing knowledge, the AKSRTF identified the following potential drivers
influencing production within the Pacific salmon life cycle in Alaska (in order of the number of life
history stages impacted from all to less) and associated priority research needs:

# 29 Warming climate and extreme events

Research to understand and quantify the effects of natural environmental variability and human
 factors on Alaska salmon distribution and abundance.

# 32 Salmon health and condition

- Research to understand the connections between freshwater and the marine environment that
   lead to pathogens or declines in thiamine levels for salmon as these changes could affect the
   ability of returning adults to successfully reach the spawning grounds, successfully spawn, the
   numbers of eggs produced and fertilized, and their ability to produce viable offspring.
- Research to understand prey quality and quantity on health and condition of salmon in marine
  and freshwater habitats.
- Research to understand the mechanism(s) behind declining size at age as these declines impact
  the amount of food available per fish, the number of eggs per female for future generations,
  and can contribute to declining run sizes.

# 42 *Predators*

- Research to understand potential conflicts between predator or endangered predators and prey (salmon).
- Research to address the role of hatchery salmon release sites has on drawing in assemblages
   of predator species that otherwise would not be present in coastal nurseries, thereby potentially
   increasing predation pressures on wild stocks that may also inhabit these nurseries.

# 48 *Marine food limitations*

- Research to understand the implications of habitat use by Alaska salmon populations at various
   levels of abundance, the productive capacity of habitats for each life stage, and the potential
   implications of density-dependent effects.
- Research to better understand the role of salmon in pelagic communities, the food availability
   for salmon and the nutritional quality of prey organisms, including harmful algal blooms, to
   better understand inter-and intra-specific competition among salmon at sea.

# 55 *Marine harvest and bycatch*

- Research to reduce bycatch, interception, and Illegal, Unreported, and Unregulated (IUU)
   fishing through improved understanding of distribution and migration patterns of Alaska
   salmon stocks to better predict and avoid incidental harvest in the migratory corridors for
   Alaska salmon including Bering Sea, Aleutian Island, and Gulf of Alaska areas and regions in
   the North Pacific where there is increased potential for IUU fishing.
- Research to improve our ability to determine the stock origin of chum and Chinook salmon
  taken in marine harvest, bycatch, and interception.
- Research to understand the frequency of occurrence and mortality rate (direct and discard)
  attributed to unobserved fishing mortality (e.g., IUU, unreported catch, incidental catch/mixed
  stocks); and once this information is known, what is the impact to the populations.
- Research to better define how all sources of marine harvest influence salmon abundance and
  how does this vary by species / region.
- Research to understand genetic diversity and fitness effects from fishing that may reduce a
   population's resilience and ability to recover from climate induced depression of population
   abundance or low productivity.

# 71 *freshwater habitat changes*

- Research to develop meaningful measures of ecosystem performance (space and time scales)
   that supports biological diversity of Alaska salmon to maintain and conserve the processes that
   confer resilience (habitat and/or genetic diversity) in face of ongoing environmental change.
- Research that improves our understanding of the impact of hatchery strays on wild salmon
   where intermingling with those stocks in freshwater has the potential to reduce genetic
   diversity, reproductive success, and resilience to climate variability and change.

# 78 freshwater harvest

Research that addresses mortality rate attributable to unobserved freshwater fishing mortality
 due to release, incomplete capture, unreported catch, and illegal fishing and how this source of
 mortality impacts the populations.

Research that documents production changes and spawning success in the high Arctic as
salmon begin to return in larger numbers to the region.

## 84 Recommended Applied Strategies to Address Priority Research Needs

## 85 Improved stock identification methods

Bevelop/improve novel stock and fish identification methods at a finer scale than is currently available from genetic mixed stock analysis. In some areas, like western Alaska, current stock groupings based on genetic distinctions cover wide geographical areas that do not allow a full understanding of the impacts of marine harvest at the finer resolution used for management and impact assessment.

# 90 Better characterization of ocean distributions and marine migration routes

91 Develop/improve/expand research to understand the migratory routes and ocean distributions for

92 western Alaska salmon to reduce bycatch potential (see recommendations by the Alaska Bycatch

93 Review Task Force report dated November 2022; Appendix 5) and interception and to understand

94 potential for impacts on their health from shifting food webs and competitive pressures.

95 *Expanded ocean ecosystem surveys* 

Develop/improve/continue ocean surveys to understand how shifts in climate and ocean conditions (climate warming/extreme events) impact the food web and health and condition of juvenile salmon populations, with the primary research goal to identify additional marine management actions, with the secondary goal of improving forecasts of short- and long-term prospects for decision makers.

100 Strategies to minimize human impacts on freshwater and coastal habitats

101 Develop/improve/expand strategies to prioritize actions that reduce human impacts on freshwater and 102 coastal ecosystems with the goal of maximizing the number, diversity, and health of wild smolts and 103 spawners.

104 *Making use of new technologies* 

Develop/improve/expand use of new technologies and advanced analytical methods for Alaska salmon
 research, including molecular identification, genomics, environmental DNA, mass marking,
 intelligent tags, salmon observation systems, and remote sensing/ autonomous vehicles.

# 108 More effective monitoring of salmon Indicator Stocks

109 Develop/improve/continue to identify indicator stocks for Chinook and chum salmon that can be 110 monitored and tracked throughout their life cycle to better understand mechanisms impacting survival

111 in marine and freshwater habitats and provide an early warning system.

112 *Improved stock assessments for in-season management* 

113 Develop/improve/continue stock assessment programs that allow for timely in season management

decisions to mitigate uncertainties in adult salmon return strength.

# 115 *Life-cycle modeling and management strategy evaluations for climate resilience*

Develop/improve/expand approaches to modeling biological impacts of climate change on full life cycle of Pacific salmon that include management strategy evaluations to test how different management actions may impact production in relation to climate scenarios, ensemble models to characterize uncertainty in climate impact projections, and ocean intelligence systems for targeted information on impacts of climate warming and extreme events on ocean ecosystems and salmon growth and health.

122 Better data management and sharing – work in progress

# 123 **Recommended Framework**

Work with Tribes/Federal/State to initiate Two-Eyed Seeing framework that embraces "learning to see from one eye with the strengths of Indigenous knowledges and ways of knowing, and from the other eye with the strengths of mainstream knowledge and ways of knowing"

127

## 129 Critical Need to Understand Shifts in Alaska Salmon Productivity

Alaska is warming at a rate two times faster than the Lower 48 contiguous states which is having a profound effect on Alaska salmon populations. Improved understanding of the mechanisms that regulate the distribution, migration and abundance of Alaska salmon will promote their conservation, allow for better projections, characterize uncertainty for production trends under climate warming, and enhance the sustainable fisheries management, food security and economic security for Alaskans.

135

These dramatic shifts in Alaska salmon productivity are occurring despite the fact that freshwater 136 habitats where Alaska salmon reside during their life history are relatively pristine, especially when 137 compared to habitats that salmon stocks encounter at lower latitudes. Existing knowledge regarding 138 Alaska salmon ecology indicates that warming in both freshwater and marine habitats is creating 139 divergent impacts on salmon species and stocks where some are responding positively to warming 140 (i.e., abundance is increasing) and others are declining in response. For example, residents of the 141 Yukon River drainage have experienced declines in returning Chinook salmon since 2001 with 142 minimal improvement and periodic crashes in chum salmon, most recently in 2021-2023. These 143 144 declines have led to the cessation of in-river commercial fisheries for Chinook salmon and severe restrictions to subsistence fishing for all salmon species, including a complete closure of subsistence 145 146 fishing in 2022.

147


148 It is critical that we take action now to understand mechanisms driving Alaska salmon 149 production and to provide a path for mitigating negative impacts. Acting now allows for a 150 response before there is further decline, making immediate action more successful and cost 151 effective than waiting until stocks become critically low or reach depleted status.

152

As such, the AKSRTF members were acutely aware of the impact the declining Chinook and chum salmon returns in western Alaska are having on culture and food security in that region. Through the AKSRTF Act, the AKSRTF formed the Arctic Yukon Kuskokwim Working Group (AYK WG) to identify priority research needs for salmon in that region. A report from the AYK WG is located on page 27. Public testimony often included recommendations for action that can be done now, such as management actions to close fisheries where western Alaska salmon are harvested as bycatch or interception. Although the AKSRTF understood it would be impossible to meet all expectations, our

- 160 commitment was to develop priority research needs that would enable decision makers at local and161 regional levels to act quickly in response to research results.
- 162
- 163 It is through this lens that the AKSRTF set out to recommend a coordinated research strategy based
- 164 on the review of existing Pacific salmon research and the identification of knowledge gaps, and applied
- research needed to better understand why Alaska salmon are experiencing increased variability and
- 166 declines.

167 **Coordinated Research Strategy** (Gravel to Gravel/Life History Figure)



<sup>168</sup> 

Salmon life begins and ends in the gravel (Gravel to Gravel – G2G) and throughout their life history they depend on freshwater and marine habitats to grow and thrive. It is through the utilization of diverse, pristine habitats found in Alaska that salmon can gain resiliency to the effects of climate warming. However, Alaska salmon are facing greater challenges than those at lower latitudes due to the pace of warming (two times faster than lower latitudes) and extreme events (marine heat waves,

174 loss of seasonal sea ice, drought, spikes in freshwater temperatures, etc.) that can have dramatic 175 impacts on their survival. Through, G2G federal agencies, Tribes, state agencies and institutions and 176 others can work together to build a strong foundation for co-stewardship, where traditional and 177 Indigenous Knowledge along with western science play important parts in support of resilient habitats 178 and communities within Alaska.

We propose a G2G assessment approach of coordinated research where individual projects, regardless 179 180 of whether they are led by state, federal, university, tribal, or non-profit entities, will share information with all other projects, and this strategy includes an intentional integration of data and information 181 182 across research projects. Prior salmon research efforts have undoubtedly enabled important 183 advancements in our knowledge and understanding of poor salmon abundance patterns across Alaska. 184 However, when each research project is advanced and understood in isolation, which is the norm, we often fail to develop a synthesized and holistic perspective across the entire salmon life cycle. 185 186 Consequently, it becomes difficult to develop an integrated and unified picture of the nature of where salmon bottlenecks are, and to clearly identify the most important drivers of these bottlenecks. 187 Perceived progress takes a long time to develop, and it becomes challenging for experts with different 188 perspectives to coalesce under a common understanding. 189

190

191 To do this, for each stock selected for study, an intensive suite of studies will be implemented concurrently over a 5 to 6-year period. Together, the suite of projects should address all relevant 192 potential drivers for salmon abundance identified in the AKSRTF report and all life stages, ideally 193 with overlap across multiple projects. In addition to data collection studies, retrospective analyses and 194 modeling efforts will be useful to "bring it all together" by consolidating data across suites of projects 195 and integrating data from separate studies. This synthesis will highlight critical factors or life stages 196 limiting salmon adult run abundance and inform potential policy and management actions. Research 197 approaches that are intensive and holistic, employing coordinated and focused examinations of all 198 potential drivers at once, have been particularly successful for identifying factors most important to 199 survival and productivity of salmon in other areas (e.g., Salish Sea Marine Survival Project), and it is 200 expected that the G2G assessment approach would be similarly successful. 201

## 202 Potential Drivers for Alaska Salmon Production

Based on a review of existing knowledge, the AKSRTF identified the following potential drivers 203 204 influencing production within in the Pacific salmon life cycle in Alaska (listed in order of the number of salmon life history stages impacted (all to less); no priority assigned): warming climate/extreme 205 events, salmon health and condition, predators, marine food limitations, marine harvest and bycatch, 206 freshwater habitat changes, and freshwater harvest. Research gaps were then identified within these 207 208 potential drivers of Alaska salmon productivity which set the stage for the AKSRTF to develop priority research needs and applied strategies (listed in the Executive Summary) to address each potential 209 210 driver.

211

# 212 Warming Climate/Extreme Events (impacts all life stages)

Alaska is warming at a rate two times faster than the lower latitudes, and this warming is affecting all 213 214 aspects of the salmon life cycle. Indirect impacts of climate-related phenomena include changes to 215 timing of salmon smolt outmigration and upriver migration, reduced fitness in response to shifts to lower quality prey as well as shifts in their ocean migration and distribution patterns. Warming is also 216 increasing the frequency of extreme events that have profound negative impacts on some species and 217 stocks of salmon depending on what freshwater systems are utilized for spawning and the migration 218 routes taken during their marine life history. Extreme events can lead to short term (days to weeks) 219 changes in water flow, oxygen levels in freshwater, and spikes in freshwater temperatures, as well as 220 seasonal changes (months) such as loss of seasonal sea and lake/river ice, and longer-term phenomena 221 (months to years) such as marine heat waves. These extreme events have direct physiological impacts 222 on salmon stocks, such as heat stress that can compromise the success of returning adult spawners or 223 juveniles in the ocean and reduce health and condition of salmon in the marine environment in 224 response to shifts to lower quality prey as well as shifts in their ocean migration and distribution 225 226 patterns.

227 Salmon Health and Condition (impacts all life stages)

Changes to the health and condition of Pacific salmon in Alaska have been noted by fishers and biologists. Across Alaska, Chinook salmon and other species of salmon have been experiencing declining size at age as well as shifts in age at maturity (to younger age). Other changes include

increased presence of ichthyophonus (a fungal-like infection) and a reduction in thiamine levels in Chinook salmon, both of which are believed to come from their marine prey. Other changes that are found for Alaska salmon include deficiencies in fat stores, particularly during early marine life history stages and potentially during adult spawning migration. All of these changes could increase mortality rates during marine life history stages and affect the ability of returning adults to reach spawning grounds and spawn successfully, and decrease their ability to produce viable offspring.

237 *Predators (impacts all life stages)* 

Predation occurs throughout the Pacific salmon life cycle, but can be difficult to assess, especially in estuarine and marine environments. Salmon sharks are estimated to consume 73 to 146 million Pacific salmon each year. Many marine mammal predators of salmon, particularly seals and sea lions, have increased in abundance in recent decades. Changes in subsistence practices along some Alaska rivers, for example, the decreased need to harvest fish such as northern pike to feed dog teams, has resulted in increasing abundance of freshwater predators in some places, as has the expansion of pike beyond its native range.

245

Socio-economic changes over since the 1970s have also affected the ecological system. As snow machines replaced dog teams for transportation, community residents kept fewer dogs and needed fewer salmon and non-salmon fish species, such as sheefish and Northern pike, to feed them. Some researchers have theorized that the decreased harvests of these piscivorous fish have led to increased predation on juvenile salmon in the fresh water environment.

# 251 *Marine Food Limitations (impacts marine life history stages)*

Many Alaskan salmon stocks are experiencing particularly poor marine survival, which could be due to marine food limitations. Pacific salmon move from rivers to sea to take advantage of better feeding and growing opportunities in the ocean. However, there are several indications that critical changes have occurred for Alaskan salmon marine feeding and growth opportunities that adversely impact their marine survival and/or their health and condition when they return to their natal spawning rivers. For instance, local and Indigenous knowledge holders are raising concerns about deteriorating fat content and health of returning salmon. Additionally, a growing body of scientific literature associates many

of these abundance or size declines with competition among salmon species, including those of hatchery origin. Therefore, it is critical to understand the mechanisms and degree to which marine food limitations may be causing poor returns of Alaskan salmon, and to understand what actions could possibly mediate these conditions.

# 263 *Marine Harvest and Bycatch (immature and maturing life stages)*

Marine harvest of Pacific salmon occurs at multiple scales (i.e., international, national, state, regional, etc.) and there are many complexities to reporting and attributing catch (limits to genetic stock identification) at these various levels. In addition, most ocean fisheries have some amount of bycatch or interception of Pacific salmon as part of the harvest process. There are challenges in tracing bycaught salmon to their stock of origin given that the various stocks intermingle in the marine environment.

# 270 Freshwater Habitat Changes (spawning adult to smolt life stages)

Much of Alaska's freshwater habitat is considered relatively pristine compared to those in lower latitudes. While intact landscapes are most likely to support biological diversity and the reliable delivery of salmon to ecosystems and people, they remain subject to large-scale drivers such as warming climate. For example, intact freshwater landscapes help to buffer environmental variability and contribute to long-term stability of salmon populations through differing responses to varying conditions (much like the stabilizing effect of asset diversity on financial portfolios), yet the buffering can be overwhelmed in times of drought or changing water tables.

278

279 In some regions of Alaska, dramatic changes to freshwater habitat have been brought on by glacial 280 recession, isostatic rebound, and tectonic forces. These changes to the landscape impact freshwater habitats in countless ways, both positive and negative, across the state. Intact habitat allows for salmon 281 populations to respond positively in some cases, such as in Glacier Bay in Southeast Alaska, where 282 283 receding glaciers have resulted in new freshwater habitats and colonization by salmon. In other cases, 284 invasive species such as *Elodea* present a significant risk to salmon streams in some regions of Alaska as the plant affects the quality of habitat for juvenile salmon. Hatchery adults also stray into streams 285 286 where wild stocks are spawning and have been known to intermingle with those stocks potentially

reducing genetic diversity, reproductive success, and resilience to climate variability and change. The presence of hatchery strays can also make it difficult to monitor escapements of wild salmon by inflating aerial and foot survey counts, and has resulted in reductions in geographic coverage of wild stock escapement indices in some areas where high hatchery stray proportions have been documented.

291

Traditional Knowledge holders often communicate that sport fishing can interfere with salmon 292 survival, both through the physical disturbances caused by sport fishers walking in streams and on 293 riverbanks, as well as through overall disrespectful behavior towards salmon often described as 294 "playing with one's food." Beavers have also increased in number in interior regions of Alaska: "...the 295 beavers, they dam the river where the spawners can't even go through the dam...seeing...lots of beaver 296 dams. There used to be no beavers in our area (Kuskokwim River)...migrating into the lower 297 streams...down to the coast now." In addition, more roads along freshwater river systems can create 298 new challenges: "A chemical sprayed on tires actually kills salmon...driving on roads...can affect the 299 salmon." 300

# 301 Freshwater Harvest (adult life stage)

The effect and magnitude of historical freshwater harvest (commercial, sport, and subsistence) on salmon populations is not well understood. Even though freshwater harvest has been reduced or eliminated in some areas because of recent declines, the productivity of stocks continues to decline, which suggests that freshwater harvest is not the primary driver on the abundance relative to other influences. The biggest information gap for subsistence harvest is in the Arctic, but, overall, this is not seen as a major influence on abundance relative to other influences.

308

309 For subsistence harvest, traditional and Indigenous Knowledge illustrates that salmon provides not only an essential food source for families, but also supports important cultural, linguistic, and family 310 311 traditions that encourage health and wellbeing within communities. Changing patterns in the harvest and use of salmon continue to drive disconcerting social changes in the region, such as reducing the 312 time that families spend together at fish camps and the resulting challenges to passing on cultural 313 knowledge between older and younger generations. "We have to respect them and keep what you do 314 315 catch very clean and handle them carefully...The animal, fish, or something we take into our home the women they take care of it right away...with no complaining...because the person who hunts will get 316

- 317 more not less...if we leave it... will not catch anymore while others are catching more. We have to 318 respect them and keep what you do catch very clean and handle them carefully." "...our people are 319 into sharing. Elders first. Families first. We never kept the first king. We shared it." "Our cultural and 320 our self-identity was giving to the Elders and sharing our catch."
- 321
- 322

## 323 Existing Knowledge

324 Because of salmon's importance to food security as well as its cultural and economic value, there is a 325 lot known about what salmon need to thrive, particularly in the freshwater phase of their life cycle. Adult salmon require the ability to move from the ocean to freshwater habitats, which must provide 326 the conditions that support the healthy development of eggs, fry, and juveniles. Collectively, 327 freshwater habitat quality can be characterized by the 4Cs: "clean, cool, complex, and connected". 328 Spawning beds must consist of clean gravel that is free of silt, or there must be sufficient movement 329 of water between the stream and its gravel bed ("hyporheic flow", or upwelling/downwelling) to 330 prevent eggs and embryos from suffocating. Streams and lakes must be free of toxic levels of heavy 331 metals, pesticides, and other pollutants. Complex habitats, such as rivers with healthy floodplains, are 332 important for fueling food webs and giving juvenile salmon the ability to move into side channels that 333 might have better feeding conditions or fewer predators. Habitat features such as undercut banks and 334 logiams provide small salmon protection from high flows and from predators, and lakes and beaver 335 ponds can provide good overwintering habitat for species such as coho salmon that spend at least a 336 full year in freshwater before going to sea. Rivers with good forest cover, intact floodplains, or active 337 338 hyporheic zones can also provide salmon the opportunity to seek out waters that are neither too cold in the winter nor too warm in the summer. Finally, young salmon must be able to migrate downstream 339 to the ocean without being blocked by dams or culverts or entrained in diversions, and returning adult 340 salmon must also be able to travel unimpeded to their spawning grounds. 341

342

Each female salmon can produce several thousands of eggs during her one chance at spawning. 343 344 Because of this, salmon can support high levels of non-industrial harvest (i.e., nearshore and in-river; commercial, sport, and subsistence) when freshwater and ocean conditions are good. They can also 345 346 rebound from population declines when the climate is favorable and freshwater habitats have not been 347 impacted by damming, logging, or floodplain development. Healthy salmon populations seem to do best when they are subject to an intermediate level of harvest. Too little harvest can allow too many 348 adults on the spawning grounds, which can reduce egg survival, and when too many juvenile salmon 349 350 are produced they can compete with each other for food. When harvest is too high, freshwater habitats suffer from the lack of gravel cleaning by spawning salmon and a dearth of nutrients that are brought 351 in from the ocean to spawning grounds by adult salmon. Very small populations can also lose genetic 352 353 diversity, which can jeopardize their ability to evolve in response to external stressors such as climate

change. These ecological concepts align well with an ethic that is shared among many Indigenous
peoples in Alaska, which is to harvest salmon when they make themselves available, take only what
is needed, and to leave salmon alone when they are not doing well.

357

The ocean is where salmon spend most of their lives, and where they put on 99% of their body weight. 358 However, this is a challenging place to study, and we only have a broad understanding of this phase 359 of their life cycle. Upon entering the ocean, salmon generally move in a counterclockwise direction 360 around the North Pacific, cycling between productive summer habitat in the Bering Sea and ice-free 361 waters of the Gulf of Alaska during the winter. Salmon feed on a variety of prey, including 362 zooplankton (e.g., copepods, krill, crab larvae) when small and moving up to squid and forage fishes, 363 such as herring, as they grow. Salmon growth and survival is the result of complex and poorly 364 understood interactions among climate, food, predators, and competitors. Large-scale climate 365 variation, such as shifts in the Aleutian Low Pressure system, control the oceanic currents, ocean 366 temperatures, and weather patterns that set up food production in the ocean - phytoplankton and 367 zooplankton - and oscillations between conditions favorable and unfavorable for salmon can be seen 368 369 throughout Alaska's history (e.g., the 1976/77 "regime shift" that was a boon for Alaska salmon fisheries). The number of hatchery salmon released into the ocean is also at an all-time high, likely 370 371 leading to reduced food for wild salmon in certain places and times.

372

373 In recent years, we have seen the emergence of extreme events such as drought, short term spikes in river temperatures, and marine heatwaves, which manifest as dramatic and persistent increases in 374 375 temperatures across broad regions of the North Pacific. Marine heat waves profoundly alter marine ecosystems, including affecting what and how much food is available for salmon and the kinds of 376 377 predators they face, and they are accompanied by hot weather over land as well, leading to river 378 temperatures high enough to weaken or kill migrating salmon even as far north as the Yukon River. 379 Extreme events such as heat waves complicate our ability to maintain optimal levels of salmon harvest, 380 because the methods for setting harvest are usually developed against some average background level 381 of natural mortality and are not well-equipped to deal with big shifts in natural mortality.

382

Following these scientific observations, we acknowledge that Western science is not the only knowledge source useful for understanding the complex relationships between salmon, their

environments, their life-cycle needs, climate, and other factors. Indigenous and traditional knowledge often goes beyond that which is directly related to ecological aspects of the natural world and includes values associated with the entire world view, such as relationship, responsibility, reciprocity, and redistribution (4 Rs). Empirical observation is critical but in ways that focus on and teach appropriate human action as an integral part of the natural world. For example, beliefs about reciprocal relationships of care between humans and fish teach culturally appropriate behavior around concepts of salmon return and conservation.

392

With regards to existing knowledge about salmon, Indigenous and traditional knowledge have existed,been passed down, and been built upon for thousands of years.

395

396 "Growing up in the village, we lived from the land, river and sea. While engaging in subsistence 397 activities, passed and taught from generation to generation, we continue to do so with great respect to 398 the environment we live in, which can be unforgiving if not taught to survive in it. They taught us how 399 to travel on different land, river, weather and sea ice conditions which can change in a heartbeat.

400

All that they taught us is woven into the fabric of our culture, to be able to survive and perpetuate the 401 life of our culture. We were taught how to relate to the environment we were born to, as well as relating 402 to other persons within our culture. For example, the great respect we have for our parents, aunts and 403 404 uncles. To share our hunting catches with widows and those that need help. Translating the nuances of our culture and life is a challenge at times, from one language to another. I think that Indigenous 405 406 knowledge better conveys all those things passed on that are deep at the core of the subsistence lifestyle we live. The permanency and perpetuity of our culture. Closely too is traditional knowledge, that 407 408 conveys the timelessness of cultural traditions and subsistence practices. To continue, unbroken, with our subsistence way of life." 409

410 (Oscar citation, per his preference)

411

This knowledge can appear in many forms and from multiple cultural traditions and is commonly derived from keen, long- term observation of and interaction with local landscapes. Ultimately, we need an intertwined and holistic approach to understanding Alaska salmon, including relationships between Alaska salmon and all living and nonliving things, that includes observations from multiple 416 knowledge types and from a variety of experiences from scientific research to generational observation417 and harvest.

418

419 "…'Traditional Knowledge,' it includes both Western and Indigenous knowledge…[there] are non-420 natives and have a lot of Traditional Knowledge regarding fish and game. So, I personally feel that 421 using the term "Traditional Knowledge" is an inclusive term which is what this Task Force was 422 required to use. Incorporating Indigenous Traditional Knowledge would benefit knowledge from the 423 locals throughout the state also using Traditional Knowledge from local fisherman regardless of 424 ethnicity." (*Jacob, cited per preference*)

425

As such, Indigenous and Traditional Knowledge is included in this report to the best of our ability. Information sources include ethnographic interviews, public testimony, and literature sources focused on the social science of traditional knowledge. (Table #) In addition, information and recommendations provided by the Arctic-Yukon-Kuskokwim Working Group, which has local members from each of the Arctic, Yukon, and Kuskokwim regions of Alaska, provided another opportunity to document traditional knowledge which will be presented later in this report.


- 432
- 433

# 434 **Productivity Trends**

## 435 North Pacific

The North Pacific Anadromous Fish Commission (NPAFC) collates Pacific salmon commercial catch 436 data and the number of hatchery salmon released into the North Pacific Ocean each year. These data 437 come from Canada, Japan, Korea, Russia and the United States, where Alaska salmon harvest is 438 separated from Washington, Oregon and California. (Note that catch numbers are an imperfect metric 439 of salmon abundance, because they also depend on fishing effort.) What is noticeable within this near 440 100-year time series is the number of Pacific salmon harvested each year by all five nations is at 441 historically high levels since the 1990s (Figure 1). Peak commercial catches of 600 million Pacific 442 salmon by all five nations occurred several times during the mid-2000s to present. Hatchery salmon 443 releases began during the 1950s, but the numbers of salmon released into the North Pacific Ocean 444 increased during the 1970s and has peaked at around 5 billion salmon each year from 1987 to present 445 (Figure 2). Overall, Japan, the United States, and Russia release the highest number of hatchery salmon 446 447 into the ocean each year when compared with Canada and Korea.


- 448
- 449
- 450
- 451



452 453 Figure 1. Commercial catch (millions of fish; 1925 to 2022) of Pacific salmon within nearshore and

rivers for Canada (light blue), Japan (dark blue), Korea (green), Russia (purple), and the United States(hlue/green)

- 455 (blue/green).
- 456
- 457
- . -
- 458
- 459



Draft Alaska Salmon Research Task Force Report - April 1, 2024

460

Figure 2. Hatchery releases of Pacific salmon (billions of fish) by Washington, Oregon, California
(WOCI; dark blue), Canada (yellow), Alaska (green), Russia (light blue), Japan (orange), and Korea
(grey). within nearshore and rivers for Canada (light blue), Japan (dark blue), Korea (green), Russia
(purple), and the United States (blue/green).

# 465 State of Alaska

The time series of salmon harvest by species and region are shown in Figures (3–7). The time period 466 shown (1959–2022) illustrates the variability in harvest during since statehood and includes the higher 467 production period starting 1976/77, the onset of increased releases of salmon from hatcheries, and 468 recent declines in productivity for some species and stocks. Chinook salmon commercial harvest 469 470 averaged around 600 thousand from the late 1950s to mid-1970s, then increased to roughly 800 thousand during the early 1980s and has since gradually declined to around 250 thousand. In general, 471 472 the downward trend in Chinook salmon commercial harvest since the mid-1980s includes dramatic 473 declines within the AYK, Central and Southeast regions and high variability in harvest within the Westward region starting around 2007. Chum salmon commercial harvest was around 5 million from 474 the late 1950s to 1980 but then increased to around 18 million through 2018. The commercial harvest 475 476 of chum salmon has recently declined to levels seen during the mid-1980s, and subsistence harvest to lowest levels on record, with the AYK region having the largest decline. Sockeye salmon commercial 477

harvest in Alaska has varied between 10 to 60 million. Commercial harvest of sockeye salmon has
been strong during recent years with proportionally higher harvest coming from Bristol Bay in the
Central Region. There is also large regional variability in sockeye salmon harvest with recent declines
in some regions and record harvests in others.

482

483 Pink salmon are characterized by considerable variability and commercial harvest has varied between 50 to 210 million annually since the mid-1980s. The lowest catches occurred during 1959 and 1978 484 and the highest catches occurred during 2013 and 2015. Much of the pink salmon harvest occurs within 485 the Gulf of Alaska regions including Southeast Alaska, Prince William Sound and Kodiak. Harvest 486 within the Bering Sea is considerably lower in comparison. Pink salmon are caught in sport and 487 subsistence fisheries, but those numbers are small compared to commercial harvest. Coho salmon 488 commercial harvest has ranged between 2.5 to nearly 10 million, following the 1976/77 shift with the 489 peak commercial harvest occurring during the early 1990s and the lowest commercial harvest 490 occurring during 2020. Much of the recent decline in coho salmon harvest is within the Southeast and 491 AYK regions. 492

493






Figure 3. Number (thousands) of Chinook salmon harvested in Alaska (1959–2022). Series include
total commercial harvest (main panel), commercial harvest for ADF&G Commercial Fisheries
Regions (lower panels), and sport and subsistence harvest through 2021 (side panels). Note change in
scale of y-axis. Data source: ADF&G, adapted from NPAFC (2023).

500





Figure 4. Number (millions) of chum salmon harvested in Alaska (1959–2022). Series include total
commercial harvest (main panel), commercial harvest for ADF&G Commercial Fisheries Regions
(lower panels), and sport and subsistence harvest through 2021 (side panels). Note change in scale of
y-axis. Data source: ADF&G, adapted from NPAFC (2023).

508



509

Figure 5. Number (millions) of sockeye salmon harvested in Alaska (1959–2022). Series include total 512 commercial harvest (main panel), commercial harvest for ADF&G Commercial Fisheries Regions 513 (lower panels), and sport and subsistence harvest through 2021 (side panels). Note change in scale of 514 y-axis. Data source: ADF&G, adapted from NPAFC (2023). 515

- 516
- 517




Figure 6. Number (millions) of pink salmon harvested in Alaska (1959-2022) as reported annually to
NPAFC. Series include total commercial harvest (main panel), commercial harvest for ADF&G
Commercial Fisheries Regions (lower panels), and sport and subsistence harvest through 2021 (side
panels). Note change in scale of y-axis. Data source: ADF&G, adapted from NPAFC (2023).

- 527
- 528




Figure 7. Number (millions) of coho salmon harvested in Alaska (1959-2022). Series include total
commercial harvest (main panel), commercial harvest for ADF&G Commercial Fisheries Regions
(lower panels), and sport and subsistence harvest through 2021 (side panels). Note change in scale of
y-axis. Data source: ADF&G, adapted from NPAFC (2023).

- 535 See Appendices 1-7
- 536
- 537
- 538

# 539 Arctic Yukon Kuskokwim Working Group Report

#### 540 EXECUTIVE SUMMARY

1. The most pronounced declines of chum salmon and Chinook salmon in Alaska have occurred in the 541 Arctic-Yukon-Kuskokwim (AYK) region, a vast and remote area dominated by the Yukon and 542 Kuskokwim rivers, and including habitat throughout Norton Sound and into the western Arctic. 543 Communities throughout this region have been intimately dependent on salmon for subsistence and 544 culture for millennia and are currently suffering immense hardship due to restrictions on fishing 545 intended to protect dwindling AYK salmon populations. The AYK Working Group (WG) of the 546 Alaska Salmon Research Task Force (AKSRTF) included 42 volunteer members (no volunteers were 547 excluded) representing a wide variety of knowledge holders, from salmon harvesters and processors 548 to agency and academic scientists, with extensive experience with salmon in this region. The goal of 549 550 the WG activities was to develop a prioritized list of research needs for understanding the causes of declines AYK and Chinook 551 recent in chum salmon salmon populations. 552

2. The AYK-WG held virtual meetings twice-monthly in the autumn of 2023 to develop a process for 553 assembling a list of potential concerns contributing to recent declines in AYK salmon and for 554 translating these into a set of prioritized research themes. The WG adopted the framework developed 555 556 by the AKSRTF that organized potential research themes around the life cycles of salmon. A variety 557 of criteria were used by WG members to establish the prioritization, including whether the research could provide new insights in the short-term and whether the knowledge derived from this research 558 had the potential to be actionable in fisheries management. There was not a consensus across diverse 559 WG members regarding the most likely causes of salmon declines, and priorities described here 560 561 represent research that a majority of WG members felt would help support resilient salmon populations. 562

3. The two top priority research themes identified by the WG were to better understand impacts of
marine harvest on AYK salmon and changes in the quantity and quality of marine food for AYK
salmon. For marine harvest, emphasis was placed on improving: 1) the sampling of marine fisheries
(state, federal, and foreign) for incidental harvest of salmon, 2) methods for identifying the stock of

origin of chum and Chinook salmon caught in these fisheries, and 3) the escapement monitoring needed to quantify the consequences for salmon populations throughout the AYK region. Research to improve understanding of changes in the quantity and quality of food for AYK salmon in marine environments included understanding climate-related changes to salmon food resources, as well as the impacts of hatchery-origin pink salmon and chum salmon, and high abundance sockeye salmon from Bristol Bay, on feeding, growth, and survival of wild AYK chum salmon and Chinook salmon.

573

4. Additional top priority research themes included understanding changes in the health of migrating 574 575 and spawning adult salmon and how climate change is affecting freshwater and marine 576 ecosystems. The health of migrating and spawning adults theme included particular emphasis on 577 understanding the interacting effects of reduced body size and physiological condition, changes in disease prevalence and parasite loads, and changes in the hydrology and water temperatures 578 579 experienced by adult fish in freshwater habitats. These interacting factors are expected to affect both 580 the survival of fish during their spawning migrations and their fitness once they have reached spawning grounds, yet these effects on population dynamics of AYK salmon are currently not clear. The climate 581 change theme emphasized interactions between changing physical features of freshwater and marine 582 ecosystems (e.g., hydrology, water temperature) and nearly all other themes described in this report. 583 Additional research topics that were also highlighted included understanding how climate change 584 affects the incidence of harmful algae blooms, changes in sea ice, and melting permafrost - all of 585 which could have important impacts on AYK salmon ecosystems. 586

587

588 6. The three lowest priority research themes were: 1) changing freshwater conditions (beyond 589 effects on spawner health), 2) historical freshwater harvest, and 3) marine and freshwater 590 predators. While considered important by many WG participants, these were not considered as high 591 a priority as the themes mentioned above. Within these research themes were topics such as estimating 592 the effects of increases in freshwater predators because of reduced harvest on these species, changes 593 in flow regimes affecting egg incubation and juvenile rearing, and increased predation by expanding 594 marine mammal populations.

595 8. The WG also identified critical activities to improve coordination of research across AYK and to 596 develop more equitable opportunities for people of this region to have more meaningful engagement 597 in the fishery science, management, and regulatory processes. In particular, the WG was united in its support to develop formal approaches to integrate Indigenous and western ways-of-knowing in the 598 management process. There was also widespread emphasis on research that explored the efficacy of 599 alternative management approaches for achieving sustainability, given the inevitable uncertainties in 600 601 data and understanding. Other emphases for improving research coordination across the AYK included developing robust and publicly accessible databases that could incorporate data collected by 602 communities, agencies, and academic institutions. Last, the WG believes that more emphasis should 603 be placed on synthetic research approaches, such as life-cycle modeling, that would provide the 604 platform for better synthesis of research that is widely spread across geography, time, and life stages 605 of AYK salmon. 606

#### 608 1. Overview of the charge to the Working Group as Part of Act

Congress passed the Alaska Salmon Research Task Force Act during December 2022 to form an 609 610 Alaska Salmon Research Task Force (AKSRTF) to characterize trends in the productivity and abundance of Pacific salmon in Alaska, identify and prioritize research needs with respect to 611 understanding increased variability or decline in Pacific salmon returns to Alaska, and to establish 612 a coordinated research strategy to address salmon returns that are in decline or experiencing increased 613 614 variability. One requirement within the Act was for the AKSRTF to establish a work group (by July 2023) focused specifically on the research needs associated with salmon returns in the Arctic-Yukon-615 616 Kuskokwim (AYK) regions of western Alaska.

# 617 2. Overview of the AYK region and its salmon

The Arctic-Yukon-Kuskokwim (AYK) region of western Alaska is located north of Bristol Bay and is dominated by the watersheds of the Kuskokwim and Yukon rivers which drain into the Eastern Bering Sea. The region also includes smaller rivers draining to Norton Sound and Kotzebue Sound, and extends into the Alaska Arctic where rivers drain to the Chukchi and Beaufort seas. The headwaters of the Yukon River extend into Canada where approximately 40% of the watershed is located (Figure 1).


Five species of anadromous Pacific salmon spawn in AYK rivers, though the most important species for fisheries and the cultures of the people living in the region are chum salmon and Chinook salmon. In the Alaska Arctic, north of Kotzebue Sound, salmon are not historically the preferred subsistence fish (although attitudes may be in transition with expanding salmon distributions); hence, research on salmon in this vast area remains low.

The AYK region includes the traditional homelands of several Alaska Native groups, with Inupiat and Yup'ik people typically living in coastal regions, and Athabascan people living in upriver regions for millennia (Langdon 2002, Wolfe and Spaeder 2009). The cultures, economies and nutritional foundation of tribes throughout the AYK region are intimately woven with salmon, particularly chum salmon and Chinook salmon. Sockeye salmon, pink salmon, and coho salmon are also harvested, along with a variety of freshwater resident species such as northern pike, burbot, and whitefish.

Fisheries in the AYK region are primarily subsistence fisheries. The region has supported in-river 635 commercial fisheries for Chinook salmon and chum salmon over the last century, and some sport 636 637 fisheries exist as well. Due to severe salmon population declines over the last two decades, there have been increasing restrictions on commercial, sport, and subsistence fisheries to protect spawning 638 populations of these species in all rivers. Chum salmon populations in the AYK region showed 639 severely depressed populations in 1999-2001, and again in 2020 - 2023. Chinook salmon populations 640 in the region have shown steady declines since the early 2000s. Closed and restricted salmon fishing 641 infringes on the opportunity for rural residents to maintain a subsistence way of life (Alaska National 642 Interest Lands Conservation Act [ANILCA] of 1980, 16 U.S.C. § 3101–3233) and opportunity to pass 643 down subsistence culture to younger generations. 644

The vast watersheds of rivers that drain the AYK region remain largely undeveloped and remote, which presents serious impediments to western science approaches to understanding the ecology of the watersheds and their salmon. Management of fishery resources is also hampered by these geographical challenges because of the inevitably high levels of uncertainty in stock assessments of fish populations, and the widely dispersed nature of fishing which makes it difficult to monitor.

The cultures, economies, and food security of people who live throughout the AYK region are being seriously impacted by declining salmon returns and the associated restrictions on fisheries. Climate change is also producing new challenges for people living in the AYK region. These changes are affecting the cultures and sustainability of AYK communities that primarily lead salmon-centered subsistence lifestyles. The AYK Working Group emphasizes the urgency of the need to take action to maintain vibrant and sustainable communities that are robust to the inevitable changes in the salmon resources that form the foundation of these communities.



658

659

Figure 1. Map of Alaska with the major watersheds and communities (black dots). The Arctic-Yukon-Kuskokwim region is located north of Bristol Bay north through Norton and Kotzebue Sounds into the Alaska Arctic. Watershed data available from the United States Geological Survey (<u>https://www.sciencebase.gov/catalog/item/5a1632bae4b09fc93dd1721f</u>). Community locations provided by the U.S. Census Bureau.

# 665 Background

The Alaska Salmon Research Task Force Act identified the need to convene an AYK Working Group (hereafter AYK WG), to address the research needs related to salmon declines in the Arctic-Yukon-Kuskokwim region. Fifteen members of the Task Force who had knowledge and expertise for this region volunteered to serve on this AYK WG. However, it was acknowledged that the information needs and expertise for the AYK WG should be far broader than those represented by Task Force

671 members alone. Nominations for public members of the working group (non-task force members) 672 were solicited in July 2023. All nominated public members who agreed to serve on the AYK WG were 673 accepted as working group members. In total, 42 individuals agreed to serve on the AYK WG, 674 including 15 Task Force members and 27 public members (Appendix 1). The AKSRTF appointed 675 Katie Howard (Alaska Department of Fish & Game) to lead the AYK WG on behalf of the task force, 676 and Daniel Schindler (University of Washington) was elected by public members of the AYK WG to 677 co-chair the working group.

A strength of the convened AYK WG was the diversity of perspectives represented by the 42 members (Appendix #). These members are knowledge holders from Kuskokwim Bay, Kuskokwim River, lower, middle and upper parts of the Yukon River, Norton Sound, North Slope, academia, federal and state management agencies, environmental and fisheries non-profits, tribal organizations, inter-tribal fish commissions, and the commercial fishing industry.

Many AYK WG members repeatedly expressed the value and need for voices at the table to equitably 683 include western science and local and Indigenous knowledge holder insights and expertise. Due to the 684 short timelines stipulated for this endeavor, the large geographic scope of the AYK region, and limited 685 686 resources to support in-person communications, the work and collaboration among working group 687 members was heavily reliant on technology and digital forms of communications (e.g., video conferences, Excel and Word documents through cloud sharing, and email). It should be 688 acknowledged that technology-heavy communications create their own inequities, particularly for a 689 690 region where internet access can be limited and computer-based information sharing may be somewhat 691 foreign to some members, particularly elders. This was a significant challenge for the AYK WG and it is recommended that, if another group is similarly convened in the future, resources and planning 692 693 be used to allow for in-person engagement as a more equitable means of communication and 694 collaboration among knowledge holders.

# 695 Working Group Process

The AYK WG met by virtual teleconference about every 2 weeks from September through December
2023 to discuss potential explanations for AYK salmon declines, other research needs in the region,
and AYK-specific research priorities. Initial meetings were an open discussion for WG members to

express ideas and concerns, which guided the co-chairs in developing structures and agendas forsubsequent meetings.

701 The AYK WG adopted the conceptual framework developed by the AKSRTF for the possible explanations of AYK salmon decline for the following themes: 1) spawner health; 2) freshwater 702 harvest; 3) freshwater predators; 4) marine predators; 5) freshwater conditions for eggs and juvenile 703 rearing and migration; 6) marine food limitation; 7) climate change; and 8) marine harvest (focused 704 705 primarily on bycatch in federal fisheries, harvest and interceptions in state-managed salmon fisheries, and illegal, unreported and unregulated foreign fisheries). It was clear from initial scoping meetings 706 707 that AYK WG members wished to express concerns and suggest research priorities that were not 708 strictly evaluating reasons for AYK salmon declines, so additional categories and discussion topics 709 were created, and research priorities were not confined to evidence for salmon declines.

Shared spreadsheets allowed AYK WG members to formalize questions and hypotheses they had within each of these different research themes. AYK WG members who found access to the shared spreadsheets challenging were encouraged to reach out to the co-chairs to ensure their input was captured on the spreadsheets by working with other AYK WG members. AYK WG members with better technology access and comfort were encouraged to work with those who found these forms of communication challenging, so that as many perspectives as possible were represented.

716 After discussing research questions and hypotheses that fell under each of the research themes 717 specified by the AKSRTF, the AYK WG discussed how best to prioritize these research themes The AYK WG agreed that each member should have the independence to assess each research priority 718 based on their own knowledge base and criteria. Examples of criteria that AYK WG members cited in 719 720 their assessments were: a) whether reducing a specific scientific uncertainty would lead to actionable 721 management changes, b) whether reducing a scientific uncertainty was a short-term or long-term goal, 722 c) whether research on a specific topic would improve synthesis of AYK salmon ecology, d) whether research on a specific topic would improve community engagement and knowledge sharing, e) would 723 724 research lead to more holistic science such as what Tribes currently use rather than the typically narrow 725 focus of western science, f) would research on a topic produce knowledge that was immediately 726 applicable, g) whether research would provide immediate information with high potential benefit, h) whether the research would benefit management of salmon ecosystems in the Arctic region of the 727

AYK, and i) whether the research would produce knowledge that would be applicable beyond theAYK region.

730 The primary goal of the AYK WG was to prioritize research needs where research could fill knowledge gaps in understanding recent AYK salmon declines. To establish these priorities, members of the AYK 731 WG were presented with the list of research questions and hypotheses developed by the group 732 (Appendix 2). As a way for individuals to express their own perspectives about how to prioritize 733 734 potential research, each AYK WG member was asked to assign a total of 20 points across all potential hypotheses or questions, with the constraint that the maximum score they could assign to an individual 735 736 question or hypothesis was 10. We received scores from the majority of AYK WG members (29 of 737 42) and these scores were summed across individual AYK WG members and across hypotheses and 738 questions to provide weights to each of the eight AKSRTF research themes. We also summarized scores by tallying the number of AYK WG members who assigned a score of at least 1 to any of the 739 740 questions or hypotheses within each of the eight AKSRTF research themes. The intention of this 741 exercise was not to treat individual hypotheses or questions as competing alternative explanations for the recent AYK salmon declines. Rather, it was a process for summarizing the variety of perspectives 742 on what were the highest priority research themes across the entire spectrum of beliefs held by 743 members of the AYK WG. Last, we ranked individual questions and hypotheses based on the total 744 number of points assigned to them by all the AYK WG members who participated in the survey. All 745 hypotheses and questions and their associated scores are provided in Appendix 2. Here we highlight 746 the top nine as these attracted the majority of attention from across the AYK WG. 747

Subsequent to AYK WG discussions, co-chairs drafted a report on behalf of the WG, with a first draft
provided to the AYK WG for review by March 11, 2024. Co-chairs revised the report based on AYK
WG member feedback and delivered a final report to the Task Force on March 21, 2024.


# 751 **3.** Overview of AYK WG general priorities for future research needs on AYK salmon

Results from the survey of AYK WG members about priorities for future research provided one mechanism for summarizing across the range of perspectives within the AYK WG. AYK WG members developed a set of questions that could be explored by future research to support resilient AYK chum salmon and Chinook salmon populations. The two specific questions that received the

highest scores across AYK WG members both deal with the availability and quality of marine food
for AYK salmon (Figure 2, purple bars). The top specific hypothesis focused on competition between
AYK salmon and hatchery pink and chum salmon, and with high abundances of sockeye salmon from
Bristol Bay. The second most popular question was concerned with whether changes in climate and
marine food webs may be limiting growth and survival of AYK salmon.

Four of the top nine questions dealt with bycatch and interceptions of AYK fish in federal, state, and foreign fisheries (Figure 2, blue bars). Three of these hypotheses focused on understanding the biological impacts of these marine harvests on AYK salmon populations. One of these questions was concerned with increasing the analytical resolution to distinguish the stock of origin of AYK fish captured in marine fisheries.

Climate change, particularly as an interacting stressor on all other drivers of change in AYK salmon populations, was one of the nine top questions discussed by the AYK WG (Figure 2, yellow bar). Two of the top nine specific hypotheses and questions focused on the implications of changes in spawner health and quality for AYK salmon populations (Figure 2, red bars). All other specific questions received less than 20 points from the scoring exercise performed by the AYK WG (Appendix 2).




771

Figure 2. Total assigned scores to the top nine individual research questions considered by the
AYK WG, colored by the associated research theme. All other questions or hypotheses received
less than 20 total points. Hypotheses and questions are colored according to: 1) Marine Food
Limitation theme (purple), 2) Marine Harvest theme (blue), 3) Climate Change theme (yellow),
and 4) Spawner Health theme (red).

777 When summing AYK WG scores across all questions within each research theme, marine harvest, 778 marine food limitation, climate change and spawner health were the top research priorities (Figure 3, 779 black bars). The highest priority research theme was to better quantify the impacts of marine harvest on AYK chum salmon and Chinook salmon stocks (Figure 3, black bars). The second highest research 780 priority identified by this approach was to understand the consequences of changes in marine food 781 limitation for AYK salmon growth and survival. The third highest priority research theme was to 782 understand the effects of a variety of stressors affecting the health and quality of migrating and 783 spawning adult salmon and how these translate into population dynamics. Understanding climate 784 change, particularly through interactions with other stressors of AYK salmon, was the fourth highest 785 priority research theme identified by the AYK WG scoring exercise. We note, however, that there 786 were elements of climate change in research priorities that were included under other themes. The 787 remaining four research themes all received some level of support for prioritization in the following 788 descending order: freshwater conditions that affect egg incubation and juvenile growth and survival, 789 freshwater harvest, marine predators, and freshwater predators. These last four categories received 790 791 only about 22% of the total weight assigned by the AYK WG to the eight research themes. The top 792 four research themes collectively received 78% of the total weight of scores from the AYK WG (Figure 3). 793

794 As a complementary way to summarize research priorities across the diverse members of the AYK WG, we also tallied the number of AYK WG members who assigned any degree of weight to each of 795 the AKSRTF research themes (Figure 3, gray bars). This method of summarizing perspectives from 796 across the AYK WG generally reinforced the priority list established by the weighted scoring method, 797 except for two notable differences. First, marine harvest and marine food limitation were ranked 798 equally as the two top research priorities. Spawner health and climate change were the next two 799 priorities and their order in the rankings did not change. Of the remaining four research themes, marine 800 predators received more than twice as much weight via this second method of ranking themes 801 compared to the first method. However, the combined weight of the lowest priority four research 802 themes was still only about 32% compared to 68% for the four top research themes (Figure 3). In the 803 sections below we discuss each of these general research themes and the distribution of support for 804 805 individual questions or hypotheses that were aggregated under each theme.






Figure 3. Summary of the AYK Working Group research priorities organized by themes established by the AKSRTF. Black bars show the distribution of relative weights assigned by WG members to hypotheses or questions that were aligned within each theme. Values are relative weights that are proportional to the sums across scores assigned to individual hypotheses and questions under each research theme. Gray bars are weights that are proportional to the number of WG members who expressed any level of weight to each of the research themes.

In the section below, in decreasing order of priority by research theme, we describe the details of individual questions or hypotheses that could guide research within each of the general themes.

815 A) Marine Harvest

Support for research quantifying and mitigating marine harvest impacts on AYK salmon was a dominant topic throughout AYK WG discussions. Within the theme of marine harvest, there were relatively high scores prioritizing research that further quantified the number of AYK salmon harvested in state, federal, and foreign marine fisheries, and improved data and methods to compare these numbers to the abundances of AYK salmon stocks as a way to estimate the biological and social consequences of marine fishery catches. Emphasis was also placed on improving the resolution of

current techniques (i.e., genetic stock identification, GSI) for estimating stock-specific impact rates of 822 823 marine harvest on individual AYK salmon stocks. Overall, there was roughly equal emphasis placed 824 on understanding interceptions in state, federal and foreign fisheries. There was recognition of the amount of effort currently being applied towards rigorously quantifying Chinook and chum salmon 825 bycatch in federal marine fisheries, though weak stock resolution in current GSI and limited coverage 826 of Chinook and chum salmon escapements throughout the AYK region place limits on the desired 827 resolution of these efforts to understand their biological impacts. The scoring was similarly high for 828 more research to understand the biological implications of both chum salmon and Chinook salmon 829 catches in state and foreign fisheries. 830

831 B) Marine Food Limitation

Research on marine food limitation for AYK chum salmon and Chinook salmon was identified as 832 833 another top priority research theme by the WG. The highest score for all individual research questions and hypotheses was in this theme. The AYK WG expressed particular support for research 834 understanding competition between AYK salmon and increasing pink salmon, chum salmon, and 835 sockeye salmon abundance from other regions, and especially in consideration of hatchery-produced 836 837 competitors. This research theme also received relatively high scores for research to determine 838 whether there were climate-induced changes in the quantity and quality of marine food for AYK salmon in the Bering Sea, and whether changes in nearshore habitat conditions were reducing the 839 survival of AYK smolts upon their migration to the nearshore ocean. 840

841 C) Spawner Health

842 Research to understand causes and consequences of adult salmon health status while migrating and 843 spawning was a relatively complex theme as WG members identified many dimensions of this problem that affect the survival of fish as they migrate from the ocean to spawning grounds, and their 844 845 subsequent success on the spawning grounds. In particular, emphasis was placed on understanding the 846 effects of changing climate and ocean conditions on en route and pre-spawn mortality of adult fish. 847 Whether changes in the incidence of parasites and diseases (such as with Ichthyophonous), and nutritional health exacerbated these effects on reproductive health and stock productivity. Warming 848 849 river temperatures with climate change was also identified as a poorly understood interactor with other

factors that affect the health of adult fish in freshwater habitats. Other concerns within this broad theme 850 851 emphasized quantification of thresholds or conditions where genetic population sizes were so low that 852 there was increased risk of extirpation of individual sub-populations in AYK watersheds, and quantifying how widespread these thresholds were surpassed at present. Also identified as an 853 important research activity was to systematically review existing in-river stock assessments to 854 determine whether there was adequate precision and accuracy to quantify *en route* mortality in the 855 large AYK rivers, particularly in the Yukon River where there is an international commitment to meet 856 an escapement goal for Chinook and chum salmon that spawn in Canadian components of the 857 watershed. Last, there was concern that little was known or acknowledged about the effects of handling 858 fish for research purposes on their stress levels and eventual success on the spawning grounds. 859

860 D) Climate Change

The AYK WG widely emphasized that the consequence of climate change in freshwater and marine 861 habitats was an overriding priority because climate change is likely modifying or amplifying nearly 862 all other stressors identified in other research themes. Beyond climate change as an amplifier of other 863 stressors, the AYK WG identified other research questions to be pursued. These include understanding 864 865 linkages between climate stressors in the ocean and in freshwater habitats, whether climate-driven 866 changes in sea ice have affected plankton phenology and potential mismatches with AYK salmon, and whether climate change has increased the incidence and intensity of harmful algal blooms that affect 867 juvenile salmon growth and survival in the ocean. There was also interest in understanding whether 868 the expansion of anadromous salmon into the Arctic was affecting the ecology of resident fishes in 869 870 rivers that historically did not have salmon in them, though this question did not receive any support in the scoring exercise. 871

872 E) Freshwater Conditions

Changing freshwater habitat conditions that affect egg incubation success, and juvenile salmon rearing and migration conditions, rated to be of intermediate priority, focused on issues related to whether changes in watershed habitat productivity and capacity were reducing the fitness and abundance of salmon smolts leaving AYK rivers. Related to this question was whether changes in hydrology may be affecting the complexity, connectivity and geomorphology of freshwater habitats in ways that affect

freshwater food webs that support juvenile salmon growth, and hydrologic effects on egg incubation conditions. Additional questions were focused on whether climate driven effects on floods, spring ice breakup, thermal regimes, and permafrost loss were contributing to a degradation of freshwater habitat for juvenile salmon. Other considerations included asking whether the expansion of beavers was altering salmon habitat in substantial ways, and whether the declines of marine-derived nutrients from declines in abundant species (e.g., pink salmon and chum salmon) were reducing the productivity of freshwater habitats.

### 885 F) Freshwater Harvest

One of the lower priority research themes was associated with current and legacy effects of harvest in 886 freshwater fisheries. The research question that received the highest score within this theme was 887 focused on understanding whether the current escapement goals were still valid given the observed 888 889 changes in the environment and the different constraints on population productivity that are expressed in freshwater versus marine ecosystems. Other important concerns were focused on understanding 890 whether there were legacy effects of historical freshwater fisheries, particularly commercial fisheries, 891 on the current demographic structure of AYK salmon populations, which have shown a pronounced 892 893 decline in average body size and age-at-maturity for Chinook salmon in particular. Other questions 894 were focused on whether large fish that become entangled in but drop out from small mesh gill nets actually survive to reproduce, as is typically assumed. While it is often assumed that the consequences 895 of managing to the higher end of an escapement goal range has the same biological consequences as 896 897 managing to achieve the bottom end of an escapement goal range, several AYK WG members believed 898 this assumption needed to be more thoroughly explored. Further, there was interest to explore the consequences for harvest on weak stocks when they mingle with dominant or highly abundant stocks 899 900 within the same river, and whether there were ways to develop more stock specificity in freshwater fisheries. Last, because of the recent proliferation of early-maturing males ("jacks") in Chinook 901 salmon, some AYK WG members believed that research should explore whether there were potential 902 903 negative consequences of harvesting these small individuals from a population, though this question 904 also received no support in the scoring exercise.

905 G) Marine and Freshwater Predators

906 The two research themes determined to be the lowest research priority by the AYK WG were 907 associated with changes in the predation rates on salmon in marine and freshwater habitats. In marine 908 habitats there was concern that increasing apex predators such as resident killer whales, seals, and salmon sharks may be reducing marine survival of AYK salmon and generating new evolutionary 909 pressures on large fish that may be related to declining body size in AYK salmon, particularly Chinook 910 911 salmon. In freshwater habitats, there was concern that piscivores may be increasing because of declining harvest of resident species (such as northern pike) that could be translating into increased 912 predation rates on juvenile salmon. Other questions were focused on whether the proliferation of 913 beavers was enhancing predation on juvenile salmon through changes in habitat that facilitate 914 predators such as pike and juvenile coho salmon, and whether climate-induced changes in freshwater 915 habitats were increasing vulnerability of juvenile salmon to freshwater predators. 916

# 917 Other research needs and priorities for improving science and management of AYK salmon

The AYK WG also identified critical activities to improve coordination of research across the AYK 918 919 and to develop more equitable opportunities for people of this region to have more meaningful 920 engagement in the fishery science, management, and regulatory processes. These concerns 921 complement similar existing efforts to prioritize research on Chinook salmon and chum salmon in the 922 AYK region by the Arctic-Yukon-Kuskokwim Sustainable Salmon Initiative (Schindler et al. 2013). In addition to discussing the eight research themes of the AKSRTF as they relate to understanding the 923 recent declines in AYK chum salmon and Chinook salmon, the AYK WG discussed several other 924 925 research needs for improving the science and management of AYK salmon. These additional themes 926 are listed and described briefly below.

927 A) Knowledge integration and alternative approaches towards fishery management

The AYK WG was united in its support to develop and explore the efficacy of formal approaches to integrate traditional, Indigenous, and western ways-of-knowing in the science and management process. At present, there is little integration of alternative sources of knowledge and perspectives about how the AYK salmon ecosystems function despite a considerable amount of knowledge held by the individuals who live throughout the region. The challenge is to find ways to weave different types of knowledge and different perspectives into a coherent framework for informing management and

conservation efforts. The AYK WG believes that exploring alternative ways of accomplishing this
integration is itself a research priority that has the distinct potential to both improve understanding of
AYK salmon and their ecosystems, and to improve integration of a variety of knowledge types to
better inform the management process.

B) Need for synthetic life-cycle approaches to improve integration

939 As AYK salmon complete their life cycles, their biology integrates across a wide variety of habitats, from freshwater streams, lakes and wetlands, to estuaries and the coastal ocean, to the Bering Sea and 940 941 Gulf of Alaska. Understanding how changes in management and the environment affect salmon 942 populations is seriously challenged by this complexity. Most research on AYK salmon has focused on individual life stages in specific habitats which, as the AYK WG expressed, has hampered scientific 943 progress towards developing a holistic understanding of the drivers of population dynamics. The AYK 944 945 WG believes that more emphasis should be placed on synthetic research approaches, such as life-cycle modeling, that would better integrate across geography, time, and life stages of AYK salmon. 946 Additional emphasis would be placed on integrating life cycle modeling, field research, traditional 947 knowledge, and management activities into a coherent framework to enhance knowledge generation 948 949 and improve management outcomes.

950

#### 951 C) Management under uncertainty

952 Fisheries management is an imperfect process that makes decisions despite incomplete understanding of ecosystems and fish populations, including their responses to management actions. These 953 954 challenges are especially acute in the AYK region of Alaska because ecosystems are so vast, remote, 955 and heterogeneous which hinders comprehensive monitoring needed to reduce key uncertainties in 956 data and models used in management. People who rely on salmon fisheries also hold a wide range of values and goals for assessing the success or failure of management decisions. Thus, there was 957 widespread support for research that evaluates trade-offs associated with various management 958 959 strategies for achieving a variety of goals for stakeholders, given the inevitable uncertainties in data and understanding. Such research would use combinations of simulation modeling and community 960 961 engagement to co-develop projects that explicitly quantify trade-offs associated with the potential risks

and rewards of alternative management strategies given uncertainties in how ecosystems function andhow they may change in the future.

964 D) Database coordination

A final emphasis for improving research coordination across the AYK included developing robust and 965 publicly accessible databases that could incorporate data collected by communities, agencies, and 966 academic institutions. This is a non-trivial task given the heterogeneity in types of data relevant to 967 salmon populations and aquatic ecosystems throughout AYK. Research is needed to develop ways to 968 both capture the vast amounts of historical data on AYK salmon and ecosystems, and to seamlessly 969 add new data streams as further research is pursued. Both serious design considerations and substantial 970 funding will be needed to accomplish this ambitious goal, but the invaluable payoff would be increased 971 transparency in research and management, and greater leveraging of data to improve ecological 972 973 understanding. Such an effort would need to proactively plan for the sustainability of such a database, and to make specific policies to acknowledge and respect data sovereignty. 974

# 975 **References**

976 Langdon, S.J. 2002. The Native People of Alaska. Greatland Graphics, Anchorage, Alaska.

977 Schindler, D., C. Krueger, P. Bisson, M. Bradford, B. Clark, J. Conitz, K. Howard, M. Jones, J.

978 Murphy, K. Myers, M. Scheuerell, E. Volk, and J. Winton. 2013. Arctic-Yukon-Kuskokwim

979 Chinook Salmon Research Action Plan: Evidence of Decline of Chinook Salmon Populations and

Recommendations for Future Research. Prepared for the AYK Sustainable Salmon Initiative
(Anchorage, AK). v + 70pp.

982 Wolfe and Spaeder. 2009. People and salmon of the Yukon and Kuskokwim drainages and Norton

983 Sound in Alaska: fishery harvests, culture changes, and local knowledge systems. American

984 Fisheries Society Symposium 70: 349-379.

985 See Appendices 8 - 9

987	Appendix 1
988	Engrossed in Senate (12/14/2022)
989	117th CONGRESS 2d Session
990	S. 3429
991	AN ACT
992 993	To establish an Alaska Salmon Research Task Force. Be it enacted by the Senate and House of Representatives of the United States of America in Congress assembled,
994	SECTION 1. SHORT TITLE.
995	This Act may be cited as the "Alaska Salmon Research Task Force Act".
996	SEC. 2. PURPOSES.
997	The purposes of this Act are—
998 999	(1) to ensure that Pacific salmon trends in Alaska regarding productivity and abundance are characterized and that research needs are identified;
1000	(2) to prioritize scientific research needs for Pacific salmon in Alaska;
1001 1002	(3) to address the increased variability or decline in Pacific salmon returns in Alaska by creating a coordinated salmon research strategy; and
1003	(4) to support collaboration and coordination for Pacific salmon conservation efforts in Alaska.
1004	SEC. 3. SENSE OF CONGRESS.
1005	It is the sense of Congress that—
1006 1007 1008 1009	(1) salmon are an essential part of Alaska's fisheries, including subsistence, commercial, and recreational uses, and there is an urgent need to better understand the freshwater and marine biology and ecology of salmon, a migratory species that crosses many borders, and for a coordinated salmon research strategy to address salmon returns that are in decline or experiencing increased variability;
1010	(2) salmon are an essential element for the well-being and health of Alaskans; and
1011	(3) there is a unique relationship between people of Indigenous heritage and the salmon they rely on

1013 SEC. 4. ALASKA SALMON RESEARCH TASK FORCE.

for subsistence and traditional and cultural practices.

- 1014 (a) In General.—Not later than 90 days after the date of enactment of this Act, the Secretary of
- 1015 Commerce, in consultation with the Governor of Alaska, shall convene an Alaska Salmon Research
- 1016 Task Force (referred to in this section as the "Research Task Force") to—
- 1017 (1) review existing Pacific salmon research in Alaska;
- 1018 (2) identify applied research needed to better understand the increased variability and declining salmon
- 1019 returns in some regions of Alaska; and
- 1020 (3) support sustainable salmon runs in Alaska.
- 1021 (b) Composition And Appointment.—
- 1022 (1) IN GENERAL.—The Research Task Force shall be composed of not fewer than 13 and not more 1023 than 19 members, who shall be appointed under paragraphs (2) and (3).
- 1024 (2) APPOINTMENT BY SECRETARY.—The Secretary of Commerce shall appoint members to the 1025 Research Task Force as follows:
- 1026 (A) One representative from each of the following:
- (i) The National Oceanic and Atmospheric Administration who is knowledgeable about salmon andsalmon research efforts in Alaska.
- 1029 (ii) The North Pacific Fishery Management Council.
- 1030 (iii) The United States section of the Pacific Salmon Commission.

(B) Not less than 2 and not more than 5 representatives from each of the following categories, at least
2 of whom shall represent Alaska Natives who possess personal knowledge of, and direct experience
with, subsistence uses in rural Alaska, to be appointed with due regard to differences in regional
perspectives and experience:

- (i) Residents of Alaska who possess personal knowledge of, and direct experience with, subsistenceuses in rural Alaska.
- 1037 (ii) Alaska fishing industry representatives throughout the salmon supply chain, including from—
- 1038 (I) directed commercial fishing;
- 1039 (II) recreational fishing;
- 1040 (III) charter fishing;
- 1041 (IV) seafood processors;

- 1042 (V) salmon prohibited species catch (bycatch) users; or
- 1043 (VI) hatcheries.

1044 (C) 5 representatives who are academic experts in salmon biology, salmon ecology (marine and
1045 freshwater), salmon habitat restoration and conservation, or comprehensive marine research planning
1046 in the North Pacific.

1047 (3) APPOINTMENT BY THE GOVERNOR OF ALASKA.—The Governor of Alaska shall appoint to
1048 the Research Task Force one representative from the State of Alaska who is knowledgeable about the
1049 State of Alaska's salmon research efforts.

- 1050 (c) Duties.—
- 1051 (1) REVIEW.—The Research Task Force shall—

1052 (A) conduct a review of Pacific salmon science relevant to understanding salmon returns in Alaska,
 1053 including an examination of—

1054 (i) traditional ecological knowledge of salmon populations and their ecosystems;

1055 (ii) marine carrying capacity and density dependent constraints, including an examination of interactions
 1056 with other salmon species, and with forage base in marine ecosystems;

- 1057 (iii) life-cycle and stage-specific mortality;
- 1058 (iv) genetic sampling and categorization of population structure within salmon species in Alaska;
- 1059 (v) methods for predicting run-timing and stock sizes;
- 1060 (vi) oceanographic models that provide insight into stock distribution, growth, and survival;
- 1061 (vii) freshwater, estuarine, and marine processes that affect survival of smolts;
- 1062 (viii) climate effects on freshwater and marine habitats;
- 1063 (ix) predator/prey interactions between salmon and marine mammals or other predators; and

1064 (x) salmon productivity trends in other regions, both domestic and international, that put Alaska salmon
 1065 populations in a broader geographic context; and

1066 (B) identify scientific research gaps in understanding the Pacific salmon life cycle in Alaska.

1067 (2) REPORT.—Not later than 1 year after the date the Research Task Force is convened, the Research 1068 Task Force shall submit to the Secretary of Commerce, the Committee on Commerce, Science, and

1069 Transportation of the Senate, the Committee on Environment and Public Works of the Senate, the

Subcommittee on Commerce, Justice, Science, and Related Agencies of the Committee on
Appropriations of the Senate, the Committee on Natural Resources of the House of Representatives,
the Subcommittee on Commerce, Justice, Science, and Related Agencies of the Committee on
Appropriations of the House of Representatives, and the Alaska State Legislature, and make publicly
available, a report—

1075 (A) describing the review conducted under paragraph (1); and

1076 (B) that includes—

- 1077 (i) recommendations on filling knowledge gaps that warrant further scientific inquiry; and
- 1078 (ii) findings from the reports of work groups submitted under subsection (d)(2)(C).
- 1079 (d) Administrative Matters.—

(1) CHAIRPERSON AND VICE CHAIRPERSON.—The Research Task Force shall select a Chairand Vice Chair by vote from among the members of the Research Task Force.

1082 (2) WORK GROUPS.—

1083 (A) IN GENERAL.—The Research Task Force—

1084 (i) not later than 30 days after the date of the establishment of the Research Task Force, shall establish

a work group focused specifically on the research needs associated with salmon returns in the AYK

1086 (Arctic-Yukon-Kuskokwim) regions of Western Alaska; and

- (ii) may establish additional regionally or stock focused work groups within the Research Task Force,as members determine appropriate.
- 1089 (B) COMPOSITION.—Each work group established under this subsection shall—
- 1090 (i) consist of not less than 5 individuals who—
- 1091 (I) are knowledgeable about the stock or region under consideration; and
- 1092 (II) need not be members of the Research Task Force; and

(ii) be balanced in terms of stakeholder representation, including commercial, recreational, and
 subsistence fisheries, as well as experts in statistical, biological, economic, social, or other scientific
 information as relevant to the work group's focus.

(C) REPORTS.—Not later than 9 months after the date the Research Task Force is convened, each
 work group established under this subsection shall submit a report with the work group's findings to
 the Research Task Force.

- 1099 (3) COMPENSATION.—Each member of the Research Task Force shall serve without compensation.
- 1100 (4) ADMINISTRATIVE SUPPORT.—The Secretary of Commerce shall provide such administrative
- 1101 support as is necessary for the Research Task Force and its work groups to carry out their duties, which
- 1102 may include support for virtual or in-person participation and travel expenses.
- (e) Federal Advisory Committee Act.—The Federal Advisory Committee Act (5 U.S.C. App.) shall
  not apply to the Research Task Force.
- 1105 SEC. 5. DEFINITION OF PACIFIC SALMON.
- 1106 In this Act, the term "Pacific salmon" means salmon that originates in Alaskan waters.
- 1107 Passed the Senate December 14, 2022.

# 1108 Appendix 2

# 1109 Alaska Salmon Research Task Force

- 1110 During the first meeting in June 2023, the AKSRTF members agreed on an approach/milestone
- timeline for the objectives provided in the ACT (Table 1).


# 1112 Table 1

Date	Approach/Milestones
June 28, 2023	Task Force Establishes Regional Teams (TF members; address ALL of ALASKA) to build EXISTING KNOWLEDGE and begin to discuss RESEARCH GAPS/NEEDS
July 27, 2023	Task Force Meeting; Establish the ARCTIC YUKON KUSKOKWIM WORKING GROUP
August 18, 2023	Begin to close our REVIEW OF EXISTING KNOWLEDGE; Final comments on DRAFT REPORT OUTLINE
October 2023	DRAFT document on EXISTING KNOWLEDGE – and initial list of RESEARCH GAPS/NEEDS (place on website for Public review)
November 2023	FINAL DRAFT EXISTING KNOWLEDGE; Continue to list RESEARCH GAPS and NEEDS (Public input)
April 2024	DRAFT FINAL REPORT; Begin one month Public Review
May 2024	FINAL DRAFT of REPORT
June 2024	FINAL REPORT

## 1113

Figure 8 illustrates the process the AKSTRF utilized to complete the objectives and prepare a Final 1114 1115 Report for the June 2024 deadline. First, the AKSRTF formed Regional Teams (Southeast, Central, Westward, and Arctic-Yukon-Kuskokwim) with the task of: 1) providing reference materials to 1116 understand salmon returns in Alaska; 2) identifying gaps in understanding the Pacific salmon life 1117 cycle in Alaska; and 3) recommendations on filling knowledge gaps that warrant further scientific 1118 inquiry. In addition, the AKSRTF formed an Arctic-Yukon-Kuskokwim (AYK) Working Group 1119 (WG) to focus on the research needs associated with salmon returns in the AYK regions of Western 1120 Alaska. The AYK WG merged the AYK Regional Team members (15) from the AKSRTF with 30 1121 other members from the Arctic, Yukon, and Kuskokwim region (see AYK WG Report section for 1122

1123 more details).



### 1124

In consideration of the ACT objectives, the AKSRTF scheduled bimonthly (every other month)
public meetings to discuss current progress and solicit feedback from the public. Public comment on
these tasks was solicited throughout the process during bimonthly meetings, online through our
AKSRTF web page and during the in person/hybrid meeting held in Anchorage, AK on November
14 and 15, 2023. The DRAFT report was also provided on the web page for Public comment and
input during mid-October to mid-November (prior to our November 14 and 15, 2023 in person
hybrid meeting in Anchorage, AK) and during the month of April 2024.

# 1132 **AKSRTF Meetings**

**Table 2** shows the dates and focus of the AKSRTF Meetings open to the public.

#### 1134

Date	Format and Primary Focus
7/27/2023	Virtual. Establish the ARCTIC YUKON KUSKOKWIM WORKING GROUP
9/19/23	Virtual. Discuss report outline and progress toward existing knowledge and research gaps
11/14-15/23	Hybrid in Anchorage, AK. Existing knowledge and gaps, Research needs and Public comment/testimony
1/25/24	Virtual. Report on progress toward DRAFT Report
3/27/24	Virtual. Report on DRAFT FINAL REPORT
5/22/24	Virtual. Comment on DRAFT FINAL REPORT and incorporation of public comments

1135

# 1137 Appendix 3

# 1138 Existing Knowledge Review

1139 Indigenous Knowledge/Traditional Knowledge

Source	Quotes
SCIA	18:38 Murkowski – "We don't fish. we don't eat." & "Multiple other
	factors, including management structures we have to look at"
SCIA	31:20 Ridley - "130,000 salmon, 1-3% prohibited species catch, but we
	know every salmon counts"
SCIA	37:20 Samuelson – "serve on all councils to represent leverage holistic
	approach, etc."
SCIA	44:10 Ulvi – "claim that Aleutian fisheries taking fish from western
	Alaska nothing new" since 1905 officially
SCIA	49:50 Winkelman – "Not just negatively affecting our culture and
	wellbeing, but our good health."
SCIA	1:05:00 Menadelook – "In my opinion, if we don't do anything we'll run
	out of salmon and/or marine mammals in 5-6 years."
SCIA	1:07:00 Menadelook – "I think that the main thing we need to do is change
	the way the salmon is managedmanaged towards maximum commercial
	yield instead of other yieldThat's what's killing salmon."
SCIA	:17:30 Ridley – "I had an Elder tell me that they lived through the Great
	Depression but didn't even know it because they had everything they
	needed."
SCIA	1:20:00 Samuelson – "My brother is the provider in my family. He had
	Covid and went out anyways to get a moose. He had no choice. The
	provider role in our communities is so important."
SCIA	1:24:30 Winkelman – "This are (Y-K Delta) has one of the highest rates
	of traditional food consumption in the entire state"
Interview	4:44 Adolph Lupie – "I'm turning 70My parents are gone. They told me
	that I'll be reaching this era when we will get no more fish on the rive
	swimming, but we have to deal with it withscience."
Interview	10:51 Adolph Lupie – "today we are hurting and suffering, sacrificing
	fish."
Interview	2:36 Evon Waska – "We came, our mom and dad, (to look) forward to the
	return of the salmon, and that's the way I grew up."
Interview	3:27 Evon Waska – "Most of the year it's winter in Alaska. To sustain us
	through the long winter monthsour parents taught us how to live the
	way of life of how important our salmon was."
Interview	12:22 Evon Waska – "Sometimes that king salmon would be the only
	dinner we had."
Interview	14:59 Evon Waska – "I'm into commercial fishing that was a great
	helpwe entered the cash economy and didn't get richUnemployment,
	lack of jobs getting heating oil, billscommercial fishing ended in
	2014that was their incomeGot to stop hardship."
	1 1

Interview	21:20 Evon Waska – "no more commercial fishing milk is now 10, maybe 10-12 bucks, maybe 15 bucks a gallonyou have to barge 'em and freight air freightmy people are going to turn to subsistence."
Public Testimony Nov. 15	2:18:36 Brooke Woods – "the crisis we're in. This is a cultural crisis."
Public Testimony Nov. 14	2:31:25 Stanislaus Sheppard – "North Pacific Fishery(ies) (Management) Council member said, "Sounds like we're in a humanitar[ian] crisis."
Public Testimony Nov. 14	2:56:25 James Nicori – "We work together. We have to work together."
Public Testimony Nov. 14	3:27:40 Virgil Umphenour – "Whenever there's no salmon (I think - hard to hear - might say famine) it affects the entire ecosystem."
Section Existing Knowledge	<b>Traditional Knowledge</b> 15:12 Adolph Lupie – "There will be a lot of mosquitoes before the salmon come in."
Existing Knowledge	
Existing Knowledge	15:26 Adolph Lupie – "Mountains…there'll be lots of snow and they'll be flooding…more water means more fish coming."
Existing Knowledge	17:55 Adolph Lupie – "salmon will come in really fast when we have a good stormy weather from the southit's bringing them in from the
	ocean."
Existing Knowledge	25:44 Adolph Lupie – Explaining Traditional Knowledge to westerners with an example – "My dad used towhen he's carpenteringhe used a stick to measure and eyeball. I told himwe could use a measuring tapebut he said that the stick measure is more accuratewhen you're doing carpentry. Experiencing with it for long time you get used to it and you can eyeball how good to cut it."
Existing Knowledge	34:17 Adolph Lupie – "The moonthe phases controls the tide"
Existing Knowledge	Public Testimony Day $1 - 2:45:40$ James Nicori – "Predicting the salmonstarting winterThey check the thickness of the ice and how much snow we have on the hillsIn the springtime they really pay attention to birds that are coming and when the birds are plentiful they are happy to have salmon this summer."
Existing Knowledge	Public Testimony Day 1 – 2:47:00 James Nicori – "Observe the storm coming inwhen we have strong west wind it pushes some of the Yukon fish into the Kuskokwim and if we have really south winds it pushes somethat are supposed to be going into the Kuskokwim into the Yukon."
Section	Research Priorities based on Traditional Knowledge
Interview	5:28 Adolph Lupie – "I wish to learn moresalmon in the ocean and our Kuskokwim River."
Interview	30:22 Adolph Lupie – In regards to salmon ecology – "How come we don't include things like photosynthesis, plants, food, light" referring to the fact that we don't look at everything, just the fish.
Interview	40:33 Adolph Lupie – "Drop the scientific protocols and integrate Traditional Knowledge."
Interview	40:53 Adolph Lupie – "Talk to more that are not fluent in English. Talk to ones that have more knowledge with their language. They have better words to describe the fish and the situation going on."

Interview	41:17 Adolph Lupie – "What I mean by (scientific) protocolssomeone once had to do research on not catching more fish while the other one was catching a lotcan we share that too?" – Look into how we can share fish that are collected for research purposes."
Interview	35:38 Evon Waska – "My suggestion is, why don't you tag 'em and see where they go…some electronic device."
Interview	36:40 Evon Waska – * "The interceptions are reaching Western Alaska." – research into interceptions of Western Alaska salmon
Public Testimony Nov. 15	2:11:46 Dan Gillikin – "Personally would love to see nothing more than a large scale program for collecting stream temperature data."
Public Testimony Nov. 15	2:17:45 Brooke Woods – "I just don't think we're looking at the physical and mental impacts of no salmon to our dietIt's essential."
Public Testimony Nov. 15	2:19:45 Brooke Woods – "My big concerns are bycatch in the pollock industryand Area M fishery."
Public Testimony Nov. 15	2:20:40 Brooke Woods – "We need to look at the marine environment and see what we can do as far as management decisions."
Public Testimony Nov. 15	2:20:55 Brooke Woods – "'…looking at the health and wellbeing of our people and what salmon means."
Public Testimony Nov. 15	2:22:50 Brooke Woods – "Pink salmon hatchery productionneeds to be addressed."
Public Testimony Nov. 15	2:32:00 Gabe Canfield – "The impacts that are happening to Alaska Pacific salmon and Alaska Pacific salmon habitat. Beyond the more well known ones some more localized human impactssuperfund sites and areas of impacted water qualityincreased road creationshuman impact(s) to critical salmon habitatshould be a priorityheard from community visits across community (Yukon River) this summer."
Public Testimony Nov. 14	2:37:00 Stanislaus Sheppard – "We're all fighting for the same thing. That's salmonWe're talking about protecting the migratory routes just like birds."
Public Testimony Nov. 14	2:42:00 James Nicori – "If it is possible in any way, create a bufferwest coast of Alaska where the salmon travels12 mile buffer zoneso trawlers and bycatch cannot be operatingCreate a buffer zone."
Public Testimony Nov. 14	2:50:25 James Nicori – "The ladies cutting the fish say, "Come look at this fish."The intestines were welded to their meat and to their bones. They can't figure out why that is happening. And sometimes growth on the bodyand they smell differently. "
Public Testimony Nov. 14	2:53:55 James Nicori – "six inches (net) can catch the bigger salmons but they hangbefore you pull in your net they come loose and they fall down into the bottom of the riverThat's salmon reproduce being wastedWe have to look at too."
Public Testimony Nov. 14	3:01:30 Charles Lean – "I think that should be a research priority. At what point do we throw in the towel and say that stocks done? It's never coming back." - referring to addressing sustainable escapement threshold
Public Testimony Nov. 14	3:04:30 Charles Lean – "I'd like to see these research efforts focus on existing science. I'd like to see synthesis papersreach some conclusionssomething you can act on."

Public Testimony Nov. 14	3:05:40 Charles Lean – "Most importantly taking action and evaluating that action towards efficacy."
Public Testimony Nov. 14	3:06:15 Charles Lean – "I think there's a lot of things that we should be looking atdrivers of the system bigger ecosystem kind of approach."
Public Testimony Nov. 14	3:07:58 Tiffany Agayar – "I was wondering if anybody has ever compiled all the data together with how they run, returning back to their streams."
Public Testimony Nov. 14	3:11:15 Tiffany Agayar – "I was just wondering if they were thinking of making a reconciliation of tagged fishwith different areas where fish are." - asking about synthesis and compilation of tagged fish data (e.g., migration)
Public Testimony Nov. 14	3:12:20 Tiffany Agayar – "I believe that tracking would help with being able to point how to and when to regulate just like they do on the Yukon."
Public Testimony Nov. 14	3:15:20 Kathleen Demientieff – "I would rather have the Native (community - not sure - indiscernible) approvewe have Traditional Knowledge."
Public Testimony Nov. 14	3:59:45 Daniel Schindler (Dan Gillikin) – "freshwater stream temperature regime"
Public Testimony Nov. 14	4:02:35 Vanessa von Biela – "We do have a lot of water temperature data from across the state one of the major problemswe're thinking about them in terms of averagesbut is that an accurate reflection of what animals are experiencing?often missing events because we're doing things like taking averages." - saying we need more detailed water temperature data
Public Testimony Nov. 14	4:04:45 Dan Gillikin – "Second, is the loss of MDN subsidies to our streamsWhat has been the total loss of biomass contributed by returning salmon in these systems over the last 10, 20, 30, 40 (years). This loss has had to of reduced the carrying capacity of these systemsCan this be correlated to juvenile growth inferred from scale analysis?"
Public Testimony Nov. 14	4:08:40 Vanessa von Biela – "What does that mean for the way we manage salmon if the productivity of the ecosystem is gonna be reduced?"
Public Testimony Nov. 14	4:09:55 Vanessa von Biela – "The one research gap that I can identify is we really don't have the funding to do all the things that we need to do."
Section Climate Change	<b>Traditional Knowledge</b> 29:30 Adolph Lupie – "Climate change…we are experiencing it and handling it…integrating the science and Taditional Knowledge."
Climate Change	Godduhn et al. 2020 – Another fisher shared similar sentiments regarding the effects of climate change on the fishery: "You have climate change, the biology of the water, the salmon ecosystem is changing, the acidity is increasing, the temperature is increasing which changes their food that they eat and have available."
Climate Change	Godduhn et al. 2020 – One of the more common themes among responses pertained to the effects of climate change. Many respondents cited late freeze-ups, early breakups, and an overall lack of snow and ice for the past several years.
Climate Change	Public Testimony Day 2 – 2:28:30 James Nicori – "One summer we hadtemperatures in the low 90s, upper 80s a couple weeks. They were

starting to have fish floating down the river...haven't reached spawning grounds."

<b>Climate Change</b>	Public Testimony Day 1 – 2:52:25 James Nicori – "Temperature was in the
	90s three or four daysOur river was too warm for those salmon to survive
	salmons were being overheated and they floated down the river."
<b>Climate Change</b>	Public Testimony Day 1 – 4:08:20 Vanessa von Biela – "we're learning

Predation

Predation

Change

Change

Change

Change

Harvest

Harvest

Harvest

Harvest

Harvest

Harvest Harvest

**Freshwater Habitat** 

**Freshwater Habitat** 

**Freshwater Habitat** 

Freshwater Habitat

Public Testimony Day 1 – 4:08:20 Vanessa von Biela – "…we're learning very quickly…that several of these salmon stocks are just becoming far less productive…with changes to climate."

Mikow et al. 2019 – Some respondents described how the sloughs and tributaries of the Kuskokwim River are constantly changing. A beaver dam or sandbar could block a free-flowing slough, creating a dead end slough: "And those are full of [salmon] carcasses. And then the predators are there: the rainbows, dollies, pike, sheefish are there, going crazy in those areas"

Public Testimony Day 1 – 3:32:18 Virgil Umphenour – "On my hunting operation where I guide out of Huslia...grayling...would be digging up the salmon eggs...and grayling ate (them)."

35:04 Adolph Lupie – "When there's high water, they (salmon) go to the river banks to eat grass and moss."

Mikow et al. 2019 – Key respondents also described their perceptions that sport fishers interfere with salmon spawning and contribute to the physical disturbance of Chinook salmon spawning habitat by walking in streams and on riverbanks.

Public Testimony Day 2 – 2:26:30 James Nicori – "... the beavers, they dam the river where the spawners can't even go through the dam...seeing...lots of beaver dams. There used to be no beavers in our area (Kuskokwim River)...migrating into the lower streams...down to the coast now."

Public Testimony Day 2 – 2:35:50 Jacob Ivanoff – "A chemical sprayed on tires actually kills...driving on roads...can affect the salmon."

13:25 Adolph Lupie – "We have to respect them and keep what you do catch very clean and handle them carefully...The animal, fish, or something we take into our home the women they take of it right away...with no complaining...because the person who hunts will get more not less...if we leave it... will not catch anymore while others are catching more.We have to respect them and keep what you do catch very clean and handle them carefully."

18:44 Adolph Lupie – "When's it's too hot we don't try to fish for subsistence."

6:59 Evon Waska – "Wood boaters...As soon as ice went out the setnet went in. King setnet...those weapons were 8.5 to 8 inches."

7:26 Evon Waska – "...our people are into sharing. Elders first. Families first. We never kept the first king. We shared it."

7:46 Evon Waska – "Our cultural and our self-identity was giving to the Elders and sharing our catch."

10:36 Evon Waska – "The moms would teach the daughters how to cut fish." 12:37 Evon Waska – "Mom and dad were boss. They would say enough of those." – meaning time to stop fishing.

Harvest	McDevitt & Koster 2021 – Many respondents attribute the decline in the use
	of fish camps to increased restrictions on fishing opportunity and an
	associated increase in fishing costs
Harvest	McDevitt & Koster 2021 – Although thousands of residents throughout the
	drainage harvest salmon each season, several factors differentiate one region
	of the river to the next. These include differences in the physical nature of
	the river through its course, species distribution and abundance, types of gear
	used by fishers, and population sizes of communities.
Harvest	McDevitt & Koster 2021 – The most common gear types for harvesting
	salmon include drift gillnets, set gillnets, fish wheels, and rod and reel.
	Although both set and drift gillnets are used drainage-wide, disparate
	physical characteristics between the three regions of the river typically
II	demand different gear types in each region.
Harvest	Godduhn et al. 2020 – These (Kuksokwim River) communities are all highly
Harvest	subsistence-dependent with strong traditions surrounding the use of salmon. Godduhn et al. 2020 –harvest of wild foods is still a primary way of life
Hai vest	for most residents.
Harvest	Godduhn et al. 2020 – Residents affirm that salmon provides not only an
	essential food source for families, but also supports important cultural,
	linguistic, and family traditions that encourage health and wellbeing within
	communities. Changing patterns in the harvest and use of salmon continue to
	drive disconcerting social changes in the region, such as reducing the time
	that families spend together at fish camps and the lack of transmission of
	cultural knowledge between older and younger generations.
Harvest	in Godduhn et al. 2020 – Limited archeological documentation in the region
	suggests that Kuskokwim River residents historically exploited consistent
	salmon runs that provided the most reliable element of the resource base,
	certainly for the last 3,000 years (Shaw 1998).
Harvest	in Godduhn et al. 2020 – Prior to Alaska statehood in 1959, commercial
	fishing in the Kuskokwim Area was regulated by quota, and subsistence
	fishing was unregulated and mostly undocumented; fishers either kept fish
	they caught for subsistence use or, if a commercial buyer were available, they apuld call the fish (light at al. 2012)
Howyost	could sell the fish (Ikuta et al. 2013). Mikow et al. 2019 – Fishers in all study communities explained that they
Harvest	maintain close communication with family and friends downstream to obtain
	news about the arrival timing of Chinook salmon.
Harvest	Mikow et al. 2019 – Almost all key respondents observed a decline in
Hai vest	Chinook salmon abundance in fishing areas near their communities, and
	some have decided to no longer target the species as a conservation measure.
Harvest	Ikuta et al. 2013 – Salmon have been a vital source of protein and a cultural
	and economic resource since time immemorial.
Harvest	Ikuta et al. 2013 – Many fishing traditions related to avoiding waste were
	described by research respondents. People do not catch more fish than can be
	processed in a timely manner, and avoid cutting in the hottest time of day,
	when the quality of the meat degrades.

Harvest	Ikuta et al. 2013 – Historical methods of harvesting salmon near Kwethluk
	include gillnets, fish spears, fish traps, and dipnets.
Harvest	Brown et al. 2012 – The most widely used resource category was fishwhich
	was also the resource most commonly harvested and the one making up the
	bulk of the total subsistence harvest.
Harvest	Public Testimony Day 2 – 2:30:00 James Nicori – "In the older daysthere
	weren't any biologists going aroundthe fish cutters, the wiveswere fish
	biologiststold us no more fishing."
Harvest	Public Testimony Day $1 - 2:40:25$ James Nicori $- \dots$ just tell us (take) what
	you can and let the rest go so we can have more to come in later years."
Harvest	Public Testimony Day 1 – 2:40:40 James Nicori – "When the salmon comes
	in fish on the first run that are comingthe first ones are mostly male and
	they go up to the headwaters for the females. Sowhen you have
	enoughthe females pass by."
Harvest	Public Testimony Day $1 - 3:58:00$ Victor Lord – "If the fishing got real low,
	people would pull their nets on their own."

1140

- **1142** *Combination of Regional Teams' References* (As of 10/2/23)
- 1143Reference Title
- 1144 2020 KRITFC Community-Based Harvest Monitoring Program Summary & Report
- 1145 2020 Takotna River Salmon Run Timing and Abundance
- 1146 2021 KRITFC Community-Based Harvest Monitoring Program Summary & Report
- 1147 2021 Kuskokwim River Salmon Situation Report
- 1148 2021 Takotna River Salmon Run Timing and Abundance
- 1149 2022 Kuskokwim River Salmon Situation Report
- 1150 2022 Takotna River Salmon Run Timing and Abundance
- 1151 A

1152 A.R. Godduhn; D.M. Runfola; C.R. McDevitt; J. Park; G. Rakhmetova; J.S. Magdanz; H.S. Cold;

- 1153 C.L. Brown. 2020. Patterns and trends of subsistence salmon harvest and use in the Kuskokwim
- 1154 River Drainage, 1990-2016. ADF&G Division of Subsistence, Technical Paper No. 468.
- Ackerman, M. W., C. Habicht, and L. W. Seeb. 2011. SNPs under diversifying selection provide
  increased accuracy and precision in mixed stock analyses of sockeye salmon from Copper River,
  Alaska. (PDF) Transactions of the American Fisheries Society 140: 865-881.
- Ackerman, M.W., W.D. Templin, J.E. Seeb, and L.W. Seeb. 2013. Landscape heterogeneity and
  local adaptation define the spatial genetic structure of Pacific salmon in a pristine environment.
- 1160 Conservation Genetics 14: 483-498.
- ADF&G 2022. A study of the interactions between hatchery and natural pink and chum salmon in
- 1162 Southeast Alaska and Prince William Sound streams Progress Synopsis May 2022 ADF&G Chinook

1163 Salmon Research Team. 2013. Chinook salmon stock assessment and research plan, 2013. Alaska

1164 Department of Fish and Game, Special Publication No. 13-01, Anchorage.

- 1165 http://www.adfg.alaska.gov/FedAidPDFs/SP13-01.pdf
- Adkison, M.D., 2002. Preseason forecasts of pink salmon harvests in southeast Alaska using
- 1167 Bayesian model averaging. Alaska Fishery Research Bulletin 9, 1–8. Agler, B.A., G.T. Ruggerone,
- 1168 L.I. Wilson, and F.J. Mueter. 2013. Historical growth of Bristol Bay and Yukon River, Alaska chum
- salmon in relation to climate and inter-and intraspecific competition. Deep-Sea Research II 94:165-170
  - 60

- Agler, B.A., Ruggerone, G.T., Wilson, L.I., Mueter, F.J., 2013. Historical growth of Bristol Bay and
  Yukon River, Alaska chum salmon (Oncorhynchus keta) in relation to climate and inter- and
  intraspecific competition. Deep Sea Research Part II: Topical Studies in Oceanography 94, 165–177.
  https://doi.org/10.1016/j. dsr2.2013.03.028
- Alaska Department of Fish and Game Staff, 2022. Preliminary Harvest Rates of Western Alaska and
  Alaska Peninsula Chum Salmon Stocks in South Alaska Peninsula Fisheries, 2022. Regional
  Information Report No. 5J23-02
- Alaska Department of Fish and Game Staff, 2023. Chignik River King Salmon Action Plan, 2023. RC
  5 Alaska Board of Fisheries. Alaska Department of Fish and Game. February 2023
- Alaska Department of Fish and Game Staff, 2023. Chignik River Sockeye Salmon Action Plan, 2023.
  RC 4 Alaska Board of Fisheries. Alaska Department of Fish and Game. February 2023
- Alexander, R.F. and K. K. English. 2022. Preliminary assessment of the Canadian and Alaskan
  Sockeye stocks harvested in the northern boundary fisheries using run reconstruction techniques,
  2009-2021. Prepared by LGL Limited, Sidney, BC, for the Pacific Salmon Commission, Vancouver,
  BC, and Fisheries and Oceans, Canada, Prince Rupert, BC. 127p.
- Alexandersdottir, M. 1987. Life history of pink salmon (Oncorhynchus gorbuscha) in Southeast
  Alaska and implications for management. Ph.D. Thesis. University of Washington, Seattle.
- Andersen, D. B., B. Retherford, and C. Brown. 2013. Climate Change and Impacts on Subsistence
  Fisheries in the Yukon River Drainage, Alaska. Fisheries Resource Monitoring Program 10-250 final
  report.
- Anderson, J.H., Faulds, P.L., Atlas, W.I. and Quinn, T.P., 2013. Reproductive success of captively
  bred and naturally spawned Chinook salmon colonizing newly accessible habitat. Evolutionary
  Applications, 6(2), pp.165-179.
- Anderson, J.H., Ward, E.J. and Carlson, S.M., 2011. A model for estimating the minimum number of
   offspring to sample in studies of reproductive success. Journal of Heredity, 102(5), pp.567-576.
- Anderton, Isaac, and Phyllis Frost. "Traditional/Local Knowledge Salmon Survey." Yukon River
  Panel Project CRE-16-02 Final Report (2002).
- Andrews, A.G, E.V. Farley, Jr., J.H. Moss, J.M. Murphy, and E.F. Husoe. 2009. Energy density and
  length of juvenile Pink salmon in the eastern Bering Sea from 2004 to 2007: a period of relatively
  warm and cool sea surface temperatures. North Pacific Anadromous Fish Commission Bulletin 5:183189.
- Araki, H. and Blouin, M.S., 2005. Unbiased estimation of relative reproductive success of different
   groups: evaluation and correction of bias caused by parentage assignment errors. Molecular Ecology,
   14(13), pp.4097-4109.

- 1205Arctic-Yukon-KuskokwimSustainableSalmonInitiativeProjectDatabase1206https://www.aykssi.org/projects/
- 1207 Arctic-Yukon-Kuskokwim Sustainable Salmon Initiative Research and Discovery Report: 2002-2010
- AYKSSI. 2013. AYK Chinook salmon research plan; evidence of decline of Chinook salmonpopulations and recommendations for future research. AYK SSI Chinook Salmon Expert Panel
- Azuma, T., 1992. Diel feeding habits of sockeye and chum salmon in the Bering Sea during thesummer. Nippon Suisan Gakkaishi 58, 2019–2025.
- Azumaya, T., Ishida, Y., 2000. Density interactions between pink salmon (Oncorhynchus gorbuscha
  and chum salmon (O. keta) and their possible effects on distribution and growth in the North Pacific
  Ocean and Bering Sea. NPAFC Bulletin 2, 165–174.

## 1215 **B**

Baker, M.R., Hollowed, A.B., 2014. Delineating ecological regions in marine systems: Integrating physical structure and community composition to inform spatial management in the eastern Bering
Sea. Deep Sea Research Part II: Topical Studies in Oceanography 109, 215–240.
https://doi.org/10.1016/j.dsr2.2014.03.001

- Baker, T.T., Wertheimer, A., Burkett, R.D., Dunlap, R., Eggers, D.M., Fritts, E.I., Gharrett, A.J.,
  Holmes, R.A., Wilmot, R.L., 1996. Status of Pacific salmon and steelhead escapements in
  Southeastern Alaska. Fisheries 21, 6–18.
- Barclay, A. W., and C. Habicht. 2019. Genetic baseline for Cook Inlet coho salmon and evaluations
  for mixed stock analysis. Alaska Department of Fish and Game, Fishery Manuscript Series No. 1919, Anchorage.
- Barclay, A. W., and E. L. Chenoweth. 2021. Genetic stock identification of Upper Cook Inlet sockeye
  salmon harvest, 2020. Alaska Department of Fish and Game, Division of Commercial Fisheries,
  Regional Information Report No. 5J21-04, Anchorage.
- Barclay, A. W., D. F. Evenson, and C. Habicht. 2019. New genetic baseline for Upper Cook Inlet
  Chinook salmon allows for the identification of more stocks in mixed stock fisheries: 413 loci and 67
  populations. Alaska Department of Fish and Game, Fishery Manuscript Series No. 19-06, Anchorage.
- 1232 Barclay, A. W., H. A. Hoyt, and C. Habicht. 2021. Genetic population structure of Chinook salmon
- 1233 from Middle and Upper Susitna River: A report to Alaska Energy Authority, Susitna-Watana
- 1234 hydroelectric project (submitted July 25, 2017). Alaska Department of Fish and Game, Division of
- 1235 Commercial Fisheries, Regional Information Report No. 5J21-02, Anchorage.
- Barclay, A. W., M. Schuster, C. M. Kerkvliet, M. D. Booz, B. J. Failor, and C. Habicht. 2019. Coded
  wire tag augmented genetic mixed stock analysis of Chinook salmon harvested in Cook Inlet marine

- sport fishery, 2014-2017. Alaska Department of Fish and Game, Fishery Manuscript No. 19-04,Anchorage.
- Barclay, A. W., M. Schuster, C. M. Kerkvliet, M. D. Booz, B. J. Failor, and C. Habicht. 2019. Coded
  wire tag augmented genetic mixed stock analysis of Chinook salmon harvested in Cook Inlet marine
  sport fishery, 2014–2017. Alaska Department of Fish and Game, Fishery Manuscript No. 19-04,
  Anchorage.
- Barclay, A. W., P. A. Crane, D. B. Young, H. A. Hoyt, and C. Habicht. 2017. Current status of genetic
  studies of coho salmon from Southcentral Alaska and evaluations for mixed-stock analysis in Cook
  Inlet. Alaska Department of Fish and Game, Fishery Manuscript Series No. 17-01, Anchorage.
- Barclay, A. W., S. Gilk-Baumer, K. Shedd, J. Botz, and C. Habicht. 2022. Genetic stock composition
  of the commercial harvest of Chinook salmon in Copper River District, 2018-2021. Alaska
  Department of Fish and Game, Fishery Data Series No. 22-35, Anchorage.
- Bataille, C.P., Brennan, S.R., Hartmann, J., Moosdorf, N., Wooller, M.J., Bowen, G.J., 2014. A
  geostatistical framework for predicting variability in strontium concentrations and isotope ratios in
  Alaskan rivers. Chemical Geology 389, 1–15. https://doi.org/10.1016/j. chemgeo.2014.08.030
- Batten, S.D., G.T. Ruggerone, and I. Ortiz. 2018. Pink salmon induce a trophic cascade in plankton
  populations in the southern Bering Sea and around the Aleutian Islands. Fisheries Oceanography
  DOI:10.111/fog.12276
- Batten, S.D., Ruggerone, G.T., Ortiz, I., 2018. Pink Salmon induce a trophic cascade in plankton
  populations in the southern Bering Sea and around the Aleutian Islands. Fisheries Oceanography.
  https:// doi.org/10.1111/fog.12276
- Baumer, J., Wadle, J. and Dublin, R. 2022 Chignik King Salmon Stock of Concern Draft Action Plan.Oral Presentation to Alaska Board of Fisheries. Alaska Department of Fish and Game.
- Beacham, T. D., M. Wetklo, L. Deng, and C. MacConnachie. 2011. Coho Salmon Population Structure
  in North America Determined from Microsatellites, Transactions of the American Fisheries Society,
  140:2, 253-270
- Beacham, T.D., Candy, J.R., Wallace, C., Urawa, S., Sato, S., Varnavskaya, N.V., Le, K.D., Wetklo,
  M., 2009. Microsatellite stock identification of Chum Salmon on a Pacific Rim basis. North American
  Journal of Fisheries Management 29, 1757–1776. https://doi.org/10.1577/M08-188.1
- Beacham, T.D., Murray, C.B., and Withler, R.E. 1989. Age, morphology, and biochemical genetic
  variation of Yukon River Chinook salmon. Transactions of the American Fisheries Society 118: 46https://doi.org/10.1577/1548-8659(1989)118<0046 :AMABGV>2.3.CO;2
- Beamish, R. J., editor. 2018 The ocean ecology of Pacific salmon and trout. American FisheriesSociety, Bethesda, Maryland.

Beaudreau, A.H., Chan, M.N., Loring, P.A., 2018. Harvest portfolio diversification and emergent
conservation challenges in an Alaskan recreational fishery. Biological Conservation 222, 268–277.
https:// doi.org/10.1016/j.biocon.2018.04.010

Beechie, T.J., C. Fogel, C. Nicol, J. Jorgensen, B. Timpane-Padgham, and P. Kiffney. 2022. How does
habitat restoration influence resilience of salmon populations to climate change. Freshwater Ecology
DOI:10.1002/ecs2.4402

Bellmore, J.R., Fellman, J.B., Hood, E., Dunkle, M.R., Edwards, R.T., 2022. A melting cryosphere
constrains fish growth by synchronizing the seasonal phenology of river food webs. Global Change
Biology 4807–4818. https://doi.org/10.1111/gcb.16273

Berkman, S.A., Sutton, T.M., Mueter, F.J., Elliott, B.W., 2021. Effects of early-life stage and
environmental factors on the freshwater and marine survival of Chinook salmon (Oncorhynchus
tshawytscha) in rivers of southeast Alaska. Fishery Bulletin 119, 201–215. https://
doi.org/10.7755/FB.119.4.1

- 1285 Bernard, D.R., 1983. Variance and bias of catch allocations that use the age composition of 1286 escapements.
- Bond, N., Overland, J.E., Spillane, M., Stabeno, P., 2003. Recent shifts in the state of the North Pacific.
  Geophysical Research Letters 30, 2–5. https://doi.org/10.1029/2003GL018597
- Botz and Sommerville (2021) Management of salmon stocks in the Copper River, 2018-2020 -- a
  report to the Alaska Board of Fisheries.

Bowen L, von Biela VR, McCormick SD, Regish AM, Waters SC, Durbin-Johnson B, Britton M,
Settles ML, Donnelly DS, Laske SM, et al. (2020) Transcriptomic response to elevated water
temperatures in adult migrating Yukon River Chinook salmon (Oncorhynchus tshawytscha). Conserv
Physiol 8: 1–22.

- Brabets, T.P., Walvoord, M.A., 2009. Trends in streamflow in the Yukon River Basin from 1944 to
  2005 and the influence of the Pacific Decadal Oscillation. Journal of Hydrology 371, 108–119.
- Bradford, M. J., J. Duncan, and J. W. Jang. 2009. Downstream migrations of juvenile salmon andother fishes in the upper yukon river. Arctic 61(3).
- Bradford, M.J., Grout, J.A., and Moodie, S. 2001. Ecology of juvenile Chinook salmon in a small nonnatal stream of the Yukon River drainage and the role of ice conditions on their distribution and
  survival. Canadian Journal of Zoology. 79(11): 2043-2054. https://doi.org/10.1139/z01-165
- Bradley, P. T., M. D. Evans, and A. C. Seitz. 2015. Characterizing the Juvenile Fish Community in
  Turbid Alaskan Rivers to Assess Potential Interactions with Hydrokinetic Devices. Transactions of
  the American Fisheries Society 144(5):1058–1069.

Braem, N. M., E. Mikow, A. Goddhun, A. R. Brenner, A. Trainor, S. J. WIlson and M. L. Kostick.
2017. Key Subistence Fisheries in Nortwest Alaska, 2012-2014. Alaska Department of Fish and Game
Division of Subsistence, Technical Paper No. 433, Fairbanks.

Briscoe, R.J., Adkison, M.D., Wertheimer, A., Taylor, S.G., 2005. Biophysical factors associated with
the marine survival of Auke Creek, Alaska, coho salmon. Transactions of the American Fisheries
Society 134, 817–828.

Brock, M and P. Coiley-Kenner. 2009. A compilation of traditional knowledge about the fisheries of
Southeast Alaska. ADF&G Division of Subsistence, Techncial Paper No. 332

Bromaghin, J.F., 2005. A versatile net selectivity model, with application to Pacific salmon and
freshwater species of the Yukon River, Alaska. Fisheries Research 74, 157–168.
https://doi.org/10.1016/j. fishres.2005.03.004

Bromaghin, J.F., 2008. Bels: Backward elimination locus selection for studies of mixture composition
or individual assignment. Molecular Ecology Resources 8, 568–571. https://doi. org/10.1111/j.14718286.2007.02010.x

Bromaghin, J.F., R.M. Nielson, and J.J. Hard. 2008. An investigation of the potential effects of
selective exploitation on the demography and productivity of Yukon River Chinook salmon. Alaska
Fisheries Technical Report Number 100. USFWS

1521 Tisheries Teenmeur Report Rumber 100. OST WS

1322 Bromaghin, Jeffrey F, Evenson, D.F., McLain, T.H., Flannery, B.G., 2011. Using a Genetic Mixture Model to Study Phenotypic Traits: Differential Fecundity among Yukon River Chinook Salmon. 1323 American Transactions Fisheries Society 235-249. https:// 1324 of the 140. 1325 doi.org/10.1080/00028487.2011.558776

Bromaghin, Jeffrey F., Nielson, R.M., Hard, J.J., 2011. A model of Chinook salmon population
dynamics incorporating size-selective exploitation and inheritance of polygenic correlated traits.
Natural Resource Modeling 24, 1–47. https://doi. org/10.1111/j.1939-7445.2010.00077.x

Brown, C.L., and M.L. Kostick, editors. 2017. Harvest and Use of Subsistence Resources in 4
Communities in the Nenana Basin, 2015. Alaska Department of Fish and Game Division of
Subsistence, Technical Paper No. 429, Fairbanks.

Brown, Caroline L., H. Cold, L. Hutchinson-Scarbrough, B. Jones, J.M. Keating, B.M. McDavid, M.
Urquia, J. Park, L.A. Sill, and T. Barnett. 2022. Alaska Subsistence and Personal Use Salmon Fisheries
2019 Annual Report. Alaska Department of fish and Game Division of Subsistence, Technical Paper
No. 490, Anchorage.

1336 Brown, R.J., 2012. Yukon Chinook Review, memo.

Brown, R.J., A. von Finster, R.J. Henszey, and J.H. Eiler. 2017. Catalog of Chinook salmon spawningareas in Yukon River Basin in Canada and United States

- Brown, R.J., Bradley, C., Melegari, J.L., 2020a. Population trends for Chinook and summer chum
  salmon in two Yukon River tributaries in Alaska. Journal of Fish and Wildlife Management 11, 377–
  400. https://doi.org/10.3996/072019-JFWM-064
- Brown, Z.W., Arrigo, K.R., 2013. Sea ice impacts on spring bloom dynamics and net primary
  production in the Eastern Bering Sea. Journal of Geophysical Research: Oceans 118, 43–62.
  https://doi.org/10.1029/2012JC008034
- Brown, Z.W., van Dijken, G.L., Arrigo, K.R., 2011. A reassessment of primary production and
  environmental change in the Bering Sea. Journal of Geophysical Research 116, C08014. https://doi.
  org/10.1029/2010JC006766
- Bryant, M.D., 2000. Estimating Fish Populations by Removal Methods with Minnow Traps in
  Southeast Alaska Streams. North American Journal of Fisheries Management 20, 923–930. https://
  doi.org/10.1577/1548-8675(2000)020<0923:EFPBRM>2.0.CO;2
- Bryant, M.D., 2009. Global climate change and potential effects on Pacific salmonids in freshwater
  ecosystems of southeast Alaska. Climatic Change 95, 169–193.
- Bryant, M.D., Everest, F.H., 1998. Management and condition of watersheds in southeast Alaska: the
  persistence of anadromous salmon. Northwest Science 72, 249–267.
- Bryant, M.D., Woodsmith, R.D., 2009. The response of salmon populations to geomorphic
  measurements at three scales. North American Journal of Fisheries Management 29, 549–559.
- Bryant, M.D., Zymonas, N.D., Wright, B.E., 2004. Salmonids on the fringe: abundance, species,
  composition, and habitat use of salmonids in high-gradient headwater streams, southeast Alaska.
  Transactions of the American Fisheries Society 133, 1529–1538.
- Bue, B.G., Molyneaux, D.B., Schaberg, K.L., 2008. Kuskokwim River chum salmon run
  reconstruction., Fishery Data Series. Alaska Department of Fish and Game, Anchorage.
- Bue, B.G., Schaberg, K.L., Liller, Z.W., Molyneaux, D.B., 2012. Estimates of the historic run and
  escapement for the Chinook salmon stock returning to the Kuskokwim River, 1976-2011. Fishery
  Data Series 12–49, 1–47.
- 1365 Bue, F.J., Hayes, S.J., Fisheries, A.D. of F. and G.D. of C., 2009. 2009 Yukon area subsistence,
- personal use, and commercial salmon fisheries outlook and management strategies., RegionalFisheries Report. Anchorage.
- Buffington, J.M., Lisle, T.E., Woodsmith, R.D., Hilton, S., 2002. Controls on the size and occurrence
  of pools in coarse-grained forest rivers. River Research and Applications 18, 507–531. https://
  doi.org/10.1002/rra.693
- Bugaev, A.B., Zavolokina, E.A., Zavarina, L.O., Kiryeev, E.H., Shubin, A.O., Ignatev, Yo.E.,
  Zolotukhia, C.F., Kaplanova, H.F., Volobuev, M.B., 2009. Origin and distribution of local stocks of

- chum salmon in the western Bering Sea based on trawl surveys of the R/V TINRO in 2004 and 20062.
- Bugaev, A.V., Myers, K.W., 2009. Stock-specific distribution and abundance of immature Chinook
  salmon in the western Bering Sea in summer and fall 2002-2004. North Pacific Anadromous Fisheries
  Commission Bulletin 5, 87–97.
- Bugaev, A.V., Zavolokina, E.A., Zavarina, L.O., Shubin, A.O., Zolotukhin, S.F., Kaplanova, N.F.,
  Volobuev, N., 2006. Identification of local stocks of chum salmon Oncorhynchus keta in the western
- **1380** Bering Sea in trawl catches of R/V TINRO in September-October, 2002 and 2003. 2.
- Buklis, L.S., 1999. A description of economic changes in commercial salmon fisheries in a region of
  mixed subsistence and market economies. Arctic 52, 40–48.
- Burkett, J. Koenings, M. Haddix, and D. Barto. 1989. Cooperative ADF&G, FRED Division/U.S.
  Forest Service lake enrichment program for Southeast Alaska. Alaska Department of Fish and Game,
- 1385 FRED Report No. 98.
- Burnside, C., and B. A. Fuerst. 2023. Chignik Management Area salmon annual management report,
  2022. Alaska Department of Fish and Game, Fishery Management Report No. 23-02, Anchorage.
- Burril, S. E., V. R. von Biela, N. Hillgruber, and C. E. Zimmerman. 2018. Energy allocation and
  feeding ecology of juvenile chum salmon (Oncorhynchus keta) during transition from freshwater to
  saltwater. Polar Biology 41(7):1–15.
- 1391 **C**
- C. L. Brown; James S. Maganz; David S. Koster; Nicole M. Braem. 2009. Subsistence Harvests in 8
  Communities in the Central Kuskokwim Drainage, 2009. Technical Paper 365.
- C. McDevitt; D. Koster; D. Runfola; M. Horne-Brine; J. Esquible-Hussion. 2020. Subsistence fisheries
  harvest monitoring report, Kuskokwim Management Area, Alaska, 2018. ADF&G Division of
  Subsistence, Technical Paper No. 467.
- C. McDevitt; D. Koster; D. Runfola; M. Horne-Brine; J. Esquible. 2021. Subsistence fisheries harvest
  monitoring report, Kuskokwim Management Area, Alaska, 2019. ADF&G Division of Subsistence,
  Techncial Paper No. 475.
- 1400 C. McDevitt; D. Koster. 2022. Subsistence fisheries harvest monitoring report, Kuskokwim Fisheries
  1401 Management Area, 2021. ADF&G Division of Subsistence, Techncial Paper No. 489.
- Carey MP, von Biela VR, Dunker A, Keith KD, Schelske M, Lean C, Zimmerman CE (2021) Egg
  retention of high-latitude sockeye salmon (Oncorhynchus nerka) in the Pilgrim River, Alaska, during
  the Pacific marine heatwave of 2014–2016. Polar Biol 44: 1643–1654.

1405 Carothers, C., Cotton, S. and K. Moerlein. 2013. Subsistence Use and Knowledge of Salmon in Barrow
1406 and Nuiqsut, Alaska. Final Report for OCS Study BOEM 2013-0015. University of Alaska Coastal
1407 Marine Institute, Fairbanks.

- 1408 Carothers, C., J. Black, S.J. Langdon and others. 2021. Indigenous peoples and salmon stewardship:1409 a critical relationship. Ecology and Society 26(1):16.
- Carothers, C., T.L. Sformo, S. Cotton, J.C. George, and P.A.H. Westley. 2019. Pacific salmon in the
  rapidly changing Arctic: Exploring local knowledge and emerging fisheries in Utqiagvik and Nuiqsut,
  Alaska Arctic 72(2):272–288
- 1412 Alaska. Arctic 72(3):273-288
- 1413 Carvalho, K.S., Smith, T.E., Wang, S., 2021. Bering Sea marine heatwaves: Patterns, trends and
  1414 connections with the Arctic. Journal of Hydrology 600, 126462. https://doi.org/10.1016/j.
  1415 jhydrol.2021.126462
- 1416 Celewycz, A.G., Wertheimer, A.C., Orsi, J.A., Lum, J.L., 1994. Nearshore distribution and residency
  1417 of pink salmon (Oncorhynchus gorbuscha) and chum salmon (O. keta) fry and their predators in Auke
  1418 Bay and Gastineau Channel, southeast Alaska. AFSC Processed Report 94-05.
- 1419 Chaloner, D.T., Martin, K.M., Wipfli, M.S., Ostrom, P.H., Lamberti, G.A., 1996. Marine carbon and
  1420 nitrogen in southeastern Alaska stream food webs: evidence from artificial and natural streams.
  1421 Canadian Journal of Fisheries and Aquatic Sciences 59, 1257–1265.
- Chasco, B., R. Hilborn, and G.T. Ruggerone. 2004. Chignik salmon studies investigations of salmon
  populations, hydrology, and limnology of the Chignik lakes, Alaska, during 2003-2004. Technical
  Report, Shool of Aquatic and Fishery Science, Fisheries Research Institute, Washington University
  Issue 0304.
- Chasco, B.E., and others. 2017. Competing tradeoffs between increasing marine mammal predation
  and fisheries harves of Chinook salmon. Nature/Scientific Reports. doi:10.1038/s41598-017-14984-8
- 1428 Cheng, W., C. Habicht, W.D. Templin, Z.D. Grauvogel, S.D. Moffitt, R.E. Brenner, R.P. Josephson,1429 and A.J. Gharrett. Population genetic structure of odd-year pink salmon from Prince William Sound
- based on a single year (2013). Alaska Hatchery Research Group Technical Document No. 14.
- 1431 Cheng, W., Curchitser, E., Ladd, C., Stabeno, P., Wang, M., 2014. Influences of sea ice on the
- 1432 Eastern Bering Sea: NCAR CESM simulations and comparison with observations. Deep Sea
- 1433 Research Part II: Topical Studies in Oceanography 109, 27–38. https://doi.
- 1434 org/10.1016/j.dsr2.2014.03.002.
- 1435 Chenoweth E. M., and K. R. Criddle. 2019. The Economic Impacts of Humpback Whale
- 1436 Depredation on Hatchery-Released Juvenile Pacific Salmon in Southeast Alaska. Marine and Coastal
- 1437 Fisheries: Dynamics, Management, and Ecosystem Science 11:62–75.

1438 Chenoweth E. M., J. M. Straley, M. V. McPhee, S. Atkinson, S. Reifenstuhl. 2017. Humpback whales
1439 feed on hatchery-released juvenile salmon.R. Soc. open sci. 4: 170180. http://dx.doi.org/10.1098/
1440 rsos.170180

1441 Chinook Stock Assessment & Research Project, Alaska Department of Fish and Game

1442 Chris McDevitt; David Koster; David Runfola; Maureen Horne-Brine; Janessa Esquible. 2021.
1443 Subsistence Fisheries Harvest Monitoring Report, Kuskokwim Fisheries Management Area, Alaska,
1444 2020. ADF&G Division of Subsistence, Technical Paper No. 483.

Cieciel, K., Farley, E.V., Eisner, L.B., 2009. Jellyfish and Juvenile Salmon Associations with
Oceanographic Characteristics during Warm and Cool Years in the Eastern Bering Sea. North Pacific
Anadromous Fish Commission Bulletin 5, 209–224.

- 1448 Clark, R., M. Willette, S. Fleischman, and D. Eggers. 2007. Biological and fishery-related aspects of
  1449 overescapement in Alaska sockeye salmon Onchorynchus nerka. Alaska Department of Fish and
  1450 Game, Special Publication No. 07-17, Anchorage.
- 1451 Cline, T.J., Ohlberger, J. & Schindler, D.E. Effects of warming climate and competition in the ocean
  1452 for life-histories of Pacific salmon. Nat Ecol Evol 3, 935–942 (2019).

1453 Connors, B., M.J. Malick, G.T. Ruggerone, P. Rand, M. Adkison, J.R. Irvine, R. Campbell, and K.
1454 Gorman. 2020. Climate and competition influence sockeye salmon population dynamics across the
1455 Northeast Pacific Ocean. Canadian Journal of Fisheries and Aquatic Sciences. 77:943-949.

Connors, B.M., Siegle, M.R., Harding, J., Rossi, S., Staton, B.A., Jones, M.L., Bradford, M.J., Brown,
R., Bechtol, B., Boherty, B., Cox, S., and Sutherland, B.J.G. 2022. Chinook salmon diversity
contributes to fishery stability and trade-offs with mixed-stock harvest. Ecological Applications 32(8):
e2709. https://doi.org/10.1002/eap.2709

- 1460 Conrad, S., Davidson, W., 2013. Overview of the 2012 Southeast Alaska and Yakutat commercial,1461 personal use, and subsistence salmon fisheries.
- 1462 Cook, M.E.A., Sturdevant, M.V., Fisheries, N., Fisheries, A., Stevens, T., 2013. Diet Composition and
  1463 Feeding Behavior of Juvenile Salmonids Collected in the Northern Bering Sea from August to October
  1464 , 2009 2011 118–126.
- Copeman, L.A., Salant, C.D., Stowell, M.A., Spencer, M.L., Kimmel, D.G., Pinchuk, A.I., Laurel,
  B.J., 2022. Annual and spatial variation in the condition and lipid storage of juvenile Chukchi Sea
  gadids during a recent period of environmental warming (2012 to 2019). Deep Sea Research Part II:
  Topical Studies in Oceanography 205, 105180. https://doi.org/10.1016/j.dsr2.2022.105180
- Courtney, M.B., Evans, M., Shedd, K.R. et al. Understanding the behavior and ecology of Chinook
  salmon (Oncorhynchus tshawytscha) on an important feeding ground in the Gulf of Alaska. Environ
  Biol Fish 104, 357–373 (2021). https://doi.org/10.1007/ s10641-021-01083-x

Cox, C.J., Stone, R.S., Douglas, D.C., Stanitski, D.M., Gallagher, M.R., 2019. The Aleutian Low-1472 1473 Beaufort Sea Anticyclone: A Climate Index Correlated With the Timing of Springtime Melt in the Cryosphere. Geophys. Res. Lett. 46. 7464-7473. https://doi. 1474 Pacific Arctic 1475 org/10.1029/2019GL083306

Coyle, K.O., Eisner, L.B., Mueter, F.J., Pinchuk, a. I., Janout, M. a., Cieciel, K.D., Farley, E.V.,
Andrews, a. G., 2011. Climate change in the southeastern Bering Sea: impacts on pollock stocks and
implications for the oscillating control hypothesis. Fisheries Oceanography 20, 139–156.
https://doi.org/10.1111/j.1365-2419.2011.00574.x

Coyle, K.O., Pinchuk, A.I., Eisner, L.B., Napp, J.M., 2008. Zooplankton species composition,
abundance and biomass on the eastern Bering Sea shelf during summer: The potential role of watercolumn stability and nutrients in structuring the zooplankton community. Deep Sea Research Part II:
Topical Studies in Oceanography 55, 1775–1791. https://doi.org/10.1016/j.dsr2.2008.04.029

Creelman, E., L. Hauser, R. Simmons, W. Templin, and L. Seeb. 2011. Temporal and geographic
genetic divergence: Characterizing sockeye salmon populations in the Chignik watershed, Alaska,
using single nucleotide polymorphisms. Transactions of the American Fisheries Society 140: 749-762.

1487 Crone, R.A., Bond, C.E., 1976. Life history of coho salmon, Oncorhynchus kisutch, in Sashin Creek,
1488 Southeastern Alaska. Fisheries Bulletin 74, 897–923.

1489 Crowell, A.L., and Arimitsu, M. 2023. Cimate change and pulse migration: intermittent Chugach Inuit
1490 occupation of glacial fjords on the Kenai Coast, Alaska. Frontiers in Environmental Archaeology 1:
1145220. doi: 10.3389/fearc.2023.1145220

Cunningham, C. J., T. A. Branch, T. H. Dann, M. Smith, J. E. Seeb, L. W. Seeb, and R. Hilborn. 2017.
A General Model for Salmon Run Reconstruction That Accounts for Interception and Differences in
Availability to Harvest. Canadian Journal of Fisheries and Aquatic Sciences. Online. doi:
10.1139/cjfas-2016-0360.

Cunningham, C.J., G.T. Ruggerone, and T.P. Quinn. 2013. Size selectivity of predation by brownbears depends on the density of their sockeye salmon prey. The American Naturalist 181(5):

Cunningham, C.J., P.A.H. Westley, and M.D. Adkison. 2018. Signals of large scale climatic drivers,
hatchery enhancement, and marine factors in Yukon River Chinook salmon survival revealed with a
Bayesian life history model. Global Change Biology DOI:10.111/ gcb.14315

1501 D

Dann, T. 2023. Genetic Stock Composition of Chum Salmon Harvested in Commercial Salmon
Fisheries of the South Alaska Peninsula, 2022. A Report to the Alaska Board of Fisheries February
2023.

Dann, T. H., A. Barclay and C. Habicht. 2012. Western Alaska Salmon Stock Identification Program
Technical Document 5: Status of the SNP baseline for sockeye salmon. (PDF 2,631 kB) Alaska

- 1507 Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 5J12-10, Anchorage.
- Dann, T. H., C. Habicht, J. R. Jasper, E. K. C. Fox, H. A. Hoyt, H. L. Liller, E. S. Lardizabal, P. A.
  Kuriscak, Z. D. Grauvogel, and W. D. Templin. 2012. Sockeye salmon baseline for the Western
  Alaska Salmon Stock Identification Program. (PDF 9,421 kB) Alaska Department of Fish and Game,
  Special Publication No. 12-12, Anchorage.
- 1513 Dann, T. H., C. Habicht, J. R. Jasper, H. A. Hoyt, A. W. Barclay, W. D. Templin, T. T. Baker, F. W.
  1514 West, and L. F. Fair. 2009. Genetic stock composition of the commercial harvest of sockeye salmon
- 1515 in Bristol Bay, Alaska, 2006-2008. Alaska Department of Fish and Game, Fishery Manuscript Series
- 1516 No. 09-06, Anchorage.
- Dann, T. H., C. Habicht, T. T. Baker, and J. E. Seeb. 2013. Exploiting genetic diversity to balance
  conservation and harvest of migratory salmon. Canadian Journal of Fisheries and Aquatic Sciences.
  70: 785-793.
- Dann, T. H., C. Habicht, W. D. Templin, L. W. Seeb, G. McKinney, and J. E. Seeb. 2018.
  Identification of genetic markers useful for mixed stock analysis of Chinook salmon in Cook Inlet,
  Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional
  Information Report 5J18-04, Anchorage.
- 1524 Dann, T.H., H. Hoyt, E. Lee, E. Fox, and M. B. Foster. 2023. Genetic Stock Composition of Chum1525 Salmon Harvested in Commercial Salmon Fisheries of the South Alaska Peninsula, 2022
- Dann, T.H., Smoker, W.W., Hard, J.J., Gharrett, A.J., 2010. Outbreeding depression after two
  generations of hybridizing Southeast Alaska coho salmon populations. Transactions of the American
  Fisheries Society 139, 1292–1305. https://doi.org/10.1577/T09-203.1
- 1529 Dann, Tyler (2021), Stock composition of subsistence harvests and total return of sockeye salmon
  1530 from the Kvichak River, Dryad, Dataset, https://doi.org/10.5061/dryad.j0zpc86fm
- Daum, D. and Flannery, B. 2011. Canadian-origin Chinook salmon rearing in nonnatal U.S. tributary
  streams of the Yukon River, Alaska. Transactions of the American Fisheries Society 140. 207-220.
  10.1080/00028487.2011.545004.
- 1534 David M. Runfola; Christopher R. McDevitt; Caroline L. Brown. 2018. Overview of the development
  1535 and implementation of the Kuskokwim River household subsistence king salmon permit system, 2018.
  1536 Alaska Department of Fish and Game Division of Subsistence, Special Publication No. -006.
- 1537 David Runfola; Loraine S. Naaktgeboren; David Koster. 2019. Inseason subsistence salmon harvest
  1538 assessments in 9 communities of the middle Kuskokwim River, 2015–2018. ADF&G Division of
  1539 Subsistence, Technical Paper No. 455.

- 1540 Davidson, B., Bachman, R., Thynes, T., Gordon, D., Piston, A., Jensen, K., Monagle, K., Walker, S.,
- 1541 2017. Annual Management Report of the 2011 Southeast Alaska Commercial Purse Seine and Drift
- 1542 Gillnet Fisheries by.
- Davis, N.C.D., 2003. Feeding Ecology of Pacific Salmon (Oncorhynchus spp.) in the Central North
  Pacific Ocean and Central Bering Sea, 1991-2000. Hokkaido University.
- Davis, N.D., Armstrong, J.L., Myers, K.W., 2004a. Bering Sea Salmon Diet Overlap in Fall 2002 and
  Potential for Interactions Among Salmon by Bering Sea Salmon Diet Overlap in Fall 2002 and
  Potential for Interactions Among Salmon.
- Davis, N.D., Fukuwaka, M., Armstrong, J.L., Myers, K.W., 2004b. Salmon Food Habits Studies in
  the Bering Sea , 1960 to Present 24–28.
- Davis, N.D., Volkov, A.V., Efimkin, A.Y., Kuznetsova, N.A., Armstrong, J.L., Sakai, O., 2009.
  Review of BASIS Salmon Food Habits Studies 197–208.
- 1552 De Robertis, A., Cokelet, E.D., 2012. Distribution of fish and macrozooplankton in ice-covered and
- 1553 open-water areas of the eastern Bering Sea. Deep Sea Research Part II: Topical Studies in
- 1554 Oceanography 65–70, 217–229. https://doi.org/10.1016/j.dsr2.2012.02.005
- 1555 Decovich, N. a, Howard, K.G., 2011. Genetic Stock Identification of Chinook Salmon Harvest on the1556 Yukon River 2010.
- 1557 Deeg, C.M., et al. 2022. Pathogens and stressors of overwintering salmon in the Gulf of Alaska. North
  1558 Pacific Anadromous Fish Commission Technical Report No. 18: 47-52.
- Denman, R.A., and G.T. Ruggerone. 1994. Effects of beaver colonization on the hydrology and
  spawning habitat of sockeye salmon in the Chignik Lakes, Alaska. Natural Resources Consultants, Inc
  report
- 1562 Denton, C., 1988. Marine survival of Chinook salmon, Oncorhynchus tshawytscha, reared at three1563 densities. Alaska Department of Fish and Game, Juneau, AK.
- 1564 Der Hovanisian, J., J. S. McPherson, E. Jones, P. Richards, R. Chapell, B. Elliott, T. Johnson, and S.
  1565 Fleischman. 2011. Chinook salmon status and escapement goals for stocks in Southeast Alaska.
  1566 Alaska Department of Fish and Game, Special Publication No. 11-19, Anchorage.
- Dickerson, B., Brinck, K., Willson, M., Bentzen, P., P, Q.T., 2005. Relative importance of salmon
  body size and arrival time at breeding grounds to reproductive success. Ecology 86, 347–352.
- Dickerson, B.R., Quinn, T.P., Willson, M.F., 2002. Body size, arrival date, and reproductive success
  of pink salmon, Oncorhynchus gorbuscha. Ethology Ecology & Evolution 14, 29–44. https://doi.
- 1571 org/10.1080/08927014.2002.9522759.

- 1572 Division of Commercial Fisheries Staff 2023. Preliminary Harvest Rates of Western Alaska and
- Alaska Peninsula Chum Salmon Stocks in South Alaska Peninsula Fisheries, 2022. A Report to theAlaska Board of Fisheries February 2023. Alaska Department of Fish and Game
- 1374 Alaska Board of Fishenes February 2025. Alaska Department of Fish and Game
- 1575 Doctor, K.K., Hilborn, R., Rowse, M., and Quinn, T. 2009. Spatial and temporal patterns of upriver
  1576 migration by sockeye salmon populations in the Wood River system, Bristol, Bay, Alaska.
  1577 Transactions of the American Fisheries Society 139: 80-91. DOI: 10.1577/T08-227.1
- 1578 Donkersloot, R., Black, J.C., Carothers, C., Ringer, D., Justin, W., Clay, P.M., Poe, M.R., Gavenus,
  1579 E.R., Voinot-Baron, W., Stevens, C. and Williams, M., 2020. Assessing the sustainability and equity
  1580 of Alaska salmon fisheries through a well-being framework. Ecology and Society 25(2):18.
- Donnellan, S. J., and A. R. Munro, editors. 2023. Run forecasts and harvest projections for 2023
  Alaska salmon fisheries and review of the 2022 season. Alaska Department of Fish and Game, Special
  Publication No. 23-10, Anchorage.
- 1584 Dube, M., B. Muldoon, J. Wilson, and K.B. Maracle. 2012. Accumulated state of the Yukon River
  1585 watershed: Part 1 Critical Review of Literature. Integrated Environmental Assessment and
  1586 Management 9(3):426-438
- 1587 Dube, M., J.E. Wilson, and J. Waterhouse. 2012. Accumulated state assessment of the Yukon River
  1588 watershed: Part II Quantitative effects-based analysis integrating western science and traditional
  1589 ecological knowledge. Integrated Environmental Assessment and Management 9(3):439-455.
- 1590 DuBois, L., Liller, Z.W., 2010. Yukon River chinook salmon aging consistency., Fishery Data Series.
  1591 Alaska Department of Fish and Game, Anchorage.
- Duffy-Anderson, J.T., Stabeno, P.J., Siddon, E.C., Andrews, A.G., Cooper, D.W., Eisner, L.B., Farley,
  E.V., Harpold, C.E., Heintz, R.A., Kimmel, D.G., Sewall, F.F., Spear, A.H., Yasumiishi, E.C., 2017.
  Return of warm conditions in the southeastern Bering Sea: Physics to fluorescence. PLoS ONE 12, 1–
  21. https://doi.org/10.1371/journal. pone.0185464
- Duncan, D. H., and A. H. Beaudreau. 2019. Spatiotemporal Variation and Size-Selective Predation on
  Hatchery- and Wild-Born Juvenile Chum Salmon at Marine Entry by Nearshore Fishes in Southeast
  Alaska. Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science 11:372–390.
- Dunmall, K.M., N.J. Mochnacz, C.E. Zimmerman, C. Lean, and J.D. Reist. 2016. Using thermal limits
  to assess establishment of fish dispersing to high-latitude and high-elevation watersheds. Can.J.Fish.
  Aquat. Sci 73:1750-1758
- 1602 E
- E.H. Mikow; B. Retherford; D.M. Runfola; D. Gonzalez. 2019. Local traditional knowledge of salmon
  freshwater ecology in the middle and upper Kuskokwim River. ADF&G Division of Subsistence,
  Technical Paper No. 450.

Ebbin, S. a., 2002. What's Up? The Transformation of Upstream-Downstream Relationships on
Alaska's Kuskokwim River. Polar Geography 26, 147–166. https://doi.org/10.1080/789610136

Echave, J.D., Manhard, C.V., Smoker, W.W., Adkison, M.D., Gharrett, A.J., 2017. Out crosses
between seasonally different segments of a Pacific salmon population reveal local adaptation.
Environmental Biology of Fishes 100, 1469–1481. https://doi.org/10.1007/s10641-017-0657-3

- 1611 Echave, K., Eagleton, M., Farley, E., Orsi, J., 2012. NOAA Technical Memorandum NMFS-AFSC-
- 1612 236 A Refined Description of Essential Fish Habitat for Pacific Salmon Within the U. S. Exclusive
   1613 Economic Zone in Alaska.
- Eggers, D.M., Bachman, R.L., Stahl, J., 2010. Stock status and escapement goals for Chilkat Lakesockeye salmon in Southeast Alaska.
- 1616 Eiler JH, Evans AN, Schreck CB (2015) Migratory patterns of wild Chinook salmon Oncorhynchus
  1617 tshawytscha returning to a large, free-flowing river basin. PLoS One 10: 1–33.
- 1618 Eiler JH, Masuda MM, Evans AN (2023) Swimming depths and water temperatures encountered by
- 1619 radio-archival-tagged Chinook Salmon during their spawning migration in the Yukon River basin.
- **1620** Trans Am Fish Soc 152: 51–74.
- 1621 Eiler JH, Masuda MM, Spencer TR, Driscoll RJ, Schreck CB (2014) Distribution, Stock Composition
  1622 and Timing, and Tagging Response of Wild Chinook Salmon Returning to a Large, Free-Flowing
  1623 River Basin. Trans Am Fish Soc 143: 1476–1507.
- 1624 Eisner, L., Hillgruber, N., Martinson, E., Maselko, J., 2012. Pelagic fish and zooplankton species
  1625 assemblages in relation to water mass characteristics in the northern Bering and southeast Chukchi
  1626 seas. Polar Biology 36, 87–113. https://doi.org/10.1007/ s00300-012-1241-0
- 1627 Eldridge, W.H., J.J. Hard, and K.A. Naish. 2010. Simulating fishery-induced evolution in chinook
  1628 salmon: the rol of gear, location, and gentic correlation among traits. Ecological Applications
  1629 20(7):1936-1948
- 1630 Elison, T.B., Schaberg, K.L., Bergstrom, D.J., 2012. Kuskokwim River Salmon Stock Status and
  1631 Kuskokwim Area Fisheries , 2012 ; a Report to the Alaska Board of Fisheries by.
- 1632 Eskelin, A., and A. W. Barclay. 2022. Eastside set gillnet Chinook salmon harvest composition in
- 1633 Upper Cook Inlet, Alaska, 2021. Alaska Department of Fish and Game, Fishery Data Series No. 22-
- 1634 06, Anchorage.
- Eskelin, T., and A. W. Barclay. 2022. Russian River early-run sockeye salmon run timing into the
  Kenai River, 2018–2020. Alaska Department of Fish and Game, Fishery Data Series No. 22-33,
  Anchorage.

Espinasse, B., Hunt, B.P.V., Finney, B.P., Fryer, J.K., Bugaev, A.V., Pakhomov, E.A., 2020. Using
stable isotopes to infer stock-specific high-seas distribution of maturing sockeye salmon in the North
Pacific. Ecology and Evolution 10, 13555–13570. https:// doi.org/10.1002/ece3.7022

1641 Estensen, J.L., Schmidt, S.N., Garcia, S., Gleason, C.M., Borba, B.M., Jallen, D.M., Padilla, A.J.,
1642 Hilton, K.M., 2018. Annual Management Report Yukon Area, 2016, Fishery Management Report
1643 No.18-14.

Evenson, D. F., C. Habicht, M. Stopha, A. R. Munro, T. R. Meyers, and W. D. Templin. 2018. Salmon
hatcheries in Alaska – A review of the implementation of plans, permits, and policies designed to
provide protection for wild stocks. Alaska Department of Fish and Game, Special Publication No. 1812, Anchorage.

Evenson, D. F., C. Habicht, M. Stopha, A. R. Munro, T. R. Meyers, and W. D. Templin. 2018. Salmon
hatcheries in Alaska — A review of the implementation of plans, permits, and policies designed to
provide protection for wild stocks. Alaska Department of Fish and Game, Special Publication No. 1812, Anchorage.

1652 **F** 

Fall, James A., Ronald T. Stanek, Brian Davis, Liz Williams, and Robert Walker. 2004. Cook Inlet
Customary and Traditional Subsistence Fisheries Assessment. Alaska Department of Fish and Game,
Division of Subsistence Technical Paper No. 285. Juneau.

- Farley, E.V., A. Starovoytov, S. Naydenko, R. Heintz, M. Trudel, C. Guthrie, L. Eisner, and J.R.
  Guyon. 2011. Implications of a warming eastern Bering Sea for Bristol Bay sockeye salmon. ICES
  Journal of Marine Science 68(6):1138-1146
- Farley, E.V., and J.H. Moss. 2009. Growth rate potential of juvenile chum salmon on the eastern
  Bering Sea shelf: an assessment of salmon carrying capacity. North Pacific Anadromous Fish
  Commission Bulletin 5:265-277.
- Farley, E.V., Jr., and M.Trudel. 2009. Growth rate potential of juvenile sockeye salmon in warmerand cooler years on the eastern Bering Sea shelf. Journal of Marine Biology 10.1155/2009/640215.
- Farley, E.V., Jr., J.M. Murphy, K. Cieciel, E.M. Yasumiishi, K. Dunmall, T. Sformo, and P. Rand.
  2020. Response of Pink salmon to climate warming in the northern Bering Sea. Deep-Sea Research II
- Farley, E.V., Jr., J.M. Murphy, M.D. Adkison, L.B. Eisner, J.H. Helle, J.H. Moss, and J. Nielsen.
  2007. Early marine growth in relation to marine-stage survival rates for Alaska Sockeye salmon.
  Fishery Bulletin 105:121-130.
- Farley, E.V., Jr., T.D. Beacham, and A.V Bugaev. 2018. Ocean ecology of sockeye salmon. Pges 319389 in R.J. Beamish, editor. The ocean ecology of Pacific salmon and trout. Amercian Fisheries
  Society, Bethesda, Maryland.

Farley, E.V., Murphy, J.M., Wing, B.W., Moss, J.H., Middleton, A., 2005. Distribution, migration
pathways, and size of western Alaska juvenile salmon along the eastern Bering Sea shelf. Alaska
Fishery Research Bulletin 11, 15–26.

Farley, E.V., Trudel, M., 2009. Growth Rate Potential of Juvenile Sockeye Salmon in Warmer and
Cooler Years on the Eastern Bering Sea Shelf. Journal of Marine Biology 2009, 1–10. https://
doi.org/10.1155/2009/640215

Feddern, M.L., E.R. Schoen, R. Shaftel, C.J. Cunningham, C. Chythlook, B.M. Connors, A.D.
Murdoch, V.R. vonBiela, B. Woods. 2023. Disciplines to understand the effects of changing climate
on Chinook salmon in the Arctic-Yukon Kuskokwim Region. Fisheries DOI: 10.1002/fish.10923.

Fellman, J.B., Hood, E., Dryer, W., Pyare, S., 2015. Stream Physical Characteristics Impact Habitat
Quality for Pacific Salmon in Two Temperate Coastal Watersheds. Plos One 10, e0132652. https://
doi.org/10.1371/journal.pone.0132652

Fellman, J.B., Nagorski, S., Pyare, S., Vermilyea, A.W., Scott, D., Hood, E., 2014. Stream temperature
response to variable glacier coverage in coastal watersheds of Southeast Alaska. Hydrological
Processes 28, 2062–2073. https://doi.org/10.1002/hyp.9742

Fergusson E, Miller T, McPhee MV, Fugate C, Schultz H (2020) Trophic responses of juvenile Pacific
salmon to warm and cool periods within inside marine waters of Southeast Alaska. Prog Oceanogr.
https://doi.org/10.1016/j.pocean.2020.102378

- Fergusson, E. A., M. V. Sturdevant, and J. A. Orsi. 2013. Trophic relationships amongjuvenile salmon during 16-year time series of climate variability in Southeast Alaska. North
- 1692 Pac Anadromous Fish Comm Tech Rep 9: 112–117

Fergusson, E., J. Murphy, and A. Gray. 2021. Southeast Alaska coastal monitoring survey: salmon
trophic ecology and bioenergetics, 2019. NPAFC Doc. 1949. 40 pp. National Oceanic and
Atmospheric Administration (NOAA), National Marine Fisheries Service (NMFS), Alaska Fisheries
Science Center, Auke Bay Laboratories, Ted Stevens Marine Research Institute (Available at
https://npafc.org).

Fergusson, E., W. Strasburger, J. Murphy, A. Piston, S. Heinl, and A. Gray. 2022. Southeast Alaska
Coastal Monitoring Survey May–July 2021. NPAFC Doc. 2021. 44 pp. National Oceanic and
Atmospheric Administration (NOAA), National Marine Fisheries Service (NMFS), Alaska Fisheries
Science Center, Auke Bay Laboratories, Ted Stevens Marine Research Institute, and Alaska
Department of Fish and Game (Available at https://npafc.org).

Finkle, H. 2023. Chignik watershed sockeye salmon run reconstruction and escapement goals. OralReport to the Alaska Board of Fisheries 2023. Alaska Department of Fish and Game.

- 1705 Finkle, H., Schaberg, K., Foster, M., and Polum, T. 2022 Review of Salmon Escapement Goals in
- the Chignik Management Area, 2020. Fishery Manuscript No. 22-05 Alaska Department of Fish andGame.
- 1708 Finkle, H., Schaberg, K., Foster, M., Wattum, M. and Polum, T. 2022 Review of Salmon
- 1709 Escapement Goals in the Alaska Peninsula and Aleutian Islands Management Areas, 2020. Fishery
- 1710 Manuscript No. 22-06 Alaska Department of Fish and Game

Finnegan, S.P., Svoboda, N.J., Schooler, S.L., and Belant, J.L. 2023. Phenological overlap of
terrestrial and marine food resources did not reduce salmon consumption by Kodiak brown bears.
Global Ecology and Conservation 45: e02506.

- Fish, T. M., and A. W. Piston. 2022. Hugh Smith Lake sockeye salmon stock assessment, 2021. Alaska
  Department of Fish and Game, Fishery Data Series No. 22-18, Anchorage.
- Flannery, B. G., P. A. Cran, J. H. Eiler, D. Beacham, N. A. DeCovich, W. D. Templin, O. L. Schleiand J. K. Wenburg. 2012. Comparison of Radiotelemetry and Microsatellites for Determining the
- Origin of Yukon River Chinook Salmon. North American Journal of Fisheries Management 32(4):
  720-730.
- Flannery, B. G., Wenburg, J.K., Gharrett, A.J., 2007. Evolution of mitochondrial DNA variation
  withina and among Yukon River chum salmon populations. Transactions of the American Fisheries
  Society 136, 902–910.
- Flannery, B.G., Wenburg, J.K., Gharrett, A.J., 2007. Variation of amplified fragment length
  polymorphisms in Yukon River chum salmon: population structure and application to mixed-stock
  analysis. Transactions of the American Fisheries Society 136, 911–925.
- Flemming, S. M., N. L. Zeiser, S. C. Heinl, C. S. Jalbert, and S. E. Miller. 2022. Stock assessment
  study of Chilkoot Lake sockeye salmon, 2020–2021. Alaska Department of Fish and Game, Fishery
  Data Series No. 22-31, Anchorage.
- Fox, E. K. C., T. D. Lawson, and M. D. Keyse. 2022. 2022 South Alaska Peninsula salmon annual
  management report and 2021 subsistence fisheries in the Alaska Peninsula, Aleutian Islands, and AtkaAmlia Islands management areas. Alaska Department of Fish and Game, Division of Commercial
  Fisheries, Fishery Management Report No. 22-32, Anchorage.
- Friedland, K D, Walker, R.V., Davis, N.D., Myers, K.W., Boehlert, G.W., Urawa, S., Ueno, Y., 2001.
  Open-ocean orientation and return migration routes of chum salmon based on temperature data from
- data storage tags. Marine Ecology Progress Series 216, 235–252.
- Frost, T.J., Yasumiishi, E.M., Agler, B.A., Adkison, M.D., McPhee, M.V., 2021. Density-dependent
  effects of eastern Kamchatka pink salmon (Oncorhynchus gorbuscha) and Japanese chum salmon (O.
- keta) on age-specific growth of western Alaska chum salmon. Fisheries Oceanography 30, 99–109.
- 1739 https://doi.org/10.1111/ fog.12505

- 1740 Fukushima, M., Smoker, W.W., 1998. Spawning Habitat Segregation of Sympatric Sockeye and Pink
- Salmon. Transactions of the American Fisheries Society 127, 253–260. https://doi.org/10.157 7/15488659(1998)127<0253:SHSOSS>2.0.CO;2
- Fukushima, M., Taylor, S.G., Smoker, W.W., 1997. Fry and smolt production of pink and sockeye
  salmon in the Auke Lake system, Southeast Alaska. Acta Hydrobiologica Sinica 21, 1–21.

1745 **G** 

- Garcia, S., Sewall, F., 2021. Fishery Data Series No . 21-05 Diet and Energy Density Assessment of
  Juvenile Chinook Salmon from Northeastern Bering Sea Trawl by.
- Garvin, M. R., C. M. Kondzela, P. C. Martin, B. Finney, J. Guyon, W. D. Templin, N. DeCovich, S.
  Gilk-Baumer, and A. J. Gharrett. 2013. Recent physical connections may explain weak genetic
  structure in western Alaskan chum salmon (Oncorhynchus keta) populations. Ecology and Evolution
  3(7): 2362-2377.
- Gazey, W.J., and K.K. English. 2000. Assessment of sockeye and pink salmon stocks in the northern
  boundary area using run reconstruction techniques, 1982-95. Canadian Technical Report of Fisheries
  and A guatic Sciences, 2220: 122p
- and Aquatic Sciences. 2320: 132p.
- Geiger, H. J. 2003. Sockeye salmon stock status and escpement goal for ReDoubt Lake in SoutheastAlaska. Alaska Department of Fish and Game, Regional Information Report No. 1J03-01.
- Geiger, H.J., Smoker, W.W., Zhivotovsky, L.A. and Gharrett, A.J., 1997. Variability of family size
  and marine survival in pink salmon (Oncorhynchus gorbuscha) has implications for conservation
  biology and human use. Canadian Journal of Fisheries and Aquatic Sciences, 54(11), pp.2684-2690.
- Geiger, H.J., Smoker, W.W., Zhivotovsky, L.A., Gharrett, A.J., 1997. Variability of family size and
  marine survival in pink salmon (Oncorhychus gorbuscha) has implications for conservation biology
  and human use. Canadian Journal of Fisheries and Aquatic Sciences 54, 2684–2690.
- Geiger, H.J., Wang, I., Malecha, P., Hebert, K., Smoker, W.W., Gharrett, A.J., 2007. What causes
  variability in pink salmon family size? Transactions of the American Fisheries Society 136, 1688–
  1698.
- Gharrett, A., Joyce, J., Smoker, W., 2013. Fine-scale temporal adaptation within a salmonid
  population: mechanism and consequences. Molecular Ecology 22, 4457–4469.
  https://doi.org/10.1111/mec.12400
- Gharrett, A.J., Shirley, S.M., 1985. A genetic examination of spawning methodology in a salmonhatchery. Aquaculture 47, 245–256.
- Gharrett, A.J., Smoker, W.W., 1991. Two generations of hybrids between even- and odd-year pink
  salmon (Oncorhynchus gorbuscha): a test for outbreeding depression? Canadian Journal of Fisheries
  and Aquatic Sciences 48, 1744–1749.

- Gilk-Baumer et al. (2017) Genetic stock composition of the commercial harvest of Chinook salmonin Copper River District, 2013-2017.
- 1776 Gilk-Baumer, S. E., S. D. Rogers Olive, D. K. Harris, S. C. Heinl, E. K. C. Fox, and W. D. Templin.
- 1777 2015. Genetic mixed stock analysis of sockeye salmon harvests in selected northern Chatham Strait
- 1778 commercial fisheries, Southeast Alaska, 2012-2014. Alaska Department of Fish and Game, Fishery
- 1779 Data Series No. 15-03, Anchorage.
- Gilk-Baumer, S., S. M. Turner, C. Habicht, and S. C. Heinl. 2013. Genetic stock identification of
  McDonald Lake sockeye salmon in selected Southeast Alaska fisheries, 2007-2009. Alaska
  Department of Fish and Game, Fishery Manuscript Series No. 13-04, Anchorage.
- Gilk, S.E., D.B. Molyneaux, T. Hamazaki, J.A. Pawluk and W.D. Templin. 2009. Biological and
  genetic characteristics of fall and summer chum salmon in the Kuskokwim River, Alaska. Pages 161in Krueger, C.C. and C.E. Zimmerman, editors. Pacific Salmon: ecology and management of
  western Alaska's populations. American Fisheries Society Symposium 70, Bethesda, MD.
- Gilk, S.E., Templin, W.D., Molyneaux, D.B., Hamazaki, T., Pawluk, J.A., 2005. Characteristics offall chum salmon Oncorhynchus keta in the Kuskokwim River drainage.
- Gilk, S.E., Wang, I.A., Hoover, C.L., Smoker, W.W., Taylor, S.G., 2004. Outbreeding depression in
  hybrids between spatially separated pink salmon, Oncorhynchus gorbuscha, populations: marine
  survival, homing ability, and variability in family size. Environmental Biology of Fishes 69, 287–297.
- Gisclair, B.R., 2009. Salmon bycatch management in the Bering Sea walleye pollock fishery: threats
  and opportunities in western Alaska. American Fisheries Society Symposium 70, 799–816.
- Graham, C. J., Sutton, T. M., Adkison, M. D., McPhee, M. V., & Richards, P. J. (2019). Evaluation
  of growth, survival, and recruitment of Chinook Salmon in Southeast Alaska Rivers. Transactions of
  the American Fisheries Society. 148, 243–259. https://doi.org/10.1002/tafs.10148
- Granath, K.L., Smoker, W.W., Gharrett, A.J., Hard, J.J., 2004. Effects on embryo development time
  and survival of intercrossing three geographically separate populations of Southeast Alaska coho
  salmon Oncorhynchus kisutch. Environmental Biology of Fishes 69, 299–306.
- Gray, A.K., Mccraney, W.T., Marvin, C.T., Kondzela, C.M., Nguyen, H.T., Guyon, J.R., 2011.
  Genetic Stock Composition Analysis of Chum Salmon Bycatch Samples from the 2007 Bering Sea
  Groundfish Fisheries. Fisheries (Bethesda).
- 1803 Grebmeier, Jacqueline M, Overland, J.E., Moore, S.E., Farley, E.V., Carmack, E.C., Cooper, L.W.,
  1804 Frey, K.E., Helle, J.H., McLaughlin, F.A., McNutt, S.L., 2006. A Major Ecosystem Shift in the
  1805 Northern Bering Sea. Science 311, 1461–1464.
- Grebmeier, Jacqueline M., Cooper, L.W., Feder, H.M., Sirenko, B.I., 2006. Ecosystem dynamics of
  the Pacific-influenced Northern Bering and Chukchi Seas in the Amerasian Arctic. Progress In
  Oceanography 71, 331–361. https://doi.org/10.1016/j.pocean.2006.10.001

- Griffiths, J.R. D.E. Schindler, L.S. Balistrieri, and G.T. Ruggerone. 2011. Effects of simultaneous
  climate change and geomorphic evolution on thermal characteristics of a shallw Alaskan lake.
  Limnology and Oceaongraphy 56(1):193-205
- 1812 Griffiths, J.R., D.E. Schindler, G.T. Ruggerone, and J.D. Bumgarner. 2014. Climate variation is
  1813 filtered differently among lakes to influence growth of juvenile sockeye salmon in an Alaskan
  1814 watershed. Oikos 0:1-12
- 1815 Guthrie, C., Nguyen, H., Guyon, J., 2014. Genetic Stock Composition Analysis of Chinook Salmon
  1816 Bycatch Samples from the 2012 Bering Sea and Gulf of Alaska Trawl Fisheries. NOAA Technical
  1817 Memorandum 33.
- 1818 Guthrie, C.M., Wilmot, R.L., 2004. Genetic structure of wild chinook salmon populations of Southeast
  1819 Alaska and northern British Columbia. Environmental Biology of Fishes 69, 81–93.
- 1820 Guthrie, C.M.G., Nguyen, H.T., Marsh, M., Guyon, J.R., 2019. Genetic Stock Composition Analysis
  1821 of Chinook Salmon Bycatch Samples from the 2017 Gulf of Alaska Trawl Fisheries.
- 1822 Guyon, J R, Guthrie III, C.M., Nguyen, H., 2010. Genetic stock composition analysis of Chinook
  1823 salmon bycatch samples from the 2007 "B" season and 2009 Bering Sea trawl fisheries., Report to the
  1824 North Pacific Fishery Management Council.
- 1825 Guyon, Jeffrey R, Kondzela, C., McCraney, T., Marvin, C., Martinson, E., 2010. Genetic Stock
  1826 Composition Analysis of Chum Salmon Bycatch Samples from the 2005 Bering Sea Groundfish
  1827 Fisheries, Report to the North Pacific Fishery Management Council.
- 1828 **H**
- Habicht, C., L. W. Seeb, K. W. Myers, E. V. Farley Jr., and J. E. Seeb. 2010. Summer-Fall distribution
  of stocks of immature sockeye salmon in the Bering Sea as revealed by single-nucleotide
  polymorphisms. Transactions of the American Fisheries Society 139(4): 1171-1191.
- Habicht, C., L.W. Seeb, K.W. Myers, E.V. Farley, and J.E. Seeb. 2010. Summer-fall distribution of
  stocks of immature sockeye salmon in the Bering Sea as revealed by single-nucleotide ploymorphisms.
  Transactions of the American Fisheries Society 139:1171-1191.
- Habicht, C., Seeb, L.W., and Seeb, J.E. 2007. Genetic and ecological divergence defines population
  structure of sockeye salmon populations returing to Bristol Bay, Alaska, and provides a toold for
  admixture analysis. Transactions of the American Fisheries Society 136:82-94. DOI: 10.1577/T06001.1
- Habicht, C., Seeb, L.W., Myers, K.W., Farley, E.V., Seeb, J.E., 2010. Summer-fall distribution of
  stocks of immature sockeye salmon in the Bering Sea as revealed by single-nucleotide polymorphisms.
  Transactions of the American Fisheries Society 139, 1171–1191.

Hagerman, G., Ehresmann, R., Shaul, L., 2018. Annual Management Report for the 2017 SoutheastAlaska/Yakutat Salmon Troll Fisheries.

Hagerman, G., Vaughn, M., Priest, J., 2021. Annual management report for the 2020 Southeast
Alaska/Yakutat salmon troll fisheries. Fishery Management Report No. 21-17.

Hagerman, G.T., D.K. Harris, J.T. Williams, et al. 2021. DRAFT-northern southeast Alaska kingsalmon stock status and action plan, 2021. RD6 Report to the Alaska Board of Fisheries.

Halas, G. and M. Cunningham. 2019. Nushagak River Chinook Salmon: Local and Traditional
Knowledge and Subsistence Harvests. Alaska Department of Fish and Game Division of Subsistence,
Technical Paper No. 453, Anchorage.

Halupka, K.C., Willson, M.F., Bryant, M.D., Everest, F.H., Gharrett, A.J., 2003. Conservation of
population diversity of Pacific salmon in southeast Alaska. North American Journal of Fisheries
Management 23, 1057–1086.

Hamazaki, T. and N. DeCovich. Application of the genetic mark-recapture technique for run size
estimation of Yukon River Chinook salmon. 2014. North American Journal of Fisheries Management
34: 276-286.

Hamazaki, T., Evenson, M., Fleischman, S.J., Schaberg, K.L., 2012. Spawner-recruit analysis and
escapement goal recommendation for Chinook salmon in the Kuskokwim River drainage.

Hamilton, K., Mysak, L.A., 1986. Possible Effects of the Sitka Eddy on Sockeye Oncorhynchus nerka)
and Pink Salmon (Oncorhynchus gorbuscha) Migration off Southeast Alaska. Canadian Journal of
Fisheries and Aquatic Sciences 43, 498–504.

Hard, J.J., Heard, W.R., 1999. Analysis of straying variation in Alaskan hatchery chinook salmon
(Oncorhynchus tshawytscha) following transplantation. Canadian Journal of Fisheries and Aquatic
Sciences 56, 578–589. https://doi.org/10.1139/f98-199

Hard, J.J. 2004. Evolution of Chinook salmon life history under size-selective harvest. In A. Hendry
and S. Sterns (eds), Evolution Illuminated: Salmon and their relatives, p. 315 – 337. Oxford, NY.

Hard, J.J., M.R. Gross, M. Heino, R. Hilborn, R.G. Kope, R. Law, and J.D. Reynolds. 2008.
Evolutionary consequences of fishing and their implications for salmon. Evolutionary Applications doi:1111.j.1752-4571.2008.00020.x

Haynie, A.C., Huntington, H.P., 2016. Strong connections, loose coupling: the influence of the Bering
Sea ecosystem on commercial fisheries and subsistence harvests in Alaska. Ecology and Society 21.
https://doi.org/10.5751/ES-08729-210406

Heard, W.R., 1978. Probable case of streambed overseeding - 1967 pink salmon, Oncorhnchus
gorbuscha, spawners and survival of their progeny in Sashin Creek, Southeastern Alaska. Fishery
Bulletin 76, 569–582.

Heard, W.R., 1998. Do hatchery salmon affect the North Pacific Ocean ecosystem ? NPAFC Bulletin1, 405–411.

Heard, W.R., 2012. Overview of salmon stock enhancement in southeast Alaska and compatibility
with maintenance of hatchery and wild stocks. Environmental Biology of Fishes 94, 273–283.
https://doi.org/10.1007/s10641-011-9855-6

Heard, W.R., Orsi, J., Heard, W.R., Taylor, S.G., Orsi, J.A., 2013. Survival and Early Marine
Migration of Enhanced Age-0 Sockeye Salmon Smolts Raised in Freshwater and Seawater at Auke
Creek, Southeast Alaska. North Pacific Anadromous Fish Comission Technical Report 9, 235–238.

Heard, W.R., Orsi, J.A., Wertheimer, A.C., Sturdevant, M.V., Murphy, J.M., Mortensen, D.G., Wing,
B.L., Celewycz, A.G., 2001. A synthesis of research on early marine ecology of juvenile Pacific
salmon in southeast Alaska. North Pacific Anadromous Fisheries Commission Technical Report 2, 3–
6.

Hebert, K.P., Goddard, P.L., Smoker, W.W., Gharrett, A.J., 1998. Quantitative genetic variation and
genotype by environment interaction of embryo development rate in pink salmon (Oncorhynchus
gorbuscha). Canadian Journal of Fisheries and Aquatic Sciences 55, 2048–2057.

Heinl, S. C., and A. W. Piston. 2009. Standardizing and automating the Southeast Alaska pink salmon
escapement index. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional
Information Report No. 1J09-06, Douglas.

Heinl, S. C., J. F. Koerner, and D. J. Blick. 2000. Portland Canal chum salmon coded wire tagging
project, 1988-1995. Alaska Department of Fish and Game, Division of Commercial Fisheries,
Regional Information Report No. 1J00-16, Juneau.

Heinl, S. C., R. L. Bachman, and K. Jensen. 2011. Sockeye salmon stock status and escapement goals
in Southeast Alaska. Alaska Department of Fish and Game, Special Publication No. 11-20, Anchorage.

Henry, E., 2021. Redoubt Lake Sockeye Monitoring and Lake Fertilization 2021 Annual Report. U.S.
Department of Agriculture, Forest Service, Sitka, Alaska.

Hillgruber, N., and C. E. Zimmerman. 2009. Estuarine Ecology of Juvenile Salmon in Western Alaska:
a Review. Pages 183–199. American Fisheries Society.

Hiroko Ikuta; Andrew R. Brenner; Anna Godduhn. 2013. Socioeconomic patterns in subsistence
salmon fisheries: historical and contemporary trends in five Kuskokwim River communities and
overview of the 2012 season. ADF&G Division of Subsistence, Technical Paper No. 382.

Hoffman, R., and T. Thynes. 2021. DRAFT-Klukshu River sockeye salmon stock status and actionplan, 2021. RC% report to the Alaska Board of Fisheries

Hoffman, S. H. 1982. Northern Southeastern Alaska pink salmon (Oncorhynchus gorbuscha) tagging
investigations, 1977-1980. Alaska Department of Fish and Game, Division of Commercial Fisheries,

- 1910 Information Leaflet No. 196, Juneau.
- 1911 Hoffman, S. H. 1983. Southern Southeastern Alaska pink salmon (Oncorhynchus gorbuscha) tagging
- 1912 investigations, 1981. Alaska Department of Fish and Game, Division of Commercial Fisheries,
- 1913 Technical Data Report No. 92, Juneau.
- Hoffman, S. H. 1983. Southern Southeastern Alaska pink salmon (Oncorhynchus gorbuscha) tagging
  investigations, 1981. Alaska Department of Fish and Game, Division of Commercial Fisheries,
  Technical Data Report No. 92, Juneau.
- Hoffman, S. H., L. Talley, and M. C. Seibel. 1983. 1982 U.S./Canada research pink and sockeye
  salmon tagging, interception rates, migration patterns, run timing, and stock intermingling in southern
  Southeast Alaska and Northern British Columbia. [in]: Final Report 1982 salmon research conducted
  in Southeast Alaska by the Alaska Department of Fish and Game in conjunction with joint U.S.Canada Interception investigations. Contract No. NASO-82-00134.
- Hoffman, Stephen H., Larry Talley, and M. C. Seibel. 1984. U.S./ Canada cooperative pink and
  sockeye salmon tagging, interception rates, migration pattern, run timing, and stock intermingling
  research in southern Southeast Alaska and northern British Columbia, 1982. Alaska Department of
  Fish and Game Technical Data Report No. 110.
- Hoffman, Stephen H., L. Talley, and M. C. Seibel. 1985. 1984 pink and chum salmon tagging, national
  contribution rates, migration patterns, run timing, and stock intermingling research in southern
  Southeast Alaska and northern British Columbia [in]: Final Report. 1984 salmon research conducted
  in Southeast Alaska by the Alaska Department of Fish and Game in conjunction with National Marine
  Fisheries Service Auke Bay Laboratory for joint U.S.-Canada Interception Studies. Contract No.
  WASC-84-00179.
- Holen, D., S.M. Hazell, and G Zimpelman, editors. 2015. The Harvest and Use of Wild Resources in
  Selected Communities of the Copper River Basin and East Glenn Highway, Alaska, 2013. Alaska
- 1934 Department of Fish and Game Division of Subsistence, Technical Paper No. 405. Anchorage.
- Holen, Davin, J.A. Fall, and R. La Vine. 2011. Customary and traditional use worksheet: salmon,
  Copper River District, Prince William Sound Management Area. Alaska Department of Game
  Division of Subsistence Special Publication No. BOF 2011-06. Anchorage.
- Hollowed, A.B., Barbeaux, S.J., Cokelet, E.D., Farley, E., Kotwicki, S., Ressler, P.H., Spital, C.,
  Wilson, C.D., 2012. Effects of climate variations on pelagic ocean habitats and their role in structuring
  forage fish distributions in the Bering Sea. Deep Sea Research Part II: Topical Studies in
  Oceanography 65–70, 230–250. https://doi. org/10.1016/j.dsr2.2012.02.008
- Hollowell, G., E. O. Otis, and E. Ford. 2023. 2022 Lower Cook Inlet area salmon annual management
  report. Alaska Department of Fish and Game, Fishery Management Report No. 23-04, Anchorage.

- 1944 Honea, J.M., Jorgensen, J.C., McClure, M.M., Cooney, T.D., Engie, K., Holzer, D., and Hilborn, R.
- 2009. Evaluating habitat effects on population status: influence of habitat restoration on spring-run
  Chinook salmon. Freshwater Biology 54 (7): 1576-1592. doi:10.1111/j.1365-2427.2009.02208.x
- Horne-brine, M.H., Warnke, D., Dubois, L., 2011. Fishery Data Series No . 11-16 Salmon Age and
  Sex Composition and Mean Lengths for the Yukon River Area , 2009 by.
- Howard, K. 2023 AYK Chum Salmon Marine Research Overview. Oral Report to Alaska Board ofFisheries. Alaska Department of Fish and Game
- Howard, K. G., K. M. Miller, and J. Murphy. 2017. Estuarine Fish Ecology of the Yukon River Delta,
  2014–2015. Page 76. Fishery Data Series, Alaska Department of Fish and Game, Anchorage, AK.
- Howard, K.G., and D.F. Evenson. 2010. Yukon River Chinook salmon comparative mesh size study.
  Fishery Data Series No. 10-92
- Howard, K.G., and V.v Biela. 2023. Adult spawners: a critical period for subarctic Chinook salmon
  in a changing climate. Global Change Biology. DOI: 10.1111/gcb.16610
- Howard, K.G., and von Biela, V. 2023. Adult spawners: a critical period for subarctic Chinook salmon
  in a changing climate. Global Change Biology 29: 1759-1773. DOI: 10.1111/gcb.16610
- Howard, K.G., Hayes, S.J., Evenson, D.F., 2009. Yukon River Chinook salmon stock status and action
  plan 2010; a report to the Alaska Board of Fisheries. Alaska Department of Fish and Game,
  Anchorage.
- Howard, K.G., J.M. Murphy, L.I. Wilson, J.H. Moss, and E.V. Farley, Jr. 2016. Size-selective
  mortality of Chinook salmon in relation to body energy after the first summer in nearshore marine
  habitats. North Pacific Anadromous Fish Commission Bulletin 6:1-11.
- Howard, K.G., Miller, K.M., Murphy, J., 2017. Estuarine fish ecology of the Yukon River Delta, 2014.
  Alaska Department of Fish and Game Fishery Data Series No. 17-16.
- Howe, E. Lance, Martin, Stephanie, 2009. Demographic Change, Economic Conditions, and
  Subsistence Salmon Harvests in Alaska 's Arctic-Yukon-Kuskokwim Region. American Fisheries
  Society Symposium 70, 433–461.
- Hunt Jr, G.L., Stabeno, P., Walters, G., Sinclair, E., Brodeur, R.D., Napp, J.M., Bond, N. a., 2002.
  Climate change and control of the southeastern Bering Sea pelagic ecosystem. Deep Sea Research Part
  II: Topical Studies in Oceanography 49, 5821–5853. https:// doi.org/10.1016/S0967-0645(02)003211
- 1974 Hunt, G.L., Coyle, K.O., Eisner, L.B., Farley, E.V., Heintz, R.A., Mueter, F., Napp, J.M., Overland,
- 1975 J.E., Ressler, P.H., Salo, S., Stabeno, J., 2011. Climate impacts on eastern Bering Sea foodwebs : a
- 1976 synthesis of new data and an assessment of the Oscillating Control Hypothesis. ICES Journal of
- 1977 Marine Science 68, 1230–1243.

- 1978 Hunt, G.L., Ressler, P.H., Gibson, G.A., De Robertis, A., Aydin, K., Sigler, M.F., Ortiz, I., Lessard,
- 1979 E.J., Williams, B.C., Pinchuk, A., Buckley, T., 2015. Euphausiids in the eastern Bering Sea: A
- 1980 synthesis of recent studies of euphausiid production, consumption and population control. Deep-Sea
- 1981 Research Part II: Topical Studies in Oceanography 134, 204–222. https://doi.org/10.1016/j.
- 1982 dsr2.2015.10.007

Huntington, H. P., S. L. Danielson, F. K. Wiese, M. Baker, P. Boveng, J. J. Citta, A. De Robertis, D.
M. S. Dickson, E. Farley, J. C. George, K. Iken, D. G. Kimmel, K. Kuletz, C. Ladd, R. Levine, L.
Quakenbush, P. Stabeno, K. M. Stafford, D. Stockwell, and C. Wilson. 2020. Evidence suggests
potential transformation of the Pacific Arctic ecosystem is underway. Nature Climate Change.

Hutchinson-Scarbrough, L. and D. Koster. 2021. Subsistence Harvest Assessment of Salmon and
Local Traditional Knowledge of Chinook Salmon in the Chignik Management Area, 2014–2016.
Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 462, Anchorage

- Hutchinson-Scarbrough, L. and Koster, D. 2021. Subsistence Harvest Assessment of Salmon andLocal Traditional Knowledge of Chinook Salmon in the Chignik Management Area, 2014-2016.
- 1992 Technical Paper No. 462 Alaska Department of Fish and Game
- Hutchinson-Scarbrough, L., D. Gerkey, G. Halas, C. Larson, L. A. Sill, J. M. Van Lanen, and M.Cunningham.

2020. Subsistence Salmon Networks in Select Bristol Bay and Alaska Peninsula Communities, 2016.
Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 459, Anchorage

Hyatt, K. D., and K. L. Mathias. 2005. Evaluation of hatchery versus wild sockeye salmon fry growth
and survival in two British Columbia Lakes. North American Journal of Fisheries Management
25:745-762.

- Hyer, K.E., Schleusner, C.J., 2005a. Chinook salmon age, sex, and length analysis from selected
  escapement projects on the Yukon River., Alaska Fisheries Technical Report Number 87.
- Hyer, K.E., Schleusner, C.J., 2005b. Chinook salmon age, sex, and length analysis from selected
  escapement projects on the Yukon River. Alaska Fisheries Technical Report Number 87 70.
- 2004 I
- Ianelli, J.N. and Stram, D.L., 2015. Estimating impacts of the pollock fishery bycatch on western
  Alaska Chinook salmon. ICES Journal of Marine Science, 72(4), pp.1159-1172.
- Ianelli, J.N., Gauvin, J., Stram, D.L., Haflinger, K., Stabeno, P., 2010. Temperature/depth datacollections on Bering Sea groundfish vessels to reduce bycatch.
- Inman, S.C., J. Esquible, M.L. Jones, W.R. Bechtol, and B. Connors. 2021. Opportunites and
  impediments for use of local data in the management of salmon fisheries. Ecology and Society 26(2):
  26

Irvine JR, Macdonald RW, Brown RJ, Godbout L, Reist JD, Carmack EC (2009) Salmon in the Arctic
and how they avoid lethal low temperatures. North Pacific Anadromous Fish Comm Bull 5: 39–50.

Ishida, Y., Azumaya, T., Fukuwaka, M., Davis, N., 2002. Interannual variability in stock abundance
and body size of Pacific salmon in the central Bering Sea Interannual variability in stock abundance
and body size of Pacific salmon in the central Bering Sea. Progress in Oceanography 55, 223–234.
https://doi.org/10.1016/ S0079-6611(02)00080-0

- 2018 **J**
- Jaenicke, H.W., Celewycz, A.G., 1994. Marine distribution and size of juvenile Pacific salmon in
  southeast Alaska and northern British Columbia. Fishery Bulletin 92, 79–90.
- Jaenicke, H.W., Celewycz, A.G., Bailey, J.E., Orsi, J.A., 1985. Paired open beach seines to study
  estuarine migrations of juvenile salmon. Marine Fisheries Review 46, 62–67.
- Jalbert, C. S., S. C. Heinl, A. Reynolds Manney, and K. R. Shedd. 2022. Commercial harvest of
  Klawock Lake sockeye salmon in the District 103 and 104 purse seine fisheries, Southeast Alaska,
  2025 2018–2021. Alaska Department of Fish and Game, Fishery Data Series No. 22-34, Anchorage.
- Jallen, D. M., Decker, S.K.S., and Hamazaki, T. 2017. Subsistence and personal use salmon harvests
  in the Alaska portion of the Yukon River drainage, 2013. Alaska Department of Fish and Game,
  Fishery Data Series No. 17-08, Anchorage.
- James S. Magdanz. 1981. Northern Bering Sea subsistence report. ADF&G Division of Subsistence,
   Technical Paper No. 4.
- Jasper J. R., C. Habicht, S. Moffitt, R. Brenner, J. Marsh, B. Lewis, E. Creelman Fox, Z. Grauvogel,
  S.D. Rogers Olive, and W. S. Grant. 2013. Source-sink estimates of genetic introgression show
  influence of hatchery strays on wild chum salmon populations in Prince William Sound, Alaska. PLoS
  ONE 8(12):e81916.
- Jasper, J. R., N. DeCovich, W. D. Templin. 2012. Western Alaska Salmon Stock Identification
  Program Technical Document 4: Status of the SNP baseline for chum salmon. (PDF 1,013 kB) Alaska
  Department of Fish and Game, Regional Information Report 5J12-09, Anchorage
- Jasper, J.R., Evenson, D.F., 2006. Length-girth, length-weight, and fecundity of Yukon River Chinook
  salmon Oncorhychus tshaytscha., Fishery Data Series. Alaska Department of Fish and Game,
  Anchorage.
- Johnson, G., Kondzela, C., Whittle, J., Miller, K., Guyon, J., 2019. Genetic Characterization of
- Juvenile Chum Salmon (Oncorhynchus keta) Migrating out of the Yukon River Delta. Technical
   Report 51–53. https://doi.org/10.23849/npafctr15/51.53.

Johnson, G.C., Stabeno, Phyllis.J., 2004. The Bering Slope Current system revisited. Journal of
Physical Oceanography 34, 384–398. https://doi.org/10.1175/1520-0485(2004)034<0384</li>
:TBSCSR>2.0.CO;2

Johnson, S.W., Murphy, M.L., Csepp, D.J., Harris, P.M., Thedinga, J.F., 2003. A survey of fish
assemblages in eelgrass and kelp habitats of southeastern Alaska. U. S. Department of Commerce.

Johnson, S.W., Thedinga, J.F., 2005. Fish use and size of eelgrass meadows in southeastern Alaska: a
baseline for long-term assessment of biotic change. Northwest Science 79, 141–155.

Johnson, T. A., S. C. Heinl, and H. J. Geiger. 2005. McDonald Lake: Stock status report. Alaska
Department of Fish and Game, Fishery Manuscript No. 05-07, Anchorage.

Jones, Bronwyn and M. Kukkonen. 2017. Local and Traditional Knowledge of Abundance of Chinook
Salmon in the Kenai River. Alaska Department of Fish and Game Division of Subsistence, Technical
Paper No. 431, Anchorage

Jones, Bronwyn and Margaret Cunningham. 2020. The Harvest and Use of Salmon by Residents of
King Salmon, Naknek, and South Naknek, Alaska, 2017 and 2018. Alaska Department of Fish and
Game Division of Subsistence, Technical Paper No. 470, Anchorage.

Jones, Bronwyn and Margaret Cunningham. 2020. The Harvest and Use of Wild Resources in Port
Heiden, Alaska, 2018. Alaska Department of Fish and Game Division of Subsistence, Technical Paper
No. 465: Anchorage.

Jones, Bronwyn E. and David Koster. 2018. Subsistence Harvests and Uses of Salmon in Tyonek,
2063 2015 and 2016. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No.
2064 439, Anchorage.

Jones, Bronwyn, Cunningham, Margaret, and Koster, David. 2019. Subsistence Harvest Assessment
and Biological Sampling of Chinook Salmon in the Togiak River Drainage. Alaska Department of
Fish and Game Division of Subsistence, Technical Paper No. 454, Anchorage.

Jones, E.L., Quinn, T.J., Van Alen, B.W., 1998. Observer Accuracy and Precision in Aerial and Foot
Survey Counts of Pink Salmon in a Southeast Alaska Stream. North American Journal of Fisheries
Management 18, 832–846. https://doi.org/10.1577/1548-8675(1 998)018<0832:OAAPIA>2.0.CO;2

Jorgensen, J.C., C. Nicol, C. Fogel, and T.J. Beechie. 2021. Identifying the potential of anadromous
salmonid habitat restoration with life cycle models. PLOS One 16(9): e0256792. https://doi.org/

2073 10.1371/journal.pone.0256792

Josephson, R., A. Wertheimer et al. 2021. Proportions of hatchery fish in escapements of summer-run
 chum salmon in southeast Alaska, 2013-2015. North American Journal of Fisheries Management. DOI
 10.1002/nafm.10580

Josephson, R., Wertheimer, A., Gaudet, D., Knudsen, E., Adams, B., Bernard, D., Heinl, S., Piston,
A., and Templin, W., 2021. Proportions of Hatchery Fish in Escapements of Summer-Run Chum
Salmon in Southeast Alaska, 2013-2015. North American Journal of Fisheries Management 41:724 738, 2021

Josephson, R., Wertheimer, A., Gaudet, D., Knudsen, E.E., Adams, B., Bernard, D.R., Heinl, S.C.,
Piston, A.W. and Templin, W.D., 2021. Proportions of hatchery fish in escapements of summer-run
Chum Salmon in Southeast Alaska, 2013–2015. North American Journal of Fisheries Management,
41(3), pp.724-738.

Joy et al. (2021) Escapement goal review of Copper and Bering Rivers and Prince William SoundPacific salmon stocks, 2020.

Joy et al. (2021) Run reconstruction, spawner-recruit analysis, and escapement goal recommendation
for Chinook salmon in the Copper River.

Joy, P. J., S. B. Haught, R. E. Brenner, S. Miller, J. W. Erickson, J. W. Savereide, and T. R. McKinley.
2090 2021. Escapement goal review of Copper and Bering Rivers and Prince William Sound Pacific salmon
stocks, 2020. Alaska Department of Fish and Game, Fishery Manuscript No. 21-02, Anchorage.

Joy, P., J. W. Savereide, M. Tyers, and S. J. Fleischman. 2021. Run reconstruction, spawner–recruit
analysis, and escapement goal recommendation for Chinook salmon in the Copper River. Alaska
Department of Fish and Game, Fishery Manuscript No. 21-01, Anchorage.

Julie Raymond-Yakoubian, Brenden Raymond-Yakoubian, Catherine Moncrieff. The incorpoation of
 traditional knowledge into Alaska federal fisheries management. Mar Policy, 78 (2017) 132-142

2097 K

K.M. Dunmall, J.D. Reist, E.C. Carmack, J.A. Babaluk, M.P. Heide-Jørgensen, and M.F. Docker.
2099 2013. Pacific Salmon in the Arctic: Harbingers of Change. In: F.J. Mueter, D.M.S. Dickson, H.P.
Huntington, J.R. Irvine, E.A.Logerwell, S.A. MacLean, L.T. Quakenbush, and C. Rosa (eds.),
Responses of Arctic Marine Ecosystems to Climate Change. Alaska Sea Grant, University of Alaska
Fairbanks. doi:10.4027/ ramecc.2013.07

Kaeriyama, M., Nakamura, M., Edpalina, R., Bower, J.R., Yamaguchi, H., Walker, R.V., and Myers,
K.W. 2004. Change in feeding ecology and trophic dynamics of Pacific salmon (Oncorhynchus spp.)
in the central Gulf of Alaska in relation to climate events. Fisheries Oceanography 13 (3): 197-207.

Kaga, T., Sato, S., Azumaya, T., Davis, N.D., Fukuwaka, M., 2013. Lipid content of chum salmon
Oncorhynchus keta affected by pink salmon O. gorbuscha abundance in the central Bering Sea.
Marine Ecology Progress Series 478, 211–221. https://doi.org/10.3354/ meps10179

Karpenko, V.I., Volkov, A.F., Koval, M.V., 2007. Diets of Pacific salmon in the Sea of Okhotsk,
Bering Sea, and Northwest Pacific Ocean. N. Pac. Anadr. Fish Comm. Bull 4, 105–116.

- Keating, J.M., D. Koster, and J.M. Van Lanen. 2020. Recovery of a Subsistence Way of Life:
  Assessments of Resource Harvests in Cordova, Chenega, Tatitlek, Port Graham, and Nanwalek,
  Alaska since the Exxon Valdez Oil Spill. Alaska Department of Fish and Game Division of
  Subsistence, Technical Paper No. 471, Anchorage.
- 2115 Khen, G.V., Zavolokin, A.V., Хен, Г.В., Заволокин, А.В., 2015. Change in the water circulation and
  2116 its implications for distribution and abundance of salmons in the western Bering Sea. TINRO News
  2117 95–115.
- Kishi, M.J., Kaeriyama, M., Ueno, H., Kamezawa, Y., 2010. The effect of climate change on the
  growth of Japanese chum salmon (Oncorhynchus keta) using a bioenergetics model coupled with a
  three-dimensional lower trophic ecosystem model (NEMURO). Deep-Sea Research Part Ii-Topical
  Studies in Oceanography 57, 1257–1265. https://doi.org/10.1016/j.dsr2.2009.12.013
- Kline, T.C., Goering, J.J., Mathison, O.A., Poe, P.H., Parker, P.L., 1990. Recycling of elements
  transported upstream by runs of Pacific salmon: I. 15-N and 13-C evidence in Sashin Creek,
  southeastern Alaska. Canadian Journal of Fisheries and Aquatic Sciences 47, 136–144.
- Knudsen, E.E., Rand, P.S., Gorman, K.B., Bernard, D.R. and Templin, W.D., 2021. Hatchery-origin
  stray rates and total run characteristics for Pink Salmon and Chum Salmon returning to prince william
  sound, Alaska, in 2013–2015. Marine and Coastal Fisheries, 13(1), pp.41-68.
- Kocan R, Hershberger P, Sanders G, Winton J (2009) Effects of temperature on disease progression
  and swimming stamina in Ichthyophonus-infected rainbow trout, Oncorhynchus mykiss Walbaum. J
  Fish Dis 32: 835–843.
- Kocan, R., Hershberger, P., 2006. Differences in Ichthyophonous prevalence and infection severity
  between upper Yukon River and Tanana River chinook salmon, Oncorhynchus tshawytscha
  (Walbaum) stocks. Journal of Fish Diseases 29, 497–503.
- Kocan, R., Hershberger, P., and Winton, J. 2004. Ichthyophoniasis: An emerging disease of Chinook
  salmon in the Yukon River. Journal of Aquatic Animal Health, 16:2, 58-72, DOI: 10.1577/H03-068.1
- Kohan, M. L., F. J. Mueter, J. A. Orsi, and M. V. McPhee. 2017. Variation in size, condition, and
  abundance of juvenile chum salmon (Oncorhynchus keta) in relation to marine factors in Southeast
  Alaska. Deep Sea research Part II, https://doi.org/10.1016/j. dsr2.2017.09.005
- Kondzela, C. M., C. M. Guthrie, S. L. Hawkins, C. D. Russell, and J. H. Helle. 1994. Genetic
  relationships among chum salmon populations in southeast Alaska and northern British Columbia.
  Canadian Journal of Fisheries and Aquatic Sciences 51(Suppl. 1):50-64.
- Kondzela, C.M., J.A. Whittle, CIT. Marvin et al. 2016. Genetic analysis identifies consistent
  proportions of seasonal life history types in Yukon River juvenile and adulst Chum salmon. North
  Pacific Anadromous Fish Commission Bulletin 6:439-450

- Kondzela, C.M., M. Garvin, R. Riley, J. Murphy, J. Moss, S. Adam Fuller, and A. Gharrett. 2009.
  Preliminary genetic analysis of juvenile Chum salmon from the Chukchi Sea and Bering Strait. North
  Pacific Anadromous Fish Commission Bulletin 5:25-27.
- 2148 Kondzela, C.M., Whittle, J.A., Vulstek, S.C., Nguyen, H.T., Guyon, J.R., 2017. Genetic stock composition analysis of Chum Salmon from the prohibited species catch of the 2015 Bering Sea 2149 trawl fishery and Gulf of Alaska groundfish fisheries. 2150 Walleye Pollock 49. https://doi.org/10.7289/V5/TM-AFSC-314 2151
- Kondzela, C.M., Whittle, J.A., Vulstek, S.C., Nguyen, H.T., Guyon, J.R., 2017. Genetic stock 2152 2153 composition analysis of Chum Salmon from the prohibited species catch of the 2015 Bering Sea Pollock trawl fishery and Gulf of Alaska groundfish fisheries. 2154 Walleve 49. https://doi.org/10.7289/V5/TM-AFSC-314 2155
- Kondzela, C.M., Whittle, J.A., Yates, D., Vulstek, S.C., Nguyen, H.T., Guyon, J.R., 2016. Genetic
  Stock Composition Analysis of Chum Salmon from the Prohibited Species Catch of the 2014 Bering
  Sea Walleye Pollock Trawl Fishery and Gulf of Alaska Groundfish Fisheries. NOAA Technical
  Memorandum NMFS AFSC i–48.
- Kondzela, C.M., Whittle, J.A., Yates, D., Vulstek, S.C., Nguyen, H.T., Guyon, J.R., 2016b. Genetic
  stock composition analysis of Chum Salmon from the prohibited species catch of the 2014 Bering Sea
  Walleye Pollock trawl fishery and Gulf of Alaska groundfish fisheries. U. S. Department of
  Commerce, NOAA Tech Memo. https://doi. org/10.7289/V5/TM-AFSC-314
- Kovach, R. P., A. J. Gharrett, and D. A. Tallmon. 2015. Genetic change for earlier migration timing
  in a pink salmon population. Proc. R. Soc. B doi:10.1098/rspb.2012.1158.
- Kovach, R.P., Ellison, S.C., Pyare, S., Tallmon, D.A., 2015. Temporal patterns in adult salmon
  migration timing across Southeast Alaska. Global Change Biology 21, 1821–1833.
- 2168 KRITFC Graphics & Explanations
- Krkosek, M., R. Hilborn, R. M. Peterman, and T. Quinn. 2011. Cycles, stochasticity and density
  dependence in pink salmon population dynamics. Proceedings of the Royal Society B 278:2060–2068.
- Krueger, Charles & Zimmerman, Christian. 2009. Pacific salmon: ecology and management ofwestern Alaska's populations.
- 2173 L
- La Vine, R., M. Kukkonen, B. Jones, and G. Zimpelman. 2013. Subsistence harvests and uses of
- 2175 wild resources in Copper Center, Slana/Nabesna Road, Mentasta Lake, and Mentasta Pass, Alaska,
- 2176 2010. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 380.
- 2177 Anchorage, Alaska.

- 2178 LaCroix, J.J., Wertheimer, A.C., Orsi, J.A., Sturdevant, M.V., Fergusson, E.A., Bond, N.A., 2009. A
- 2179 top-down survival mechanism during early marine residency explains coho salmon year-class strength
- 2180 in southeast Alaska. Deep-Sea Research Part II: Topical Studies in Oceanography 56, 2560–2569.
- 2181 https://doi.org/10.1016/j. dsr2.2009.03.006
- Ladd, C., 2014. Seasonal and interannual variability of the Bering Slope Current. Deep Sea Research
  Part II: Topical Studies in Oceanography 109, 5–13. https://doi.org/10.1016/j.dsr2.2013.12.005
- Landingham, J., Sturdevant, M., Brodeur, R., 1998. Feeding habits of juvenile Pacific salmon in
  marine waters of southeastern Alaska and northern British Columbia. Fishery Bulletin 96, 285–302.
- Langdon, S. 2021. K'iis Xaadas relations with sockeye salmon: contemporary efforts at constructing
  a neo-traditional regime of stewardship. Maritime Studies (20) 157-173
- Langdon, S.J. Traditional knowledge and harvesting of salmon by Huna and Hinya Tlingit. 2006. Final
  Report, Fisheries Information Service Porject 02-104
- Langdon, S.J. Traditional knowledge and harvesting of salmon by Huna and Hinya Tlingit. 2006. Final
   Report, Fisheries Information Service Project 02-104
- Langdon, S.J., 2006. Tidal pulse fishing: Selective traditional Tlingit salmon fishing techniques on the
  west coast of the Prince of Wales Archipelago., in: Traditional Ecological Knowledge and Natural
  Resource Management. University of Nebraska Press, pp. 21–46.
- Langdon, S.J., 2007. Sustaining a relationship: inquiry into the emergence of a logic of engagement
  with salmon amon the Southern Tlingits, in: Native Americans and the Environment: Perspectives on
  the Ecological Indian. University of Nebraska Press.
- Langdon, S.J., 2018. Approaching Leviathan : Efforts to Establish Small-Scale , Community Based
  Commercial Salmon Fisheries in Southeast Alaskan Indigenous Communities, in: Fisheries, Quota
  Management and Quota Transfer, MARE Publication Series. Springer International Publishing, pp.
  197–214.
- Larsen, C. F., R. J. Motyka, J. T. Freymueller, K. A. Echelmeyer, and E. R. Ivins. 2005. Rapid
  viscoelastic uplift in southeast Alaska caused by post-Little Ice Age glacial retreat. Earth and Planetary
  Science Letters 237 (2005) 548–560.
- Larsen, C. F., R. J. Motyka, A. A. Arendt, K. A. Echelmeyer, and P. E. Geissler. 2007. Glacier changes
  in southeast Alaska and northwest British Columbia and contribution to sea level rise. JOURNAL OF
  GEOPHYSICAL RESEARCH, VOL. 112, F01007, doi:10.1029/2006JF000586.
- Larson W. A., L. W. Seeb, M. V. Everett, R. K. Waples, W. D. Templin, and J. E. Seeb. 2014.
  Genotyping by sequencing resolves shallow population structure to inform conservation of Chinook
  salmon (Oncorhynchus tshawytscha). Evolutionary Applications 7(3): 355-369.

Larson, W. A., F. M. Utter, K. W. Myers, W. D. Templin, J. E. Seeb, C. M. Guthrie III, A. V. Bugaev,
and L. W. Seeb. 2013. Single-nucleotide polymorphisms reveal distribution and migration of Chinook
salmon (Oncorhynchus tshawytscha) in the Bering Sea and North Pacific Ocean. Canadian Journal of
Fisheries and Aquatic Sciences 70: 128-141.

Larson, W. A., J. E. Seeb, C. E. Pascal, W. D. Templin, and L. W. Seeb. 2014. Single-nucleotide
polymorphisms (SNPs) identified through genotyping-by-sequencing improve genetic stock
identification of Chinook salmon (Oncorhynchus tshawytscha) from western Alaska. Canadian
Journal of Fisheries and Aquatic Sciences 71(5):698-708.

Larson, W., Utter, F., Myers, K.W., Templin, W.D., Seeb, J.E., Guthrie III, C.M., Bugaev, A.V., Seeb,
L.W., 2013. Single-nucleotide polymorphisms reveal distribution and migration of Chinook salmon
(Oncorhynchus tshawytscha) in the Bering Sea and North Pacific Ocean. Canadian Journal of
Fisheries and Aquatic Sciences 70, 128–141. https://doi.org/dx.doi.org/10.1139/cjfas-2012-0233
Published

Larson, W.A., F.M. Utter, K.W. Myers, W.D. Templin, J.E. Seeb, C.M. Guthrie III, A.V. Bugaev, and
L.W. Seeb. 2013. Single-nucleotide ploymorphisms reveal distribution and migration of Chinook
salmon in the Bering Sea and North Pacific Ocean. Canadian Journal of Fisheries and Aquatic Sciences
70:128-141.

LaVine, R., M.J. Lisac and P. Coiley-Kenner. 2007. Traditional ecological knowledge of 20th century
ecosystems and fish populations in the Kuskokwim Bay Region. U.S. Fish and Wildlife Service,
Office of Subsistence Management, Fisheries Resource Monitoring Program (Project no. FIS 04 –
351) Anchorage, Alaska.

2232 Lee, E., Dann, T., Hoyt, H., 2021. Yukon River Chinook Genetic Baseline Improvements.

Leon, J.M., McPhee, M.V., 2013. Freshwater Growth and Recruitment in Two Western Alaskan
Populations of Chinook Salmon. NPAFC Technical Report 9, 229–231.

2235 Levi, T., Allen, J.M., Bell, D., Joyce, J., Russell, J.R., Tallmon, D.A., Vulstek, S.C., Yang, C., Yu,

D.W., 2019. Environmental DNA for the enumeration and management of Pacific salmon. Molecular
 Ecology Resources 19, 597–608. https://doi.org/10.1111/1755-0998.12987

Lewis B., W. S. Grant, R. E. Brenner, and T. Hamazaki. 2015. Changes in size and age of Chinook
salmon Oncorhynchus tshawytscha returning to alaska. PLoS ONE 10(6): e0130184. 17 pp.
doi:10.1371/journal.pone.0130184.

Lewis, B.A., Zadina, T.P.C.N.-RIR. 1J. 2001. 39, 2001. The history of the subsistence and commercial
fisheries, stock assessment and enhancement activities, and watershed disturbances in the Klawock
Lake drainage on Prince of Wales Island.

Liller, Z.W., Brodersen, A.R., Clark, J.N., 2013. Regional Information Report 3A13-01 Salmon Age
, Sex , and Length Catalog for the Kuskokwim Area , 2010 and 2011 Annual Report for Project 10303 by.

Litzow, M. a, Mueter, F.J., Hobday, A.J., 2014. Reassessing regime shifts in the North Pacific:
incremental climate change and commercial fishing are necessary for explaining decadal-scale
biological variability. Global change biology 20, 38–50. https://doi.org/10.1111/gcb.12373

- Litzow, M.A., 2017. Indications of hysteresis and early warning signals of reduced community
  resilience during a Bering Sea cold anomaly. Marine Ecology Progress Series 571, 13–28.
- Loewen, M., and L. Henslee. 2017. The 2016 Chignik River sockeye salmon smolt outmigration.
  Alaska Department of Fish and Game, Fishery Data Series No. 17-11, Anchorage.
- Loring, P. a, Gerlach, C., 2010. Food Security and Conservation of Yukon River Salmon: Are We
  Asking Too Much of the Yukon River? Sustainability 2, 2965–2987.
  https://doi.org/10.3390/su2092965
- Lum, J.L., Fair, L., 2018. Chilkat River and King Salmon River King Salmon Stock Status and ActionPlan, 2018.
- 2259 M
- MacNeil, M.A., Graham, N.A.J., Cinner, J.E., Dulvy, N.K., Loring, P.A., Jennings, S., Polunin,
  N.V.C., Fisk, A.T., McClanahan, T.R., 2010. Transitional states in marine fisheries: adapting to
  predicted global change. Philosophical Transactions of the Royal Society B: Biological Sciences 365,
  3753–3763.
- Magdanz, J. S. and C. Utermohle. 1997. The subsistence salmon fishery in the Norton Sound, Port
  Clarence and Kotzebue Districts, 1994. Alaska Department of Fish and Game, Division of Subsistence
  Technical Paper No. 237, Juneau.
- Magdanz, J. S., E Trigg, A. Ahmasuk, P. Nanouk, D. S. Koster and K. R. Kamletz. 2005. Patterns and
  trends in subsistence salmon harvests, Norton Sound and Port Clarence, 1994-2003. Alaska
  Department of Fish and Game, Division of Subsistence Technical Paper No. 294, Juneau.
- Magdanz, J. S., H. Smith, N. Braem and P. Fox and D. S. Koster. 2011. Patterns and trends in
  subsistence fish harvests, Northwest Alaska, 1994-2004. Alaska Department of Fish and Game,
  Divsion of Subsistence Technical Paper No. 366, Kotzebue.
- Magdanz, J. S., S. Tahbone, A. Ahmasuk, D. S. Koster and B. L. Davis. 2007. Customary trade and
  barter in subsistence fish in the Seward Peninsula area, Alaska. Alaska Department of Fish and Game,
  Division of Subsistence Technical Paper No. 328, Juneau.
- Magdanz, J. S., S. Tahbone, K. R. Kamletz. and A. Ahmasuk. 2005. Subsistence salmon fishing by
  residents of Nome, Alaska, 2001. Alaska Department of Fish and Game, Division of Subsistence
  Technical Paper No. 274, Juneau.

- Mahnken, C., Ruggerone, G., Waknitzl, W., Flagg, T., 1998. A Historical Perspective on Salmonid
  Production from Pacific Rim Hatcheries. North Pacific Anadromous Fish Commission Bulletin 1, 38–
  53.
- Malick, M. J., M. D. Adkison, A. C. Wertheimer. 2009. Variable effects of biological and
  environmental processes on coho salmon survival in southeast Alaska. Trans Am Fish Soc 138: 846860.
- Malison et al. (2016), Do beaver dams reduce habitat connectivity and salmon productivity in
  expansive river floodplains? PeerJ 4:e2403; DOI 10.7717/peerj.2403
- Manhard, C. V., J. E. Joyce, W. W. Smoker, and A. J. Gharrett. 2016. Ecological factors influencing
  lifetime productivity of pink salmon (Oncorhynchus gorbuscha) in an Alaskan stream. Can. J. Fish.
  Aquat. Sci. 74: 1325–1336.
- Manhard, C.V., Adkison, M.D., Hard, J.J., Smoker, W.W., Gharrett, A.J., 2018. Local adaptation of
   phenology revealed in outcrosses between spawning segments of a salmonid population. Molecular
   Ecology 27, 4698–4710. https://doi.org/10.1111/mec.14908
- Manhard, C.V., Joyce, J.E., Gharrett, A.J., 2017. Evolution of phenology in a salmonid population: a
   potential adaptive response to climate change. Canadian Journal of Fisheries and Aquatic Sciences 74,
   1519–1527. https://doi.org/10.1139/cjfas-2017-0028
- Manhard, C.V., Joyce, J.E., Smoker, W.W., Gharrett, A.J., 2017. Ecological factors influencing
  lifetime productivity of pink salmon (Oncorhynchus gorbuscha) in an Alaskan stream. Canadian
  Journal of Fisheries and Aquatic Sciences 74, 1325–1336. https://doi.org/10.1139/cjfas-2016-0335
- Manhard, C.V., Joyce, J.E., Smoker, W.W., Gharrett, A.J., 2017. Ecological factors influencing
  lifetime productivity of pink salmon (Oncorhynchus gorbuscha) in an Alaskan stream. Canadian
  Journal of Fisheries and Aquatic Sciences 74, 1325–1336. https://doi.org/10.1139/cjfas-2016-0335
- Manishin, K.A., Cunningham, C.J., Westley, P.A. and Seitz, A.C., 2021. Can late stage marine
  mortality explain observed shifts in age structure of Chinook salmon?. Plos one, 16(2), p.e0247370.
- Manishin, K.A., Cunningham, C.J., Westley, P.A.H., Seitz, A.C., 2021. Can late stage marine
  mortality explain observed shifts in age structure of Chinook salmon? PLoS ONE 16, 1–14. https://doi.
  org/10.1371/journal.pone.0247370
- Mantua, N.J., N.G. Taylor, G.T. Ruggerone. K.W. Myers, et al. 2009. The salmon MALBEC Project:
  A North Pacific-scale study to support salmon conservation planning. North Pacific Anadromous Fish
  Commission Bulletin 5:333-354.
- Manzer, J.I. 1964. Preliminary observations on the vertical distribution of Pacific salmon (Genus
  Oncorhynchus) in the Gulf of Alaska. Journal of the Fisheries Board of Canada 21(5): 891-903. https://
  doi.org/10.1139/f64-086

- 2313 Marchioni, M. A., J. F. Fall, B. Davis, and G. Zimpleman. 2016. Kodiak City, Larsen Bay and Old
- 2314Harbor: An Ethnographic Study of Traditional Subsistence Salmon Harvests and Uses. Alaska
- 2315 Department of Fish and Game Division of Subsistence, Technical Paper No. 418, Anchorage."
- Marston, B., and A. Frothingham. 2022. Upper Cook Inlet commercial fisheries annual management
  report, 2021. Alaska Department of Fish and Game, Fishery Management Report No. 22-16,
  Anchorage.
- Martin, D.J., Whitmus, C.J., Hachmeister, L.E., Volk, E.C., Schroder, S.L., 1987. DISTRIBUTION
  AND SEASONAL ABUNDANCE OF JUVENILE SALMON AND OTHER FISHES IN THE
  YUKON DELTA b.
- Martinson, E.C., J.H. Helle, D.L. Scarnecchia, and H.H. Stokes. 2008. Density-dependent growth of
  Alaska sockeye salmon in relation to climate-ocean regimes, poulation abundance, and body size,
  1925 to 1998. Marine Ecology Progress Series 370:1 18.
- Maschmann, G.F., 2011. Abundance and Run Timing of Adult Pacific Salmon in the East Fork
  Andreafsky River, Yukon Delta National Wildlife Refuge, Alaska Number, Alaska Fisheries Data
  Series.
- Matarese, A.C., Blood, D.M., Picquelle, S.J., Benson, J.L., 2003. Atlas of abundance and distribution
  patterns of ichthyoplankton from the northeast Pacific Ocean and Bering Sea ecosystems based on
  research conducted by the Alaska Fisheries Science Center (1972-1996). NOAA Professional Paper
  NMFS 1.
- McConnell, C.J., Westley, P.A.H., McPhee, M.V., 2018. Differences in fitness-associated traits
  between hatchery and wild chum salmon despite long-term immigration by strays. Aquaculture
  Environment Interactions 10, 99–113. https://doi.org/10.3354/aei00261
- McCraney, W.T., E.V. Farley, C.M. Kondzela, S.V. Naydenko, A.N. Starovoytov, and J.R. Guyon.
  2012. Genetic stock identification of overwintering chum salmon in the North Pacific Ocean.
  Environmental Biology of Fishes 94:663-668
- McGregor, A.J., Van Alen, B.W., Alen, V., 1987. ABUNDANCE , AGE , AND SEX
  COMPOSITIONS OF CHINOOK , SOCKEYE , COHO , AND CHUM SALMON CATCHES AND
  ESCAPEMENTS I N SOUTHEAST ALASKA IN BY : Commissioner. ADF&G Technical Data
  Report 200.
- McKinley, T., N. DeCovich, J. W. Erickson, T. Hamazaki, R. Begich, and T. L. Vincent. 2020. Review
  of salmon escapement goals in Upper Cook Inlet, Alaska, 2019. Alaska Department of Fish and Game,
  Fishery Manuscript No. 20-02, Anchorage.
- McKinney, G.J., Barry, P.D., Pascal, C., Seeb, J.E., Seeb, L.W., McPhee, M.V., 2022. A New
  Genotyping-in-Thousands-by-Sequencing Single Nucleotide Polymorphism Panel for Mixed-Stock
  Analysis of Chum Salmon from Coastal Western Alaska. North American Journal of Fisheries
  Management 42, 1134–1143. https://doi.org/10.1002/ nafm.10805

McMillan, J.R., B. Morrison, N. Chambers, G. Ruggerone, L. Bernatches, J. Stanford, and H. Neville.
2023. A global synthesis of peer-reviewed research on the effects of hatchery salmonids on wild
salmonids. Fisheries Management and Ecology DOI:10.1111/ fme.12643

McPhee, M., Siegel, J., Adkison, M., 2019. Is a Warming Bering Sea Leading to Smaller Chinook
Salmon? NPAGC Technical Report 1,5 117–119. https://doi.org/10.23849/npafctr15/117.119.

McPhee, M.V., Leon, J.M., Wilson, L.I., Siegel, J.E., Agler, B.A., 2016. Changing growth and
maturity in western Alaskan chinook salmon, Oncorhynchus tshawytscha, Brood Years 1975-2005.
North Pacific Anadromous Fish Commission Bulletin 6, 307–327.

McPhee, M.V., Tappenbeck, T.H., Whited, D.C., Stanford, J.A., 2009a. Genetic diversity and
population structure in the Kuskokwim River drainage support the recurrent evolution hypothesis for
sockeye salmon life histories. Transactions of the American Fisheries Society 138, 1481–1489.

McPhee, M.V., Zimmerman, M.S., Beacham, T.D., Beckman, B.R., Olsen, J.B., Seeb, L.W., Templin,
W.D., 2009. A hierarchical framework to identify influences on Pacific salmon population abundance
and structure in the Arctic-Yukon-Kuskokwim region. American Fisheries Society Symposium 70,
1177–1198.

Meka, J.M., Zimmerman, C.E., Heintz, R.A., Wang, S.W., 2005. Body condition and feeding ecology
of Kuskokwim River chum salmon (Oncorhynchus keta) during freshwater outmigration 1–61.

Meredith, B.L., N.D. Frost, K.S. Reppert, and G.T. Hagerman. 2021. DRAFT-Unik and Chickamin
King salmon stock status and action plan, 2021. RC4 Report to the Alaska Board of Fisheries

Mikow, E., B. Retherford, A. Goddhun and M. L. Kostick. 2016. Exloring the Subsistence Fisheries
of Point Lay and Wainwright, Alaska. Alaska Department of Fish and Game Division of Subsistence,
Technical Paper No. 419, Fairbanks.

- Mikow, E.H et al. 2019. Local Traditional Knowledge of Salmon Freshwater Ecology in the Middleand Upper Kuskokwim River
- Miller, K., Shaftel, R., Bogan, D., 2020. Diets and Prey Items of Juvenile Chinook (Oncorhynchus tshawytscha) and Coho Salmon (O. kisutch) on the Yukon Delta. U.S. Dep. Commer., NOAA Tech.
  Memo. NMFS-AFSC-410 1–54. https://doi.org/10.13140/ RG.2.2.20435.60961
- Miller, K.B. and Weiss, C.M. 2023. Disentangling population level differences in juvenile migration
   phenology for three species of salmon on the Yukon River. Journal of Marine Science and Engineering
   11, 589. https:// doi.org/10.3390/jmse11030589

Miller, S. E., J. M. Murphy, S. C. Heinl, A. W. Piston, E. A. Fergusson, R. E. Brenner, W. W.
Strasburger, and J. H. Moss. 2022. Southeast Alaska pink salmon forecasting models. Alaska
Department of Fish and Game, Fishery Manuscript No. 22-03, Anchorage.

Miller, S.E., Adkison, M., Haldorson, L., 2012. Relationship of water column stability to the growth,
condition, and survival of pink salmon (Oncorhynchus gorbuscha) in the northern coastal Gulf of
Alaska and Prince William Sound. Canadian Journal of Fisheries and Aquatic Sciences 69, 955–969.
https://doi.org/10.1139/f2012-031

- Minobe, S., 2002. Interannual to interdecadal changes in the Bering Sea and concurrent 1998 / 99
  changes over the North Pacific Progress in Oceanography 55: 45–64.
- Molyneaux, D.B., Folletti, D.L., Brannian, L.K., Roczicka, G., 2005. Fishery Data Series No . 05-45
  Age, Sex, and Length Composition of Chinook Salmon from the 2004 Kuskokwim River Subsistence
  Fishery Final Report for Project 04-353 USFWS Office of Subsistence Management by.
- Molyneaux, D.B., Folletti, D.L., Brodersen, A.R., 2008. Salmon age, sex and length catalog for the
  Kuskokwim area, 2007., Regional Information Report. Alaska Department of Fish and Game, Division
  of Commercial Fisheries, Anchorage, AK. Moncrieff, C., Brown, C., Sill, L., 2009. Natural indicators
  of salmon abundance and timing, Yukon River., AYK Sustainable Salmon Initia-tive Final Report.
  Bering Sea Fisherman's Association, Anchorage.
- Mortensen, D., Wertheimer, A., Taylor, S., Landingham, J., 2000. The relation between early marine
  growth of pink salmon, *Oncorhynchus gorbuscha*, and marine water temperature, secondary
  production, and survival to adulthood. Fishery Bulletin 98, 319–335.
- Moss, J.H., J.M. Murphy, E.V. Farley, Jrl., L.B. Eisner, and A.G. Andrews. 2009. Juvenile pink and
  chum salmon distribution, diet, and growth in the northern Bering and Chukchi seas. NPAFC Bulletin
  5:191-196.
- Mossop, B. and Bradford, M.J. 2011. Importance of large woody debris for juvenile Chinook salmon
  habitat in small boreal forest streams in the upper Yukon River basin, Canada. Canadian Journal of
  Forest Research 34(9): 1955-1966. https://doi.org/10.1139/x04-066
- Mueter, F. J., B. Planque, G. L. Hunt, I. D. Alabia, T. Hirawake, L. Eisner, P. Dalpadado, M. Chierici,
  K. F. Drinkwater, N. Harada, P. Arneberg, and S.-I. Saitoh. 2021. Possible future scenarios in the
  gateways to the Arctic for Subarctic and Arctic marine systems: II. prey resources, food webs, fish,
  and fisheries. ICES Journal of Marine Science 78(9):3017–3045.
- Mueter, F.J., Litzow, M. a, 2008. Sea ice retreat alters the biogeog-raphy of the Bering Sea continental
  shelf. Ecological applications : a publication of the Ecological Society of America 18, 309–20. Mundy,
  P.R., and D.F. Evenson. 2011. Environmental controls of phenology of high-latitude Chinook salmon
  populations of the Yukon River, North America, with application to fishery managment. ICES Journal
  of Marine Science 68(6):1155-1164.
- Munro, A. R., C. Habicht, T. H. Dann, D. M. Eggers, W. D. Templin, M. J. Witteveen, T. T. Baker,
  K. G. Howard, J. R. Jasper, S. D. Rogers Olive, H. L. Liller, E. L. Chenoweth, and E. C. Volk. 2012.
  Harvest and harvest rates of chum salmon stocks in fisheries of the Western Alaska Salmon Stock
  Identification Program (WASSIP), 2007–2009. (Very Large PDF 57,581 kB) Alaska Department of
  Fish and Game, Special Publication No. 12-25, Anchorage

Munro, A. R. 2023. Summary of Pacific salmon escapement goals in Alaska with a review of
escapements from 2014 to 2022. Alaska Department of Fish and Game, Fishery Manuscript No. 2301, Anchorage.

- Murphy, J. M., E.A. Fergusson, A. Piston, S. Heinl, A. Gray, and E. Farley. 2019. Southeast Alaska
  pink salmon growth and harvest forecast models. North Pacific Anadromous Fish Commission
  Technical Report No. 15: 75-91.
- Murphy, J., Garcia, S., Dimond, J., Moss, J., Sewall, F., Strasburger, W., Lee, E., Dann, T., Labunski,
  E., T.Zeller, Gray, A., Waters, C., Jallen, D., Nicolls, D., Conlon, R., Cieciel, K., Howard, K., Harris,
  B., N.Wolf, Farley, E., 2021. Northern Bering Sea surface trawl and ecosystem survey cruisereport,
  2019. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-423, 124 p.
- Murphy, J., Howard, K., Eisner, L., Andrews, A., Templin, W., Guthrie, C., Cox, K., Farley, E., 2013.
  Linking Abundance, Distribution, and Size of Juvenile Yukon River Chinook Salmon to Survival in
  the Northern Bering Sea. North Pacific Anadromous Fish Commission 9, 25–30.
- Murphy, J., S. Garcia, J. Dimond, J. Moss, F. Sewall, et al. 2021. Northern Bering Sea surface trawl
  and ecosystem survey cruise report, 2019. NOAA Tech Memo NMFS-AFSC-423
- Murphy, J.M., Farley, E.V., Ianelli, J.N., Stram, D.L., 2016a. Distribution, Diet, and Bycatch of
  Chum Salmon in the Eastern Bering Sea 219–234. https://doi.org/10.23849/npafcb6/219.234.Abstract
- Murphy, J.M., Howard, K.G., Gann, J.C., Cieciel, K.C., Templin, W.D., and Guthrie III, C.M. 2017.
  Juvenile Chinook salmon abundance in the northern Bering Sea: implications for future returns and
  fisheries in the Yukon River. Deep Sea Research Part II: Topical Studies in Oceanography 135: 156167. https://doi.org/10.1016/j. dsr2.2016.06.002.
- Murphy, J.M., K.G. Howard, J.C. Gann, K.C. Cieciel, W.D. Templin, and C.M. Guthrie. 2017.
  Juvenile Chinook salmon abundance in the northern Bering Sea: implications for future returns and
  fisheries in the Yukon River. Deep-Sea Research II 135:156-167
- Murphy, J.M., W.D. Templin, E.V. Farley, Jr., and J.E. Seeb. 2009. Stock-structured distribution of
  western Alaska and Yukon juvenile Chinook salmon (Oncorhynchus tshawytscha) from United States
  BASIS surveys, 2002–2007. N. Pac. Anadr. Fish Comm. Bull. 5: 51–59.
- Murphy, M.L., 1984. Primary production and grazing in freshwater and intertidal reaches of a coastal
  stream, southeast Alaska. Limnology and Oceanography 29, 805–815.
- Murphy, M.L., Heifetz, J., Thedinga, J.F., Johnson, S.W., Koski, K.V., 1989. Habitat utilization by
  juvenile Pacific salmon (Oncorhynchus) in the glacial Taku River, southeast Alaska. Canadian Journal
  of Fisheries and Aquatic Sciences 46, 1677–1685.
- 2451 Myers, K., Irvine, J., Logerwell, E., Urawa, S., Naydenko, S., Zavolokin, A., Davis, N., 2016.
- 2452 Pacific Salmon and Steelhead: Life in a Changing Winter Ocean. NPAFC Bull. 6, 113–138.
- 2453 https://doi.org/10.23849/npafcb6/113.138

2454 Myers, K., R. Walker, N. Davis, J. Armstrong, W. Fournier, N. Mantua, and J. Raymond-Yakoubian.
2455 2010. Climate-Ocean Effects on AYK Chinook Salmon. Page 249. SAFS-UW -1003, School of
2456 Aquatic and Fishery Sciences, University of Washington, Seattle.

- Myers, K.W., Davis, N.D., Walker, R.V., Armstrong, J.L., 2006. Migration studies of salmon in the
  Bering Sea. Final Report, NOAA contract No. NA17RJ1232 AM021. University of Washington.
- Myers, K.W., R.V. Walker, N.D. Davis, J.L. Armstrong, and M. Kaeriyama. 2009. High seas
  distribution, biology, and ecology of AYK salmon: direct information from high seas tagging
  exeriments, 1954 2006. American Fisheries Society Symposium 70:201-239.
- 2462 Myers, K.W., Walker, R.V., Davis, N., 2001. Ocean Distribution and Migration Patterns of Yukon2463 River Chinook Salmon.
- Myers, K.W., Walker, R.V., Davis, N.D., Armstrong, J.L., and Kaeriyama, M. 2009. High seas
  distribution, biology, and ecology of Arctic-Yukon-Kuskokwim salmon: direct information from high
  seas tagging experiments, 1954-2006. American Fisheries Society Symposium 70:201-239.
- 2467 N
- Nagasawa, K. 2023. Catch of coho salmon (Oncorhynchus kisutch) infected with the freshwater
  parasite Salvelinema walkeri (Nematoda: Cystidicolidae) in the Gulf of Alaska in the early winter.
  Species Diversity 28: 141-146. DOI: 10.12782/specdiv.28.141
- 2471 Nagasawa, K. 1998. Predation by salmon sharks on Pacific salmon in the North Pacific Ocean. North
  2472 Pacific Anadromous Fish Commission Bulletin 1:419-433.
- 2473 Nakatani, R. E., G. J. Paulik, and R. Van Cleve. 1975. Pink salmon (Oncoryhnchus gorbuscha) tagging
  2474 experiments in S. E. Alaska, 1938–1942 and 1945. National Oceanic and Atmospheric Administration,
  2475 National Marine Fisheries Service, Special Scientific Report, Fisheries Series 686, Seattle.
- 2476 National Research Council. 2005. Developing a Research and Restoration Plan for Arctic-Yukon2477 Kuskokwim (Western Alaska) Salmon. Washington, DC: The National Academies Press. https://
  2478 doi.org/10.17226/11080.
- Naves, L.C., M.F. Turek, and W.E. Simeone. 2010. Subsistence– personal use salmon harvest,
  Southeast–Yakutat Management Region, 1996–2006. Alaska Department of Fish and Game Division
  of Subsistence Technical Paper No. 350, Anchorage.
- NBTC (Northern Boundary Technical Committee). 2005. Stock Composition Estimates and individual
  stock assignments based on genetic microsatellites and scale patterns for test mixtures of Alaskan and
  Canadian sockeye salmon. TCNB (05)-2.
- Neilsen, J.L., G.T. Ruggerone. 2004. Top-down and bottom-up linkages among climate, growth,
  competition, and production of sockeye salmon populations in Bristol Bay, Alaska, 1955-2000. PICES
  13th Annual Meeting book of abstracts. 24

- 2488 Neuswanger, Jason R, Wipfli, M.S., Evenson, M.J., Hughes, N.F., Rosenberger, A.E., 2015. Low
- productivity of Chinook salmon strongly correlates with high summer stream discharge in two
  Alaskan rivers in the Yukon drainage. Canadian Journal of Fisheries and Aquatic Sciences 72, 1125–
  1137.
- Neuswanger, Jason R., Wipfli, M.S., Evenson, M.J., Hughes, N.F., Rosenberger, A.E., Jonsson, B.,
  2015. Low productivity of Chinook salmon strongly correlates with high summer stream discharge
  in two Alaskan rivers in the Yukon drainage. Canadian Journal of Fisheries and Aquatic Sciences 1–
  13. https://doi.org/10.1139/ cjfas-2014-0498
- Nielsen, J.L., and G.T. Ruggerone. 2009. Climate change and a dynamic ocean carrying capacity:
  growth and survival of Pacific salmon at sea. American Fisheries Society Symposium 71:77-96.
- Nielsen, J.L., G.T. Ruggerone, and C.E. Zimmerman. 2012. Adaptive strategies and life history
  characteristics in a warming climate: salmon in the Arctic? Environ Biol Fish 96:1187-1226
- North Pacific Anadromous Fish Commission. 2023. The status and trends of Pacific salmon and
  steelhead trout stocks with linkages to their ecosystem. N. Pac. Anadr. Fish Comm. Tech. Rep. 19.
  2502 256 pp. https://doi.org/10.23849/LOEX7610 NPFMC LKTKS Search Engine
- 2503 <u>https://lktks.npfmc.org/</u>
- 2504 0
- O'Neill, D., 2012. The fall of the Yukon kings, in: Banerjee, S. (Ed.), Arctic Voices: Resistance at the
  Tipping Point.
- Ohlberger, J., D.E. Schindler, R.J. Brown, J.M.S. Harding, M.D. Adkison, A.R. Munro, L. Horstmann,
  and J. Spaeder. 2020. The reproductive value of large females: consequences of shifts in demographic
  structure for puoulation reproductive potential in Chinook salmon. Canadian Journal of Fisheries and
  Aquatic Sciences dx.doi.org/10.1139/cjfas-2020-0012
- Ohlberger, J., E.J. Ward., R.E. Brenner, M.E. Hunsicker, et al. 2022. Non-stationary and interactive
  effects of climate and competition on pink salmon productivity. Global Change Biology.
  DOI:10.1111/gcb.16049.
- Oke, K.B., Mueter, F., Litzow, M.A., 2019. Warming leads to opposite patterns in weight-at-age for
   young versus old age classes of Bering Sea walleye pollock 1–39.
- Olsen, J. B., P. A. Crane, B. G. Flannery, K. Dunmall, W. D. Templin, and J. K. Wenburg. 2010.
  Comparative landscape genetic analysis of three Pacific salmon species from subarctic North America.
  Conservation Genetics 12: 223-241.
- 2519 Olsen, J. B., T. D. Beacham, M. Wetklo, L. W. Seeb, C. T. Smith, B. G. Flannery, J. K. Wenburg.
- 2520 2010. The influence of hydrology and waterway distance on population structure of Chinook salmon
  2521 Oncorhynchus tshawytscha in a large river. Journal of Fish Biology 76: 1128-1148.

Orsi, J. A., A. C. Wertheimer, M. V. Sturdevant, E. A. Fergusson, D. G. Mortensen, and B. L. Wing.
2004. Juvenile chum salmon consumption of zooplankton in marine waters of southeastern Alaska: a
bioenergetics approach to implications of hatchery stock interactions. Reviews in Fish Biology and
Fisheries 14 :335-359.

Orsi, J. A., and H. W. Jaenicke. 1996. Marine distribution and origin of prerecruit Chinook salmon
Oncorynchus tshawytscha, in southeastern Alaska. Fishery Bulletin 94:482-497

Orsi, J. A., M. V. Sturdevant, J. M. Murphy, D. G. Mortensen, and B. L. Wing. 2000. Seasonal habitat
use and early marine ecology of juvenile Pacific salmon in Southeastern Alaska. North Pacific
Anadromous Fish Commission Bulletin No. 2: 111–122.

Orsi, J., A. Wertheimer, M. Sturdevant, E. Fergusson, and B. Wing. 2009. Insights from a 12-year
biophysical time series of juvenile Pacific salmon in Southeast Alaska: the Southeast Alaska Coastal
Monitoring Project (SECM). Alaska Fisheries Science Center Quarterly Report July–September 2009.

Orsi, J., Fergusson, E., Joyce, J.E., 2013. Connecting the "Dots" Among Coastal Ocean Metrics and
Pacific Salmon Production in Southeast Alaska, 1997-2012.

Orsi, J.A., Fergusson, E.A., Sturdevant, M.V., Wing, B.L., Wertheimer, A.C., Heard, W.R., 2009.
Annual survey of juvenile salmon, ecologically-related species, and environmental factors in the
marine waters of southeastern Alaska, May - August 2008. NPAFC Doc. 1181, 72 pp.

Orsi, J.A., Wertheimer, A.C., 1995. Marine vertical distribution of juvenile Chinook and coho salmon
in southeastern Alaska. Transactions of the American Fisheries Society 124, 159–169.

Oslund, S., S. Ivey, and D. Lescanec. 2020. Area Management Report for the sport fisheries of
northern Cook Inlet, 2017–2018. Alaska Department of Fish and Game, Fishery Management Report
No. 20-04, Anchorage.

Otis, E. O., G. J. Hollowell, and E. G. Ford. 2018. Observations of pink salmon hatchery proportions
in selected Lower Cook Inlet escapements, 2014?2017. Alaska Department of Fish and Game, Special
Publication No. SP18-11, Anchorage.

- Otis, E. O., J. W. Erickson, C. Kerkvliet, and T. McKinley. 2016. A review of escapement goals for
  salmon stocks in Lower Cook Inlet, Alaska, 2016. Alaska Department of Fish and Game, Fishery
  Manuscript Series No. 16-08, Anchorage.
- Ovando, D., Cunningham, C., Kuriyama, P., Boatright, C. & Hilborn, R. (2022) Improving forecasts
  of sockeye salmon (Oncorhynchus nerka) with parametric and nonparametric models. Canadian
  Journal of Fisheries and Aquatic Sciences, 99, 1-13.
- Oxman, D.S., Smoker, W.W., Gharrett, A.J., 2013. Developmental progression of gill rakers as a post hatch developmental marker in pink salmon, Oncorhynchus gorbuscha. Environmental Biology of
   Fishes 96, 677–689. https://doi.org/10.1007/s10641-012-0058-6

2556 **P** 

Paige, A. et al. 2009. Local knowledge, harvest patterns, and community uses of salmon in Wrangell,
Alaska. ADF&G Division of Subsistence, Techncial Paper No. 323.

Paige, A. W., S. Churchill, N. Ratner, M. Turek, and P. Coiley-Kenner. 2009. Local knowledge,
harvest patterns, and community use of salmon in Wrangell, Alaska. Alaska Department of Fish and
Game, Division of Subsistence Technical Paper No. 323, Juneau.

Parker-Stetter, S.L., Horne, J.K., Farley, E.V., Barbee, D.H., Andrews, A.G., Eisner, L.B., Nomura,
J.M., 2013. Summer distributions of forage fish in the eastern Bering Sea. Deep-Sea Research Part II:
Topical Studies in Oceanography 94, 211–230. https://doi.org/10.1016/j.dsr2.2013.04.022

Pella, J.J., Geiger, H.J., 2009. Sampling considerations for estimating geographic origins of Chinook
salmon bycatch in the Bering Sea pollock fishery., Special Publication No. SP 09-08. Alaska
Department of Fish and Game, Anchorage.

- Peltz, L., and J. P. Koenings. 1989. Evidence for temperature limitation of juvenile sockeye salmon,
  Oncorynchus nerka, growth in Hugh Smith Lake, Alaska. FRED Reports No. 90
- Pestal, G., C. J. Schwarz and R. A. Clark. 2020. Taku River Sockeye Salmon Stock Assessment
  Review and Updated 1984-2018 Abundance Estimates. Pacific Salmon Comm. Tech. Rep. No. 43:
  118 p.
- Petrou, E. L., J. E. Seeb, L. Hauser, M. J. Witteveen, W. D. Templin, L. W. Seeb. 2014. Fine-scale
  sampling reveals distinct isolation by distance patterns in chum salmon (Oncorhynchus keta)
  populations occupying a glacially dynamic environment. Conservation Genetics 15(1): 229-243.

Petrou, E. L., L. Hauser, R. S. Waples, J. E. Seeb, W. D. Templin, D. Gomez-Uchida and L. W. Seeb.
2013. Secondary contact and changes in coastal habitat availability influence the nonequilibrium
population structure of a salmonid (Oncorhynchus keta). Molecular Ecology 22(23): 5848-5860.

- Piston, A. W. 2021. District 104 purse seine fishery harvest pattern analysis. Pacific Salmon Comm.
  Tech. Rep. No. 44: 127 p.
- Piston, A. W., and S. C. Heinl. 2012. Hatchery Chum Salmon Straying in Southeast Alaska, 2011.
  Alaska Department of Fish and Game, Fishery Data Series No. 12-45, Anchorage.
- Piston, A. W., and S. C. Heinl. 2012. Hatchery Chum Salmon Straying Studies in Southeast Alaska,
  2008–2010. Alaska Department of Fish and Game, Fishery Manuscript Series No. 12-01, Anchorage.

Piston, A. W., S. C. Heinl, H. J. Geiger, and T. A. Johnson. 2006. Hugh Smith Lake sockeye salmon
adult and juvenile studies, 2003-2005. Alaska Department of Fish and Game, Fishery Data Series
No. 06-51, Anchorage.

- Piston, A.W., and S.C. Heinl. 2020. Chum salmon stock status and escapement goals in southeastAlaska through 2019. Spec Pub No 20-10 ADFG
- Piston, A.W., and S.C. Heinl. 2020. Pink salmon stock status and escapement goals in southeast Alaska
  through 2019. Spec Pub No 20-09 ADFG
- 2592 Poetter, A.D., Aaron, T., 2017. Annual Management Report, Kuskokwim Area, 2016.
- Polum, T. 2023. Report on selected sport fisheries of the Alaska Peninsula–Aleutian Islands
  Management Area, 2012–2021. Alaska Department of Fish and Game, Fishery Management Report
  No. 23-01, Anchorage.
- Polum, T., M. Witteveen, M. Stratton, and M. Evans. 2019. Report on selected sport fisheries of the
  Kodiak Management Area, 2008–2017. Alaska Department of Fish and Game, Fishery Management
  Report No. 19-04, Anchorage.
- Porter, T. J., S. W. Schoenemann, L. J. Davies, E. J. Steig, S. Bandara, and D. G. Froese. 2019. Recent
  summer warming in northwestern Canada exceeds the Holocene thermal maximum. Nature
  Communications 10(1):1631.
- Priest, J., S.C. Heinl, and L.D. Shaul. 2021. Coho salmon stock status in southeast Alaska: a review
  of trends in productivity, harvest, and abundance through 2019. Pacific Salmon Commission Technical
  Report No. 45.
- Punt, A.E., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., Haddon, M., 2016. Management
  strategy evaluation: best practices. Fish and Fisheries 17, 303–334. https://doi.org/10.1111/faf.12104
- 2607 **R**
- Ramos, J. and R. Mason. Traditional Ecological Knowledge of Tlingit People Concerning the SockeyeSalmon Fishery of the Dry Bay Area
- Ransbury, S. R., N. L. Zeiser, J. A. Bednarski, S. C. Heinl, C. S. Jalbert, and S. E. Miller. 2021. Stock
  assessment study of Chilkat Lake and River sockeye salmon, 2017–2020. Alaska Department of Fish
  and Game, Fishery Manuscript Series No. 21-06, Anchorage.
- Ratner, N.C. and J.A. Dizard. 2006. Local knowledge, harvest patterns, and community use of sockeye
  salmon in Hoonah, Alaska. ADF&G Division of Subsistence, Technical Paper No. 307.
- Ratner, N.C., et al. 2006. Local knowledge, customary practices, and harvest of sockeye salmon from
  the Klawock and Sarkar rivers, Prince of Wales Island, Alaska. ADF&G Division of Subsistence,
  Technical Paper No. 308.
- Raymond-Yakoubian, J., B. Raymond-Yakoubian, C. Moncrieff. 2017. The incorporation of
  traditional knowledge into Alaska federal fisheries management. Marine Policy 78:132-142.

Raymond-Yakoubian, B., Raymond-Yakoubian, J. 2015. "Always taught not to waste": Traditional
Knowledge and Norton Sound/ Bering Strait Salmon Populations." 2015 Arctic-Yukon-Kuskokwim
Sustainable Salmon Initiative Project 1333 Final Product. Kawerak. Inc. Social Science Program:
Nome, AK.

Raymond-Yakoubian, J., 2009. Climate-Ocean Effects on Chinook Salmon: Local Traditional
Knowledge Component. Final report to the Arctic Yukon Kuskokwim Sustainable Salmon Initiative
for project 712. Kawerak. Inc. Social Science Program: Nome, AK.

Ream, J. T. and J. Merriam. 2017. Local and traditional knowledge of Stikine River Chinook salmon:
a local perspective on a vital commercial, sport, and subsistence fish. Alaska Department of Fish and
Game Division of Subsistence, Technical paper No. 430, Anchorage.

Ream, J.T. and J. Merriam. 2017. Local and Traditional Knowledge of Stikine River Chinook Salmon:
a Local Perspective on a Vital Commercial, Sport, and Subsistence Fish. ADF&G Division of
Subsistence, Technical Paper No. 430.

Reese, C., Hillgruber, N., Sturdevant, M., Wertheimer, A., Smoker, W., Focht, R., 2009. Spatial and
temporal distribution and the potential for estuarine interactions between wild and hatchery chum
salmon (Oncorhynchus keta) in Taku Inlet, Alaska. Fishery Bulletin 107, 433–450.

Reimer, A. M., and N. A. DeCovich. 2020. Susitna River Chinook salmon run reconstruction and
escapement goal analysis. Alaska Department of Fish and Game, Fishery Manuscript No. 20-01,
Anchorage.

Rich, W. H. 1927. Salmon-tagging experiments in Alaska, 1924 and 1925. Bulletin of the United
States Bureau of Fisheries 42:109–146.

Rich, W. H., and A. J. Suomela. 1929. Salmon-tagging experiments in Alaska, 1926. Bulletin of the
United States Bureau of Fisheries 43(Part 2):71–104.

Riddell, B.E., K.G. Howard, and A.R. Munro. 2022. Salmon returns in the North Pacific in relation to
expedition observations (and next steps). In L. Fitzpatrick, tech. editor. Virtual Conference on Winter
Ecology of Pacific Salmon and Results from the Two Gulf of Alaska Expeditions, N. Pac. Anadr. Fish
Comm. Tech. Rep. 18: 115–139. (Available at https://npafc.org)

2647 Riddell, B.E., K.G. Howard, and A.R. Munro. 2022. Salmon returns in the North Pacific in relation

to expedition observations (and next steps). In L. Fitzpatrick, tech. editor. Virtual Conference on

2649 Winter Ecology of Pacific Salmon and Results from the Two Gulf of Alaska Expeditions, N. Pac.

2650 Anadr. Fish Comm. Tech. Rep. 18: 115–139. (Available at https://npafc.org)

Riddell, B.E., K.G. Howard, and A.R. Munro. 2022. Salmon returns in the North Pacific in relation to
expedition observations (and next steps). In L. Fitzpatrick, tech. editor. Virtual Conference on Winter
Ecology of Pacific Salmon and Results from the Two Gulf of Alaska Expeditions, N. Pac. Anadr. Fish

2654 Comm. Tech. Rep. 18: 115–139. (Available at https://npafc.org)

Riffe, R., Mercer, B., 2006. Effects of habitat and predator-prey interactions on stocked sockeye fry
in Tatsamenie Lake (No. No. 06-02), Fishery Manuscript. Alaska Department of Fish and Game,
Anchorage.

Rogers Olive, S.D., E.K.C. Fox, and S.E. Gilf-Baumer. 2018. Genetic baseline for mixed stock
analyses of sockey salmon harvested in SEAK for Pacific salmon treaty applications, 2018. Fishery
Manuscript no. 18-03

Rogers, D.E., and G.T. Ruggerone. 1993. Factors affecting marine growth of Bristol Bay sockeyesalmon. Fisheries Research 18:89-103.

- Ruggerone, G.T. 2003. Rapid natural habitat degradation and consequences for sockeye salmon
  production in the Chignik Lakes system, Alaska. SAFS-UW-0309, 133pgs.
- Ruggerone, G.T. 2004. Pre-season forecast of sockeye salmon migration timing in Bristol Bay, Alaska
  based on oceanographic and biological variables. North Pacific Research Board Report 39pgs

Ruggerone, G.T., and B.M. Connors. 2015. Productivity and life history of sockeye salmon in relation
to competition with pink and sockeye salmon in the North Pacific Ocean. Canadian Journal of
Fisheries and Aquatic Sciences 72:818-833.

- Ruggerone, G.T., and D.E. Rogers. 1992. Predation on sockeye salmon fry by juvenile coho salmon
  in the Chignik Lakes, Alaska: Implications for salmon management. North American Journal of
  Fisheries Management 12:87-102.
- Ruggerone, G.T., and J.L. Nielsen. 2004. Evidence for competitive dominance of Pink salmon over
  other salmonids in the North Pacific Ocean. Reviews in Fish Biology and Fisheries 14:371-390.
- Ruggerone, G.T., and J.L. Nielsen. 2009. A review of growth and survival of salmon at sea in response
  to competitioni and climate change. American Fisheries Society Symposium 70:241-265.

2677 Ruggerone, G.T., B.A. Agler, B.M. Connors, E.V. Farley J.R. Irvine et al. 2016. Pink and sockeye
2678 salmon interactions at sea and their influence on forecast error of Bristol Bay sockeye salmon. North
2679 Pacific Anadromous Fish Commission Bulletin 6:349 - 361.

- Ruggerone, G.T., B.A. Agler, B.M. Connors, E.V. Farley, Jr., et al. 2016. Pink and sockeye salmon
  interactions at sea and their influence on forecast error of Bristol Bay sockeye salmon. North Pacific
  Anadromous Fish Commission Bulletin 6:349-361.
- Ruggerone, G.T., B.A. Agler, J.L. Nielsen. 2012. Evidence for competition at sea between Norton
  Sound chum salmon and Asian hatchery chum salmon. Environmental Biology of Fishes 94:149-163.

Ruggerone, G.T., B.A. Agler, L. Wilson, and E.V. Farley, Jr. 2013. Size-selective mortality of Bristol
Bay sockeye smolts in relation to smolt characteristics, ocean conditions, and sockeye salmon
productivity. North Pacific Anadromous Fish Commision Technical Report 9:210-213.

Ruggerone, G.T., Connors, B.M., 2015. Productivity and life history of sockeye salmon in relation to
competition with pink and sockeye salmon in the North Pacific Ocean. Canadian Journal of Fisheries
and Aquatic Sciences 72, 818–833. https://doi.org/dx.doi. org/10.1139/cjfas-2014-0134

Ruggerone, G.T., E. Farley, J. Nielsen, and P. Hagen. 2005. Seasonal marine growth of Bristol Bay
sockeye salmon in relation to competition with Asian pink salmon and the 1977 ocean regime shift.
Fishery Bulletin 103:355-370.

- Ruggerone, G.T., J.L. Nielsen, and J. Bumgarner. 2007. Linkages between Alaskan sockeye salmon
  abundance, growth at sea, and climate, 1955 2002. Deep-Sea Research II 54:2776-2793.
- Ruggerone, G.T., J.R. Irvine, and B. Connors. 2021. Did recent heatwaves and record high pink
  salmon abundance lead to a tipping point that caused record declines in North Pacific salmon
  andubnance and harvest in 2020? North Pacific Anadromous Fish Commission, Technical Report
  17:78-82.
- Ruggerone, G.T., M. Zimmermann, K.W. Myers, J.L. Nielsen, and D.E. Rogers. 2003. Competition
  between Asian pink salmon and Alaskan sockeye salmon in the North Pacific Ocean. Fisheries
  Oceanography 12(3):209-219.
- Ruggerone, G.T., Nielsen, J.L., Agler, B.A., 2009. Linking marine and freshwater growth in western
  Alaska Chinook salmon Oncorhynchus tshawytscha. Journal of Fish Biology 75, 1287–1301.
- Ruggerone, G.T., Nielsen, J.L., and Agler, B.A. 2009. Climate, growth and population dynamics of
  Yukon River Chinook salmon. North Pacific Anadromous Fish Commission Bulletin 5: 279–285
- Ruggerone, G.T., R. Hanson, and D.E. Rogers. 2000. Selective predation by brown bears foraging on
  spawning sockeye salmon. Canadian Journal of Zoology 78:974-981.
- Ruggerone, G.T., R.M. Peterman, B. Dorner, and K.W. Myers. 2010. Magnitude and trends in
  abundance of hatchery and wil pink salmon, chum salmon, and sockeye salmon in the North Pacific
  Ocean. Marine and Coastal Fisheries: Dynamics, Managment, and Ecosystem Science 2:306-328.
- Ruggerone, G.T., Zimmermann, M., Myers, K.W., Nielsen, J.L., Rogers, D.E., 2003. Competition
  between Asian pink salmon (Oncorhynchus gorbuscha) and Alaskan sockeye salmon (O. nerka) in the
  North Pacific Ocean. Fisheries Oceanography 12, 209–219.
- 2715 Ruggerone, G.T., A.M. Springer, G.B. van Vliet, B. Connors, J.R. Irvine, L.D. Shaul, M.R. Sloat,
- and W.I. Atlas. 2023. From diatoms to killer whatles: iimpacts of pink salmon on North Pacific
  ecosystems. Marine Ecology Progress Series 719:1-40.
- Runfola, D.M. and D. Koster. 2019. Inseason estimation of subsistence salmon fishing effort and
  harvest in the lower Kuskokwim River, 2015–2018. Alaska Department of Fish and Game, Division
  of Subsistence Technical Paper No. 449, Fairbanks.

- Runfola, D.M., H. Ikuta, A.R.Brenner, J.J. Simon, J. Park, D. S. Koster, and M. Kostick. 2017. Bethel
  subsistence, 2012: wild resource harvests and uses, land use patterns, and subsistence economy in the
  hub community of the Yukon–Kuskokwim Delta, Alaska. Alaska Department of Fish and Game
  Division of Subsistence, Technical Paper No. 393, Fairbanks.
- Russell et al. (2021) Prince William Sound area salmon fisheries -- a report to the Alaska Board of
  Fisheries, 2021.
- Russell, C. W. 2023. North Alaska Peninsula commercial salmon annual management report, 2022.
  Alaska Department of Fish and Game, Fishery Management Report No. 23-03, Anchorage.
- 2729 **S**
- Salomone, P.G., K.R. Courtney, G.T. Hagerman, P.A. Fowler, and P.J. Richards. 2021. DRAFTStikine River and Andrew Creek King salmon stock status and action plan, 2021. RC& report to the
  Alaska Board of Fisheries
- 2733 Sato, S., Moriya, S., Azumaya, T., Nagoya, H., 2009. Stock Distribution Patterns of Chum Salmon in
- the Bering Sea and North Pacific Ocean during the Summer and Fall of 2002 2004. North Pacific
- 2735 Anadromous Fish Commission Bulletin 5, 29–37.
- Scannell, H., J. Botz, K. Gatt, J. Morella, J. Buza, and R. Ertz. 2023. 2021 Prince William Sound area
  finfish management report. Alaska Department of Fish and Game, Fishery Management Report No.
  23-06, Anchorage.
- Schaberg, K., T. McKinley and H. Finkle. 2023 Review of Salmon Escapement Goals in the Alaska
  Peninsula, Aleutian Islands and Chignik Management Areas. Oral Report to the Alaska Board of
  Fisheries 2023. Alaska Department of Fish and Game Report.
- Schaberg, K.L., Liller, Z.W., Molyneaux, D.B., 2010. Fishery Data Series No . 10-32 A Mark –
  Recapture Study of Kuskokwim River Coho , Chum , Sockeye , and Chinook Salmon , 2001 2006
  Final Report for Project FIS 04-308 by.
- Schaberg, K.L., Liller, Z.W., Molyneaux, D.B., Bue, B.G., Stuby, L., 2012. Estimates of Total Annual
  Return of Chinook Salmon to the Kuskokwim River, 2002 2007 by, Fishery Data Series.
- Scheuerell, M.D., Hilborn, R., Ruckelshaus, M.H., Bartz, K.K., Lagueux, K.M., Haas, A.D., and
  Rawson, K. 2006. The Shiraz model: a tool for incorporating anthropogenic effects and fish-habitat
  relationships in conservatino planning. Canadian Journal of Fisheries and Aquatic Sciences 63: 15961607. doi:10.1139/F06-056
- Schindler, D., C. Krueger, P. Bisson, M. Bradford, B. Clark, J. Conitz, K. Howard, M. Jones, J.
  Murphy, K. Myers, M. Scheuerel, E. Volk, and J. Winton. 2013. Arctic -Yukon-Kuskokwim Chinook
  Salmon Research Action Plan: Evidence of Decline of Chinook Salmon Populations and
  Recommendations for Future Research. Page 70. AYK Sustainable Salmon Initiative, Anchorage, AK.

Schindler, D., Krueger, C., Bisson, P., Bradford, M., Clark, B., Conitz, J., Howard, K., Jones, M.,
Murphy, J., Myers, K., Scheuerell, M., Volk, E., and Winton, J. 2013. Arctic-Yukon-Kuskokwim
Chinook salmon research action plan: evidence of decline of Chinook salmon populations and
recommendations for future research. Prepared for the AYK Sustainable Salmon Initiative
(Anchorage, AK). v + 70 pp.

Schoch, G.C., Albert, D.M., Shanley, C.S., 2014. An Estuarine Habitat Classification for a Complex
Fjordal Island Archipelago. Estuaries and Coasts 37, 160–176. https://doi.org/10.1007/s12237-0139622-3

Schuster, M., M. D. Booz, and A. W. Barclay. 2021. Chinook salmon sport harvest genetic stock and
biological compositions in Cook Inlet salt waters, 2014–2018. Alaska Department of Fish and Game,
Fishery Manuscript No. 21-04, Anchorage.

Schuster, M., M. D. Booz, and A. W. Barclay. 2021. Chinook salmon sport harvest genetic stock and
biological compositions in Cook Inlet salt waters, 2014–2018. Alaska Department of Fish and Game,
Fishery Manuscript No. 21-04, Anchorage.

Schwanke, C. J., and M. J. Piche. 2023. Run timing and spawning distribution of Copper River
Chinook salmon, 2019–2021. Alaska Department of Fish and Game, Fishery Data Series No. 23-14,
Anchorage.

Seeb, J. E., C. Habicht, W. D. Templin, J. B. Shaklee, L. W. Seeb, and F. M. Utter. 1999. Allozyme
and mtDNA variation describe ecologically important genetic structure of even-year pink salmon
inhabiting Prince William Sound, Alaska. Ecology of Freshwater Fish 8: 122-140.

Seeb, L. W., A. Antonovich, M. Banks, T. Beacham, R. Bellinger, S. Blankenship, M. Campbell, N.
DeCovich, J. C. Garza, C. Guthrie, T. Lundrigan, P. Moran, S. Narum, J. Stephenson, J. Supernault,
D. Teel, W. D. Templin, J. K. Wenburg, S. Young, and C. T. Smith. 2007. Development of a
standardized DNA database for Chinook salmon. Fisheries 32(11):540–552

Seeb, L. W., C. Habicht, E. V. Farley, Jr., J. E. Seeb, and F. M. Utter. 2011. Single-Nucleotide
polymorphic genotypes reveal patterns of early juvenile migration of sockeye salmon in the eastern
Bering Sea. Transactions of the American Fisheries Society 140: 734-748.

Seeb, L. W., C. Habicht, W. D. Templin, K. E. Tarbox, R. Z. Davis, L. K. Brannian, and J. E. Seeb.
2000. Genetic diversity of sockeye salmon (Oncorhynchus nerka) of Cook Inlet, Alaska, and its
application to restoration of populations affected by the Exxon Valdez oil spill. Transactions of the
American Fisheries Society 129: 1223-1249.

Seeb, L. W., N. A. DeCovich, A. W. Barclay, C. T. Smith, and W. D. Templin. 2009. Timing and
origin of Chinook salmon stocks in the Copper River and adjacent ocean fisheries using DNA
markers. (PDF 916 kB) Alaska Department of Fish and Game, Fishery Data Series No. 09-58,
Anchorage.

Seeb, L. W., P. A. Crane, C. M. Kondzela, R. L. Wilmot, S. Urawa, N. V. Varnavskaya, and J. E.
Seeb. 2004. Migration of Pacific Rim chum salmon on the high seas: insights from genetic data.
Environmental Biology of Fishes 69: 21-36.

Seeb, L. W., W. D. Templin, S. Sato, S. Abe, K. Warheit, J. Y. Park, and J. E. Seeb. 2011. Single
nucleotide polymorphisms across a species- range: implications for conservation studies of Pacific
salmon. Molecular Ecology Resources 11: 195-217.

Seeb, L. W., W. D. Templin, S. Sato, S. Abe, K. Warheit, J. Y. Park, and J. E. Seeb. 2011. Single
nucleotide polymorphisms across a species' range: implications for conservation studies of Pacific
salmon. Molecular Ecology Resources, 11:195–217. doi: 10.1111/j.1755-0998.2010.02966.x

Seeb, L.W., J.E. Seeb, C. Habicht, E.F. Farley, Jr., and F.M. Utter. 2011. Single-nucleotide
polymorphic gentoypes reveal patterns of early juvenile migration of sockeye salmon in the eastern
Bering Sea. Transactions of the American Fisheries Society 140(3):734-748.

Seitz, A.C., Courtney, M.B., Evans, M.D. and Manishin, K., 2019. Pop-up satellite archival tags reveal
evidence of intense predation on large immature Chinook salmon (Oncorhynchus tshawytscha) in the
North Pacific Ocean. Canadian Journal of Fisheries and Aquatic Sciences, 76(9), pp.1608-1615.

Sergeant, C.J., Bellmore, J.R., Bellmore, R.A., Falke, J.A., Mueter, F.J., Westley, P.A.H., 2023.
Hypoxia vulnerability in the salmon watersheds of Southeast Alaska. Science of The Total
Environment 896, 165247. https://doi.org/10.1016/j.scitotenv.2023.165247

Sergeant, C.J., Bellmore, J.R., McConnell, C., Moore, J.W., 2017. High salmon density and low
discharge create periodic hypoxia in coastal rivers. Ecosphere 8, e01846.
https://doi.org/10.1002/ecs2.1846

Shanley, C.S., Albert, D.M., 2014. Climate Change Sensitivity Index for Pacific Salmon Habitat in
Southeast Alaska. PLoS ONE 9, e104799. https://doi.org/10.1371/journal.pone.0104799

Shaul, L. D., J. A. Bednarski, J. T. Williams, and B. W. Elliott. 2019. Stock status and review of factors
affecting coho salmon returns and escapements in Southeast Alaska. Alaska Department of Fish and
Game, Regional Information Report No. 1J19-12, Douglas.

Shaul, L. D., R Ericksen, K. Crabtree, and J. Lum. 2013. Beyond the estuary: an extension of the
nomad life history strategy in coho salmon. North Pacific Anadromous Fish Commission Technical
Report No. 9:171–175.

Shaul, L.D., and H.J. Geiger. 2016. Effects of climate and competition for off shore prey on growth,
survival, and reproductive potential of coho salmon in Southeast Alaska. N. Pac. Anadr. Fish Comm.
Bull. 6: 329-347. doi: 10.23849/npafcb6/329.347

Shedd, K. R., D. L. Leonard, and J. V. Nichols. 2022. Mixed stock analysis of Chinook salmon
harvested in Southeast Alaska commercial troll and sport fisheries, 2019. Alaska Department of Fish
and Game, Fishery Data Series No. 22-20, Anchorage

Shedd, K. R., E. A. Lescak, C. Habicht, E. E. Knudsen, T. H. Dann, H. A. Hoyt, D. J. Prince, and W.
D. Templin. 2022. Reduced relative fitness in hatchery-origin Pink Salmon in two streams in Prince
William Sound, Alaska. Evolutionary Applications, 15(3): 429–446.
https://doi.org/10.1111/eva.13356

Shedd, K. R., T. H. Dann, H. A. Hoyt, M. B. Foster, and C. Habicht. 2016. Genetic baseline of North
American sockeye salmon for mixed stock analyses of Kodiak Management Area commercial
fisheries, 2014–2016. (PDF 3,362 kB) Alaska Department of Fish and Game, Fishery Manuscript
Series No. 16-03, Anchorage.

Shedd, K., Foster, M., Wattum, M., Polum, T., Witteveen, M., Stratton, M., Dann, T., Hoyt, H., and
Habicht, C. 2016. Genetic Stock Composition of the Commercial and Sport Harvest of Chinook
Salmon in Westward Region, 2014-2016, Fishery Manuscript Series No. 16-11

Shedd, K., T.H. Dann, C. Habicht, and W.D. Templin. 2014. Defining reproductive success: whichfish count? Alaska Hatchery Research Program Technical Document No. 1.

Shedd, K., T.H. Dann, C. Habicht, and W.D. Templin. 2014. Parentage SNP selection - SEAK chum.
Alaska Hatchery Research Program Technical Document No. 2.

2840 Shedd, K.R., Lescak, E.A., Habicht, C., Knudsen, E.E., Dann, T.H., Hoyt, H.A., Prince, D.J. and

2841 Templin, W.D., 2022. Reduced relative fitness in hatchery-origin Pink Salmon in two streams in

**2842** Prince William Sound, Alaska. Evolutionary Applications, 15(3), pp.429-446.

Siddon, E. 2022. Ecosystem Status Report 2022: Eastern Bering Sea, Stock Assessment and Fishery
Evaluation Report, North Pacific Fishery Management Council, 1007 W. 4rd Ave., Suite 400,
Anchorage, AK 99501.

Siddon, E.C., Kristiansen, T., Mueter, F.J., Holsman, K.K., Heintz, R. a, Farley, E.V., 2013. Spatial
match-mismatch between juvenile fish and prey provides a mechanism for recruitment variability
across contrasting climate conditions in the eastern Bering Sea. PloS one 8, e84526.
https://doi.org/10.1371/journal.pone.0084526

Siegel, J.E., Adkison, M.D., McPhee, M.V., 2018. Changing maturation reaction norms and the effects
of growth history in Alaskan Chinook salmon. Marine Ecology Progress Series 595, 187–202.
https://doi.org/10.3354/meps12564

2853 Siegel, J.E., McPhee, M.V., Adkison, M.D., 2017. Evidence that marine temperatures influence

2854 growth and maturation of western Alaskan Chinook Salmon Oncorhynchus tshawytscha. Marine and
2855 Coastal Fisheries 9, 441–456. https://doi.org/10.1080/19425120. 2017.1353563

Sigler, M., Hollowed, A., Holsman, K., Zador, S., Haynie, A., Himes-Cornell, A., Mundy, P., Davis, 2856 2857 S., Duffy-Anderson, J., Gelatt, T., Gerke, B., Stabeno, P., 2016. Alaska Regional Action Plan for the Southeastern Technical Bering Sea. NOAA Memorandum **NMFS** AFSC i–50. 2858 http://dx.doi.org/10.7289/V5/TM-AFSC-336 2859

Sigler, M., Renner, M., Danielson, S., 2011. Fluxes, fins, and feathers: relationships among the Bering,
Chukchi, and Beaufort seas in a time of climate change. Oceanography- 24, 250–265.

Sigler, M.F., Mueter, F.J., Bluhm, B.A., Busby, M.S., Cokelet, E.D., Danielson, S.L., Robertis, A.D.,
Eisner, L.B., Farley, E.V., Iken, K., Kuletz, K.J., Lauth, R.R., Logerwell, E.A., Pinchuk, A.I., 2016.
Late summer open water zoogeography of the northern Bering and Chukchi seas. Deep Sea Research
Part II: Topical Studies in Oceanography. https://doi.org/10.1016/j.dsr2.2016.03.005

- Sill et al. (2019) Copper River Chinook salmon -- the intersection of commercial fisheries and thesubsistence way of life in Cordova, Alaska.
- Sill, L. A. and J. M. Van Lanen. 2022. Local and traditional knowledge of Chilkat Chinook Salmon.
  Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 463, Anchorage.
- Sill, L. A., G. Halas, and D. Koster. 2019. Copper River Chinook Salmon: The Intersection ofCommercial
- Fisheries and the Subsistence Way of Life in Cordova, Alaska. Alaska Department of Fish and GameDivision of Subsistence, Technical Paper No. 444, Anchorage.
- Sill, L.A. and J.M. Van Lanen. 2022. Local and traditional knowledge of Chilkat Chinook salmon.
  ADF&G Division of Subsistence, Technical Paper No. 463.
- Simeone, W. E., and E. McC. Valentine, 2007. Ahtna knowledge of long-term changes in salmon runs
  in the Upper Copper River drainage, Alaska. Alaska Department of Fish and Game, Division of
  Subsistence Technical Paper No. 324. Juneau.
- Simeone, William E. and James Kari. 2002. Copper River Subsistence Evaluation 2000 & Traditional
  Knowledge Project, Part One. Alaska Department of Fish and Game, Division of Subsistence, Final
  Report No. FIS 00-040, Anchorage, Alaska.
- Sisk, J., 1991. The Southeastern Alaska Salmon Industry : Historical Overview and Current Status 1–
  15.
- Sloat, M.R., Reeves, G.H., Christiansen, K.R., 2016. Stream network geomorphology mediates
  predicted vulnerability of anadromous fish habitat to hydrologic change in southeast Alaska. Global
  Change Biology 604–620. https://doi.org/10.1111/gcb.13466
- Smith, C.T., R. J. Nelson, C. C. Wood, and B. F. Koop. 2001. Glacial biogeography of North American
  Coho salmon (Oncorhynchus kisutch). Molecular Ecology 10: 2775-2785.
- Smith, C.T., Templin, W.D., Seeb, J.E., Seeb, L.W., 2005. Single Nucleotide Polymorphisms Provide
  Rapid and Accurate Estimates of the Proportions of U.S. and Canadian Chinook Salmon Caught in
  Yukon River Fisheries. North American Journal of Fisheries Management 25, 944–953.
  https://doi.org/10.1577/m04-143.1

Smoker, W.W., Gharrett, A.J., Stekoll, M.S. and Taylor, S.G., 2000. Genetic variation of fecundity
and egg size in anadromous pink salmon Oncorhynchus gorbuscha Walbaum. Alaska Fishery
Research Bulletin, 7(1), pp.44-50.

Smoker, W.W., Gharrett, A.J., Stekoll, M.S., 1998. Genetic variation of return date in a population of
pink salmon: a consequence of fluctuating environment and dispersive selection? Alaska Fishery
Research Bulletin 5, 46–54.

Smoker, W.W., Gharrett, A.J., Stekoll, M.S., Joyce, J.E., 1994. Genetic analysis of size in an
anadromous population of pink salmon. Canadian Journal of Fisheries and Aquatic Sciences 51, S9–
S15.

Somerville, M. A., and T. R. Hansen. 2021. Fishery management report for the recreational, personal
use, and subsistence fisheries of the Upper Copper/Upper Susitna Management Area, 2019. Alaska
Department of Fish and Game, Fishery Management Report No. 21-07, Anchorage.

Springer, A.M., McRoy, C.P., 1993. The paradox of pelagic food webs in the northern Bering S e a o
HI . Patterns of primary production. Continental Shelf Research 13, 575–599. https://doi.
org/10.1016/0278-4343(93)90095-f

Springer, A.M., McRoy, C.P., Turco, K.R., 1989. The paradox of pelagic food webs in the northern
Bering Sea—II. Zooplankton communities. Continental Shelf Research 9, 359–386. https://doi.
org/10.1016/0278-4343(89)90039-3

Springer, A.M., van Vliet, G.B., 2014. Climate change, pink salmon, and the nexus between bottomup and top-down forcing in the subarctic Pacific Ocean and Bering Sea. Proceedings of the National
Academy of Sciences 111, E1880–E1888. https://doi.org/10.1073/ pnas.1319089111

St. Saviour, A., A. W. Barclay, and N. Logelin. 2020. Northern Cook Inlet Chinook salmon marine
harvest stock composition, 2016–2017. Alaska Department of Fish and Game, Fishery Data Series
No. 20-27, Anchorage.

Stachura, M.M., Essington, T.E., Mantua, N.J., Hollowed, A.B., Haltuch, M.A., Spencer, P.D.,
Branch, T.A., Doyle, M.J., 2014. Linking Northeast Pacific recruitment synchrony to environmental
variability. Fisheries Oceanography 23, 389–408. https://doi.org/10.1111/fog.12066

2920 State of Alaska Salmon and People (SASAP) <u>https://alaskasalmonandpeople.org/</u>

Stopha, M. 2012. An evaluation of the Port Graham salmon hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J12-28, Anchorage.

Stopha, M. 2012. An evaluation of the Trail Lakes salmon hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J12-21, Anchorage.

Stopha, M. 2013. An evaluation of the Armin F. Koernig salmon hatchery for consistency with
statewide policies and prescribed management practice. Alaska Department of Fish and Game,
Division of Commercial Fisheries, Regional Information Report 5J13-11, Anchorage.

Stopha, M. 2013. An evaluation of the Cannery Creek salmon hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J13-06, Anchorage.

Stopha, M. 2013. An evaluation of the Eklutna salmon hatchery for consistency with statewide policies
and prescribed management practice. Alaska Department of Fish and Game, Division of Commercial
Fisheries, Regional Information Report 5J13-02, Anchorage.

Stopha, M. 2013. An evaluation of the Gulkana salmon hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J13-05, Anchorage.

Stopha, M. 2013. An evaluation of the Main Bay salmon hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J13-07, Anchorage.

Stopha, M. 2013. An evaluation of the Solomon Gulch salmon hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J13-04, Anchorage.

Stopha, M. 2013. An evaluation of the Wally Noerenberg Hatchery for consistency with statewide
policies and prescribed management practice. Alaska Department of Fish and Game, Division of
Commercial Fisheries, Regional Information Report 5J13-10, Anchorage.

Stopha, M. and J. Musslewhite. 2012. An evaluation of the Tutka Bay Lagoon salmon hatchery for
consistency with statewide policies and prescribed management practices. Alaska Department of Fish
and Game Division of Commercial Fisheries, Regional Information Report 5J12-05, Anchorage.

Stram, D.L., Ianelli, J.N., 2009. Eastern Bering Sea Pollock Trawl Fisheries: Variation in Salmon
Bycatch over Time and Space. American Fisheries Society Symposium 70, 827–850.

Stram, D.L., Ianelli, J.N., 2015. Evaluating the efficacy of salmon bycatch measures using fisherydependent data. ICES Journal of Marine Science 72, 1173–1180. https://doi.org/10.1093/icesjms/
fsu168

Sturdevant, M. V., E. Fergusson, N. Hillgruber, C. Reese, J. Orsi, R. Focht, A. Wertheimer, and B.
Smoker. 2011. Lack of trophic competition among wild and hatchery juvenile chum salmon during
early marine residence in Taku Inlet, Southeast Alaska. Environmental Biol. Fish. DOI
10.1007/s10641-011-9899-7.

Sturdevant, M. V., R. Brenner, E. A. Fergusson, J. A. Orsi, and W. R. Heard. 2013. Does Predation
by Returning Adult Pink Salmon Regulate Pink Salmon or Herring Abundance? North Pacific
Anadromous Fish Commission Technical Report No. 9: 153-164.

Sturdevant, M.V., Fergusson, E., Hillgruber, N., Reese, C., Orsi, J., Focht, R., Wertheimer, A.,
Smoker, B., 2012. Lack of trophic competition among wild and hatchery juvenile chum salmon during
early marine residence in Taku Inlet, Southeast Alaska. Environmental Biology of Fishes 94, 101–
116. https://doi.org/10.1007/ s10641-011-9899-7

Sturdevant, M.V., Sigler, M.F. and Orsi, J.A., 2009. Sablefish predation on juvenile Pacific salmon in
the coastal marine waters of Southeast Alaska in 1999. Transactions of the American Fisheries
Society, 138(3), pp.675-691.

Su, Z., Adkison, M.D., 2002. Optimal in-season management of pink salmon (Oncorhynchus gorbuscha) given uncertain run sizes and seasonal changes in economic value. Canadian Journal of Fisheries and Aquatic Sciences 59, 1648–1659. https://doi.org/10.1139/ f02-133

2973 **T** 

Tadokoro, K., Ishida, Y., Davis, N.D., Ueyanagi, S., Sugimoto, T., 1996. Change in chum salmon
(Oncorhychus keta) stomach contents associated with fluctuation of pink salmon (O. gorbuscha)
abundance in the central subarctic Pacific and Bering Sea. Fisheries Oceanography 5, 89–99.

Tarpey, C. M., J. E. Seeb, G. J. McKinney, W. D. Templin, A. V. Bugaev, S. Sato, and L. W. Seeb.
2017. SNP data describe contemporary population structure and diversity in allochronic lineages of
pink salmon (Oncorhynchus gorbuscha). Canadian Journal of Fisheries and Aquatic Sciences Online
15 August 2017. doi:10.1139/ cjfas-2017-0023.

Templin, W. D., J. E. Seeb, J. R. Jasper, A. W. Barclay, and L. W. Seeb. 2011. Genetic differentiation
of Alaska Chinook salmon: the missing link for migratory studies. Molecular Ecology Resources 11:
226-246.

- 2984 Templin, W.D., 2001. The history of propagation and transportation of Chinook salmon
- 2985 Oncorhynchus tshawytscha stocks at hatcheries in Southeast Alaska, 1972 1998. Regional

2986 Information Report 5J01-05, 48 pp.

Templin, W.D., Seeb, J.E., Jasper, J.R., Barclay, A.W., Seeb, L.W., 2011. Genetic differentiation of
Alaska Chinook salmon: the missing link for migratory studies. Molecular ecology resources 11 Suppl
1, 226–46. https://doi.org/10.1111/j.1755-0998.2010.02968.x

Templin, W.D., Seeb, J.E., Jasper, J.R., Barclay, A.W., Seeb, L.W., 2011. Genetic differentiation of
Alaska Chinook salmon: the missing link for migratory studies. Molecular ecology resources 11 Suppl
1, 226–46. https://doi.org/10.1111/j.1755-0998.2010.02968.x

Thedinga, J.F., Wertheimer, A.C., Heintz, R.A., Maselko, J.M., Rice, S.D., 2000. Effect of stock,
coded-wire tagging, and transplant on straying of pink salmon (Oncorhynchus gorbuscha) in
southeastern Alaska. Canadian Journal of Fisheries and Aquatic Sciences 57, 2076–2085.

Theriault, V., Moyer, G.R., Jackson, L.S., Blouin, M.S. and Banks, M.A., 2011. Reduced reproductive
success of hatchery coho salmon in the wild: insights into most likely mechanisms. Molecular
Ecology, 20(9), pp.1860-1869.

Tiegs, S.D., Chaloner, D.T., Levi, P., Rüegg, J., Tank, J.L., Lamberti, G.A., 2008. TIMBER
HARVEST TRANSFORMS ECOLOGICAL ROLES OF SALMON IN SOUTHEAST ALASKA
RAIN FOREST STREAMS. Ecological Applications 18, 4–11. https://doi.org/10.1890/07-0655.1

Tiernan A., T. Elison, T. Sands, and J. Head. 2022. Overview of the Bristol Bay commercial salmon
fishery, 2019–2022: a report to the Alaska Board of Fisheries. Alaska Department of Fish and Game,
Special Publication No. 22-17, Anchorage.

Trainor et al. 2019. Local and Traditional Knowledge of Freshwater Life Stages of Chinook and CHum
Salmon in Anvik, Huslia, Allakaket, and Fort Yukon

Trainor et al. 2021. How Subsistence Salmon Connects Households and Communities: an explorationof Salmon Production and Exchange Networks in Three Communities on the Yukon River, 2018-2019

Trudel, M., Thiess, M.E., 2007. Regional Variation in the Marine Growth and Energy Accumulation
of Juvenile Chinook Salmon and Coho Salmon along the West Coast of North America. American
Fisheries Society Symposium 57, 205–232.

3012 Twardek WM, Lapointe NWR, Cooke SJ (2022) High egg retention in Chinook Salmon
3013 Oncorhynchus tshawytscha carcasses sampled downstream of a migratory barrier. J Fish Biol 100:
3014 715–726.

3015 Twardek, W.M. 2022. Evaluating the consequences of physical barriers on fish during long-distance3016 upstream migrations through rivers. Thesis Carleton University.

3017 U

3018 University of Washington Alaska Salmon Program https://alaskasalmonprogram.org/

3019 United States and Canada. 2006. Potential causes of size trends in Yukon River Chinook salmon3020 populations. Regional Information Report No. 3A06-07. Yukon River Joint Technical Committee.

3021 Urawa, S., Nagasawa, K., Margolis, L., Moles, A., 1998. Stock identification of chinook salmon
3022 (Oncorhynchus tshawytscha) in the north Pacific Ocean and Bering Sea by parasite tags. North Pacific
3023 Anadromous Fisheries Commission Bulletin 1, 199–204.

Urawa, S., Sato, S., Crane, P.A., Agler, B., 2009. Stock-specific ocean distribution and migration of
chum salmon in the Bering Sea and North Pacific Ocean. NPAFC Bulletin 5, 131–146.

3026 Utter, F.M., McPhee, M.V., and Allendorf, F.W. 2009. Population genetics and the management of 3027 Arctic-Yukon-Kuskokwim salmon populations. American Fisheries Society Symposium 70: 97-123.

3028 V

Vallion, A.C., Wertheimer, A.C., Heard, W.R., Martin, R.M., 1981. Summary of data and research
pertaining to the pink salmon population at Little Port Walter, Alaska, 1964-80. NWAFC Processed
Report 81-10.

Vanessa R. von Biela, Bowen, L., McCormick, S.D., Carey, M.P., Donnelly, D.S., Waters, S., Regish,
A.M., Laske, S.M., Brown, R.J., Larson, S., Zuray, S., and Zimmerman, C.E. 2020. Evidence of
prevalent heat stress in Yukon River Chinook salmon. Canadian Journal of Fisheries and Aquatic
Sciences. 77(12): 1878-1892. https://doi.org/10.1139/cjfas-2020-0209

3036 Various resources from Indigenizing Salmon Management and other work

Vega, S. L., J. M. Head, T. Hamazaki, J. W. Erickson, and T. R. McKinley. 2022. Review of salmon
escapement goals in Bristol Bay, Alaska, 2021. Alaska Department of Fish and Game, Fishery
Manuscript No. 22-07, Anchorage.

Vega, S.L., T.M> Sutton, and J.M. Murphy. 2016. Marine-entry timing and growth rates of juvenile
Chum salmon in Alaska waters of the Chukchi and northern Bering seas. Deep-Sea Research II

Volkov, A.F. 2022. Appendicularia in the Bering, Okhotsk, Chuckchi Seas and North Pacific and their
significance for feeding nekton. Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr., 2022,
vol. 202, no. 2, pp. 390–408. DOI: 10.26428/1606-9919-2022-202- 390-408. EDN: BXOLJN.

von Biela VR, Sergeant CJ, Carey MP, Liller Z, Russell C, Quinn-Davidson S, Rand PS, Westley
PAH, Zimmerman CE (2022) Premature Mortality Observations among Alaska's Pacific Salmon
During Record Heat and Drought in 2019. Fisheries 47: 157–168.

Vulstek, S. C., J. R. Russell, J. E. Joyce, and A. K. Gray. 2022. 2017 Auke Creek research station
report: data summary and historical tredns from 1980 to 2017. NOAA Technical Memorandum
NMFS-AFSC-436.

3051 W

Wadle, J. and Baumer, J. 2023. Chignik River Sockeye Salmon Stock of Concern Action Plan. OralReport to Alaska Board of Fisheries. Alaska Department of Fish and Game

Walker, R., Myers, K., Davis, N.D., Aydin, K.Y., Friedland, K.D., Carlson, H.R., Boehlert, G.W.,
Urawa, S., Ueno, Y., Anma, G., 2000. Diurnal variation in thermal environment experienced by
salmonids in the North Pacific as indicated by data storage tags. Fisheries Oceanography 9, 171–186.

Walsey, V., Brewer, J., 2018. Managed out of existence: over-regulation of Indigenous subsistence
fishing of the Yukon River. GeoJournal 83, 1169–1180. https://doi.org/10.1007/s10708-018-9879-y

Wang, I.A., Leder, E.H., Smoker, W.W., Gharrett, A.J., 2006. Timing of development during epiboly
in embryos of second-generation crosses and backcrosses between odd- and even-broodyear pink
salmon, Oncorhynchus gorbuscha. Environmental Biology of Fishes 75, 325–332.

Wang, J., Zhang, J., Watanabe, E., Ikeda, M., Mizobata, K., Walsh, J.E., Bai, X., Wu, B., 2009. Is the
dipole anomaly a major driver to record lows in arctic summer sea ice extent? Geophysical Research
Letters 36, 1–5. https://doi.org/10.1029/2008GL036706

Wang, M., Overland, J.E., Stabeno, P., 2012. Future climate of the Bering and Chukchi Seas projected
by global climate models. Deep-Sea Research Part II: Topical Studies in Oceanography 65–70, 46–
57. https://doi.org/10.1016/j.dsr2.2012.02.022

WAPLES, R., 2009. Conserving the evolutionary legacy of Arctic-Yukon-Kuskokwim salmon. in
Krueger, Charles & Zimmerman, Christian. 2009. Pacific salmon: ecology and management of
western Alaska's populations. pp 125–139.

Wechter, M. E., B. R. Beckman, A. G. Andrews III, A. H. Beaudreau, and M. V. McPhee. 2017.
Growth and condition of juvenile chum and pink salmon in the northeastern Bering Sea. Deep Sea
Research Part II: Topical Studies in Oceanography 135:145–155.

Wechter, M.E., B.R. Beckman, A.G. Andrews, A.H. Beaudreau, and M.V. McPhee. 2017. Growth
and condition of juvenile chum and pink salmon in the northeastern Bering Sea. Deep-Sea Research
II 135:145-155.

Weingartner, T.J., Danielson, S.L., Royer, T.C., 2005. Freshwater variability and predictability in the
Alaska Coastal Current. Deep-Sea Research Part II: Topical Studies in Oceanography 52, 169–191.
https://doi.org/10.1016/j.dsr2.2004.09.030

Weitkamp, L. A., J. A. Orsi, K. W. Myers, and R. C. Francis. 2011. Contrasting Early Marine Ecology
of Chinook Salmon and Coho Salmon in Southeast Alaska: Insight into Factors Affecting Marine
Survival, Marine and Coastal Fisheries, 3:1, 233-249, DOI:10.1080 /19425120.2011.588919.

Welch, D.W., A.D. Porter, and E.L. Rechisky. 2020. A synthesis of the coast-wide decline in survival
of West Coast Chinook salmon. Fish and Fisheries DOI:10.1111/faf.12514.

Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2013. Forecasting pink salmon
harvest in Southeast Alaska from juvenile salmon abundance and associated biophysical parameters:
2012 returns and 2013 forecast. North Pacific Anadromous Fish Commission Document 1486. 23 pp.

Wessel, M.L., Smoker, W.W., Joyce, J.E., 2006. Variation of morphology among juvenile chinook
salmon of hatchery, hybrid, and wild origin. Transactions of the American Fisheries Society 135, 333–
340. https://doi.org/10.1577/T04-078.1

Whittle, J.A., Kondzela, C.M., Nguyen, H.T., Hauch, K., Cuadra, D., Guyon, J.R., 2018. Genetic stock
composition analysis of Chum Salmon from the prohibited species catch of the 2016 Bering Sea

3093 Walleye Pollock trawl fishery and Gulf of Alaska groundfish fisheries. 56. 3094 https://doi.org/10.7289/V5/TM-AFSC-314

Whittle, J.A., Kondzela, C.M., Watson, J.T., Barry, P.D., Nguyen, H.T., Yasumiishi, E.M., Nicolls,
D., Larson, W.A., 2021a. Genetic Stock Composition Analysis of Chum Salmon from the Prohibited
Species Catch of the 2018 Bering Sea Walleye Pollock Trawl Fishery and Gulf of Alaska Groundfish
Fisheries. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-, 81 p.

- Williams, D.L., Shelden, C.A., 2011a. Fishery Data Series No . 11-49 Kogrukluk River Salmon
  Studies , 2010 by.
- Williams, D.L., Shelden, C.A., 2011b. Fishery Data Series No . 11-49 Kogrukluk River Salmon
  Studies , 2010 by.
- Williams, L., P. Coiley-Kenner, and D. Koster. 2010. Subsistence harvests and uses of salmon, trout,
  and char in Akhiok, Larsen Bay, Old Harbor, Ouzinkie, and Port Lions, Alaska, 2004 and 2005. Alaska
- 3105 Department of Fish and Game, Division of Subsistence, Technical Paper No. 329, Anchorage.
- Wilson, L. 2023. Alaska salmon fisheries enhancement annual report 2022. Alaska Department of Fish
  and Game, Division of Commercial Fisheries, Regional Information Report No. 5J23-04, Juneau.
- 3108 Wilson, L., 2023. Alaska Salmon Fisheries Enhancement Annual Report 2022. RIR No. 5J23-04
- Wipfli, M.S., 1997. Terrestrial invertebrates as salmonid prey and nitrogen sources in streams:
  contrasting old-growth and young-growth riparian forests in southeastern Alaska, U.S.A. Canadian
  Journal of Fisheries and Aquatic Sciences 54, 1259–1269. https://doi.org/10.1139/f97-034
- Wipfli, M.S., Gregovich, D.P., 2002. Export of invertebrates and detritus from fishless headwater
  streams in southeastern Alaska: Implications for downstream salmonid production. Freshwater
  Biology 47, 957–969. https://doi.org/10.1046/j.1365-2427.2002.00826.x
- Wipfli, M.S., Hudson, J., Caouette, J., 1998. Influence of salmon carcasses on stream productivity :
  response of biofilm and benthic macroinvertebrates in southeastern Alaska, 1511, 1503–1511.
- Wipfli, M.S., Hudson, J.P., Chaloner, D.T., Caouette, J.P., 1999. Influence of salmon spawner
  densities on stream productivity in Southeast Alaska. Canadian Journal of Fisheries and Aquatic
  Sciences 56, 1600–1611. https://doi.org/10.1139/f99-087
- Witherell, D., Ackley, D., Coon, C., 2002. An Overview of Salmon Bycatch in Alaska Groundfish
  Fisheries. Alaska Fishery Research Bulletin 9, 53–64.
- Witteveen, M. J., and K. Shedd. 2016. Chinook salmon genetic sampling along the Alaska Peninsula
  and adjacent areas results, 2012–2014. Alaska Department of Fish and Game, Fishery Data Series No.
  16-25, Anchorage.

Wolfe, R.J., and Spaeder, J. 2009. People and salmon of the Yukon and Kuskokwim drainages and
Norton Sound in Alaska: fishery harvests, culture change, and local knowledge systems. American
Fisheries Society Symposium 70: 349-379.

Woodgate, R. a., Aagaard, K., Weingartner, T.J., 2006. Interannual changes in the Bering Strait fluxes
of volume, heat and freshwater between 1991 and 2004. Geophysical Research Letters 33, 2–6.
https://doi.org/10.1029/2006GL026931

Woody, C. A., editor. 2018. Bristol Bay Alaska: natural resources of the aquatic and terrestrial
ecosystems. J. Ross Publishing, Plantation, Florida. 589 pp.

Woody, C. A., J. Olsen, J. Reynolds, and P. Bentzen. 2000. Temporal variation in genotypic and
phenotypic traits of two sockeye salmon populations. Transactions of the American Fisheries Society
129: 1031-1043.

3136 Y

Yasumiishi, E.M., Criddle, K.R., Hillgruber, N., Mueter, F.J., Helle, J.H., 2015. Chum salmon (
Oncorhynchus keta ) growth and temperature indices as indicators of the year-class strength of age-1
walleye pollock (Gadus chalcogrammus) in the eastern Bering Sea. Fisheries Oceanography 24, 242–
256. https://doi.org/10.1111/ fog.12108

Yasumiishi, E.M., E.V. Farley, G.T. Ruggerone, B.A. Agler, and L.I. Wilson. 2016. Trends and factors
influencing the length, compensatory growth, and size-selective mortality of juvenile Bristol Bay,
Alaska, sockeye salmon at sea. Marine and Coastal Fisheries 8(1):315-333.

Yasumiishi, E.M., E.V. Farley, Jr., J.Maselko, K.Y. Aydin, et al. 2019. Differential north-south
response of juvenile Chinook salmon marine growth to ecosystem change in the eastern Bering Sea,
1974 - 2010. ICES Journal of Marine Science, doi:10.1093/icesjms/ fsz166

Yasumiishi, E.M., Farley, E.V., Ruggerone, G.T., Agler, B.A., and Wilson, L.I. 2016. Trends and
factors influencing the length, compensatory growth, and size-selective mortality of juvenile Bristol
Bay, Alaska, sockeye salmon at sea. Marine and Coastal Fisheries: Dynamics, Management, and
Ecosystem Science 8:315-333. DOI: 10.1080/19425120.2016.1167793

3151 YRJTC. 2006. Potential causes of size trends in Yukon River Chinook salmon populations. RIP No.3152 3A06-07

#### 3153 Z

Zimmermann, H.H., et al. 2023. Marine ecosystem shifts with deglacial sea-ice loss inferred from
ancient DNA shotgun sequencing. Nature Communications: 14, Article number: 1650. https://doi.
org/10.1038/s41467-023-36845-x

- 3157 Zuray, S., Kocan, R., Hershberger, P., 2012. Synchronous Cycling of Ichthyophoniasis with Chinook
- 3158 Salmon Density Revealed during the Annual Yukon River Spawning Migration. Transactions of the
- 3159
   American Fisheries Society 141, 615–623. https://doi.org/10.108 0/00028487.2012.683476
- 3160 *Public comments from the Web Site here (during April 2024)*

#### 3161 Appendix 4

#### 3162 THE 2023–2027 NPAFC SCIENCE PLAN Primary Goal and Research Objectives

The primary goal of the 2023–2027 Science Plan is to: "Establish a research framework to develop a mechanistic understanding of the impact of changing climate on salmon abundance and distribution trends in the North Pacific Ocean." (1) Improve knowledge of the relative biomass, distribution, migration, and fitness of Pacific salmon in the ocean (Present Knowledge); (2) Understand causes and anticipate changes in the production of Pacific salmon and the marine ecosystems producing them (Forward Action).

- 3169 Improved understanding of the mechanisms that regulate the distribution and abundance of Pacific
- 3170 salmon will promote the conservation of anadromous populations in the North Pacific Ocean, allow
- 3171 for better projections, or at least include realistic uncertainty given climate change, of Pacific salmon
- 3172 production trends in the future, and enhance the sustainable fisheries management, food security, and
- 3173 economic security in member nations.
- The timing of the NPAFC 2023–2027 Science Plan overlaps with the proposed implementation of
- 3175 Basin-scale Events to Coastal Impacts (BECI; 2021–2030). It is anticipated that a BECI science plan
- 3176 will be finalized at the PICES Annual Meeting during fall 2023.

#### 3177 NPAFC Research Themes

- 3178 (1) Status of Pacific salmon and steelhead trout (Present Knowledge);
- 3179 (2) Pacific salmon and steelhead trout in a changing North Pacific Ocean (Forward Action);
- 3180 (3) New technologies;
- 3181 (4) Management systems;
- 3182 (5) Integrated information systems.

#### 3183 Theme 1. Status of Pacific Salmon and Steelhead Trout (Present Knowledge)

#### 3184 *Outcome: The present status of salmon and their environments is documented and reported.*

The purpose of this theme is to document and effectively report on the present status of salmon and 3185 their habitats. The NPAFC collates annual statistics on catch, escapement, and hatchery releases of 3186 Pacific salmon around the Pacific Rim. There is an ongoing need to maintain and improve 3187 monitoring of spawning escapement, catch, smolt production and other biological information for 3188 3189 potential use in the projecting salmon return strength or ocean survival. Long-term time series are particularly valuable in understanding linkages between climate and Pacific salmon production. Data 3190 on hatchery fish should be maintained separately from data on wild fish as much as practicable. 3191 3192 Biological information such as age composition of a population, body size, fecundity and egg size are monitored whenever feasible. 3193

121

#### 3194 (1-1) Status of Key Salmon Populations

- 3195 *Monitor Key Populations*—Continue reporting on ongoing monitoring programs for key 3196 salmon populations and identify new sampling opportunities. Identify additional key 3197 populations that can be monitored to provide status information for co-existing salmon 3198 populations and their ecosystems.
- 3199Stock Assessments—Monitor current and emerging stock assessment methods in cooperation3200with partners potentially including ICES, NASCO and the Pacific Salmon Commission.
- *Report on Status of Salmon in the NPO*—Report annually on the Status of Salmon in the
   NPO. Consider utilizing the Interactive Mapping System developed within the NPAFC
   Working Groupon Salmon Marking. Could be northern hemispheric in scope in cooperation
   with Atlantic and Arctic partners.
- 3205 *Data Quality*—Improve the quantification and documentation of uncertainty associated with 3206 existing and new data time series and maintain wild and hatchery salmon data separately in 3207 the timeseries.
- New Baseline Information—Provide a data review and annual Pacific salmon hatchery and 3208 wild abundance data updates to Ruggerone and Irvine (2018). These methods could be drawn 3209 from those described in Ruggerone et al. (2010) and Ruggerone and Irvine (2018) and 3210 adapted as needed. These data would be reviewed and provided by NPAFC member 3211 countries as part of the WGSA annual workplans. These data would be managed and 3212 warehoused by the NPAFC, similar to the catch and hatchery release statistics: 3213 https://npafc.org/statistics/. The long-term goal will be to make it possible for each party to 3214 easily estimate the annual wild and hatchery abundance and biomass of salmon in the North 3215 Pacific Ocean. 3216

#### 3217 (1-2) Monitor Salmon in the Ocean

- 3218 Gathering information on the marine ecology of Pacific salmon is critical to our understanding of how climate variability impact ecosystem function, salmon fitness, 3219 distribution, migration and survival. Anadromous salmon migrate in the ocean to maximize 3220 their growth and survival. Their seasonal migration and distribution patterns are stock 3221 3222 specific, and fundamental migration routes may be genetically fixed. Increasing information on seasonal ocean migration and distribution of key salmon populations contributes to: 3223 planning effective ocean monitoring surveys, better climate modelling and projecting, better 3224 management to avoid incidental salmon bycatch, and efficient enforcement activities to 3225 protect salmon in the ocean. 3226
- 3227 Therefore, the recommendation is to:
- Continue integrated ecosystem marine survey monitoring activities currently
   conducted by Parties within respective exclusive economic zones and the Convention
   area to collect observations on the biological and physical oceanographic

- 3231 characteristics and observations on size-at-age, external traits, gonads, health and
  3232 condition (e.g., energy density/lipid, thiamine deficiency, parasites and diseases),
  3233 stomach contents, and potential population impacts.
  3234 Monitor northward expansion of salmon into Arctic regions (e.g. northern Bering)
- Theme 2. Pacific Salmon and Steelhead Trout in a Changing North Pacific Ocean (Forward Action)
- 3238 *Outcome: The effects of natural environmental variability and human factors affecting salmon* 3239 *distribution and abundance are understood and quantified.*

Climate change may result in significant variability and overall declines in the carrying capacity and usable habitat (distribution) of Pacific salmon in the North Pacific Ocean, potentially leading to expanded use of the Arctic Ocean, at least seasonally. An improved understanding of linkages between environmental changes and Pacific salmon production will help to plan for the economic consequences of these changes. The objectives are to understand and quantify the effects of environmental variability and anthropogenic factors affecting salmon distribution and abundance, and to project future changes with improved models.

3247 (2-1) Pacific Salmon Distribution/Migration, Climate and Ocean Changes

Sea; Chukchi Sea; Beaufort Sea).

3235

- In recent years, there have been shifts in the distribution of salmon in northern regions, but some declines at the southern edges of their distribution along the Asian and North American continents. These geographical shifts in salmon abundance may be related to climate-induced changes in habitat/environments operating at regional and local scales. What are the relevant mechanisms influencing shifts spatial distribution and migration? What is driving Pacific salmon movement into the Arctic?
- 3254 (2-2) Pacific Salmon Density Dependence, Carrying Capacity, Climate and Ocean Changes
- With the potential of limited food resources in the ocean, it is important to understand the implications of habitat use by Pacific salmon populations at various levels of abundance, the productive capacity of habitats for each life stage, and the potential implications of density dependent effects.
- There is a need to understand odd/even year differences in survival/growth of salmon species thathas been correlated with pink salmon abundance. Is this a top-down effect? Or are there otherexplanations that may help explain this correlation?•
- There is a need for more comprehensive studies on the role of salmon in pelagic communities, thefood availability for salmon and the nutritional quality of prey organisms.•
- 3264 Understand inter-and intra-specific competition among salmon at sea.
- 3265 (2-3) Pacific Salmon Critical Periods, Climate and Ocean Changes

- Variation in the early marine survival of Pacific salmon has been hypothesized to have a major role in determining the numbers of adults that return to spawn. However, there has been limited evidence to support this hypothesis. We need to understand the causes of mortality at each stage of the salmon life cycle and evaluate whether any particular life history period is critical.
- 3271 *Ocean Entry*—Juvenile abundance, timing and body size at ocean entry may be important 3272 parameters that are critical to understanding and quantifying mortality at sea. Examine how 3273 these parameters are associated with salmon survival or brood year strength.
- 3274 *Growth*—Increased energy efficiency for growth of juveniles in the early marine period may 3275 be a key to their survival and optimization of hatchery production.
- 3276 *Prey Organisms*—Identify which prey organisms are important for salmon growth at each
  3277 stage and region, and examine if the abundance of prey organisms limits salmon production.
- 3278 *Salmon Health*—Examine effects of pathogens and stressors on the growth and survival of salmon in the ocean.

#### 3280 (2-4) Modelling the Future for Salmon

- Reliable projection models of future salmon distribution, abundance and survival is important for sustainable resource management and for projecting future variations in production due to changing climate. Researchers and analysts should consider developing statistical models as well as ecosystem models coupled with biophysical models to estimate the impact of climate change on salmon populations, and to create future scenarios for salmon distribution and abundance. •
- Explain the unequal stock/species specific response of Pacific salmon to climate change.
   E.g., why are Asian pink salmon and Bristol Bay sockeye thriving under contemporary
   conditions while other species/stocks are not doing as well?
- Model projections of impacts of climate change on salmon production and make progress in
   understanding unexplained variability in salmon abundance, migration, growth, size-at-age,
   and survival.
- 3293 Theme 3. New Technologies
- 3294 *Outcome: New technologies and analytical methods are advanced and applied to salmon research.*

Novel stock and fish identification methods including new molecular techniques, hatchery mass marking, and intelligent tags continue to be developed, and these tools are integral to comprehensive and cost-effective monitoring and mechanistic studies to facilitate the formulation of effective models predicting the distribution and abundance of salmon populations. Although considerable progress has been made in both the basic understanding of population differentiation of mixed marine salmonid assemblages and in genetic research technologies, this knowledge is still insufficient to understand the spatial distribution of different populations in the ocean and the

differences in their responses to changing environmental conditions. Implementing genetic methods
 to differentiate mixed marine salmonid assemblages and to expand the database of reference samples
 are increasingly needed.

- *Molecular Identification*—Develop effective molecular techniques and baselines to identify
   the geographical origin of individual fish/population.
- Genomics—Use genomic technology for the rapid assessment of the physiological health
   status and cause of the condition of salmon.
  - *Environmental DNA (eDNA)* —Develop eDNA methods for the rapid and non-lethal estimation of salmon distribution and potentially abundance.
- *Mass Marking*—Develop mass marking techniques to identify hatchery salmon in mixtures
   of populations. Thermal otolith marking is a successful tool for mass marking, but available
   mark patterns are limited.
- Intelligent Tags—Develop tagging methods to investigate the habitat conditions, predators
   and navigation mechanism of salmon migrating in the ocean.
- Salmon Observation Systems—Improve tracking technologies to increase knowledge of stock-specific patterns of migration and survival at each life stage.
- *Remote Sensing/Autonomous Vehicles*—Application of remote sensing technologies such
   satellite imagery and ocean gliders/saildrones outfitted with sensors such as acoustics and
   new camera technologies to understand changes in the biophysical environment experienced
   by salmon.
- 3322 The NPAFC should focus on the following new tools and activities to improve salmon identification:
- Conduct additional pink salmon genetic baseline studies to address questions regarding GSI and range expansion.
  - Expand our understanding of eDNA methods, appropriate use and application.
- Develop and standardize Pacific salmon genetic data and analysis methods for a
   comprehensive coastwide genetic baseline database.

#### 3328 Theme 4. Management Systems

3309

3310

3325

#### 3329 *Outcome: Information required for effective management systems is available and applied to* 3330 *enforcement activities, social systems, and salmon movements into the Arctic.*

- The objective is to provide scientific advice that effectively informs management systems including their cultural and socioeconomic aspects. Enforcement of NPAFC's convention measures that
- prohibit directed fishing for anadromous fish within the Convention Area is the responsibility of
- NPAFC's Committee on Enforcement (ENFO). The CSRS is increasingly playing a role in
- providing information on the probable distribution of Pacific salmon at different times of the year,
- and therefore likely locations of illegal, unreported, and unregulated (IUU) fishing.

- Climate change adaptation is a social process, and this is one of the key challenges facing salmon,
- ecosystems and humans moving forward. At this time, we allocate very little research effort towards
- the sociology of adaptation process in societies. The role of NPAFC to form collaborations on
- salmon, will need to consider how shifting salmon distributions and abundances, among other
- 3341 species, may disrupt the NPAFC role and connections moving forward. This involves shifting away
- from status quo ways of managing fisheries and salmon, with transformational shifts in our
- 3343 management systems.

With a northward shift in salmon distributions given climate change, the impacts on coastal communities will be large and will require NPAFC to engage with other fields of knowledge, such as the social sciences, to facilitate connections, agreements, collaborations between Nations. The absence of this field of expertise in NPAFC represents a critical gap that will be required to meet the

3348 huge challenges of climate change.

#### 3349 Theme 5. Integrated Information Systems

Outcome: Freely available information systems mobilize and synthesize historic and current data
about salmon and their environment.

3352 It is essential that salmon and ecosystem data are readily available for researchers and for machine learning AI applications. Therefore, the goal is to build upon data mobilization approaches 3353 developed during the IYS. This includes: (1) the development of data mobilization approaches that 3354 ensure adherence to the FAIR principles with data Findable, Accessible, Interoperative and Re-3355 usable; (2) the application of a "federated" approach to integrating data sets from the respective 3356 parties in common agreed to data formats, e.g. the Global Ocean Observatory System standards 3357 (GOOS); (3) improve the ability to share information and collaborate on research efficiently using a 3358 3359 modern web-based framework. Data assembled as part of the other themes are to be linked in a central data system. 3360

**Cooperative Research Approaches and Implementation of the Science Plan** Pertinent 3361 3362 approaches to cooperative research under the 2023–2027 Science Plan will include compilation and synthesis of existing data and metadata to generate and test specific hypotheses, integration of 3363 3364 ecological monitoring research in the ocean using research vessels and/or remote sensing, conceptual and quantitative modeling, process-oriented field and laboratory studies, and retrospective analyses. 3365 Scientific results from cooperative studies using these approaches will progressively reduce major 3366 gaps in knowledge with respect to the research themes, as well as make significant contributions to 3367 BECI research in collaboration with other relevant partners such as ENFO, PICES, ICES and 3368 NASCO. New scientific information will also contribute to effective enforcement activities by 3369 3370 member nations to protect Pacific salmon from IUU fishing in the Convention Area. Progress on research themes or issues of the 2023–2027 Science Plan will be reviewed annually during the 3371 3372 NPAFC Annual Meeting. Symposia, workshops, and other science meetings will be scheduled during this time as appropriate. 3373

3374

3375 Appendix 5

#### 3376 ALASKA BYCATCH REVIEW TASK FORCE FINAL REPORT — NOVEMBER 2022

3377 Salmon Recommendations

Much of the salmon research identified was similar for both the Bering Sea/Aleutian Island and the
Gulf of Alaska. Listed below is the research identified for Western Alaska salmon and research which
is unique for the Gulf of Alaska.

- 3381 Western Alaska Salmon
- **3382** *Research Goals*
- Research to improve our ability to determine the stock of origin of chum and Chinook salmon taken as bycatch.
- Research to reduce bycatch through improved understanding of distribution and migration of
   Western Alaska chum and Chinook salmon stocks migration patterns to better predict and
   therefore avoid bycatch "hot spots" in the BSAI region.ng.
- Research that helps us understand the relative importance of particular mechanisms for drivingabundance of Western Alaska Chinook and chum.
- a) Improved information on marine migration patterns
- i. The projects AFSC mentioned that Sabrina Garcia (Chinook salmon) and Wes Larson
   (chum salmon) are leading in the Bering Sea: Model ocean distribution and migration of AK
   Chum and Chinook salmon stocks in the Bering Sea to predict distribution and hotspots.
- ii. A tagging project of immature chum salmon in the North Pacific Ocean to help usunderstand their destination, timing, and maturity.
- iii. A synthesis of marine migration information from fishery-dependent data sources, marinesurveys, and tagging studies, and how these patterns have changed with a changing climate.
- **b)** Improved information on the characteristics of fishery catches.
- i. There are still improvements that can be made in the ability to assess age, and specifically
   stock-specific age of Chinook and chum salmon caught in any marine fisheries.
- 3401 c) Improved information to help understand fishery impacts
- i. Improved Adult Equivelant (AEQ) modeling through 'stock specific' chinook and chum
  salmon bycatch. Particularly for western Alaska chum salmon, AEQ analyses are limited by:

- age classification data gaps in adult chum abundance across all of the western Alaska
   stock group. Studies that improve the ability to estimate abundance of all chum salmon in
   the western Alaska stock reporting group. Continued genetics work is needed.
- the ability to break up that reporting group. This might be remedied by using
  technologies that go beyond genetic assignment alone (use of pathogens, stable isotopes,
  etc.)
- 3410 Research that can provide an additional (non-adult) abundance estimate
- 3411 This will be really powerful for helping triangulate which life stages are most important for
- 3412 determining good or poor productivity. The committee recommends that research should span the
- 3413 life-cycle of the salmon species.
- a) Understand critical survival periods for western Alaska salmon through integrated ecosystem
   assessment surveys including expansion of the northern Bering Sea pelagic trawl survey into the
- 3416 near shore waters north of the Yukon River including Norton Sound.
- 3417 i. Similar research is being planned in the southern Bering Sea to have a more comprehensive3418 assessment of Western Alaska Chinook and chum.
- **3419 NOTE**: Neither of these projects are funded beyond 2023.
- ii. Ecosystem indicators: summer sea temperature, phytoplankton/zoo plankton community structure;
  salmon and pelagic fish catch per unit effort, distribution, energy density for fitness, size, stomach
  contents. These indicators are being utilized to understand climate impact on the northern Bering Sea
  ecosystem, fish fitness and survival. The recent information from the northern Bering Sea pelagic
  trawl survey suggests that the marine heat wave within the NBS during 2016 to 2019 negatively
  affected juvenile Chum salmon fitness (shift to low quality prey, increased metabolic rates due to
  higher SST), likely leading to high winter mortality. The data suggest that Chinook salmon
- 3427 abundance is impacted by factors affecting them in freshwater and early marine residence.
- **b)** Studies that help understand how ocean/climate conditions impact future runs
- i. Marine pelagic trawl surveys in the northern and southern Bering Sea can help us address this (see above).
- ii. NOAA and ADF&G are collaborating on using International Year of the Salmon (IYS) catches
  and samples to examine immature AYK chum salmon in the North Pacific Ocean during winter.
- 3433 (This is not yet funded.)
- iii. Immature salmon surveys (like the IYS surveys) in the Bering Sea and North Pacific Ocean.
  There is currently no funding support for charter vessel to conduct the survey, collecting and
  processing samples or paying for gear and supplies.
- 3437 c) Studies that help us understand the role of diet, health, and disease on the survival and spawning
  3438 success of Western AK Chinook and chum

**i.** Understanding vectors of Ichthyophonus infection for Yukon Chinook salmon, and whether it is

3440 causing significant en route mortality during the spawning migration

3441 **ii.** Understanding diet, nutrition, and condition of Western AK Chinook and chum stocks at juvenile

- 3442 (marine pelagic trawl surveys in the northern and southern Bering Sea see above), immature (IYS
- 3443 surveys, industry catches, etc.), and adult life stages (returning samples from lower river test
- 3444 fisheries- pilot work started for Yukon Chinook, but only funded through 2022).

#### 3445 Gulf of Alaska Chinook Salmon

Conduct annual genetic and spatial assessment of Gulf of Alaska (GOA) Chinook salmon. This recommendation is intended to include, in addition to the genetic assessment that is currently taking place, that efforts should be made to produce estimates of both the spatial and temporal bycatch of Alaska stocks of Chinook salmon, as well as characterizations of the age, sex and size of the bycatch of Chinook salmon identified as stocks of Alaska origin. If further progress can be made towards identifications of stock of origin of Alaska Chinook salmon taken as bycatch, that too should be pursued.

- **a)** Studies that help us understand the relative role of marine interceptions and bycatch.
- **i.** Improved information on marine migration patterns and its relation to fishery locations and timing.
- Extend the distribution and timing projects using bycatch data in the Bering Sea to include the western GOA.
- **ii.** Improved demographic information that will enable assessment of stock specific impacts.
- •Collect samples to improve demographic information such as stock, age, sex, size and maturity for
  Chinookand chum salmon caught in any marine fisheries.
- •Improved information to help understand fishery impacts through AEQ or similar analyses.
- b) Research that can provide an additional (non-adult) abundance estimate. This is useful for helpingtriangulate which life stages are most important for determining productivity.
- i. Juvenile salmon surveys: a survey occurs annually in the eastern GOA to monitor SoutheastAlaska salmonstocks (Southeast Coastal Monitoring project).
- •ADF&G will pilot a juvenile salmon survey in the western Gulf of Alaska in 2023. This will align with surveys in the
- northern and southern Bering Sea and Southeast Alaska to give a comprehensive assessment ofAlaska Chinook and chum salmon early in the marine life stages.
- Note: neither the GOA nor the Bering Sea projects are funded beyond 2023
- 3470 c) Studies that help us understand how ocean/climate conditions impact future runs.

- 3471 i. Marine pelagic trawl surveys in the Bering Sea and Gulf of Alaska (including western/central
- 3472 Alaska and SEAK surveys).
- 3473 ii. Immature salmon surveys (like the IYS surveys) in the Bering Sea, Gulf of Alaska, and North3474 Pacific Ocean.
- 3475
- 3476

#### 3477 Appendix 6

3478 Agencies and Non-profits that support Alaska salmon research/information

#### 3479 Alaska Bycatch Advisory Council

3480 https://www.adfg.alaska.gov/index.cfm? adfg=bycatchtaskforce.main

On March 10, 2023, the Commissioner of Fish and Game established the Bycatch Advisory Council

to advise the department on ways to implement the recommendations contained within the final

3483 report of the Alaska Bycatch Review Task Force (see Appendix 4).

#### 3484 U.S. Fish and Wildlife Service – Gravel to Gravel Initiative

3485 https://www.fws.gov/page/gravel-to-gravel-keystone-iniative

With Gravel to Gravel investments, the U.S. Fish and Wildlife Service is actively supporting and 3486 3487 funding a variety of projects that will ensure safe, resilient, and equitable futures for our people, salmon, land, and waters. We are working to shape this Initiative with local and regional partners, 3488 including the Tanana Chiefs Conference, Association of Village Council Presidents, Kawerak, Inc., 3489 3490 the Kuskokwim Intertribal Fish Commission, the Yukon River Intertribal Fish Commission, the Bureau of Land Management, USGS, National Park Service, the Yukon River Drainage Fisheries 3491 Association, State of Alaska, and nonprofit partners like Trout Unlimited. Importantly, the initiative 3492 is not a one-and-done effort. Gravel to Gravel-funded projects will build upon previous work and 3493 3494 partnerships, while catalyzing the future of our service in Alaska – leveraging new funding, and strengthening fresh relationships, as we continue our work in serving Alaska's people, ecosystems, 3495 3496 and wildlife.

#### 3497 U.S. DOI Bureau of Land Management (BLM) – Gravel to Gravel Keystone Initiative

3498 https://www.blm.gov/programs/aquatic-resources/alaska/ gravel-gravel-keystone-initiative

The Department of the Interior is investing more than \$16 mil-lion over the next four years from President Biden's **Bipartisan Infrastructure Law** to improve the resilience of ecosystems and salmon in Alaska's Yukon, Kuskokwim and Norton Sound region as part of the Gravel to Gravel Keystone Initiative. This initial multi-million-dollar investment serves as a catalyst for additional investments in Gravel to Gravel, which is made up of three elements:

- 3504 Investments to improve resiliency of Pacific salmon
- 3505 Renewed commitment to strengthening relationships through co-stewardship
- **3506** Responding to ecosystem threats to food security.

While the BLM is working across all three elements of Gravel to Gravel, we are heavily focused on
improving watershed resiliency through assessment and restoration. And we, in the BLM are doing
what we can where we can with the provided funding to make a positive and significant impact for

3510 the communities and ecosystems of the Yukon, Kuskokwim, and Norton Sound region.

#### **NOAA Fisheries – Alaska Fisheries Science Center** 3511

https://www.fisheries.noaa.gov/alaska/ecosystems/alaska-ecosystem-monitoring-and-assessment 3512

3513 Pacific salmon play an important role in Alaska's marine ecosystems and are a valuable commercial,

recreational, and subsistence resource. NOAA Fisheries scientists forecast salmon harvests, assess 3514

the impact of commercial fisheries on salmon, and evaluate how salmon populations respond to 3515 environmental changes. The information we provide helps managers make science-based decisions

- 3516 to ensure sustainable fish populations, fisheries, and fishing communities.
- 3517
- 3518 **Pacific Salmon Commission**
- https://www.psc.org/ 3519

The Pacific Salmon Commission is the body formed by the govern-ments of Canada and the United 3520

States in 1985 to implement the Pacific Salmon Treaty. It is our shared responsibility to conserve the 3521

Pacific Salmon in order to achieve optimum production and to divide the harvests so that each 3522 country reaps the benefits of its investment in salmon management. The Pacific Salmon Commission 3523

oversees two Endowment Funds established in 1999 to support projects in Canada and the United 3524 States that develop improved information for resource manage-ment; rehabilitate and restore marine 3525

3526 and freshwater salmon habitats; and, enhance wild stock production through low tech-nology

techniques. 3527

#### 3528 **Alaska Department of Fish and Game**

https://www.adfg.alaska.gov/ 3529

The Alaska Department of Fish and Game maintains active and comprehensive management and 3530 3531 research programs to ensure fish and wildlife populations are "utilized, developed, and maintained on the sustained yield principle," in accordance with Alaska's Constitution. 3532

#### Salmon Ocean Ecology Program 3533 •

https://www.adfg.alaska. gov/index.cfm? adfg=salmonoceanecology. main#:~:text=Our% 3534 20Mission,salmon%20in% 20the%20marine% 20environment. 3535

The Salmon Ocean Ecology Program supports statewide salmon fisheries management through the 3536 assessment and monitoring of salmon in the marine environment. Our goals are to understand the 3537 3538 marine life of Alaskan salmon, use this information to assist fishery management decision-making, and help answer pressing questions about marine processes that influence the abundance and 3539 characteristics of our salmon populations. 3540

- 3541 Alaska Sustainable Salmon Fund
- 3542 http://www.akssf.org/

3543 The Alaska Sustainable Salmon Fund (AKSSF) program, administered by the Alaska Department of Fish and Game, manages the State of Alaska's allocations from the federal Pacific Coastal 3544

- 3545 Salmon Recovery Fund (PCSRF). PCSRF was established by Congress in 2000 to protect, restore,
- and conserve Pacific salmon and steelhead populations and their habitats. AKSSF allocates its funds
- annually through competitive calls for pro-posals (CFPs) that open in the spring (usually mid-April).
- 3548 Eligible projects conserve habitat, restore habitat, monitor subsistence salmon populations,
- 3549 investigate the causes of Chinook declines, and conduct climate impact studies to identify resilient
- 3550 salmon habitat. Please see the AKSSF Objectives document for more information.

#### 3551 North Pacific Anadromous Fish Commission

3552 <u>https://www.npafc.org/</u>

The North Pacific Anadromous Fish Commission (NPAFC) is an international inter-governmental 3553 organization established by the Convention for the Conservation of Anadromous Stocks in the North 3554 Pacific Ocean. The Convention was signed on February 11, 1992, and took effect on February 16, 3555 1993. The member countries are Canada, Japan, the Republic of Korea, the Russian Federation, and 3556 the United States of America. As defined in the Convention, the primary objective of the NPAFC is 3557 to promote the conservation of anadromous stocks in the Convention Area. The Convention Area is 3558 the international waters of the North Pacific Ocean and its adjacent seas north of 33° North beyond 3559 the 200-mile zones (exclusive economic zones) of the coastal States. For the purposes of NPAFC, 3560 3561 anadromous fish include Pacific salmon and steelhead trout. Anadromous stocks are the stocks of these species that migrate into the Convention Area. 3562

#### 3563 Scientific Research

3564 The Convention authorizes fishing for anadromous fish in the Convention Area for scientific purposes under national and joint research programs approved by the NPAFC. The taking of 3565 3566 anadromous fish for scientific purposes must be consistent with the needs of the research program and provisions of the Convention and be reported to the Commission. Scientific research is 3567 conducted under the Science Plan, which may include investigations on species ecologically related 3568 to anadromous stocks. The member countries cooperate in collecting, reporting, and exchanging 3569 3570 biostatistical data, biological samples, fisheries data, and organizing scientific communications, such as seminars, workshops, exchanges of scientific personnel, and publications. The members provide 3571 3572 catch, enhancement, and other technical information and material pertaining to areas adjacent to the Convention Area from which anadromous stocks migrate into the Convention Area. 3573

#### 3574 Yukon River Drainage Fisheries Association

3575 <u>https://yukonsalmon.org/</u>

A wide array of issues affect Yukon River fisheries. Therefore YRDFA concentrates its work in a number of program areas to achieve its mission. These include: Policy Advocacy - A wide range of agencies and boards impact Yukon River management, from the State of Alaska Board of Fish to the federal North Pacific Management Council and the international Yukon River Salmon Agreement. YRDFA participates in dialogues with all these agencies, representing the interests of Yukon River communities in state, federal, and international forums. Conservation & Restoration - YRDFA works to protect wild salmon stocks and the habitats upon which they depend. Through biological

- research and participation in management, YRDFA works on behalf of Yukon River fishers.
- 3584 Cultural Preservation Traditional ecological knowledge is a vital source of information about
- 3585 salmon populations and a rich part of Yukon River cultures.

3586 YRDFA documents Local and Traditional Knowledge (LTK) about the salmon, the river, and the people. YRDFA works to incorporate LTK into Yukon fisheries management and works to protect 3587 the subsistence rights that are the foundation of Alaskan Native culture. Economic Opportunity -3588 Inhabitants of communities in the Yukon River drainage possess valuable knowledge and skills that 3589 can be indispensable to the success of local projects. YRDFA strengthens fishing communities by 3590 bringing jobs and training to communities, and increasing participation in fisheries management. 3591 3592 YRDFA also works to increase market opportunities and values for Yukon River salmon. Information Sharing - YRDFA is the only consistent forum for ongoing dialogue and information-3593 sharing between all parties with interests in the Yukon River fishery. YRDFA plays a key role in 3594 keeping fishermen and women informed and relaying their opinions to managers. YRDFA brings 3595 together fishers from throughout the watershed to reach consensus on policy positions that are good 3596

3597 for the salmon, the people, and the river.

#### 3598 Yukon River Inter-Tribal Fish Commission

3599 <u>https://www.tananachiefs.org/tribal-resources-stewardship-program/fish-commission/</u>

The Yukon River Inter-Tribal Fish Commission (YRITFC) with Tanana Chiefs Conference (TCC)
was founded in 2014 when Yukon River Tribes came together in St. Mary's and formed the Fish
Commission in response to low king salmon returns. YRITFC works with a variety of partners to
oversee 28 federally recognized villages.

#### 3604 Yukon Delta Fisheries Development Association

3605 <u>https://ydfda.org/</u>

Our mission is to create a self-sustaining, independent fishing company that will create income andemployment opportunities for Yukon Delta residents.

#### 3608 Kuskokwim River Inter-Tribal Fish Commission

- 3609 <u>https://www.kuskosalmon.org/</u>
- 3610 Thirty-three federally-recognized Tribes working together toward unified salmon co-management,
- 3611 research, and monitoring as we protect Kuskokwim salmon and traditional ways of life. Formed in
- 2015, the Kuskokwim River Inter-Tribal Fish Commission (KRITFC) works to develop a
- 3613 meaningful role for tribes and rural residents engaged in the management of Kuskokwim fisheries
- 3614 from the headwaters to the sea.
- 3615 Arctic Yukon Kuskokwim Sustainable Salmon Initiative
- 3616 <u>https://www.aykssi.org/</u>

- 3617 In response to salmon declines, Bering Sea Fishermen's Association and regional Native
- 3618 organizations (Association of Village Council Presidents, Kawerak, Inc., and Tanana Chiefs
- 3619 Conference) joined with state and federal agencies to create the AYK SSI, a proactive science-based
- 3620 program working cooperatively to identify and ad-dress the critical salmon research needs facing
- this region. The AYK SSI is the largest example of co-management of research-funding addressing
- 3622 salmon within the Pacific Rim and one of the largest, most successful programs of its kind in North3623 America. OUR MISSION is to collaboratively develop and implement a comprehensive research
- America. OUK MISSION is to conadoratively develop and implement a comprehens.
- 3624 plan to understand the causes of the declines and recoveries of AYK salmon.

#### 3625 Bristol Bay Regional Seafood Development Association

3626 <u>https://www.bbrsda.com/</u>

Our mission is to maximize the value of the Bristol Bay fishery for the benefit of our members. One
of the primary activities of the BBRSDA is funding (actually co-funding, in most cases) projects
designed to strengthen our fishery. Across a broad range of disciplines – economic research, qualityimprovement, fisheries science and marketing, among others – these programs are where most of our
members' annual 1% assessment goes. BBRSDA's participation in approved projects – as well as
the responsibilities of funded entities – are defined in contracts with the grantees (ADFG, research
institutions, fishermen, service providers, municipalities, etc.).

#### 3634 Prince William Sound Science Center

3635 <u>https://pwssc.org/</u>

In Prince William Sound, wild and hatchery-bred pink and chum salmon are important commercial 3636 3637 fisheries. Pink salmon is the largest of any commercial fishery and is serviced primarily by the purse seine fishery. Commercial, recreational and subsistence harvests of salmon profoundly affect the 3638 economic and cultural fabric of Prince William Sound communities. Coho, sockeye, Chinook, pink, 3639 and chum salmon support valuable fisheries in the region. The economic impact of these fisheries is 3640 critical to many small coastal communities here, and around the globe. Yet, the interactions between 3641 wild and hatchery fish are little understood. Our research focuses primarily on two species of Pacific 3642 3643 salmon found in Alaska, both of which evolved from their ancient rainbow trout ancestors. They start their lives as freshwater fish, then change and develop the ability to live and grow in the ocean 3644 3645 where they mature. As native fish evolve and interact with hatchery fish, there are inevitable impacts. We seek to understand those impacts in order to help maintain the unique identity and 3646 health of every species. 3647

- 3648 Alaska Sealife Center
- 3649 <u>https://www.alaskasealife.org/</u>

*Our Science Mission* The overall goal of our Science Program is to develop an under-standing of the
 role of marine mammals, birds and fish in the arctic and subarctic marine ecosystems, and to generate
 scientific knowledge relevant to resource management and policy. Our projects focus on Alaska
 marine life and environments, but reach globally with international collaborations. The Center's

- unique geo-graphic location, marine cold water research facilities, live animal collections, andspecialized staff allows us to use a combination of experimental and field research to:
- Investigate physiological and ecological processes affecting marine animal population
   dynamics.
- Conduct controlled experiments to understand factors affecting reproductive success and
   fitness in marine species.
- Monitor marine animal responses to environmental variability and stressors.
- Evaluate human impacts on our marine environment and animal populations.
- Develop tools to support recovery and restoration of marine resources.

#### 3663 University of Washington Alaska Salmon Program

#### 3664 <u>https://alaskasalmonprogram.org/</u>

Our program focuses on all aspects of the ecology and evolution of Pacific salmon in the watersheds of western Alaska, the Bering Sea, and the Gulf of Alaska. We actively pursue discovery science in an era of rapid global change to produce data and knowledge for managing and conserving regional ecosystems and their fisheries, and provides insights relevant to fisheries and watershed management across the globe. Our educational mission provides research opportunities for undergraduate and graduate students, and we seek to engage with regional K-12 programs, other citizens, and management agencies to improve our collective understanding of these remarkable ecosystems.

#### 3672 Salmon and People Project

#### 3673 <u>https://global.si.edu/projects/salmon-and-people-project</u>

**Project Highlights** The Kenai Lowlands region of the Kenai Peninsula in south central Alaska 3674 covers roughly 9400 square km. The watersheds of this region support abundant salmon that 3675 underpin sport fisheries and millions of dollars in economic activity related to commercial salmon 3676 fisheries. The food security and cultural identity of many residents all depend on abundant and 3677 reliable salmon populations. Over the past 15+ years, the Smithsonian Environmental Research 3678 Center's Dennis Whigham has collaborated with Coowe Walker of the Kachemak Bay National 3679 Estuarine Research Reserve (KBNERR), Ryan King of Baylor University (BU) and Mark Rains of 3680 the University of South Florida (USF). The collaboration has resulted in research that shows the 3681 3682 important ecological relationships between el-ements of the landscape like local plants and how many young salmon there are in the streams. The continued existence of resilient salmon 3683 populations, partic-ularly on lands like the Kenai Lowlands that lack state or federal conservation 3684 status, will require Alaskans to decide what invest-ments they need to make to ensure they can 3685 3686 sustainably support the goods and services provided by the landscape.

#### 3687 State of Alaska's Salmon and People

3688 <u>https://alaskasalmonandpeople.org/</u>

The mission of the State of Alaska's Salmon and People project is to create an equitable decision making platform for all stakeholders through information synthesis, collaboration, and stakeholder
 engagement.

#### 3692 Center for Salmon and Society

3693 <u>https://www.uaf.edu/cfos/research/center-for-salmon-society/index.php</u>

The Center for Salmon and Society seeks to engage all salmon-connected Alaskans in objective
forums to foster dialogue using the best available science to identify trade-offs inherent in complex
natural resource decisions.

- **3697 The Salmon Project**
- 3698 https://salmonproject.org/

3699 The Salmon Project is a celebration of wild salmon's place at the heart of Alaskan life, and the

- 3700 diverse ways it is present in our values, our culture, and our landscape.
- 3701 *About*

From 2012 to 2019, The Salmon Project shone a spotlight on the role salmon has in all Alaskans' lives, reinforcing our culture and identities, and showing how our individual choices affect this incredible resource. By telling the story of our shared Salmon Love, we laid a foundation for a statewide movement to ensure that Alaskans' lives will always be salmon lives. Our project moved out of its active stage at the end of 2019. Many of our projects and initiatives are evergreen.

3707 Our Vision

- 3708 To create a future where Alaskans are united in an:
- 3709 Awareness of the economic, environmental, social and cultural importance of salmon for ourselves,
- and for all Alaskans, including those whose connection to salmon is different than our own.
- **Understanding** of the challenges facing Alaska's wild salmon resource.
- 3712 **Commitment** to collective decisions and personal actions that will ensure future generations of
- 3713 Alaskans live with an abundance of wild salmon.

#### 3714 Alaska Fish Habitat Partnerships

- 3715 <u>https://www.alaskafishhabitat.org/</u>
- 3716 *Our Philosophy* Working together to protect, maintain, restore and enhance fish habitat throughout
- 3717 Alaska. *Our History* Alaska's first partnership to be formally recognized by the National Fish
- 3718 Habitat Partnership board was the Mat-Su Basin Salmon Habitat Partnership in 2006, followed by
- the Southwest Alaska Partnership in 2008, the Kenai Peninsula Partnership in 2010, and the

- 3720 Southeast Alaska Partnership in 2014. The Western Native Trout Initiative and Pacific Lamprey
- **3721** Partnership serve larger geographies that include Alaska.
- 3722 *Alaska's Partnerships* Rainbow Trout at Fish Creek, Matanuska-Susitna Valley, Alaska.
- 3723 USFWS/K.Mueller Operating under the banner of the National Fish Habitat Partner-ship,
- Alaska's recognized fish habitat partnerships are working on behalf of Alaska's wild, native fish and
- their habitats. These six recognized partnerships are part of a national network of locally-driven,
  voluntary, and non-regulatory collaboratives. Active partnerships made up of diverse interests are
- 3720 voluntary, and non-regulatory conaboratives. Active partnerships made up of diverse interests are
   3727 increasingly necessary to sustain Alaska's locally and globally important fisheries especially in
- geographic areas where habitat overlays a mosaic of private, state, tribal and federal lands and
- 3729 threats to fish habitat are at play.
- **Basin-Scale Events to Coastal Impacts (BECI)**

### 3731 <u>https://beci.info/</u>

BECI (Basin Scale Events to Coastal Impacts) is an ambitious project to develop an international

ocean intelligence system for the North Pacific Ocean that uses enhanced observations, numerical

modeling, and data analytics infrastructure to provide timely and targeted information on the impacts

of current and future climate events on ocean ecosystems. BECI was proposed by the North Pacific

- Anadromous Fish Commission (NPAFC) and the North Pacific Marine Science Organization
- (PICES) which was endorsed by the United Nations Decade of Ocean Science and SustainableDevelopment (UNDOS) in 2021.
- 2021 Development (010000) in 2021

## **3739** North Pacific Research Board

3740 <u>https://nprb.org/</u>

Supporting Research in the North Pacific Alaska supports some of the most diverse and abundant 3741 marine ecosystems in the world. Species large and small depend upon the state's unique marine 3742 environments for survival from coastal kelp beds to the undercarriage of multi-year Arctic sea ice. 3743 For coastal communities and those that utilize Alaska marine resources, this dependence extends 3744 beyond food or survival to traditional, cultural, and economic values that are shared by many—even 3745 to those outside the state. Alaska generates more than 40% of the commercial fish landings in the 3746 3747 United States. It is no wonder that with over 44,000 miles of coastline stretching from the frigid, exposed waters of the Beaufort Sea to the sheltered narrows of the Inside Passage, a shared 3748 responsibility and knowledge is required to support the long-term health of these ecosystems. 3749 Understanding Alaska's marine ecosystems is a collaborative effort. Using science and local 3750 traditional knowledge, scientists from all over the world are studying the oceanography, plants, and 3751 3752 animals of the North Pacific, Bering Sea, and the Arctic Ocean through funding support by the North 3753 Pacific Research Board (NPRB). NPRB's mission is to build a clear understanding of these ecosystems thereby enabling effective management and sustainable use of marine resources. 3754

## **3755 Department of Interior, Office of Subsistence Management**

3756 <u>https://www.doi.gov/subsistence/osm</u>

The Office of Subsistence Management is housed within the U.S. Fish and Wildlife Service, and provides administrative support to the Federal Subsistence Board and the Federal Subsistence Regional Advisory Councils. The staff of the Office of Subsistence Management includes fish and wildlife biologists, anthropologists, technical and administrative staff, and liaisons to the Alaska Department of Fish and Game and the Alaska Native community. The staff provides support for the regulatory process and the Fisheries Resource Monitoring Program.

#### 3763 Arctic Beaver Observations Network

3764 <u>https://sites.google.com/alaska.edu/a-bon/</u>

The Arctic Beaver Observation Network (A-BON) is a group of scientists, indigenous groups, land managers, and local observers who are concerned about the expansion of beaver populations into Arctic landscapes. This collaboration began in 2020 and assembles a broad range of perspectives from Alaska, Canada, Europe, and Russia to coordinate research and observations related to beaver

colonization of the Arctic and the impacts it is having on ecosystems and people.

3770

3771

3772

#### 3773 Appendix 7 – Alaska Salmon Research Task Force Members and Affiliations

#### 3774 Andrew Munro, North Pacific Fishery Management Council

3775 Dr. Andrew Munro is a fisheries scientist and statewide stock assessment scientist for the Alaska Department of Fish and Game. He has been a member of the North Pacific Fishery Management 3776 Council's Scientific and Statistical Committee (SSC) since 2019. He has expertise in fish biology, 3777 stock assessment, and salmon. Munro is also well-versed in the Council's history with Pacific 3778 salmon conservation efforts in the federal fisheries, and the management needs for understanding the 3779 Pacific salmon life cycle in Alaska. Munro also chairs a Working Group on Stock Assessment for 3780 3781 the North Pacific Anadromous Fish Commission. He also participates on technical panels for the Arctic-Yukon-Kuskokwim Sustainable Salmon Initiative. He has a PhD in Fish and Wildlife 3782 Biology from Montana State University, an MS in Biology, with an emphasis on Marine and 3783 Freshwater Ecology from East Stroudsburg University, and a BS in Biology, with a concentration in 3784 ecology and a minor in chemistry, from Ursinus College in Collegeville, Pennsylvania. 3785

#### 3786 Ed Farley (Chair), NOAA Fisheries, Alaska Fisheries Science Center

Dr. Farley has extensive experience and expertise in marine ecology of Pacific salmon. He also 3787 commercially fished in Bristol Bay from 1982 to 1997. Farley holds a PhD in Fisheries from the 3788 University of Alaska Fairbanks. His dissertation focused on early marine ecology of Bristol Bay 3789 sockeye salmon to better understand mechanisms in their marine life history that impact pro-duction. 3790 Farley has worked for NOAA Fisheries at the Alaska Fisheries Science Center since 1997. He is 3791 currently the program manager of the Ecosystem Monitoring and Assessment Program. The program 3792 is focused on understanding the impacts of climate change on ecosystems and fish fitness and 3793 survival. He developed and implemented the Alaska Fisheries Science Center's salmon early marine 3794 3795 ecology surveys in the eastern Bering Sea in 1999. The data from these surveys are used to forecast adult returns of Yukon River Chinook salmon and to understand how the rapidly warming Bering 3796 Sea is impacting early marine growth and survival of Bristol Bay sockeye salmon and western 3797 Alaska Chum and Chinook salmon. He is the chair of the Science Sub Committee of the North 3798 3799 Pacific Anadromous Fish Commission (NPAFC). This sub committee is charged with conservation of Pacific salmon in the North Pacific Ocean. Since 1997, Farley has developed and led NPAFC 3800 3801 efforts to understand production dynamics of Bering Sea salmon stocks (Asian and North American) through the Bering Aleutian Salmon International Survey (BASIS) Program and to understand 3802 winter ecology of Pacific salmon through the International Year of the Salmon Program. Farley was 3803 3804 the lead author on the ocean ecology of sockeye salmon chapter in the Ocean Ecology of Pacific 3805 Salmon and Trout. This document was published in 2018. It is where much of the NPAFC marine research has been summarized. 3806

#### **Bill Templin**, Alaska Department of Fish and Game

Bill Templin is currently the chief fishery scientist for salmon at the Alaska Department of Fish
and Game, Division of Commercial Fisheries. In this capacity, he is responsible for overseeing the
division's statewide salmon research and stock assessment programs and helping ensure that
research is well integrated with fisheries management. He has 29 years of experience conducting and
overseeing fisheries research on commercially important fish and shellfish species in Alaska. For

nine of these years, he served as the principal geneticist, in charge of the department's genetics

- 3814 program. During this time, he has represented the State of Alaska in various statewide, national and
- 3815 international settings including the Pacific Salmon Commission, North Pacific Anadromous Fish
- **3816** Commission, and the Intergovernmental Consultative Committee on Fisheries.

#### 3817 Andrew Piston, Pacific Salmon Commission

Andrew "Andy" Piston currently supervises the Southeast Alaska pink and chum salmon research 3818 programs, Ketchikan area sockeye salmon research programs, and other regional salmon stock 3819 assessment projects for the Alaska Department of Fish and Game. He is responsible for monitoring 3820 the escapement, production, survival and harvest patterns, and overall health of Southeastern 3821 Alaska's pink and chum salmon stocks. He is also responsible for developing recommendations and 3822 scientific advice for managers, the Alaska Board of Fisheries, Pacific Salmon Commission, and 3823 other organizations. He works cooperatively with NOAA Fisheries to implement the Southeast 3824 Coastal Monitoring Project and to produce joint pre-season pink salmon harvest forecasts. He has 3825 been involved with salmon research projects in Southeast Alaska since 1994. He has served on the 3826 Northern Boundary Panel of the Pacific Salmon Commission since 2016, when he was appointed by 3827 the Governor of Alaska. Previously, Piston served as the co-chair of the Northern Boundary 3828

- **3829** Technical Committee. He has been a technical committee member since 2010.
- **3830 Oscar Evon,** *Native Village of Kwigillingok*

Mr. Evon was born and raised in Kwigillingok, Alaska. He is a subsistence fisherman and member
of the Native Village of Kwigillingok. He is the Director of Regional Affairs for the Coastal Villages
Region Fund. He served as a board member from 2000-2009, eventually becoming Board of
Directors president and chair. He also acted as the fund's director of programs. Evon's previous
roles include Tribal administrator and COVID-19 coordinator for the Native Village of
Kwigillingok, office manager for Alaska Moravian Bible Seminary, and community outreach
coordinator for E3 Alaska. He serves on the North Pacific Fishery Management Council's Salmon

3838 Bycatch Committee.

#### 3839 Jacob Ivanoff (vice Chair), Native Village of Unalakleet

Mr. Ivanoff is a resident of Unalakleet, Alaska. He has personal knowledge of, and direct experience
with, subsistence harvest and uses of salmon in rural Alaska. He is highly educated, receiving
knowledge from Elders and others about salmon and through various positions he has held
professionally over the years. Ivanoff is affiliated with the Native Village of Unalakleet, with the
Tribe, and with other salmon-related entities in the Bering Strait region. Ivanoff is chair of the

- Alaska Department of Fish and Game's South Norton Sound Advisory Committee.
- **3846** Karla Jensen, *Native Village of Pedro Bay.*

Ms. Jensen is a Services Specialist 1 with the Pedro Bay Village Council. She is a Board member for the United Tribes of Bristol Bay. The Board of Directors consists of representatives from each of United Tribes of Bristol Bay's 15 member Tribal governments. United Tribes of Bristol Bay is a Tribal consortium working to protect the traditional Yup'ik, Dena'ina, and Alutiig ways of life in Southwest

Alaska that depend on the pristine Bristol Bay watershed and all it sustains, most notably Bristol Bay's wild salmon. As a political division of our member Tribal governments, their work is primarily focused in three areas: Tribal consultation with government agencies on issues affecting the Alaska Native way of life; grassroots organizing in the local, statewide, and national arena; and youth empowerment and organizing in the Bristol Bay region.

#### **3856 Caroline Brown,** *Alaska Department of Fish and Game*

Ms. Brown is the statewide Subsistence Research Director. She is responsible for coordinating all 3857 ethnographic research and policy recommendations on subsistence practices for the Subsistence 3858 Section of Alaska Department of Fish and Game. Prior to this role, Brown was the Northern Regional 3859 Program Manager from 2017-2020 and the lead subsistence resource specialist for interior Alaska 3860 from 2002-2017. Brown holds an MA degree in anthropology from the University of Chicago where 3861 she was also a PhD candidate. Over the last 20 years, Brown has conducted multiple projects involving 3862 the documentation and analysis of local, traditional and Indigenous knowledge throughout Alaska. 3863 She also serves as the alternate U.S. co-chair on the U.S./Canada Yukon River Panel. 3864

#### **3865 Justin Leon**, *Kuskokwim River Inter-Tribal Fish Commission*

Mr. Leon serves as the Fisheries Biologist for the Kuskokwim River Inter-Tribal Fish Commission. 3866 Before this, he served as the Senior Tribal Climate Resilience Liaison for the Alaska Region for the 3867 Native American Fish and Wildlife Society. Through his roles as the Alaska Tribal Liaison with 3868 Native American Fish and Wildlife Society and Fisheries Biologist with the Kuskokwim River Inter-3869 Tribal Fish Commission, Leon has garnered experience working with Alaska Native Tribes and 3870 Tribal citizens, and bridging local and Indigenous Knowledge with western science. Leon has a BS 3871 from the University of Georgia in wildlife management. He moved to Alaska after graduating in 3872 2008 and has made Alaska home since then. He obtained his MS in fisheries from the University of 3873 Alaska Fairbanks, where he focused on Chinook salmon in the Yukon and Kuskokwim rivers. After 3874 graduate school, he spent 10 years as a fishery biologist for the Alaska Department of Fish and 3875 Game. As a fishery biologist, he worked with crab and salmon research, and wild fisheries stock 3876 3877 management in Northwest Alaska, the Alaskan interior, and the Aleutian Islands.

#### 3878 Michelle Stratton, Alaska Marine Conservation Council/Fisherman

3879 Ms. Stratton works for the Alaska Marine Conservation Council. She was born and raised in Palmer, Alaska, and grew up set netting for salmon with her family on the west side of Cook Inlet. She began 3880 her career as a technician for the Alaska Department of Fish and Game before earning her MS degree 3881 in Fisheries Science from the University of Alaska Fairbanks. At the same time, she was working eight 3882 years as an ADF&G fisheries biologist. In her position at Alaska Marine Conservation Council, 3883 3884 Stratton devotes much of her time toward fisheries research and education, helping build connections between Alaska's fishing communities and the scientific processes that support them. As a lifelong 3885 subsistence hunter and fisherman, Stratton has a passion for fisheries biology and its role in sustaining 3886 the thriving food systems and wild places that she has lived within most of her life. She also is an 3887 owner-operator of a set net site on the south end of Kodiak Island. 3888

#### **3889 Mike Flores,** *Charter Boat Fisherman*

3890 Mr. Flores has over 30 years of experience owning and operating a large charter fishing business on
3891 the Kenai Peninsula. He has recently completed service on the Alaska Bycatch Task Force. Mr. Flores
3892 is serving on the Charter Halibut Management Committee of the North Pacific Fisheries Management
3893 Council. He is familiar with the processes and procedures of Alaska State boards and committees.

## **3894 Austin Estabrooks,** *At-sea Processors Association*

3895 Mr. Estabrooks is a natural resource analyst for the At-sea Processors Association (APA). He has worked on various aspects of salmon bycatch mitigation undertaken by the pollock catcher processor 3896 (CP) fleet operating in the Bering Sea. He is the primary author of the Incentive Plan Agreement 3897 3898 (IPA) under which the CP fleet operates to meet the objectives of Amendment 91 and 110. He is responsible for monitoring in-season by catch of both Chinook and chum to help identify hot-spots 3899 for avoidance as well as working closely with Auke Bay geneticists to identify longer term spatio-3900 temporal trends of chum salmon stock distributions as reflected in the bycatch. He has conducted at-3901 3902 sea experiments with salmon lights and salmon excluder devices using deploy and retrieve cameras to quantify escapement. Through the Pollock Conservation Cooperative Research Center, Mr. 3903 3904 Estabrooks also advises APA on funding extensive research on Alaska salmon. This includes recent projects developing species distribution models for Chinook salmon and investigating Yukon chum 3905 salmon early life history. Prior to APA, he spent nearly five years in the Bering Sea and Gulf of 3906 3907 Alaska working as a North Pacific Groundfish observer, where he collected systematic genetic samples of salmon bycatch in the pollock fishery. 3908

#### **3909 Tom Carpenter**, *Commercial Fisherman*

Mr. Tom Carpenter hails from Cordova, Alaska. In 2022, he was appointed by Alaska Governor Mike
Dunleavy to serve on the Alaska Board of Fisheries. He is retired from the U.S. Coast Guard and has
participated in various fisheries throughout his career. He has served as a crewman on a seiner and a
gillnetter before buying his own boat. He also purchased and operated a sporting goods store in
Cordova. Carpenter has served for over 22 years on the Copper River/ Prince William Sound Advisory
Committee.

#### **3916 Steve Reifenstuhl** *retired*, *Northern Southeast Regional Aquaculture Association*

Mr. Reifenstuhl has over 45 years of experience in Alaska salmon fisheries management, research
(salmon biology and ecology, post-secondary education), salmon habitat restoration (cooperative
projects with U.S. Fish and Wildlife Service) and salmon hatchery production. Among his various
roles and accomplishments, he was a founding board member of Alaska's Salmon Hatchery/ Wild
research program, served on the S.E. Regional Advisory Council, North Pacific Research Board
Advisory Panel and was a board member of United Fishermen of Alaska. He recently retired as general
manager of the Northern Southeast Regional Aquaculture Association.

#### **3924** Megan McPhee, University of Fairbanks, Alaska

3925 Dr. Megan McPhee is an associate professor at the College of Fisheries and Oceans, University of
3926 Alaska, Fairbanks, located at the Juneau Fisheries Center. She is a fisheries ecologist who focuses on
3927 the ecology, evolutionary biology, and population structure of Pacific salmon. Relevant research

- topics include marine ecology of chum salmon, connections between climate, growth rate, and
- age/size at maturity in western Alaska Chinook salmon and Southeast Alaska steelhead; effects of
- 3930 competition on growth of western Alaska chum salmon in the North Pacific, genetic stock
- identification of western Alaska chum salmon, and hatchery-wild interactions in Southeast Alaska.
- 3932 She sits on the steering committees of the Southeast Alaska Fish Habitat Partnership and the
- International Year of the Salmon. She is also an associated faculty with the Tamamta program atUAF, which seeks to elevate the role of Indigenous knowledge in fisheries education and research.
- **3935** Megan Williams, Arctic Program, Ocean Conservancy/ University of Alaska Fairbanks

Dr. Megan Williams is a fisheries scientist with Ocean Conservancy. Her education and professional 3936 experiences have focused on fisheries interactions with apex predators and predator ecology in Alaska. 3937 She has worked extensively on bycatch issues and climate readiness in fisheries management at both 3938 state and federal levels since 2010. Her current work focuses on integrating western science and 3939 3940 Traditional/Indigenous Knowledge to understand cumulative threats to salmon and rural communities and to identify climate resilient solutions. She also serves as the chair of the Alaska Scientific Review 3941 Group that advises NOAA Fisheries and the U.S. Fish and Wildlife Service on the best available 3942 science and subsistence information to be included in annual Marine Mammal Stock Assessment 3943 Reports in Alaska. 3944

**3945 Tommy Sheridan,** *University of Alaska Fairbanks* 

Mr. Sheridan is the associate director for the University of Alaska Fairbanks (UAF) Alaska Blue 3946 Economy Center. He is also currently serving as community site coordinator for the Alaska Regional 3947 Collaboration for Innovation and Commercialization (ARCTIC) Program through UAF's Alaska 3948 Center for Energy and Power to establish Cordova, Alaska as a Community Innovation Hub. He spent 3949 3950 eight years working for Northern Southeast Regional Aquaculture Association in the salmon hatchery industry, six years working for Alaska Department of Fish and Game as a commercial salmon fisheries 3951 manager, and three years working for Silver Bay Seafoods in the seafood processing sector. He has 3952 graduate level education in fisheries and fisheries management, and has taught undergraduate fisheries 3953 3954 courses in both Alaska and Oregon. He has served as a board member for Prince William Sound Aquaculture Corporation, Alaska Fisheries Development Foundation, Prince William Sound Science 3955 3956 Center, and Sitka Sound Science Center. He was appointed to the Governor's Alaska Bycatch Review Task Force (ABRT) in 2021, and currently serves on the North Pacific Anadromous Fish Commission 3957 as one of two US Commissioners. 3958

#### **3959** Noëlle Yochum, Alaska Pacific University/ Trident Seafoods

Dr. Noëlle Yochum is affiliated faculty with Alaska Pacific University and is the Senior Manager of
Fishing Innovation and Sustainability for Trident Seafoods. Prior to this, Dr. Yochum led the
Conservation Engineering group at the Alaska Fisheries Science Center (NOAA Fisheries). In both
capacities, Dr. Yochum's focus is on collaborative research with industry and scientific partners to
find innovative ways to evaluate and mitigate incidental impacts of fishing, including bycatch,
bycatch mortality, and effects on fish habitat. This work is done through field and laboratory
research to understand fish behavior and to improve fishing gear design and practices. In addition to

work in Alaska and the U.S. west coast, she has conducted related work on the U.S. east coast andabroad.

#### **3969** Katie Howard, Alaska Department of Fish and Game

3970 Dr. Kathrine "Katie" Howard is a statewide fishery scientist with the Alaska Department of Fish and3971 Game and serves as the Salmon Ocean Ecology Lead. She holds a PhD and an MS degree in zoology

3971 Game and serves as the Samon Ocean Ecology Lead. She holds a FilD and an WS degree in zoology 3972 from the University of Hawaii and a BS in biology and a BA in English from the University of

3973 Idaho. Katie began at ADF&G in 2009 as the Yukon Area Research biologist and was quickly

3974 promoted to the Arctic-Yukon-Kuskokwim regional research biologist before becoming the AYK

3975 fisheries scientist and eventually holding a statewide scientist position overseeing the Salmon Ocean

3976 Ecology Program. Katie has been involved with multiple initiatives and projects that include

3977 collaborative marine surveys to assess juvenile Chinook and chum salmon stocks, North Pacific

3978 Anadromous Fish Commission, and International Year of the Salmon.

3979

# Appendix 8. Names and primary affiliations of the Arctic-Yukon-Kuskokwim Working Group of the AKSRTF.

Name	Primary Affiliation
AKSRTF Members	
Andrew Munro	Alaska Department of Fish and Game
Austin Estabrooks	At-Sea Processors Association
Bill Templin	Alaska Department of Fish and Game
Caroline Brown	Alaska Department of Fish and Game
Ed Farley	NOAA Fisheries, Alaska Fisheries Science Center
Jacob Ivanoff	Native Village of Unalakleet
Justin Leon	Kuskokwim River Inter-Tribal Fish Commission
Katie Howard	
(co-chair of WG)	Alaska Department of Fish and Game
Megan McPhee	University of Alaska Fairbanks
Megan Williams	Ocean Conservancy
Michelle Stratton	Alaska Marine Conservation Council/Fisherman
Noelle Yochum	Trident Seafoods
Oscar Evon	Coastal Villages Region Fund (CVRF)
Steve Reifenstuhl	Salmon Biologist/Consultant (Retired)
Tom Carpenter	Commercial Fisherman
Public Members	
Adolph Lupie	Lower Kuskokwim River

Andy Bassich	Upper Yukon River
Bill Alstrom	Lower Yukon River
Brooke Woods	Yukon River
Charlie Lean	Norton Sound
Charlie Wright	
Courtney Weiss	Yukon Delta Fisheries Development Association (YDFDA)
Curry Cunningham	University of Alaska – Fairbanks
Dan Gillikin	Native Village of Napaimute, Aniak, AK
Daniel Schindler	
(co-chair of WG)	University of Washington
Hannah Heimbuch	Commercial Fisher
James Nicori	Kuskokwim region
Jennifer Hooper	Association of Village Council Presidents, Yukon-Kuskokwim Delta Region
Jessica Black	Associate vice chancellor for rural, community education and Native education. University of Alaska - Fairbanks
Joe Spaeder	Arctic-Yukon-Kuskokwim Sustainable Salmon Initiative
Mark McNeley	Native Village of Nelson Lagoon
Martin Andrew	Kuskokwim region
Marvin Okitkun	Kotlik, lower Yukon Delta
Michelle Quillin	Tanana Chiefs Conference, Fairbanks
Patrick Barry	NOAA Alaska Fisheries Science Center

Ragnar Alstrom	Executive Director, Yukon Delta Fisheries Development Association (YDFDA)
Renae Ivanoff	NSEDC's Fisheries Research and Development Director
Scott Gende	National Park Service
Serena Fitka	Yukon Delta Fisheries Development Association (YDFDA)
Todd Sformo	North Slope Borough - Department of Wildlife Management
Vanessa von Biela	U.S. Geological Survey, Alaska Science Center
Virgil Umphenour	Upper Yukon River

3982

3983 Appendix 9. List of all hypotheses and research questions considered by the AYK WG for understanding the causes of recent declines in

3984 AYK chum salmon and Chinook salmon. Individual hypotheses and questions were developed as components of each of the major

3985 research themes (possible explanations) specified by the AKSRTF. Individual scores are the total number of points allocated to each

3986 hypothesis or question by the AYK WG, which were summed within each theme (total score = TS) to develop weights for prioritizing

3987 research across the AKSRTF research themes.

Research Theme: Possible explanation for AYK salmon decline (TS = total score)	Individual score	Specific question or hypothesis
Spawner Health TS = 102	28	Quantify causes, magnitude, and consequences of <i>en route</i> and pre-spawn mortality. What are interactions with climate and changing ocean conditions? Roles of disease and parasites
	23	Improved understanding of spawner quality (age, sex ratios, size, genetic divesity, nutritional and health condition of spawners) and how changes to spawner quality impact reproductive success and stock productivity
	9	Identifying vectors of salmon disease, and conditions leading to changes in disease prevalence and intensity (e.g., environmental conditions and/or changes to migratory patterns)
	13	Interactive effects of multiple stressors on spawning adults during their freshwater migration (e.g., climate conditions)
	3	Sublethal impacts, such as straying or other behavior changes, during poor migration conditions (i.e. warm temperatures, low water) on migrating fish
	19	Identifying conditions/thresholds where genetic population size is so critically low that stocks are at particular risk for extirpation, and assessing whether any existing stocks meet those criteria

	6	Programmatic review to determine whether inriver stock assessments have adequate precision and accuracy for estimating abundance
Freshwater Harvest (Commercial and Sport) TS = 35	9	What are legacy effects of historical FW fisheries (e.g., gear effects, harvest rates) on current population status and demographics?
	6	What is role of unobserved fishery-mortality (e.g., drop-outs from small mesh nets) on mortality rates, fishery selection, and ultimately population productivity?
	4	What is the extent that FW harvest can further depress weak stock components while targeting productive stocks (related to management under uncertainty). Can we improve stock resolution and harvest specificity?
	12	Are the current BEG's/SEG's spawner recruit relationships used to estimate surplus fish available for harvest still valid in light of new theories about where mortality may be occurring i.e. fresh vs marine waters and does it matter?
	4	Ecological consequences of harvesting to achieve the upper limits of escapement goal range, compared to lower limits of escapement goal range?
	0	Is there any down side (related to future recruitment) to focusing harvest on "Jack" Chinook salmon?
Freshwater	9	Are changes in freshwater predators reducing FW survival of juvenile salmon?
Predators TS = 19	1	Do changing environmental conditions (temp, flow, turbidity) have an influence on predation success and/or the abundance and size of predators?
	3	How do beavers indirectly affect predation on juvenile salmon by altering habitat?
	0	Are changes in harvest of FW predators changing predation on juvenile salmon?
	6	Do climate-induced changes in coho salmon alter predation regimes on other salmon in FW
Marine Predators	12	What is the role of changes in marine apex predators (orcas, sharks, sea lions) on demographic structure (eg. body sizes and ages) and abundance of AYK salmon? What are abundance trends, diets and spatial distribution of predator populations?

TS = 24	5	What are impacts of marine predators on pre-adult and smolt life stages?
15 - 24		
	7	Are there interactions between changing climate and marine predators that affect ecology of AYK salmon?
Freshwater conditions for eggs and juvenile rearing and migration TS = 51	14	Are changes in watershed habitat productivity and capacity reducing fitness and abundance of smolts leaving watersheds?
	15	Is the hydrology (discharge, magnitude, duration, bank full flows) changing and what is the effect? Loss of complexity, disconnected habitats, channel morphology. Effects on FW food webs and growth and survival of juvenile salmon. Effects on egg incubation?
	6	Have changes in marine-derived nutrients (MDN) from declining salmon populations, changes in fish and wildlife management, reduced the productivity of freshwater nursery habitats?
	5	Is the spread of beavers into areas that had not previously had beavers changing the quality and quantity of habitat for salmon?
	0	How do fires, permafrost degradation, and localized logging in riparian zone for firewood influence habitat value?
	11	Climate effects on floods, spring breakup, thermal refuges, flows, temperatures, permafrost ?
Marine food	47	Are changing marine food webs and climate reducing quantity and quality of food for smolts?
limitation TS = 131	65	Is competition with hatchery pinks and chums, Bristol Bay sockeye, reducing marine survival of AYK salmon?
	19	Are changes in nearshore habitat conditions affecting smolt survival of AYK salmon
Climate change (freshwater and marine) TS = 62	35	Interactions between climate change and all other factors
	0	What are the impacts of increasing presence of salmon in the high Arctic on resident non-salmon species?
	8	Are large scale climate variation linked between freshwater and marine ecosystems in the AYK region? If so how do we separate them, do we need to?

	16	Climate-driven changes in sea ice and impacts for plankton phenology (match/mismatch)
Marine Harvest	42	What are 1) catch rates and 2) stock composition of fish in state, federal and foreign fisheries? How do these compared to escapement estimates in AYK watersheds. Need more effective monitoring of all 3 components
TS = 156	49	Is bycatch in US federal fisheries causing declines of AYK salmon?
	38	Are interceptions in other Alaska state fisheries causing declines of AYK salmon?
	15	Is foreign IUU (illegal, unregulated, unreported) fisheries causing declines of AYK salmon?
	21	Are genetic markers adequate for understanding impacts of bycatch and interceptions on population dynamics?