

Revised Table 3-19 Chum salmon bycatch rates (chum salmon per mt of pollock) by genetic cluster area in the early and late period B season pollock fishery, 2011 through 2022

	<i>Cluster area 1</i>		<i>Cluster area 2</i>		<i>Cluster area 3</i>		<i>Cluster area 4</i>	
	Early	Late	Early	Late	Early	Late	Early	Late
2011	0.53	0.46	0.18	0.34	0.16	0.24	0.13	0.07
2012	0.04	0.23	0.02	0.11	0.01	0.09	0.001	0.004
2013	0.42	0.64	0.14	0.35	0.02	1.09	0.01	0.03
2014	0.23	0.46	0.56	1.02	0.33	0.40	0.04	0.39
2015	0.15	0.88	0.08	0.69	0.16	1.00	0.04	0.12
2016	0.31	0.50	0.40	0.67	0.66	0.40	0.05	0.81
2017	0.59	0.13	1.21	0.48	0.45	0.24	0.30	0.41
2018	0.45	0.16	0.97	1.72	0.96	0.64	0.01	0.05
2019	0.46	0.43	0.45	0.63	0.84	0.19	0.02	0.87
2020	0.13	0.64	0.73	1.86	0.06	1.30	0.05	0.38
2021	1.02	0.07	8.98	0.25	0.13	0.61	0.01	0.06
2022	0.34	1.12	0.11	1.96	0.04	0.77	0.02	0.92
Average	0.39	0.48	1.15	0.84	0.32	0.58	0.06	0.34

Source: NMFS Alaska Region Catch Accounting System; ChumRates_YrTempGrpcorrected