

**Special Publication No. 17-07**

---

---

**Protocol for Estimating Age of Weathervane Scallops  
*Patinopecten caurinus* in Alaska**

by

**Chris Siddon**

**Quinn Smith**

**Kevin McNeel**

**Dion Oxman**

and

**Kenneth Goldman**

---

---

March 2017

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



## Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the *Système International d'Unités* (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

<b>Weights and measures (metric)</b>		<b>General</b>		<b>Mathematics, statistics</b>	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H <sub>A</sub>
gram	g			base of natural logarithm	<i>e</i>
hectare	ha			catch per unit effort	CPUE
kilogram	kg	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	coefficient of variation	CV
kilometer	km			common test statistics	(F, t, $\chi^2$ , etc.)
liter	L	at	@	confidence interval	CI
meter	m	compass directions:		correlation coefficient (multiple)	R
milliliter	mL	east	E	correlation coefficient (simple)	r
millimeter	mm	north	N	covariance	cov
		south	S	degree (angular)	°
<b>Weights and measures (English)</b>		west	W	degrees of freedom	df
cubic feet per second	ft <sup>3</sup> /s	copyright	©	expected value	<i>E</i>
foot	ft	corporate suffixes:		greater than	>
gallon	gal	Company	Co.	greater than or equal to	≥
inch	in	Corporation	Corp.	harvest per unit effort	HPUE
mile	mi	Incorporated	Inc.	less than	<
nautical mile	nmi	Limited	Ltd.	less than or equal to	≤
ounce	oz	District of Columbia	D.C.	logarithm (natural)	ln
pound	lb	et alii (and others)	et al.	logarithm (base 10)	log
quart	qt	et cetera (and so forth)	etc.	logarithm (specify base)	log <sub>2</sub> , etc.
yard	yd	exempli gratia (for example)	e.g.	minute (angular)	'
		Federal Information Code	FIC	not significant	NS
<b>Time and temperature</b>		id est (that is)	i.e.	null hypothesis	H <sub>0</sub>
day	d	latitude or longitude	lat or long	percent	%
degrees Celsius	°C	monetary symbols (U.S.)	\$, ¢	probability	P
degrees Fahrenheit	°F	months (tables and figures): first three letters	Jan, ..., Dec	probability of a type I error (rejection of the null hypothesis when true)	$\alpha$
degrees kelvin	K	registered trademark	®	probability of a type II error (acceptance of the null hypothesis when false)	$\beta$
hour	h	trademark	™	second (angular)	"
minute	min	United States (adjective)	U.S.	standard deviation	SD
second	s	United States of America (noun)	USA	standard error	SE
		U.S.C.	United States Code	variance	
<b>Physics and chemistry</b>		U.S. state	use two-letter abbreviations (e.g., AK, WA)	population	Var
all atomic symbols				sample	var
alternating current	AC				
ampere	A				
calorie	cal				
direct current	DC				
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

***SPECIAL PUBLICATION NO. 17-07***

**PROTOCOL FOR ESTIMATING AGE OF WEATHERVANE SCALLOPS  
*PATINOPECTEN CAURINUS* IN ALASKA**

by

Chris Siddon, Quinn Smith, Kevin McNeel, Dion Oxman, and Kenneth Goldman  
Alaska Department of Fish and Game, Division of Commercial Fisheries

Alaska Department of Fish and Game  
Division of Sport Fish, Research and Technical Services  
333 Raspberry Road, Anchorage, Alaska, 99518-1565

March 2017

The Special Publication series was established by the Division of Sport Fish in 1991 for the publication of techniques and procedures manuals, informational pamphlets, special subject reports to decision-making bodies, symposia and workshop proceedings, application software documentation, in-house lectures, and became a joint divisional series in 2004 with the Division of Commercial Fisheries. Special Publications are intended for fishery and other technical professionals. Special Publications are available through the Alaska State Library, Alaska Resources Library and Information Services (ARLIS) and on the Internet: <http://www.adfg.alaska.gov/sf/publications/>. This publication has undergone editorial and peer review.

*Chris Siddon*

*Alaska Department of Fish and Game, Division of Commercial Fisheries,  
1255 W. 8<sup>th</sup> Street, Juneau, AK 99801*

*Quinn Smith*

*Alaska Department of Fish and Game, Division of Commercial Fisheries,  
803 3<sup>rd</sup> Street, Douglas, AK 99824*

*Kevin McNeel*

*Alaska Department of Fish and Game, Division of Commercial Fisheries,  
PO Box 115526, Juneau, AK 99811*

*Dion Oxman*

*Alaska Department of Fish and Game, Division of Commercial Fisheries,  
PO Box 115526, Juneau, AK 99811*

*Kenneth Goldman*

*Alaska Department of Fish and Game, Division of Commercial Fisheries,  
3298 Douglas Place, Homer, AK 99603*

*This document should be cited as follows:*

*Siddon, C., Q. Smith, K. McNeel, D. Oxman, and K. Goldman. 2017. Protocol for estimating age of weathervane scallops *Patinopecten caurinus* in Alaska. Alaska Department of Fish and Game, Special Publication No. 17-07, Anchorage.*

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

**If you believe you have been discriminated against in any program, activity, or facility please write:**

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

**The department's ADA Coordinator can be reached via phone at the following numbers:**

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

**For information on alternative formats and questions on this publication, please contact:**

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2375

# TABLE OF CONTENTS

	<b>Page</b>
LIST OF FIGURES .....	iii
LIST OF APPENDICES .....	iii
ABSTRACT .....	1
INTRODUCTION .....	1
Age Estimation Methods .....	1
Error and Quality Control .....	3
METHODS .....	5
Shell Collection .....	5
Age Estimation .....	6
Quality Control .....	9
Reference Collection .....	9
Training new readers .....	9
Examining Bias and Precision .....	10
Addressing Bias and Poor Precision .....	11
Data Management .....	12
REFERENCES CITED .....	13
APPENDIX A: DIFFERENCES IN SHELL APPEARANCE ACROSS THE STATE OF ALASKA .....	15
APPENDIX B: EXAMPLES OF DAMAGED SHELLS .....	17
APPENDIX C: FIRST ANNULUS MEASUREMENTS .....	19
APPENDIX D: TABLES OF SYMMETRY EXAMPLES .....	23
APPENDIX E: DATA COLLECTION FIELDS .....	25

## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
1. A: Weathervane scallop aged by the color band pair and circuli methods at 9 years. B: Three axes from which age assessment counts are conducted. ....	2
2. Magnified image of scallop shell circuli showing area of compression and annulus. ....	3
3. Two 9-year-old scallops from Management Area D (Yakutat) in 2006 emphasizing the differences in growth within an area through different shell heights and to document observer codes.....	4
4. A: Scallop aged at 10 years using the circuli method. B: Scallop aged at 9 years using the circuli method, showing 2 false checks in the shell which can also be seen using the color band method. ....	6
5. Twelve-year-old albino scallop shell from commercial fishery Management Area D in 2006.....	8
6. Example of annuli on scallop shell auricle.....	8

## LIST OF APPENDICES

<b>Appendix</b>	<b>Page</b>
A1. Example images of scallops from 5 management areas across the state of Alaska showing differences in shell appearance. Map of areas is provided in Appendix C2. ....	16
B1. Examples of shell damage illustrating which shells can be aged and which cannot. ....	18
C1. First annulus measurements (mm) along the primary aging axis from scallops collected by ADF&G fishery observer program and research surveys.....	20
C2. Management Areas of collection. ....	21
D1. Table A shows no bias in results; ages are randomly distributed across the main diagonal of the table.....	24
D2. Table B shows asymmetry across the main diagonal after age 14, indicating specimens in the circled area need to be re-aged. ....	24
E1. Data fields to be entered by age readers. ....	26

## ABSTRACT

Age determination is an essential component to many fisheries stock assessments that provide the foundation for sound fisheries management. Although stock assessment models that include age data are generally an improvement over length-based models, errors in age data can have serious repercussions on fisheries management. Thus, a standardized methodology and formal quality control measures are essential. Here we present a standardized protocol to assess the age of weathervane scallops *Patinopecten caurinus* collected by the Alaska Department of Fish and Game's statewide scallop research and observer programs. These methods and analyses also provide measures and thresholds to test for precision (percent agreement and mean CV) and bias (Bowker's and Evans-Hoenig tests of symmetry) in age estimates, as well as quality control mechanisms for a long-term scallop age assessment program.

Key words: weathervane scallops, *Patinopecten caurinus*, Alaska, age assessment, precision, bias

## INTRODUCTION

Determining the age of individual organisms provides the foundation for many quantitative fisheries stock assessments; age determination models are more informative and precise than non-age-based models. However, ages can be difficult to estimate due to differences in biology (life spans, growth rates, and environments) and estimation procedure (structure processing and band identification; Campana 2001). Potential error in estimated ages can cause bias in stock assessments and potentially lead to mismanagement of fisheries (Heifetz et al. 1998; Campana 2001). Such bias can deleteriously affect estimates of growth rates, stock recruitment, year class strength (cohort abundance), fishing mortality, and ultimately yield (Kimura 1990; Lai and Gunderson 1987; Heifetz et al. 1998). The methods and error of ages estimated for weathervane scallops *Patinopecten caurinus* caught across Alaska has yet to be assessed. Assuming currently used sampling designs for weathervane scallops are appropriate, this report will set a procedure for estimating and assessing ages. The common methods to minimize and assess age estimation error that will be used in this report include (1) having a standard protocol, (2) using separate individuals (age readers) to estimate ages to evaluate individual error, (3) using a standard set of specimens (reference collection) to train or evaluate individuals, (4) formally training new age readers, and (5) establishing repeatable tests for precision and bias.

## AGE ESTIMATION METHODS

Common protocols used to estimate the age of bivalves such as geoduck clams, Arctic quahogs, and scallops include sectioning shells and producing acetate peels (Shaul and Goodwin 1982; Ropes 1984; Fiori and Morsán 2004; Lomovasky et al. 2008) as well as visually estimating age by counting rings visible on the surface of the shell (MacDonald and Bourne 1987). Sectioning shells involves the use of a precision diamond blade saw to remove sections of the shell, mounting sections onto slides, and sanding and polishing the resulting slide-mounted sections. Producing acetate peels involves cutting shells using tile or diamond blade saws, etching the resulting surface with acid (e.g., hydrochloric acid), and using acetone while pressing the acetate into the etched surface to transfer the annuli (Ropes 1984). These methods are typically time intensive and require potentially expensive laboratory equipment, but are necessary for species in which annual banding is not discernable on the external surface of the shell. For species where annual bands or circuli are visible from the surface, such as the weathervane scallops, developing estimates directly from the surface are more efficient and cost effective.

Estimating the age of scallops using counts of annuli visible on the surface of shell is well established. While the earliest studies were completed on Atlantic scallops such as *Placopecten*

*magellanicus* and *Pecten maximus* (Stevenson and Dickie 1954), age estimates of weathervane scallops in Alaska waters have been conducted since the late 1960s (Hennick 1970). The enumeration of annuli (dark bands) is the most efficient and cost effective method, and can be aided using slightly more time consuming methods (e.g., identifying compressed circuli under magnification) or labor and equipment intensive processing (sectioning or preparing acetate peels). Further, several studies have supported the hypothesis that these annuli are formed once a year (Merrill et al. 1966; MacDonald and Thompson 1985; MacDonald and Bourne 1987; Smith et al. 2001; Hart and Chute 2009). However, there is evidence that these bands may not form annually in all species (Chute et al. 2012), and further validation of the annual nature is needed.

Annuli on the surface of weathervane scallops are described as bands or “rings” of different color or texture observed under reflected light (Figure 1A; Ropes and Jearld 1987, Gustafson and Goldman 2012, Chute et al. 2012, Spafard and Rosenkranz 2014). This sequential light/dark banding pattern, visible on weathervane scallop shells using reflected light from the umbo (Figure 1B) to the edge, is in response to seasonal growth trends where the dark band (theoretically representing a slow growth cycle) is considered the annulus. Small textured ridges within these bands (circuli) are also distinguishable and can be used to identify individual annuli (Figure 2; Spafard and Rosenkranz 2014). Annuli can also be counted on the auricle (Figure 1B) and have been used to assist in determining age when the dark band and circuli methods proved ineffective on shells, like those that are heavily worn (R. Burt, ADF&G, pers. comm.). The annulus determined by the dark band, circuli, or auricle method is ultimately what is counted to estimate age, and the age assigned to individual annuli are estimated from the umbo (earliest growth) out to the outer edge (Figure 1; CARE 2006; Gustafson and Goldman 2012).




Figure 1.–A: Weathervane scallop aged by the color band pair and circuli methods at 9 years. Arrows point to annuli. B: Three axes from which age assessment counts are conducted. All axes start at the umbo and go to the shell edge.






Figure 2.—Magnified image of scallop shell circuli showing area of compression and annulus.

## **ERROR AND QUALITY CONTROL**

There are 2 main sources of error estimating ages: (1) data and specimen mishandling, and (2) incorrect identification of annuli. The former can be mitigated through formal procedures and careful data collection, and can be detected through similar quality control procedures as incorrect age estimates. The latter can be mitigated through formal annuli criteria, training, and continual reader assessment which will be discussed below. Errors in age estimates can be detected through evaluation of a reader's ability to detect annuli, analysis of repeat estimates between readers, and using a reference collection.

Formal annuli criteria consists of objective measurement ranges for specific annuli and definitions of features that are commonly mistaken as annuli. These features are referred to as false checks and can be present in all age estimation methods. For weathervane scallops, a false check is an irregularity, crack, or shock line on the scallop shell surface. Merrill et al. (1966) found that serious developmental disturbances caused by injury or stress can result in the formation of a *shock ring* on scallop shells, which can mask or cause problems in discerning true annuli. Scallops in unfished or lightly fished areas tend to show few shock rings whereas those from heavily fished areas tend to have such rings with considerably higher frequency (Merrill et al. 1966). Typically, shock rings/false checks can be distinguished from annuli because they do not leave a continuous mark across the entire width of the shell, unlike annuli which can be tracked continuously along the entire span of the shell's surface (Spafard and Rosenkranz 2014). In-depth descriptions of annuli characteristics and false checks are needed to develop precise age estimates and to train new readers.

An effective tool for training and evaluating readers is a reference collection (Campana 2001). A reference collection is a standard set of prepared structures where the ages are known (through

tagging or research corroboration), or the ages are at least developed through laboratory consensus. Ideally, reference specimens should represent all variation in annuli characteristic used to estimate age, and should generally account for geographic range, stock, sex, size, and brood years. This is especially true for weathervane scallops as growth potentially varies considerably both within (Figure 3), and among geographic locations (Appendix A1). The development and continual use of reference collections is effective for insuring the accuracy and precision of age estimates. These evaluations can ensure that all readers are estimating structure ages similarly and that individual readers' criterion is not changing over time (reader drift).

A way to evaluate the error of ages and readers using either references or estimated ages is multiple reader comparisons. These repeated age estimates by different individuals allows for the statistical analysis of between-reader precision and bias (accuracy between readers), and assessment of the quality of the age estimates produced. Although it is important to assess precision, bias (i.e., disagreement) is more deleterious, because it has systematic impacts on model estimates and the effect of the error produced cannot be limited through increasing sample sizes (Campana 2001). Further, Campana and Jones (1992) and Hoenig et al. (1995) state that estimates of precision are only of interest and worth conducting if there is no evidence of bias. Therefore, comparisons of individual readers with other trained individuals are important especially in the absence of references that are of known age.



Figure 3.—Two 9-year-old scallops from Management Area D (Yakutat) in 2006 emphasizing the differences in growth within an area through different shell heights and to document observer codes. Fishery observer codes on sticky notes in image are: D = management area; S = scallop; 06 = year; Code C2 = missing 2 annuli at shell margin where shell height measurement was taken.

To evaluate bias and precision for quality control, both statistical and graphical analyses are used (Campana 2001). Common statistical tests for bias are the Evans-Hoenig and Bowker tests of symmetry (Bowker 1948; Hoenig et al. 1995), and graphical tests that include bias plots.

Precision is commonly measured using calculated average percent error, coefficient of variation (CV), and percent agreement (Campana 2001). Both bias and precision are discussed in more detail in the *Examining Bias and Precision* section. Assessing and tracking bias and precision provides for a standardized measure of data quality and ultimately results in more precise data. However, precision does not guarantee accuracy (i.e., reflect the true or absolute age) and should never be used as a substitute for accuracy (Campana 2001; Goldman et al. 2012).

The quality of age data ultimately relies on a solid protocol that ensures data are checked and potentially corrected in a consistent manner. Although general methods pertain to all organisms being aged, species-specific methods are required due to the difference structures and life history (e.g., fish otoliths and bivalve shells). In this report we define the standard method for estimating ages of weathervane scallops, ensuring proper data management and dissemination, testing for and improving accuracy and precision, and provide suggestions for future research to improve age estimates.

## **METHODS**

The methods for collecting representative and accurate age information to inform stock assessment can be broken down into 4 main categories: shell collection, age determination, quality control, and data management.

### **SHELL COLLECTION**

Shells may be collected from multiple sources using a variety of sampling programs depending on the suitability to the project. Currently, shells are collected from the ADF&G preseason survey (Smith et al. 2016) and the ADF&G Scallop Observer Program (ADF&G 2016). Although sampling regimes may vary, certain steps are necessary to collect and prepare shells for aging.

The top, or left valve (hereafter referred to as a shell) will be collected for aging because the annuli (dark banding pattern) are more visible and the circuli are more distinct on this shell. The bottom shell is subject to excessive wear due to resting on the sea floor and lacks the characteristic color changes of seasonal growth which are needed to estimate age. The difficulty in distinguishing annuli from the bottom shell could result in a systematic bias in scallop age estimates (Spafard and Rosenkranz 2014).

Shells must be intact enough that annuli counts may be made along multiple axes (see Age Estimation section below); age may be estimated from any shell that is not significantly damaged. For example, shells with broken hinges are acceptable because that portion of the shell is not used for age estimation. Additionally, shells with a broken margin may be acceptable if the damage is minimal (i.e., minor chips at places along the margin), but are unacceptable if the damage is considerable (e.g., the entire margin is chipped away or crushed and potential annuli can no longer be counted). Crushed shells should not be collected because this kind of damage prohibits an acceptable estimation of age. See Appendix B1 for examples of these classifications of damage.

To prepare shells for age estimation, all epifauna must be removed by scraping and brushing the surface with a 10% bleach solution. Shells should be subsequently dried and labelled with the appropriate identifying information including unique specimen numbers, location, date of collection, and any other data pertinent to the project. Detailed methods describing the collection

and preparation of scallop shells for age estimation purposes by the ADF&G Statewide Observer Program are documented in their Observer Program Manuals (ADF&G 2016).

## AGE ESTIMATION

To create the most accurate and precise data for stock assessment, age estimation protocols must be standardized and consistently applied. Consequently, the methods used to identify annuli, count annuli, and ensure that false checks are not counted must be consistent within and between readers. Due to its simplicity and speed, the standard method used to estimate scallop age will be the enumeration of alternating light and dark bands from the surface of the shell. However, it may be necessary to use the variation in circuli densities observed under magnification to determine annuli location (described below). The standard method of assessing annuli on weathervane scallop shells will be as follows:

1. Examine the outside of the shell from the umbo to the outer margin for any false checks or cracks. Dark bands that do not traverse across the entire shell represent a false check (Figure 4A), and therefore should not be counted as annuli. Cracks typically appear as a light band and will often show irregularities not observed in the dark bands that represent annual growth; consequently, these should also not be considered annuli (Figure 4B).




Figure 4.–A: Scallop aged at 10 years using the circuli method. The false checks observed between ages 3 and 4 and between ages 5 and 6 can also be seen using the color band aging method. False checks are marked in pencil and arrows on specimen, and ages are recorded with pencil. B: Scallop aged at 9 years using the circuli method, showing 2 false checks (denoted by arrows) in the shell which can also be seen using the color band method. Both cracks can be denoted by the abrupt color change to a white line (unlike an annulus which is composed of dark circuli). The crack between ages 2 and 3 also shows an irregularity from the crack in an area on the right side of the shell (encircled in a solid line). The second false check from a crack between ages 4 and 5 in the dashed circle shows where the crack crosses over (or blends) into the fifth annulus.

2. Because the outer margin of the shell can be damaged during collection, count the annuli along 3 different axes to ensure that annuli at the outer margin of shell are included, and cracks and false checks are not included in the assessed age (Figure 1B). Make the first or



primary axis count perpendicular to the hinge, starting at the umbo and moving along the straight-line height measurement to the outer margin of the shell. Make the second and third axes between 30 and 45 degrees from the umbo to an undamaged area at the outer margin on either side of the primary axis (Figure 1B).

3. Locate the first annulus, and use calipers to measure the distance from the umbo to the first annulus along the primary axis. For shells on which the dark band representing the first annulus is difficult to identify, take following steps:
  - a) Backlight the shell using a small lamp or something similar to provide greater contrast.
  - b) Use reflected light on the top of the shell under a magnifier or stereo-microscope to locate the first area of compressed circuli closest to the umbo (Figure 1B). If unsure about the presence or absence of an annulus, use the first annulus measurements provided in Appendix C1 to assist in determining if the first annulus can be located. Changing the orientation of the shell or adjusting the angle of the light can provide contrast for identifying the ridges of the circuli.
  - c) If the first annulus cannot be located using either of these methods, examine the auricle to see if the first annulus is visible.
  - d) If the first annulus cannot be located by any of the methods described above, and if the first visible annulus lies beyond the maximum distance of any first annuli measured for a shell from that management area (Appendix C1), count the remaining annuli and add 1 year to include the missing first annulus in the age of that shell.
4. Once the first annulus has been located, start with that annulus and count annuli along each of the 3 transects to the outermost edge of the shell. Special care must be taken when identifying annuli located near the outer margin of the shell because distances between annuli can decrease significantly as one moves away from the umbo, and shell damage in this area can obscure annuli. Use a magnifier to identify annuli near and at the outer margin of the shell. The maximum count from the 3 axes will be recorded as the final age for the shell. If shells are collected in the winter after January 1 or prior to the completion of the annulus (e.g., spring), count additional growth beyond the last visible annulus as another year due to application of the international January 1 birthdate (CARE 2006; Matta and Kimura 2012).
5. If the entire shell is heavily worn, or if conditions that otherwise obscure annuli from view are encountered (e.g, albinism; Figure 5), the reader will use the circuli method to estimate age.
  - a. Using a lighted magnifier or a stereo-microscope (2.5–40X), identify areas where circuli are spread apart or compressed as you move along the shell from the umbo towards the shell edge. Mark the far edge of the compressed circuli area as the annulus (Figure 2).
6. If the entire shell is heavily worn and annuli cannot be determined from dark bands or circuli on the main plane of the shell, check the auricles to determine if age can be assessed from that location using the same methods outlined above (Figure 6).
7. If the age of a specimen cannot be estimated using the above criteria, do not use the sample.




Figure 5.–Twelve-year-old albino scallop shell from commercial fishery Management Area D (Yakutat) in 2006. Fishery observer codes on sticky notes in image are D = management area; S = scallop; 06 = year.




Figure 6.–Example of annuli on scallop shell auricle. Arrows point to annuli.

## **QUALITY CONTROL**

Quality control for the ADF&G scallop age assessment program will be maintained by following similar standard practices used by the Alaska Fisheries Science Center, which has a long-standing (since 1983), consistent, and successful approach for quality control in age assessment practices (Kimura and Lyons 1991; Kimura and Anderl 2005; Matta and Kimura 2012).

### **Reference Collection**

Due to the different appearances (Appendix A1), growth rates, and associated anomalies that can be encountered on shells while estimating the age of scallops from different beds across Alaska, a reference collection will be created for each ADF&G management area. The current weathervane scallop reference collection is incomplete, and only consists of specimens collected from 2 locations. Until such time as a reference collection of 200 shells exists for all fishing areas, the current reference collection will serve for training and quality control purposes.

The reference collection will be used for quality control maintenance for experienced age readers and to train new age readers. Each year, prior to aging individuals, experienced age readers will use the reference collection to assess their precision and bias by estimating the ages of 3 sets of 20 individuals from each management area that they will be aging that year. Their results will be tested (see below) and must show a CV less than or equal to 10% and no bias before any field samples can be processed. If precision (CV) is greater than 10%, the reader will re-examine the age estimation protocol and read the discrepant samples from the reference collection again to test precision. If tests of symmetry show bias, the reader will review the protocol and re-estimate the age of samples within the age range of the bias, and run tests of symmetry again.

### **Training new readers**

Standardized training of new age readers and qualitative examination of their age estimation of reference specimens will help ensure standardization and repeatability among readers. The training of new age readers will be conducted as follows:

1. After thorough examination of this report, an experienced age reader will introduce the criteria used to identify and differentiate among annuli, cracks, and false checks to the new age reader.
2. New age readers will examine specimens in the reference collection to practice identifying annuli; at this time trainees will have access to the age data associated with each specimen to learn annuli criteria.
3. The new age reader will be given 3 sets of 20 specimens from 1 or multiple targeted management areas within the reference collection and produce age estimations without access to the reference age data (i.e., conduct blind reads). These age estimates will be tested for precision (CV) and bias (tests of symmetry).
4. Once an acceptable level of precision is achieved ( $CV \leq 10\%$ ) and tests of symmetry show no bias exists, the new age reader will conduct a blind read of the entire reference collection for the appropriate management area and their results will be tested for precision and bias. If the results show good precision and no bias, the trainee is ready to age. If the results show poor precision ( $CV > 10\%$ ), the new age reader will be required to re-age the entire collection again, until results are improved. If the results show bias, the new age reader will work with an experienced reader to improve application of annuli criteria until a test of symmetry shows differences between the reader and the reference collections are random.

## Examining Bias and Precision

Contingency table tests of symmetry will be used to determine if bias exists between age readers (Hoenig et al. 1995; Evans and Hoenig 1998; Campana 2001). These are designed to test the hypothesis that an  $m \times m$  contingency table (where  $m$  is the maximum age in the table) containing 2 categories (e.g., 1 set of ages provided by 2 age readers) is symmetric about the main diagonal of the table (Bowker 1948). The Bowker's (1948) test for differences using each off diagonal, but Evans and Hoenig (1998) provided a modification to the test where age differences from each side of the main diagonal of the table are pooled to enhance the ability to detect bias in data sets with small sample sizes. Both tests of symmetry will be used to provide the best ability to detect bias regardless of sample size. The test statistic follows a chi-square distribution for both Bowker (1948) and Evans and Hoenig (1998), and is defined as:

$$\chi^2 = \sum_{i=1}^{m-1} \sum_{j=i+1}^m \frac{(n_{ij} - n_{ji})^2}{(n_{ij} + n_{ji})} \quad (1)$$

where  $n_{ij}$  is the observed frequency in the  $i$ th row and  $j$ th column and  $n_{ji}$  is the observed frequency in the  $j$ th row and the  $i$ th column. The degrees of freedom ( $df$ ) for Bowker's (1948) test equals the number of comparisons across (on opposing sides of) the main diagonal of the table. The  $df$  for the Evans-Hoenig (1998) test is equal to the maximum difference between assigned ages that occurred between readers or methods and complete agreement (zero difference). The chi-square critical value will be determined based on the number of degrees of freedom at  $\alpha = 0.05$ . If the chi-square value from the test of symmetry is less than the critical value (i.e.,  $p$  values are not significant), the test suggests that there is not significant evidence of systematic bias. If the chi-square result is greater than or equal to the critical value (i.e.,  $p$  values are significant), then bias is present.

Tests of precision for comparing between-reader ages will include percent agreement (PA), percent agreement  $\pm 1$  yr (PA  $\pm 1$ yr), and the CV (see Chang 1982 and Campana 2001). While there is no absolute rule published for what is an acceptable mean CV for aging studies (Morrison et al. 1998; Campana 2001), Campana (2001) stated that 5–10% serves as a good reference value for many fishes aged by counting annuli in otoliths and vertebrae. Further, Kilada et al. (2007) reported similar age precision for smoothcockle (*Serripes groenlandicus*), reporting a mean CV of 4.68%. The between-reader percent agreement and CVs from Kilada et al. (2007) are similar to between-reader percent agreement and CVs reported for numerous groundfish species (Kimura and Lyons 1991; Kimura and Anderl 2005). To allow for the effects of the variable shape, shell wear, color, and look of scallop shells from different locations across Alaska, the maximum threshold of 10% will be used for mean CVs generated by the ADF&G scallop aging program and will be calculated as such:

$$CV_j = 100 \times \frac{\sqrt{\sum_{i=1}^R \frac{(X_{ij} - X_j)^2}{R-1}}}{X_j} \quad (2)$$

where  $CV_j$  is the age precision estimate for the  $j$ th scallop,  $X_{ij}$  is the  $i$ th scallop age of the  $j$ th scallop,  $X_j$  is the mean age of the  $j$ th scallop, and  $R$  is the number of final ages. The number of readers will typically be 2; however, if a third reader independently provides an age on a sample,



the  $R$  for that sample's CV calculation will be 3. Multiplying by 100 makes the CV value a percent. The  $CV_j$ s will then be averaged across all scallops to produce a mean CV, which will be compared to the 10% threshold. If the mean CV is greater than 10%, the precision of the sample will be considered too low.

### **Addressing Bias and Poor Precision**

Once age readers have completed their review of the reference collection and the results show acceptable precision and no bias, 2 readers (a primary reader and a test reader) will proceed to age collected shells. The primary reader will age all shells in a sample. The test reader will then independently read a 20% subsample without knowledge of the primary reader's estimated ages. Statistics for bias and precision (consistent with the above assessments) will be conducted on the between-reader results to ensure there is no bias or precision problems with the primary reader's estimated ages.

If no bias is detected with the test of symmetry and the CV for precision between the primary reader and the test reader is not greater than 10%, the original age estimated by the primary reader will be used for management.

If bias is detected, the following steps will be used:

1. The ages where between-reader bias is occurring will be assessed by examining the symmetry table and looking at ages surrounding the main diagonal in the table to see which samples need to be re-aged. The symmetry table will show the ages where readers are either over- or underaging samples (Appendix D2). An age range where the bias is occurring will be assessed from the table and those samples within the age range will be re-read by both readers and statistical analyses for bias will be run again.
2. If bias still exists, the primary and test reader will review all samples within the bias age range to determine ages by consensus. If no consensus can be reached on a specimen, a third reader will perform a blind read; if no 2 out of the 3 readers can agree on an assessed age, the specimen will not be aged.
3. Once the final ages of the first subsample are determined, a second randomly selected 20% subsample will be read by both readers to ensure the bias has been rectified.

If bias is not present but the CV is higher than 10%, the following steps will be used:

1. The primary and test reader will independently re-age all samples where their age estimates differed, and the CV will be calculated again. If the CV is still higher than 10%, the primary and test reader will work together to review all samples in question to assess differences and ensure that the criteria for identifying annuli is being followed appropriately.
2. If protocols were not followed, the ages for the entire sample must be aged again by a trained age reader.
3. If protocols were followed, the final ages may be determined during this consultation though consensus. If no consensus can be reached, a third reader will perform a blind read on the specimen. If no 2 out of the 3 readers can agree on an assessed age, the specimen will not be aged.

## **DATA MANAGEMENT**

All age data will be uploaded, managed, and housed in the ADF&G Mark, Tag and Age Laboratory's relational database, and will be available to ADF&G researchers and managers statewide. Date fields to be collected are in Appendix E1.

## REFERENCES CITED

- ADF&G (Alaska Department of Fish and Game). 2016. Scallop observer training and deployment manual. Alaska Department of Fish and Game, Division of Commercial Fisheries, Kodiak. [http://kodweb.fishgame.state.ak.us/index/Wiki:Observer\\_Program:Scallop\\_Observer\\_Program:Scallop\\_Observer\\_Manual:Archive](http://kodweb.fishgame.state.ak.us/index/Wiki:Observer_Program:Scallop_Observer_Program:Scallop_Observer_Manual:Archive); (Accessed February 27, 2017. URL only available from within the State of Alaska computer system).
- Bowker, A. H. 1948. A test for symmetry in contingency tables. *Journal of the American Statistical Association* 43:572–574.
- Campana, S. E. 2001. Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *Journal of Fish Biology* 59:197–242.
- Campana, S. E., and C. M. Jones. 1992. Analysis of otolith microstructure data. Pages 73–100 [In] D. K. Stevenson and S. E. Campana, editors. *Otolith microstructure examination and analysis*. Canadian Special Publication of Fisheries and Aquatic Science 117.
- CARE (Committee of Age Reading Experts). 2006. Manual on generalized age determination. <http://care.psmfc.org/docs/CareManual2006.pdf> (Accessed February 27, 2017).
- Chang, W. Y. B. 1982. A statistical method for evaluating the reproducibility of age determination. *Canadian Journal of Fisheries and Aquatic Sciences* 39:1208–1210.
- Chute, A. S., S. C. Wainright, and D. R. Hart. 2012. Timing of shell ring formation and patterns of shell growth in the sea scallop *Placopecten magellanicus* based on stable oxygen isotopes. *Journal of Shellfish Research* 31(3):649–662.
- Evans, G. T., and J. M. Hoenig. 1998. Testing and viewing symmetry in contingency tables, with application to readers of fish ages. *Biometrics* 54:620–629.
- Fiori, S. M., and E. M. Morsán. 2004. Age and individual growth of *Mesodesma mactroides* (Bivalvia) in the southernmost range of its distribution. *ICES Journal of Marine Science* 61:1253–1259.
- Goldman, K. J., G. M. Cailliet, L. J. Natanson, and A. Andrews. 2012. Assessing the age and growth of Chondrichthyan fishes. Pages 423–451 [In] J. Carrier, J. A. Musick, and M. Heithaus, editors. *The biology of sharks and their relatives*. Second Edition. CRC Press, Boca Raton, FL.
- Gustafson, R. L., and K. J. Goldman. 2012. Assessment of weathervane scallops in Kamishak Bay and at Kayak Island, 2004 through 2012. Alaska Department of Fish and Game, Fishery Data Series No. 12-62, Anchorage.
- Hart, D. R., and T. S. Chute. 2009. Verification of Atlantic sea scallop (*Placopecten magellanicus*) shell growth rings by tracking cohorts in fishery closed areas. *Canadian Journal of Fisheries and Aquatic Sciences* 66:751–758.
- Heifetz, J., D. Anderl, N. E. Maloney, and T. L. Rutecki. 1998. Age validation and analysis of aging error from marked and recaptured sablefish, *Anoplopoma fimbria*. *Fishery Bulletin* 97:256–263.
- Hennick, D. P. 1970. Reproductive cycle, size at maturity, and sexual composition of commercially harvested weathervane scallops (*Patinopecten caurinus*) in Alaska. *Journal of Fisheries Research Board of Canada* 27(11):2112–2119.
- Hoenig, J. M., M. J. Morgan, and C. A. Brown. 1995. Analyzing differences between two age determination methods by tests of symmetry. *Canadian Journal of Fisheries and Aquatic Sciences* 52:364–368.
- Kilada, R. W., D. Roddick, and K. Mombourquette. 2007. Age determination, validation, growth and minimum size of sexual maturity of the Greenland smoothcockle (*Serripes groenlandicus*, Bruguiere, 1789) in eastern Canada. *Journal of Shellfish Research* 26(2):443–450.
- Kimura, D. K., and D. M. Anderl. 2005. Quality control of age data at the Alaska Fisheries Science Center. *Marine Freshwater Research* 56:783–789.
- Kimura, D. K., and J. J. Lyons. 1991. Between-reader bias and variability in the age-determination process. *Fishery Bulletin* 89:53–60.

## REFERENCES CITED (Continued)

- Kimura, D. K. 1990. Approaches to age-structured separable sequential population analysis. *Canadian Journal of Fisheries and Aquatic Sciences* 47:2364–2374.
- Lai, H. L., and D. R. Gunderson. 1987. Effects of aging errors on estimates of growth, mortality and yield per recruit for walleye pollock (*Theragra chalcogramma*). *Fisheries Research* 5:287–302.
- Lomovasky, B. J., M. Lasta, M. Valiñas, M. Bruschetti, P. Ribeiro, S. Campodónico, and O. Iribarne. 2008. Differences in shell morphology and internal growth pattern of the Patagonian scallop *Zygochlamys patagonica* in the four main beds across their SW Atlantic distribution range. *Fisheries Research* 89:266–275.
- MacDonald, B. A., and R. J. Thompson. 1985. Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. I. growth rates of shell and somatic tissue. *Marine Ecology Progress Series* 25:279–294.
- MacDonald, B. A., and N. F. Bourne. 1987. Growth, reproductive output, and energy partitioning in weathervane scallops, *Patinopecten caurinus*, from British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* 44:152–160.
- Matta, M. E., and D. K. Kimura, editors. 2012. Age determination manual of the Alaska Fisheries Science Center age and growth program. NOAA Professional Paper NMFS 13. US Department of Commerce, Seattle WA. <http://spo.nwr.noaa.gov/pp13.pdf> (accessed February 27, 2017).
- Merrill, A. S., J. A. Posgay, and F. E. Nichy. 1966. Annual marks on shell and ligament of sea scallop (*Placopecten magellanicus*). *Fishery Bulletin* 65:299–311.
- Morrison, A. K., S. G. Robertson, and D. C. Smith. 1998. An integrated system for producing fish aging: Image analysis and quality assurance. *North American Journal of Fisheries Management* 18:587–598.
- Ropes, J. W. 1984. Procedures for preparing acetate peels and evidence validating the annual periodicity of growth lines formed in the shells of ocean quahogs, *Arctica islandica*. *Marine Fisheries Review* 46:27–35.
- Ropes, J. W., and A. Jearld Jr. 1987. Age determination in bivalves. Pages 517–526 [In] R. C. Summerfelt and G. E. Hall, editors. *The age and growth of fish*. Iowa State University Press, Ames, IA.
- Shaul, W., and L. Goodwin. 1982. Geoduck (*Panopea generosa*: Bivalvia) age determined by internal growth lines in the shell. *Canadian Journal of Fisheries and Aquatic Sciences* 39:632–636.
- Smith, Q., B. Williams, and R. Burt. 2016. Statewide weathervane scallop survey operational plan, 2016 through 2018. Alaska Department of Fish and Game, Alaska Department of Fish and Game, Regional Operational Plan ROP.CF.1J.2016.07, Juneau.
- Smith, S. J., E. L. Kenchington, M. J. Lundy, G. Robert, and D. Roddick. 2001. Spatially specific growth rates for sea scallops (*Placopecten magellanicus*). [In] G. H. Kruse, N. Bez, A. Booth, M. W. Dorn, S. Hills, R. N. Lipcius, D. Pelletier, C. Roy, S. J. Smith, and D. Witherell, editors. *Spatial processes and management of marine populations*. 17<sup>th</sup> Lowell Wakefield Symposium. Alaska Sea Grant College Program AK-SG-01-02.
- Spafard, M. A., and G. E. Rosenkranz. 2014. Age and growth of weathervane scallops *Patinopecten Caurinus* from the Alaska statewide scallop fishery, 1996–2013. Alaska Department of Fish and Game, Fishery Manuscript Series No. 14-03, Anchorage.
- Stevenson, J. A. and L. M. Dickie. 1954. Annual growth rings and rate of growth of the giant scallop, *Placopecten magellanicus* (Gmelin) in the Digby area of the Bay of Fundy. *Journal of Fisheries Research Board of Canada* 11:660–671.

**APPENDIX A: DIFFERENCES IN SHELL APPEARANCE  
ACROSS THE STATE OF ALASKA**







Appendix A1.—Example images of scallops from 5 management areas across the state of Alaska showing differences in shell appearance. Map of areas is provided in Appendix C2.



## **APPENDIX B: EXAMPLES OF DAMAGED SHELLS**



Appendix B1.—Examples of shell damage illustrating which shells can be aged and which cannot.

	
<p>Uninjured top valve</p>	<p>Cracked—Can be aged</p>
	
<p>Broken Margin—Can be aged</p>	<p>Broken Margin—Cannot be aged</p>
	
<p>Broken Margin—Cannot be aged</p>	<p>Crushed—Cannot be aged</p>




## **APPENDIX C: FIRST ANNULUS MEASUREMENTS**

Appendix C1.—First annulus measurements (mm) along the primary aging axis from scallops collected by ADF&G fishery observer program and research surveys. See Appendix C2 for a map showing management areas of collection.

Area	District	Mean	SE	Minimum	Maximum	Sample Size	Collected From	Years
D	Yakutat	21.7	0.17	10	44	1069	Observer program	2003–2015
D	District 16	22.0	0.93	14	32	30	Observer program	2006–2015
E	Kayak Island	25.4	0.07	6.6	53.6	8637	Research survey	1996–2014
H	Kamishak	29.6	0.08	7.6	52.8	9299	Research survey	1996–2015
KNE	Northeast	25.8	0.18	12	54	1396	Observer program	1996–2015
KSH	Shelikof	30.2	0.16	12	55	2395	Observer program	1999–2015
KSEM	Semidistrict Islands	27.3	0.86	18	34	27	Observer program	1996
KSW	Southwest	23.4	0.50	14	49	178	Observer program	2011–2015
M	West Chignik	21.9	0.84	16	28	17	Observer program	2008
M	Central	29.4	0.59	21	39	60	Observer program	1998–2006
M	Unimak Bight	20.6	0.40	13	38	137	Observer program	2012–2015
O	Dutch Harbor	23.7	0.35	15	52	262	Observer program	2008–2015
Q	Bering Sea	25.1	0.39	15	45	285	Observer program	1999–2015


Appendix C2.-Management Areas of collection.






## **APPENDIX D: TABLES OF SYMMETRY EXAMPLES**

Appendix D1.–Table A shows no bias in results; ages are randomly distributed across the main diagonal of the table.



Appendix D2.–Table B shows asymmetry (bias towards overestimation) across the main diagonal after age 14, indicating specimens in the circled area need to be re-aged.



## **APPENDIX E: DATA COLLECTION FIELDS**

Appendix E1.–Data fields to be entered by age readers.

Primary Reader	Test Reader
Field Data (ties age samples back to databases)	Test reader's name
Event ID or Haul ID	Date of test read
Scallop or shell ID number	Method used to determine first annulus location
F&G Region	Assessed age
Species code	Test reader sample size
Shell height (mm)	Test reader comments
<hr/>	
Lab data	
Primary reader's name	
Date of primary read	
Shell condition code (0-5):	
0 = undamaged	
1 = broken hinge	
2 = broken margin	
3 = cracked	
4 = punctured	
5 = crushed	
Method used to determine first annulus location	
Distance from umbo to first annulus (mm)	
Assessed age	
Selected for second read (y/n)?	
Primary reader sample size	
Primary reader comments	
<hr/>	
Precision and Bias Analysis:	
Between reader percent agreement	
Between-reader percent agreement $\pm$ 1 yr	
Between-reader CV (coefficient of variation)	
<hr/>	
Test of symmetry results:	
chi-square statistics	
Degrees of freedom (Bowker test)	
Degrees of freedom (Evans-Hoenig test)	

*Note:* Event ID and scallop ID # are identifiers used on surveys and Kamishak fishery; Haul ID and shell ID #are identifiers used in observer program.